District Level Variation in Hypertension Epidemiology in India and Influence of Social Determinants:

National Family Health Survey-5

Shore title: Hypertension in India: NFHS-5

Rajeev Gupta[1]

Kiran Gaur[2]

Suresh Sharma[3]

Raghubir S Khedar[1]

Rajinder K Dhamija[4]

From: [1]Department of Preventive Cardiology & Medicine, Eternal Heart Care Centre & Research Institute, Jaipur, India; [2]Department of Statistics, Mathematics and Computer Science, Government SKN Agriculture University, Jobner, Jaipur, India. [3]Population Research Centre, Ministry of Health & Family Welfare, Institute of Economic Growth, Delhi University Enclave, New Delhi, India. [4]Department of Neurology, Institute of Human Behavior and Allied Sciences, New Delhi, India.

Correspondence: Dr Rajeev Gupta, Department of Preventive Cardiology & Medicine, Eternal Heart Care Centre and Research Institute, Jagatpura Road, Jawahar Circle, Jaipur 302017 India. Phones: +91-141-5174000; FAX: +91-141-5174001; Email: rajeevgg@gmail.com; drrajeev.gupta@eternalheart.org;

ABSTRACT

BACKGROUND: Enumeration of state and district-level variation in hypertension prevalence in India

and to evaluate the influence of social determinants.

METHODS: We used data from the Fifth National Family Health Survey (NFHS-5) from 707 districts

and 825,954 participants (women 724,115, men 101,839 men) on prevalence of hypertension

defined according to standard criteria. Data on multiple social determinants were also obtained from

NFHS-5 report.

RESULTS: Age-standardized prevalence of hypertension was 22.4% (women 21.3%, men 24.0%) with

the highest prevalence in women and men, respectively, in Sikkim (34.5 and 41.6%) and Punjab (31.2

and 37.7%) and lowest in Rajasthan (15.4 and 17.9%) and Ladakh (15.7 and 17.4%). Prevalence was

more in western and southern Indian districts. High prevalence of hypertension in the young (<30y)

was observed in northeastern and northern states. District-level hypertension prevalence correlated

negatively with multi-dimensional poverty index (R^2 women 0.299, men 0.245) and positively with

female literacy (women 0.165, men 0.134). Among women, districts with the highest availability of

electricity, clean water, sanitation, clean cooking fuels, healthcare service delivery and better

nutrition were associated with more hypertension on univariate and multivariate analyses (p<0.05).

CONCLUSIONS: The study shows significant geographical variation in hypertension prevalence in

India. Hypertension is more in men with high prevalence of premature hypertension. Better district-

level development (less poverty, more literacy) and healthcare services are associated with greater

hypertension prevalence in women.

KEY WORDS: Hypertension; Raised blood pressure; Epidemiology; Non-communicable diseases;

Social determinants.

SUMMARY TABLE

What is known about the topic:

• Significant state-level variation in hypertension prevalence in India has been reported but

district-level variation is not known.

• Social determinants are important in hypertension but not well studied, especially in

women.

What this study adds:

• The study shows a significant district-level variation with greater hypertension prevalence in

southern and western India.

• Hypertension among the young, <30 years, is more in less developed districts.

• Social determinants of hypertension in women are less poverty, more literacy and

availability of healthcare services.

INTRODUCTION

World Health Organization (WHO) and others have reported that high blood pressure (BP) is the most important risk factor for mortality and disease burden in both women and men, globally and in India.[1-4] Studies have reported significant inter-country variation in hypertension prevalence with the highest prevalence rates in Eastern European, Central Asian and Eastern Asian countries.[1,3,5,6] Within-country variation has also been reported from several large countries- China, USA, Brazil, Indonesia, Pakistan, UK- and Europe.[7-13] A county (district) level variation has been reported from the USA and UK.[14,15] Studies from USA, UK and China have reported greater hypertension prevalence in less developed regions and counties.[7,8,12] Risk factors for hypertension are social and biological.[16] Social determinants of hypertension include low-quality urban and rural infrastructure, low socioeconomic status, social disorganization, unemployment, adverse work environment, illiteracy, adverse early life events, subnormal maternal and child health, etc.[16,17] It has been reported that macrolevel and micro-level variations in social, lifestyle and biological factors are responsible for difference in hypertension prevalence.[16]

Previous national studies in India- District Level Health Survey/Adult Health Survey (DLHS-4/AHS-3)[18] and Fourth National Family Health Survey (NFHS-4)[19] have reported significant state-level variation in hypertension prevalence. It has been reported that macro-level social factors such as greater urbanization, better human development index, more literacy and raised body mass index (BMI) are associated with this variation.[19] This contrasts with developed countries where hypertension is more in lesser developed and deprived regions and communities.[12,15] India has undergone rapid socio-economic progress in recent years. To correlate variation in hypertension prevalence at the district (county) level with various social factors we performed the present study. We used district-level data from Fifth National Family Health Survey (NFHS-5)[20] on hypertension prevalence and performed correlation with multiple social determinants of health.

METHODS

Data from NFHS-5 are available at the National Family Health Survey website at

http://rchiips.org/nfhs/factsheet_NFHS-5.shtml.[20] NFHS is a large-scale, multi-round survey

conducted in a representative sample of households throughout India. Multiple rounds of the survey

have been conducted since the first survey in 1992-93. The survey provides state and national

information for India on fertility, infant and child mortality, the practice of family planning, maternal

and child health, reproductive health, nutrition, anemia, utilization and quality of health and family

planning services. Each successive round of the NFHS has had two specific goals: a) to provide

essential data on health and family welfare needed by the Ministry of Health and Family Welfare and

other agencies for policy and program purposes, and b) to provide information on important

emerging health and family welfare issues. Ethics clearance has been obtained by the Institutional

Ethics Committee of Indian Institute of Population Sciences, Mumbai, India. Technical assistance for

the NFHS has been provided by ICF International (Virginia, USA) and other organizations on specific

issues. The funding for different rounds of NFHS has been provided by USAID, DFID, the Bill and

Melinda Gates Foundation, UNICEF, UNFPA, and Ministry of Health and Family Welfare, Government

of India. In NFHS-5, similar to the previous iterations, the sample was designed to provide data of all

key indicators at the national and state levels as well as estimates for most key indicators at the

district level for almost all the districts of the country.[20]

The total sample size in NFHS-5 was 610,000 households in India. This was estimated to

produce reliable data for each district of the country. The sample selection process was similar to

the previous NFHS studies. The rural sample was selected through a 2-stage sampling with villages as

the primary sampling unit (PSU) at the first stage selected with probability proportional to size,

followed by a random selection of 22 households in each PSU at the second stage. In urban areas,

there was also a 2-stage sampling design with Census Enumeration Blocks (CEB) selected at the first

stage and a random selection of 22 households in each CEB at the second stage. In the second stage

in both rural and urban areas, households were selected after conducting a complete mapping and

household listing operation in the selected first-stage units. NFHS-5 fieldwork for India was

conducted in two phases, phase one from 17 June 2019 to 30 January 2020 and phase two from 2

January 2020 to 30 April 2021 by 17 Field Agencies and gathered information from 636,699

households, 724,115 women, and 101,839 men in 705 districts of the country.

The biomarker questionnaire included measurements of height, weight, waist and hip

circumference, hemoglobin levels and finger-stick blood for additional testing in a laboratory for

women aged 15-49 years and men aged 15-54 years. Blood pressure and random blood glucose

were measured in all women and men aged 15 years and more in contrast to NFHS-4.[19]

Questionnaire information and biomarkers were collected after informed consent from each

participant. In the NFHS-5 the BP of eligible respondents was measured using an OMRON Blood

Pressure Monitor to determine the prevalence of hypertension. BP measurements for each

respondent were taken on three separate occasions and the readings were recorded in the

Biomarker Questionnaire with an interval of 5 minutes between readings. Respondents whose

systolic BP was >130 mm Hg or diastolic BP > 85 mm Hg were considered to have elevated blood

pressure readings and were encouraged to see a doctor for a full evaluation. Hypertension was

diagnosed when systolic BP was ≥140 mmHg and/or diastolic BP ≥90 mmHg or when the participant

was taking medicine to control BP and reported in per cent (%).

Statistical analyses:

Data sheets were downloaded from the NFHS-5 website. Additional data were obtained

from the Population Research Centre, Institute of Economic Growth, Delhi University, New Delhi,

India. Data were transferred to MS Excel sheets and descriptive analyses were performed using this

program. Hypertension prevalence rates are reported separately for women and men and have been

age-adjusted to the population of India. A χ^2 test was used to determine the significance of women-

men differences. Hypertension prevalence rates for various districts of the country (n=705) were

also obtained from the NFHS-5 website and reported separately for women and men. State-level

mean, medians and 25-75 interquartile intervals (IQR) for hypertension were calculated. Univariate

correlation of district hypertension prevalence with social determinants (Supplementary Table 1)

was performed using polynomial regression and graphs were computed in MS Excel. For multivariate

correlation, we used SPSS statistical package.

Role of funding sources:

No direct funding was obtained for writing this report. National Family Health Surveys are

funded by various grants from the Government of India.

142   **RESULTS**

NFHS-5 enrolled 724,115 women and 101,839 men >15 years of age (total of 825,954) with

response rates more than 90%.[20] Data for multiple socioeconomic, lifestyle and clinical variables

were obtained and recorded. National age-adjusted prevalence (%) of hypertension among women

was 21.3% and men 24.0% (overall 22.4%). Age-adjusted prevalence (%) of hypertension among

women and men in different states of India is in Table 1. There is significant state-level variation in

hypertension prevalence. In women the highest prevalence is in Sikkim (34.5%), Punjab (31.2%) and

Goa (27.5%) and the lowest in Rajasthan (15.4%), Ladakh (15.7%) and Bihar (15.9%). In men the

highest prevalence is in Sikkim (41.6%), Punjab (37.7%) and Manipur (33.2%) while the lowest is in

Ladakh (17.4%), Rajasthan (17.9%) and Bihar (18.4%) (Figure 1). Hypertension prevalence in various

states and districts of India in women and men is shown as a heat map (Figure 2). High prevalence is

observed in southern and eastern Indian states. There is a significant district-level variation with

greater prevalence in most western and southern districts in women and western, southern and

eastern districts in men. Absolute prevalence rates for each district among women and men are in

Supplementary Tables 2-34. We also calculated median (IQR) hypertension prevalence in each state

according to the district-level prevalence and prevalence rates for women and men shown in

Supplementary Figure 1.

Age-group stratified hypertension prevalence among women and men for the whole country

and all the states are in Table 2. There is a significant age-associated escalation in its prevalence in all

the states of the country (Mantel-Haenszel χ^2 for trend, p <0.001). Higher prevalence of

hypertension in the young (<30 years) is observed among men and women in most North-Eastern

Indian states (Arunachal Pradesh, Assam, Meghalaya, Nagaland, Sikkim and West Bengal), Jammu &

Kashmir, Ladakh and Goa (Supplementary Figure 2). On the other hand, high prevalence of

hypertension among the elderly ($\geq$60yr) is observed in Delhi, Goa, Karnataka, Kerala, Punjab, Sikkim

and Telangana.

Data on multiple social determinants were obtained from various government databases,

NFHS-5 and others.[19,20] District-level multidimensional poverty index is negatively associated with

the prevalence of hypertension in both women and men with polynomial R^2 of 0.299 and 0.245,

respectively (Figure 3a). Hypertension prevalence is positively associated with district-level female

literacy levels in both women and men with polynomial R^2 of 0.165 and 0.134, respectively (Figure

3b). Data on social determinants in NFHS-5 have been primarily obtained for women, therefore, we

determined variation of hypertension prevalence in women according to them. Correlation of

district-level hypertension prevalence among women with various social, healthcare delivery and

clinical determinants are in Figure 4. Increasing hypertension prevalence among women correlates

significantly with declining poverty, better literacy, sanitation, and electricity, and availability of

clean fuel for cooking (Figure 4a). Greater hypertension prevalence is also observed in district with

lower childhood marriages and teenage pregnancies and greater vaccination coverage, modern

contraceptive usage and availability of skilled delivery services (Figure 4b). Presence of childhood

overweight, adult overweight (BMI $\geq$25) and adult hyperglycemia is also associated with greater

hypertension prevalence (Figure 4c).

To identify important social determinants among women we classified various

socioeconomic-, environmental-, healthcare service-, womankind specific-, and nutritional variables

into tertiles and determined hypertension prevalence in each of them (Table 3). Analyses of social

determinants show that in districts with lower poverty and illiteracy and the highest prevalence of

availability of electricity, clean water, sanitation and clean cooking fuels, the prevalence of

hypertension is the highest. Availability of better healthcare service delivery (higher availability of

skilled childbirth, use of modern contraception and full vaccination) and better nutrition (lower

anemia, childhood stunting or wasting, overweight in children and adults) is also associated with

greater prevalence of hypertension. We also calculated univariate and multivariate regression

coefficients to identify important variables (Table 4). On univariate analysis. There are multiple

factors associated with district level hypertension prevalence including socioeconomic and

environmental variables (multidimensional poverty index, illiteracy, access to electricity, clean

drinking water and sanitation, tobacco use), healthcare service delivery parameters and nutritional

factors. However, on multivariate analyses risk factors that continue to be significant are low female

illiteracy (standardized b -0.12), improved drinking water (0.11), sanitation (0.19), modern

contraceptive use (0.20) and overweight or obesity in women (0.44) (p<0.05).

DISCUSSION

Fifth National Family Health Survey (NFHS-5) is the most representative and comprehensive

hypertension epidemiology study in India. The study shows that there is substantial variation in

hypertension prevalence across states and districts of the country. The prevalence of hypertension is

more in better-developed states and districts. Hypertension in the young (<30 y) is more in the

northeastern and northern states of the country. Greater hypertension prevalence in more

developed districts (lower multidimensional poverty, and better literacy, healthcare services and

nutritional status) in India, contrasts with developed countries where hypertension is more in lesser

developed locations and states. These findings suggest that macrolevel cardiovascular disease risk

factor transition associated with demographic and epidemiological shifts is still evolving in India.

Geographic variation in hypertension prevalence has been reported from all the large

countries of the world. [7-15] There are significant difference in prevalence rates at state and district-

levels. In China, greater hypertension prevalence has been reported from less developed western

and south-eastern regions;[7] in USA, hypertension prevalence is greater in southern and south-

eastern states and less developed counties;[8] and in the UK hypertension is more in deprived

counties of northern region.[12] Global Burden of Disease (GBD) Study sub-national collaborators have

reported greater hypertension related mortality in deprived regions of Brazil.[9] On the other hand, in

lower middle income countries such as India, Indonesia and Pakistan, greater hypertension

prevalence and hypertension related cardiovascular mortality have been reported from better

developed states and locations.[10,11,19] Country-level differences also exist in hypertension In Europe

and Asia. In Europe hypertension prevalence is significantly greater in less developed countries of

Central and Eastern Europe,[13] while in Asia the prevalence of hypertension is significantly lower in

high income countries of Eastern Asia and high in Central and Southern Asian countries.[21] Positive

association of hypertension in India with better development suggests that the risk factor transition

that happens with social development has not yet occurred in India.[23] There is, on the other hand,

evidence that at an individual level, hypertension may be more among illiterate rural and slum-

dwelling populations,[23,24] and there is a significant association of low socioeconomic and educational

status with greater cardiovascular mortality.[25,26] In the present study we have shown a positive

association of human development with hypertension in India and more studies are needed to

identify the importance of individual and macrolevel social determinants.

An important finding in the present study is a high prevalence of premature hypertension,

especially in lesser developed states of northeastern and northern India. Higher prevalence of

hypertension at young age in India, compared to countries of North and South America and Europe

has been previously reported in the DLHS-4/AHS study.[18] Northeastern Indian states have a very

proportion of tribal population and a high prevalence of hypertension among these populations has

also been reported.[27] Hypertension in the young is an important problem among less developed

societies in Africa and among African Americans in USA.[28] There is a significant association of

premature hypertension with stroke in these communities. Million Death Study in India reported a

significantly greater stroke burden and mortality in the eastern and north-eastern states of the

country.[29] This corresponds to the presence of premature hypertension in these regions. More

studies are required to elucidate reasons for greater hypertension and stroke burden among the

Sub-Himalayan regions of India.

The study has several strengths- this is the one of the first studies that has evaluated

hypertension prevalence in each district of the country. The DLHS-4/AHS study also evaluated

hypertension in multiple districts but two important states were missing.[18] IN NFHS-4 the sample

size was restricted to the young and middle aged and did not provide data on elderly which have

greater hypertension.[30] The ICMR-INDIAB study also obtained data on hypertension in multiple

states but sampling was based on state-level data and not at the district level.[31] We have also

identified districts with greater hypertension prevalence. This is important for policy-making as these

districts should be the initial focus for hypertension control initiatives.[32] Identification of districts

with premature hypertension is also important as deployment of primary prevention strategies in

these regions (salt, alcohol and tobacco control, healthy foods). Limitations of the study include a

skewed female-male ratio in the recruitment of the sample (NFHS reports are focused on women's

health factors), single-day measurement of BP which is not recommended to confirm the diagnosis

of hypertension, use of older definition of hypertension (SBP $\geq$140 and/or DBP $\geq$90 mmHg) instead of

more recently suggested criteria (SBP $\geq$130 and/or DBP $\geq$80 mmHg). In the present study we do not

yet have data on hypertension awareness, awareness and control status and this is also a limitation.

Data on pre-hypertension, which is more prevalent than hypertension in India, and should be the

focus of prevention strategies, are not yet available. There is a need for more primary data on

hypertension determinants to explain regional variations including social and cultural determinants

of physical activity and diet. Data on district-level availability and access to healthcare and

availability of drug-therapies to individuals, especially women, with hypertension are not available.

Finally, all these data need replication for external validation using data from concurrent studies

including the ICMR-INDIAB and other national surveys.

Conclusions:

It has been argued that phase transition in the human progress has been brought about by

tectonic shifts in development with technological progress in every corner of the globe. This has

lifted most societies out of Malthusian trap but aspects of local geography, culture and institutions

are also important in this transition.[33] Our study shows that the escape from poverty and illiteracy

among women is associated with greater hypertension prevalence in India and forecasts a short-

term scenario with more hypertension with better social development. We hope that a

socioeconomic tipping point shall arrive soon where hypertension is more in lower socioeconomic

status individuals.[34] India and most low-middle income and low-income countries have significantly

higher adverse cardiovascular events and mortality from hypertension.[35] Reversal of this trend is

possible using well-tried evidence-based interventions at the level of population and individuals

cross the life course. The creation of an egalitarian society with a focus on various components of

inequality,[34] availability of non-communicable disease-focused healthcare,[36] and population

empowerment,[36] would lead to lower hypertension among the more developed communities and

regions of the country.

CONTRIBUTORS

RG and KG conceptualized the study. Data analyses and estimate generation were done by KG and

SS. Maps were generated by KG. RG drafted the first iteration of the manuscript. RG, KG, SS and RKD

made substantial contributions to critical review, editing and revision of the manuscript. All the

authors approved the final version of the manuscript. RG, KG and SS accessed and verified the data

underlying the study. All authors had full access to all the data in the study and had final

responsibility for the decision to submit for publication.

DECLARATION OF INTERESTS:

All the authors declare no competing interests. KG is an employee of Government of Rajasthan and

SS and RKD are employed by Government of India.

DATA SHARING:

All the data used for the manuscript are freely available at the National Family Health Survey

website at http://rchiips.org/nfhs/.

ACKNOWLEDGEMENTS:

No specific funding was available for this article.

296   **LEGENDS TO TABLES**

Table 1: Hypertension prevalence in Indian states in NFHS-5 in women and men >15 years. Criteria

for diagnosis of hypertension were systolic BP $\geq$140 mmHg and/or diastolic BP $\geq$90 mmHg or on

medical treatment.

Table 2: Age-group specific hypertension prevalence in Indian states in women and men.

Table 3: Hypertension prevalence (%, 95% CI) in women according to tertiles of various social

indicators.

Table 4: Univariate and multivariate regression analysis (standardized b) of district-level

hypertension prevalence (women) with various socioeconomic and other factors.

LEGENDS TO FIGURES

Figure 1: Hypertension prevalence among women and men in various states of India in NFHS-5.

Figure 2: Heat map showing state-level prevalence (Figure 2a) and district-level prevalence (Figure

2b) of hypertension among women and men in NFHS-5.

Figure 3: Inverse correlation of district-level hypertension prevalence in women and men with

increasing multidimensional poverty index (Figure 3a) and positive correlation of district-level

hypertension prevalence in women and men with increasing literacy (Figure 3b).

Figure 4 (a-c): Correlation of district-level hypertension prevalence in women with various (a)

social, (b) healthcare delivery, and (c) clinical determinants.

REFERENCES

1.   World Health Organization. Invisible numbers: the true extent of non-communicable diseases

and what to do about them. Geneva. World Health Organization. 2022.

2.   Global Burden of Disease Study 2019 Risk Factor Collaborators. Global burden of 87 risk factors

in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of

Disease Study 2019. *Lancet*. 2020; 396:1223-1250.

3.   Global Burden of Diseases CVD Collaborators. Global burden of cardiovascular diseases and risk

factors, 1990-2019: Update from the Global Burden of Disease 2019 Study. *J Am Coll Cardiol*.

2020; 76:2982-3021.

4.   Gupta R, Xavier D. Hypertension: the most important non-communicable disease risk factor in

India. *Indian Heart J*. 2018; 70:565-572.

5.   NCD Risk Factor Collaboration (NCD-RiSC). Worldwide trends in blood pressure from 1975 to

2015: a pooled analysis of 1479 population-based measurement studies with 19.1 million

participants. *Lancet*. 2017; 389:37-55.

6.   NCD Risk Factor Collaboration. Worldwide trends in hypertension prevalence and progress in

treatment and control from 1990 to 2019: A pooled analysis of 1173 population-representative

studies with 104 million participants. *Lancet*. 2021; 398:957-980.

7.   Li Y, Wang L. Feng X, Zhang M, Huang Z, Deng Q, Zhou M, Astell-Burt T, Wang L. Geographical

variations in hypertension prevalence, awareness, treatment and control in China: Findings from

a nationwide and provincially representative survey. *J Hypertens*. 2018; 36:178-187.

8.   Centres for Disease Control and Prevention. Hypertension maps and data sources: prevalence of

hypertension, 2017. Available at: https://www.cdc.gov/bloodpressure/maps_data.htm. Accessed

7 March 2023.

9.   Nascimento BR, Brant LCC, Yadgir S, Oliveira GMM, Roth G, Glenn SD, Mooney M, Naghavi M,

Passos VMA, Duncan BB, Silva DAS, Malta DC, Ribeiro ALP. Trends in prevalence, mortality and

morbidity associated with high systolic blood pressure in Brazil from 1990 to 2017: estimates

from the Global Burden of Disease 2017 (GBD 2017) study. *Pop Health Metrics*. 2020; 18:e17.

10. GBD 2019 Indonesia Subnational Collaborators. The state of health in Indonesia's provinces,

1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet Glob*

Health. 2022; 10:e1632-1645.

11. GBD 2019 Pakistan Collaborators. The state of health in Pakistan and its provinces and

territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet*

Glob Health. 2023; 11:e229-243.

12. British Heart Foundation. Heart and circulatory disease statistics 2022. Available at:

https://www.bhf.org.uk/what-we-do/our-research/heart-statistics/heart-statistics-

publications/cardiovascular-disease-statistics-2022. Accessed 7 March 2023.

13. Timmis A, Townsend N, Gale CP, Torbica A, Lettino M, Petersen SE, Mossialo EA, Maggioni AP,

Kazakiewicz D, May HT, Smedt DD, Flather M, Zuhlke L, Beltrame JF, Huculeci R, Tavazzi L,

Hindricks G, Bax J, Casedei B, Achenbach S, Wright L, Vardas P. European Society of Cardiology:

cardiovascular disease statistics 2019. *Eur Heart J*. 2020; 40:12-85.

14. Vaughan AS, Coronado F, Casper M, Loustalot F, Wright JS. County level trends in hypertension

related cardiovascular mortality: United States, 2000 to 2019. *J Am Heart Assoc*. 2022;

11:e024785.

15. Hughes A. Hypertension prevalence estimates in England, 2017: Estimated from the Health

Survey for England. London. Public Health England. 2020.

16. Hall ME, Hall JE, Whelton PK. Epidemiology, pathophysiology and treatment of hypertension. In:

Fuster V, Narula J, Vaishnava P, et al. Editors. Fuster & Hurst's: The Heart, 15[th] Ed. New York:

McGraw-Hill. 2022;189-232.

17. Marmot M, Wilkinson R. Social Determinants of Health, 2[nd] Ed. Oxford. Oxford University Press.

2006.

18. Geldsetzer P, Manne-Goehler J, Theilmann M, Davies JI, Awasthi A, Vollmer S, Jaacks LM,

Barnighausen T, Atun R. Diabetes and hypertension in India: a nationally representative study of

1.3 million adults. *JAMA Intern Med*. 2018; 178:363-372.

19. Gupta R, Gaur K, Ram CVS. Emerging trends in hypertension epidemiology in India. *J Hum*

Hypertens. 2019. 33:575-587.

20. National Family Health Survey of India. Available at: http://rchiips.org/nfhs/. Accessed 7

March19023.

21. Gupta R, Gaur K. Hypertension epidemiology in India: Recent nationwide studies and comparison

with Asian countries. In: Ponde CK, Mohanan PP. Editors. *Cardiology 2022: Hypertension*.

Bengaluru. Microlabs Ltd. 2022. 235-244.

22. Subramanian S, Ambade M, Kumar A, Chi H, Joe W, Rajpal S, Kim R. Progress on Sustainable

Development Goal indicators in 707 districts of India: a quantitative mid-line assessment using

the National Family Health Surveys, 2016 and 2021. *Lancet Reg Health SE Asia*. 2023;100155.

23. Gupta R, Gupta VP, Ahluwalia NS. Educational status, coronary heart disease and coronary risk

factor prevalence in a rural population of India. *BMJ*. 1994; 309:1332-1336

24. Gupta R, Kaul V, Agrawal A, Guptha S, Gupta VP. Cardiovascular risk according to educational

status in India. *Prev Med*. 2010; 51:408-411.

25. Rosengren A, Smyth A, Rangarajan S, Ramasundarahettige C, Bangdiwala SI, AlHabib KF, Avezum

A, Bengtsson Boström K, Chifamba J, Gulec S, Gupta R, Igumbor EU, Iqbal R, Ismail N, Joseph P,

383 Kaur M, Khatib R, Kruger IM, Lamelas P, Lanas F, Lear SA, Li W, Wang C, Quiang D, Wang Y,

384 Lopez-Jaramillo P, Mohammadifard N, Mohan V, Mony PK, Poirier P, Srilatha S, Szuba A, Teo K,

385 Wielgosz A, Yeates KE, Yusoff K, Yusuf R, Yusufali AH, Attaei MW, McKee M, Yusuf S..

386 Socioeconomic status and risk of cardiovascular disease in 20 low-income, middle-income, and

387 high-income countries: the Prospective Urban Rural Epidemiology (PURE) study. *Lancet Glob*

388 *Health.* 2019; 7:e748-760.

389 26. Joseph P, Kutty VR, Mohan V, Kumar R, Mony P, Vijayakumar K, Islam S, Iqbal R, Kazmi K, Rahman

390 O, Yusuf R, Anjana RM, Mohan I, Rangarajan S, Gupta R, Yusuf S. Cardiovascular disease,

391 mortality and their associations with modifiable risk factors in a multinational South Asia cohort:

392 A PURE substudy. *European Heart J.* 2022; 43:2831-2840.

393 27. Hazarika C, Babu BV. Prevalence of hypertension in Indian tribal population: a systematic review

394 and meta-analysis. *J Racial Ethn Health Disparities.* 2023; EPub. doi: 10/1007/s40615-023-01532-

395 6.

396 28. Zhou B, Perel P, Mensah GA, Ezzati M. Global epidemiology, health burden and effective

397 interventions for elevated blood pressure and hypertension. *Nature Rev Cardiol.* 2021; 18:785-

398 802.

399 29. Ke C, Gupta R, Xavier D, Prabhakaran D, Mathur P, Kalkonde YV, Kolpak P, Suraweera W, Jha P;

400 Million Death Study Collaborators.. Divergent trends in ischemic heart disease and stroke

401 mortality in India from 2000 to 2015: A nationally representative mortality survey. *Lancet Global*

402 *Health.* 2018; 6:e914-923.

403 30. Prenissl J, Manne-Goehler J, Jaacks LM, Prabhakaran D, Awasthi A, Bischops AC, Atun R,

404 Bärnighausen T, Davies JI, Vollmer S, Geldsetzer P. Hypertension screening, awareness,

405 treatment and control in India: a nationally representative cross sectional study among

406 individuals aged 15-49 years. *PLoS Med.* 2019; 16:e1002801.

31. Anjana RM, Unnikrishnan R, Deepa M, Pradeepa R, Tandon N, Das AK, Joshi S, Bajaj S, Jabbar PK,

Das HK, Kumar A, Dhandhania VK, Bhansali A, Rao PV, Desai A, Kalra S, Gupta A, Lakshmy R,

Madhu SV, Elangovan N, Chowdhury S, Venkatesan U, Subashini R, Kaur T, Dhaliwal RS, Mohan

V; ICMR-INDIAB Collaborative Study Group. Metabolic non-communicable disease health report

of India: the ICMR-INDIAB national cross-sectional study (ICMR-INDIAB 17). *Lancet Diabetes*

Endocrinol. 2023; 11:474-489.

32. Kaur P, Kunwar A, Sharma M, Durgad K, Gupta S; India Hypertension Control Initiative

collaboration; Bhargava B. The India Hypertension Control Initiative: early outcomes in 26

districts across 5 states of India, 2018-2020. *J Hum Hypertens*. 2023; 37:560-567.

33. Galor O. Metamorphosis. In: *The Journey of Humanity: The Origins of Wealth and Inequality*.

Dublin. Penguin Random House. 2022;85-100.

34. Piketty T. Toward a democratic, ecological and multicultural socialism. In: *A Brief History of*

Equality. Cambridge, MA. Harvard University Press. 2022; 226-244.

35. Walli-Attaei M, Rosengren A, Rangarajan S, Breet Y, Abdul-Razak S, Sharief WA, Alhabib KF,

Avezum A, Chifamba J, Diaz R, Gupta R, Hu B, Iqbal R, Ismail R, Kelishadi R, Khatib R, Lang X, Li S,

Lopez-Jaramillo P, Mohan V, Oguz A, Palileo-Villanueva LM, Poltyn-Zaradna K, R SP, Pinnaka LVM,

Serón P, Teo K, Verghese ST, Wielgosz A, Yeates K, Yusuf R, Anand SS, Yusuf S; PURE

investigators. Metabolic, behavioural, and psychosocial risk factors and cardiovascular disease in

women compared with men in 21 high-income, middle-income, and low-income countries: an

analysis of the PURE study. *Lancet* 2022; 400:811-821.

36. Gupta R, Wood DA. Primary prevention of ischemic heart disease: populations, individuals, and

healthcare professionals. *Lancet*. 2019; 394:685-696.

Women

State	Value
Sikkim	35.8
Punjab	31.5
Kerala	31.4
Goa	27.5
Puducherry	27.5
Telangana	26.1
Karnataka	25.5
Arunachal Pradesh	25.3
Tamil Nadu	24.4
Delhi	24.1
Andhra Pradesh	24
Chhattisgarh	23.9
Nagaland	22.6
Uttarakhand	22.1
Manipur	21.9
Maharashtra	21.8
Odisha	21.8
Madhya Pradesh	21.1
Haryana	20.7
Himachal Pradesh	20.4
West Bengal	20.2
Jammu & Kashmir	20
Tripura	19.5
Gujarat	19.4
Uttar Pradesh	18.4
Assam	18.2
Jharkhand	17.6
Meghalaya	17.4
Bihar	16
Rajasthan	15.1
Mizoram	14.1

Men

State	Value
Sikkim	46.4
Punjab	36.2
Delhi	33.8
Kerala	33.5
Puducherry	32.2
Uttarakhand	31.9
Arunachal Pradesh	31.6
Telangana	30.3
Tamil Nadu	30.2
Nagaland	30
Manipur	29.4
Andhra Pradesh	28.5
Karnataka	27.9
Chhattisgarh	27.7
Goa	26.7
Haryana	25.2
Odisha	25
Maharashtra	23.7
Madhya Pradesh	23.2
Himachal Pradesh	23
Assam	22
Jharkhand	22
Tripura	21.4
Uttar Pradesh	21.4
Mizoram	20.7
West Bengal	20.4
Meghalaya	20
Jammu & Kashmir	18.7
Gujarat	18.2
Rajasthan	17.3
Bihar	16.7

Hypertension Prevalence Women (%)

Hypertension Prevalence Men (%)

Value

- 33 - 42
- 28 - 32
- 23 - 27
- 19 - 22
- 15 - 18

Hypertension Prevalence Women (%)

Hypertension Prevalence Men (%)

Value

- 43 - 50
- 36 - 42
- 29 - 35
- 22 - 28
- 15 - 21
- 10 - 14

Social Determinants of Health

Women Specific Healthcare Service Delivery Parameters

Women Specific Clinical Parameters