Antigenic drift and subtype interference shape A(H3N2) epidemic dynamics in the United States

Amanda C Perofsky[1,2*], John Huddleston[3], Chelsea Hansen[1,2], John R Barnes[4], Thomas Rowe[4], Xiyan Xu[4], Rebecca Kondor[4], David E Wentworth[4], Nicola Lewis[5], Lynne Whittaker[5], Burcu Ermetal[5], Ruth Harvey[5], Monica Galiano[5], Rodney Stuart Daniels[5], John W McCauley[5], Seiichiro Fujisaki[6], Kazuya Nakamura[6], Noriko Kishida[6], Shinji Watanabe[6], Hideki Hasegawa[6], Sheena G Sullivan[7], Ian G Barr[7], Kanta Subbarao[7], Florian Krammer[8,9], Trevor Bedford[2,3,10,11], Cécile Viboud[1]

Affiliations

1. Fogarty International Center, National Institutes of Health, United States
2. Brotman Baty Institute for Precision Medicine, University of Washington, United States
3. Vaccine and Infectious Disease Division, Fred Hutchinson Cancer Center, United States
4. Virology Surveillance and Diagnosis Branch, Influenza Division, National Center for Immunization and Respiratory Diseases (NCIRD), Centers for Disease Control and Prevention (CDC), United States
5. WHO Collaborating Centre for Reference and Research on Influenza, Crick Worldwide Influenza Centre, The Francis Crick Institute, United Kingdom
6. Influenza Virus Research Center, National Institute of Infectious Diseases, Japan
7. WHO Collaborating Centre for Reference and Research on Influenza, The Peter Doherty Institute for Infection and Immunity, Department of Microbiology and Immunology, The University of Melbourne, The Peter Doherty Institute for Infection and Immunity, Australia
8. Center for Vaccine Research and Pandemic Preparedness (C-VaRPP), Icahn School of Medicine at Mount Sinai, United States
9. Department of Pathology, Molecular and Cell-Based Medicine, Icahn School of Medicine at Mount Sinai, United States
10. Department of Genome Sciences, University of Washington, United States
11. Howard Hughes Medical Institute, Seattle, United States

*Correspondence to amanda.perofsky@nih.gov

Abstract

Influenza viruses continually evolve new antigenic variants, through mutations in epitopes of their major surface proteins, hemagglutinin (HA) and neuraminidase (NA). Antigenic drift potentiates the reinfection of previously infected individuals, but the contribution of this process to variability in annual epidemics is not well understood. Here we link influenza A(H3N2) virus evolution to regional epidemic dynamics in the United States during 1997—2019. We integrate phenotypic measures of HA antigenic drift and sequence-based measures of HA and NA fitness to infer antigenic and genetic distances between viruses circulating in successive seasons. We estimate the magnitude, severity, timing, transmission rate, age-specific patterns, and subtype dominance of each regional outbreak and find that genetic distance based on broad sets of epitope sites is the strongest evolutionary predictor of A(H3N2) virus epidemiology. Increased HA and NA epitope distance between seasons correlates with larger, more intense epidemics, higher transmission, greater A(H3N2) subtype dominance, and a greater proportion of cases in adults relative to children, consistent with increased population susceptibility. Based on random forest models, A(H1N1) incidence impacts A(H3N2) epidemics to a greater extent than viral evolution, suggesting that subtype interference is a major driver of influenza A virus infection dynamics, presumably via heterosubtypic cross-immunity.

Introduction

Influenza viruses continually accumulate genetic changes in epitopes of two major surface proteins, hemagglutinin (HA) and neuraminidase (NA), in a process known as "antigenic drift." Though individual hosts develop long-lasting immunity to specific influenza virus strains after infection, antigenic drift helps the virus to escape immune recognition, leaving previously exposed hosts susceptible to reinfection and necessitating the regular updates to the antigens included in the influenza vaccine [1]. While antigenic drift aids immune escape, prospective cohort studies and modeling of surveillance data also indicate that reinfection by antigenically homologous viruses occurs on average every 1-4 years, due to the waning of protection over time and antigenic drift [2,3].

Among the influenza virus types that routinely co-circulate in humans (A and B), type A viruses, particularly subtype A(H3N2), experience the fastest rates of antigenic evolution and cause the most substantial morbidity and mortality [4-7]. Seasonal influenza A viruses (IAV) cause annual winter epidemics in temperate zones of the Northern and Southern Hemispheres and circulate year-round in tropical regions [8]. Influenza A epidemic burden fluctuates substantially from year to year [9], and there is much scientific interest in disentangling the relative roles of viral evolution, prior immunity, human behavior, and climatic factors in driving this seasonal variability. Climatic factors, such as humidity and temperature, have been implicated in the seasonality and timing of winter outbreaks in temperate regions [10-14], while contact and mobility patterns contribute to the seeding of new outbreaks and geographic spread [10,15-19]. A principal requirement for the recurrence of epidemics is a sufficient and continuous source of susceptible individuals, which is determined by the degree of cross-immunity between the surface antigens of currently circulating viruses and functional antibodies elicited by prior infection or vaccination in a population.

Because mutations to the HA1 region of the HA protein are considered to drive the majority of antigenic drift [20,21], influenza virus genetic and antigenic surveillance have focused primarily on HA, and official influenza vaccine formulations prescribe the amount of HA [22]. Yet, evidence for the effect of HA drift on influenza epidemic dynamics remains conflicting. Theoretical and empirical studies have shown that HA drift between currently circulating viruses and the previous season's viruses is expected to cause earlier, larger, more severe, or more synchronized epidemics; however, the majority of these studies were limited to the pre 2009 influenza pandemic period [6,17,23-28]. Information on HA evolution has been shown to improve forecasts of seasonal influenza dynamics in Israel and the United States [29,30], but recent research has also found that HA evolution is not predictive of epidemic size in Australia [31] or epidemic timing in the United States [16]. A caveat is that many of these studies used binary indicators to study seasonal antigenic change, defined as seasons in which circulating viruses were antigenically distinct from the vaccine reference strain [16,17,24,31,32]. This may obscure epidemiologically relevant patterns, as positive selection in HA and NA is both episodic and continuous [6,32-37]. Past research has also typically focused on serological and sequence-based measures of viral evolution in isolation, and the relative importance of these two approaches in predicting epidemic dynamics has not been systematically assessed. Further, to the best of our knowledge, the epidemiologic impact of NA evolution has not been explored.

There has been recent recognition of NA's role in virus inhibiting antibodies and its potential as a vaccine target [38-40]. Though antibodies against NA do not prevent influenza infection, NA immunity attenuates the severity of infection by limiting viral replication [41-46], and NA-specific antibody titers are an independent correlate of protection in both field studies and human challenge trials [47-49]. Lastly, the phenomenon of interference between influenza A subtypes, modulated by immunity to conserved T-cell epitopes [50-52], has long been debated [53,54]. Interference effects are most pronounced during pandemic seasons, leading to troughs or even replacement of the resident subtype in some pandemics [55], but the contribution of heterosubtypic interference to annual dynamics is unclear [2,56-59].

Here, we link A(H3N2) virus evolutionary dynamics to epidemiologic surveillance data in the United States
over the course of 22 influenza seasons prior to the coronavirus disease (COVID-19) pandemic,
considering the full diversity of viruses circulating in this period. We analyze a variety of antigenic and
genetic markers of HA and NA evolution against multiple indicators characterizing the epidemiology and
disease burden of annual outbreaks. We find a signature of both HA and NA antigenic drift in surveillance
data, with a more pronounced relationship in epitope change rather than the serology-based indicator,
along with a major effect of subtype interference. Our study has implications for surveillance of
evolutionary indicators that are most relevant for population impact and for prediction of influenza burden
on inter-annual timeframes.

Results

Our study focuses on the impact of A(H3N2) virus evolution on seasonal epidemics from seasons 1997-
1998 to 2018-2019 in the US; whenever possible, we make use of regionally disaggregated indicators
and analyses. We start by identifying multiple indicators of influenza evolution each season based on
changes in HA and NA. Next, we compile influenza virus subtype-specific incidence time series for US
Department of Health and Human Service (HHS) regions and estimate multiple indicators characterizing
influenza A(H3N2) epidemic dynamics each season, including epidemic burden, severity, intensity,
type/subtype dominance, timing, and the age distribution of cases. We then assess univariate
relationships between indicators of evolution and epidemic characteristics. Lastly, we measure the
relative importance of viral evolution, heterosubtypic interference, and prior immunity in predicting
regional A(H3N2) epidemic dynamics, using multivariable regression models and random forest models.

Indicators of influenza A(H3N2) evolution

We characterized seasonal patterns of genetic and antigenic evolution among A(H3N2) viruses
circulating from 1997 to 2019, using HA and NA sequence data shared via the Global Initiative on Sharing
Avian Influenza Data (GISAID) EpiFlu database [60] and ferret hemagglutination inhibition (HI) assay data
shared by the WHO Global Influenza Surveillance and Response System (GISRS) Collaborating Centers
in London, Melbourne, Atlanta, and Tokyo. Prior to constructing phylogenetic trees, we subsampled
sequences to representative sets of 50 viruses per month, with preferential sampling for North American
sequences. Although our study is US-focused, we used a global dataset because US-collected
sequences and HI titers were sometimes sparse during the earlier seasons of the study. Time-resolved
phylogenies of HA and NA genes are shown in Figure 1.

Our choice of evolutionary indicators builds on earlier studies that found hemagglutination inhibition (HI)
phenotype or HA sequence data beneficial in forecasting seasonal influenza virus evolution [35,61-63] or
annual epidemic dynamics [27,29,30] (Table 1). Historically, HI serological assays were considered the
gold standard for measuring immune cross-reactivity between viruses, yet measurements are available
for only a subset of viruses. To overcome this limitation, we used a computational approach that maps HI
titer measurements onto the HA phylogenetic tree to infer antigenic phenotypes [35,63]. Importantly, this
model infers the antigenicity of virus isolates that lack HI titer measurements, which comprise the majority
of HA sequences in GISAID. Our sequence-based measures of drift counted substitutions at epitope sites
in the globular head domains of HA and NA, identified through monoclonal antibody escape or protein
crystal structure: 129 sites in HA epitope regions A to E [21,64-67], 7 sites adjacent to the HA receptor
binding site (RBS) [68], and 223 or 53 sites in NA epitope regions A to C [34,69].

We included other indicators of viral fitness for HA and NA, including the number of substitutions at non-
epitope sites (mutational load) [35,61] and the average rate of phylogenetic branching in a season (local
branching index, LBI) [35,62]. We also calculated the Shannon entropy of LBI values, which considers the
richness and relative abundances of viral clades with different growth rates. Lastly, we counted the
number of substitutions at epitope sites in the HA stalk domain (stalk footprint distance) [70]. Although the
majority of the antibody-mediated response to HA is directed to the immunodominant HA head,

antibodies towards the highly conserved immunosubdominant stalk domain of HA are widely prevalent in
older individuals, although at low levels [71-73]. We considered stalk footprint distance to be our "control"
metric for drift, given the HA stalk evolves at a significantly slower rate than the HA head [70].

To measure antigenic distances between consecutive seasons, we calculated mean genetic distances at
epitope sites or mean $\log_2$ titer distances from HI titer measurements (Figure 1), between viruses
circulating in the current season t and the prior season t-1 year (one season lag) or two prior seasons ago
t-2 years (two season lag). These time windows generated seasonal antigenic distances consistent with
empirical and theoretical studies characterizing transitions between H3 or N2 antigenic clusters
[6,32,35,55,62,74], with H3 epitope distance and HI $\log_2$ titer distance, at two-season lags, and N2
epitope distance, at one-season lags, capturing expected "jumps" in antigenic drift during key seasons
that have been previously associated with major antigenic transitions [32], such as the seasons
dominated by A/Sydney/5/1997-like strains (SY97) (1997-1998, 1998-1999, 1999-2000) and the 2003-
2004 season dominated by A/Fujian/411/2002-like strains (FU02) (Figures S1-S2). Prior studies explicitly
linking antigenic drift to epidemic size or severity also support a one-year [6] or two-year time window of
drift [26,27]. Given that protective immunity wanes after 1-4 years, we would also expect these
timeframes to return the greatest signal in epidemiological surveillance data.

We measured pairwise correlations between seasonal indicators of HA and NA evolution to assess their
degree of concordance. As expected, we found moderate-to-strong associations between HA epitope
distance and HI $\log_2$ titer distance and HA RBS distance and HI $\log_2$ titer distance (Figure S1-S3).
Consistent with prior serological studies [39,75,76], epitope distances in HA and NA were not correlated
(one-season lag: Spearman's $\rho = 0.25$, $P = 0.26$; two-season lag: $\rho = 0.15$, $P = 0.5$; Figures S2-S4).
Seasonal diversity of HA and NA LBI values was negatively correlated with NA epitope distance (Figure
S3), suggesting that selective sweeps follow the emergence of drifted variants.

164 Associations between A(H3N2) evolution and epidemic dynamics

We explored relationships between viral evolution and variation in A(H3N2) epidemic dynamics from
seasons 1997-1998 to 2018-2019, excluding the 2009 A(H1N1) pandemic, using syndromic and virologic
surveillance data collected by the US CDC and WHO.

We estimated weekly incidences of influenza A(H3N2), A(H1N1), and B in 10 HHS regions by multiplying
the influenza-like illness (ILI) rate – the proportion of outpatient encounters for ILI, weighted by regional
population size – by the regional proportion of respiratory samples testing positive for each influenza
type/subtype (percent positive) [57,77]. We combined pre-2009 seasonal A(H1N1) viruses and
A(H1N1)pdm09 viruses as A(H1N1) and the Victoria and Yamagata lineages of influenza B viruses as
influenza B. Weekly incidences of influenza A(H3N2), A(H1N1), and type B, averaged across the 10 HHS
regions, are shown in Figure 2. Weekly regional incidences, which show variability in the timing and
intensity of annual epidemics, are shown in Figure 2 and Figure S5. Based on these incidence time
series, we measured indicators of epidemic burden, intensity, severity, subtype dominance, timing, and
age-specific patterns during each non-pandemic season and assessed their univariate relationships with
each indicator of HA and NA evolution, which we describe in turn below. Seasonal characteristics of
A(H3N2) epidemic dynamics were based on epidemic size, defined as the cumulative weekly incidence;
peak incidence, defined as the maximum weekly incidence; excess mortality attributable to A(H3N2), an
indicator of epidemic severity; transmissibility, defined as the maximum time-varying effective
reproductive number, effective Rt; and epidemic intensity, defined as the inverse Shannon entropy of the
weekly incidence distribution (i.e., the sharpness of the epidemic curve). See methods and Table 2 for
details on all epidemic metrics and Figure S6 for pairwise correlations between metrics.

Two sequence-based measures based on broad sets of epitope sites exhibited stronger relationships with
seasonal epidemic burden and transmissibility than the serology-based measure, HI $\log_2$ titer distance.
Both H3 epitope distance ($t - 2$) and N2 epitope distance ($t - 1$) correlated with increased epidemic size

(linear models, LMs: H3, adjusted $R^2 = 0.37$, P = 0.03; N2: $R^2 = 0.26$, P = 0.08) and peak incidence (LMs,
H3: $R^2 = 0.4$, P = 0.02; N2: $R^2 = 0.33$, P = 0.04) and higher effective Rt (generalized linear models, GLMs:
H3, $R^2 = 0.38$, P = 0.05; N2, $R^2 = 0.32$, P = 0.03) (regression results: Figure 3, Spearman's correlations:
Figure S7). HI $\log_2$ titer distance ($t - 2$) exhibited positive but non-significant associations with different
measures of epidemic impact (Figure 3, Figure S7). Seasonal diversity in the growth rates of circulating
lineages in the current t or prior season ($t - 1$) had strong negative correlations with effective Rt (GLMs,
H3 ($t - 1$): $R^2 = 0.49$, P = 0.009; N2, t: $R^2 = 0.46$, P = 0.006) and epidemic intensity (Beta GLMs, H3 ($t -$
1): $R^2 = 0.45$, P = 0.003; N2, t: $R^2 = 0.51$, P = 0.001) (Figures S7-S8). Seasonal mean LBI exhibited
similar but slightly weaker correlations with effective Rt and epidemic intensity. Pneumonia and influenza
excess mortality attributable to A(H3N2) also increased with H3 epitope distance, though this relationship
was not statistically significant (Figure S9). The remaining indicators of viral evolution, including H3 and
N2 non-epitope distance (mutational load), H3 RBS distance, and H3 stalk footprint distance had weak,
non-significant correlations with the different measures of epidemic impact (Figure S7).

We explored whether evolutionary changes in A(H3N2) may predispose this subtype to dominate
influenza virus circulation in a given season. A(H3N2) subtype dominance – the proportion of influenza
positive samples typed as A(H3N2) – increased with H3 epitope distance ($t - 2$) and N2 epitope distance
($t - 1$) (Beta GLMs, H3: $R^2 = 0.32$, P = 0.05; N2: $R^2 = 0.34$, P = 0.03; Figure 4, Figure S7). Figure 4
illustrates this relationship at the regional level across two seasons in which A(H3N2) was nationally
dominant, but where antigenic change differed. In 2003-2004, we observed widespread dominance of
A(H3N2) viruses after the emergence of the novel antigenic cluster, FU02 (A/Fujian/411/2002-like
strains). In contrast, there was substantial regional heterogeneity in subtype circulation during 2007-2008,
a season in which A(H3N2) viruses were antigenically similar to those from the previous season. Patterns
in type/subtype circulation across all influenza seasons in our study period are shown in Figure S10. As
observed for the 2003-2004 season, widespread A(H3N2) dominance tends to coincide with major
antigenic transitions (e.g., A/Sydney/5/1997 (SY97) seasons, 1997-1998 to 1999-2000;
A/California/7/2004 (CA04) season, 2004-2005), though this was not universally the case (e.g.,
A/Perth/16/2009 (PE09) season, 2010-2011).

Next, we tested for associations between A(H3N2) evolution and epidemic timing, including onset week,
defined as the winter changepoint in incidence [16], and peak week, defined as the first week of
maximum incidence; spatiotemporal synchrony, measured as the variation (standard deviation, s.d.) in
regional onset and peak timing; and epidemic speed, including seasonal duration and the number of
weeks from onset to peak (Table 2, Figure S11). Seasonal duration increased with H3 or N2 LBI diversity
in the current t or prior season ($t - 1$) (Gamma GLMs, H3, t: $R^2 = 0.6$; P = 0.001; N2, t: $R^2 = 0.6$; P =
0.002; Figures S11-S12), while the number of days from epidemic onset to peak shortened with
increasing N2 epitope distance ($t - 1$) (Gamma GLM, $R^2 = 0.31$, P = 0.04; Figure S11, Figure S13). Onset
and peak timing tended to be earlier in seasons with increased H3 and N2 antigenic novelty, but
correlations were not statistically significant (Figure S14). A(H3N2) evolution did not correlate with the
degree of spatiotemporal synchrony across HHS regions.

Lastly, we considered the effects of antigenic change on the age distribution of outpatient ILI cases, with
the expectation that the proportion of cases in children would decrease in seasons with greater antigenic
novelty, due to drifted variants' increased ability to infect more immunologically experienced adults [7,78].
Consistent with this hypothesis, N2 epitope distance from prior seasons was negatively correlated with
the fraction of cases in children aged < 5 years (LMs, one-season lag: $R^2 = 0.29$, P = 0.1; two-season lag:
$R^2 = 0.59$, P = 0.003) and individuals aged 5-24 years (one-season lag: $R^2 = 0.38$, P = 0.04; two-season
lag: $R^2 = 0.17$, P = 0.18) and negatively correlated with the fraction of cases in adults aged 25-64 years
(one-season lag: $R^2 = 0.36$, P = 0.05; two-season lag: $R^2 = 0.49$, P = 0.01) and ≥65 years (one-season
lag: $R^2 = 0.39$, P = 0.01; two-season lag: $R^2 = 0.33$, P = 0.05) (Figures S15-S16). As observed in Gostic
et al. [78], H3 epitope distance ($t - 2$) had negative but non-significant associations with the fraction of
cases in children and positive but non-significant associations with the fraction of cases in adult age
groups (Figures S15-S16).

Effects of heterosubtypic viral interference on A(H3N2) epidemic burden and timing

We investigated the effects of influenza type/subtype interference – proxied by influenza A(H1N1) and B epidemic size – on A(H3N2) incidence during annual outbreaks. Across the entire study period, we observed moderate-to-strong, non-linear relationships between A(H1N1) epidemic size and A(H3N2) epidemic size (GLM, $R^2 = 0.65$, $P = 0.01$), peak incidence ($R^2 = 0.66$, $P = 0.02$), and excess mortality (all age groups and ≥ 65 years, $R^2 = 0.57$, $P = 0.01$) (Figure 5, Figure S17), wherein A(H3N2) epidemic burden and excess mortality decreased as A(H1N1) incidence increased. A(H1N1) epidemic size was also significantly correlated with A(H3N2) effective Rt, exhibiting a negative, approximately linear relationship (GLM, $R^2 = 0.45$, $P = 0.01$) (Figure 5). A(H3N2) epidemic intensity was negatively associated with A(H1N1) epidemic size, but this relationship was not statistically significant (Beta GLM, $R^2 = 0.21$, $P = 0.15$). Influenza B epidemic size was not significantly correlated with any A(H3N2) epidemic metrics (Figure 5, Figure S17).

The internal gene segments NS, M, NP, PA, and PB2 of A(H3N2) viruses and pre-2009 seasonal A(H1N1) viruses share a common ancestor [79] whereas A(H1N1)pdm09 viruses have a combination of gene segments derived from swine and avian reservoirs that were not reported prior to the 2009 pandemic [80,81]. Because pre-2009 seasonal A(H1N1) viruses and A(H3N2) are more closely related, seasonal A(H1N1) viruses may limit the circulation of A(H3N2) viruses to a greater extent than A(H1N1)pdm09 viruses. As a sensitivity analysis, we measured correlations between A(H1N1) incidence and A(H3N2) epidemic metrics separately for pre- and post-2009 pandemic time periods. Relationships between different A(H3N2) epidemic metrics and A(H1N1) epidemic size were broadly similar for both periods, with slightly stronger correlations observed during the pre-2009 period (Figure S18).

We compared A(H3N2) epidemic timing across A(H3N2) and A(H1N1) dominant seasons, which we defined as when $\geq 70\%$ of influenza A positive samples are typed as A(H3N2) or A(H1N1)/A(H1N1)pdm09, respectively. We applied a strict threshold for subtype dominance because seasons with $< 70\%$ samples of one IAV subtype tended to have greater geographic heterogeneity in circulation, resulting in regions with dominant subtypes that were not nationally dominant. A(H3N2) epidemic onsets and peaks occurred, on average, three weeks earlier in A(H3N2) dominant seasons (Wilcoxon test, $P < 0.0001$). In A(H1N1) dominant seasons, regional A(H3N2) epidemics exhibited greater heterogeneity in epidemic timing (onset s.d.: H3 dominant seasons, 12.4 weeks versus H1 dominant seasons, 16.3 weeks; peak s.d., H3 dominant seasons, 13.3 weeks versus H1 dominant seasons, 22.6 weeks; Wilcoxon tests, $P < 0.0001$) and were significantly shorter in duration compared to A(H3N2) dominant seasons (median duration: H3 dominant seasons, 29 weeks versus H1 dominant seasons, 21 weeks; Wilcoxon test, $P < 0.0001$).

We applied a wavelet approach [82] to weekly time series of type/subtype-specific incidences to measure more fine-scale differences in the relative timing of type/subtype circulation (Figure S19). A(H3N2) incidence preceded A(H1N1) incidence during most seasons prior to 2009 and during the two seasons in which A(H1N1)pdm09 was dominant, potentially because A(H3N2) viruses are more globally prevalent and migrate between regions more frequently than A(H1N1) viruses [7]. There was not a clear relationship between the direction of seasonal phase lags and A(H1N1) epidemic size (LM, $R^2 = 0.23$, $P = 0.1$; Figure S19). A(H3N2) incidence led influenza B incidence in all influenza seasons (positive phase lag), irrespective of influenza B epidemic size (LM, $R^2 = 0.05$, $P = 0.5$; Figure S19).

The relative impacts of viral evolution, heterosubtypic interference, and prior immunity on A(H3N2) epidemic dynamics

We implemented conditional inference random forest models to assess the relative importance of viral evolution, type/subtype co-circulation, prior population immunity, and vaccine-related parameters in predicting regional A(H3N2) epidemic metrics (Figure 6). We limited viral evolutionary indicators to H3 epitope distance ($t - 2$), N2 epitope distance ($t - 1$), HI $\log_2$ titer distance ($t - 2$), and H3 and N2 LBI

diversity in the current and prior season, due to weaker or non-significant correlations between the other
evolutionary metrics and epidemic burden (Figure S7). To account for potential type or subtype
interference, we included A(H1N1) epidemic size (A(H1N1) or A(H1N1)pdm09) and B epidemic size in the
current and prior season and the dominant IAV subtype in the prior season. We included A(H3N2)
epidemic size in the prior season as a proxy of natural prior immunity to A(H3N2). To account for vaccine-
induced immunity, we considered four categories of predictors and included estimates for the current and
prior seasons: seasonal vaccination coverage among adults (18-49 years coverage × ≥ 65 years
coverage), adjusted A(H3N2) vaccine effectiveness (VE), a combined metric of vaccination coverage and
A(H3N2) VE (18-49 years coverage × ≥ 65 years coverage × VE), and H3 and N2 epitope distance
between currently circulating strains and the US vaccine reference strain. We could not include a
predictor for vaccination coverage in children or consider clade-specific VE estimates, because data were
not available for most seasons in our study. We did not predict excess mortality attributable to A(H3N2),
due to data limitation (one national estimate per season) and omitted models predicting epidemic timing,
due to weak or non-significant correlations between timing-related measures and most indicators of viral
evolution (Figure S11). Lastly, we could not separate our analysis into pre- and post-2009 pandemic
periods due to small sample sizes.

Based on variable importance scores, A(H1N1) epidemic size in the current season was the most
informative predictor of A(H3N2) epidemic size and peak incidence, followed by H3 epitope distance, and
the dominant IAV subtype in the previous season or N2 epitope distance (Figure 6). For A(H3N2) subtype
dominance, the highest ranked predictors were H3 epitope distance, N2 epitope distance, and the
dominant IAV subtype in the previous season (Figure 6). We note that we did not include A(H1N1)
epidemic size as a predictor in this model, due to its confounding with the target variable. For models of
A(H3N2) effective Rt and epidemic intensity, we observed less discernable differences in variable
importance scores across the set of candidate predictors (Figure 6). For the model of effective Rt, N2 LBI
diversity in the current season, A(H1N1) epidemic size in the current season, and N2 epitope distance
between circulating strains and the vaccine strain were the highest ranked variables, while the most
important predictors of epidemic intensity were H3 and N2 LBI diversity in the current season and adult
vaccination coverage in the current and prior season. Variable importance rankings from LASSO (least
absolute shrinkage and selection operator) regression models were qualitatively similar to those from
random forest models, with A(H1N1) epidemic size in the current season, H3 and N2 epitope distance,
and the dominant IAV subtype in the prior season consistently retained across the best-tuned models of
epidemic size, peak incidence, and subtype dominance (Figure S20). Vaccine-related parameters and H3
antigenic drift (either H3 epitope distance or HI $\log_2$ titer distance) were retained in the best-tuned LASSO
models of effective Rt and epidemic intensity (Figure S20).

We measured correlations between observed values and model-predicted values at the HHS region level.
Among our various epidemic metrics, random forest models produced the most accurate predictions of
A(H3N2) subtype dominance (ρ = 0.94, regional range = 0.8 – 0.98), peak incidence (Spearman's ρ =
0.91, regional range = 0.73 – 0.95), and epidemic size (ρ = 0.9, regional range = 0.73 – 0.94), while
predictions of effective Rt and epidemic intensity were less accurate (ρ = 0.8, regional range = 0.65 –
0.91; ρ = 0.78, regional range = 0.63 – 0.91, respectively) (Figure 7). Random forest models tended to
underpredict most epidemic targets in seasons with substantial H3 antigenic transitions, in particular the
SY97 cluster seasons (1998-1999, 1999-2000) and the FU02 cluster season (2003-2004) (Figure 7).

For epidemic size and peak incidence, seasonal predictive error – root-mean-square error (RMSE) across
all regional predictions in a season – increased with H3 epitope distance (size, Spearman's ρ = 0.51, P =
0.02; peak, ρ = 0.61, P = 0.007) and N2 epitope distance (size, ρ = 0.43, P = 0.06; peak, ρ = 0.46, P =
0.04). For models of epidemic intensity, seasonal RMSE increased with N2 epitope distance (ρ = 0.62, P
= 0.006) but not H3 epitope distance (ρ = 0.07, P = 0.8) (Figures S21-S22). The RMSE of effective Rt and
subtype dominance predictions were not significantly correlated with H3 or N2 epitope distance (Figures
S21-22).

To further refine our set of informative predictors, we performed multivariable regression with the top 10 ranked predictors from each random forest model and used Bayesian Information Criterion (BIC) to select the best fit model for each epidemic metric, allowing each metric's regression model to include up to three independent variables. This additional step of variable selection demonstrated that models with few predictors fit the observed data relatively well (epidemic size, adjusted R^2 = 0.69; peak incidence, adj. R^2 = 0.63; effective Rt, adj. R^2 = 0.65; epidemic intensity, adj. R^2 = 0.75), except for subtype dominance (adj. R^2 = 0.48) (Table 3). The set of variables retained after model selection were similar to those with high importance rankings in random forest models and LASSO regression models, with the exception that HI log_2 titer distance, rather than H3 epitope distance, was included in the minimal models of effective Rt and epidemic intensity.

Discussion

Antigenic drift between currently circulating influenza viruses and the previous season's viruses is expected to confer increased viral fitness, leading to earlier, larger, or more severe epidemics. However, prior evidence for the impact of antigenic drift on seasonal influenza outbreaks is mixed. Here, we systematically compare experimental and sequence-based measures of A(H3N2) evolution in predicting regional epidemic dynamics in the United States across 22 seasons, from 1997 to 2019. We also consider the effects of other co-circulating influenza viruses, prior immunity, and vaccine-related parameters, such as coverage and effectiveness, on A(H3N2) incidence. Our findings indicate that evolution in both major surface proteins – hemagglutinin (HA) and neuraminidase (NA) – contributes to variability in epidemic magnitude across seasons, though viral fitness appears to be secondary to subtype interference in shaping annual outbreaks.

The first question of this study sought to determine which metrics of viral fitness have the strongest relationships with A(H3N2) epidemic burden and timing. Among our set of candidate evolutionary predictors, genetic distances based on broad sets of epitope sites (HA = 129 sites; NA = 223 epitope sites) had the strongest, most consistent associations with A(H3N2) epidemic size, transmission rate, severity, subtype dominance, and age-specific patterns. Increased epitope distance in both H3 and N2 correlated with larger epidemics and increased transmissibility, with univariate analyses finding H3 distance more strongly correlated with epidemic size, peak incidence, transmissibility, and excess mortality, and N2 distance more strongly correlated with epidemic intensity (i.e., the "sharpness" of the epidemic curve) and subtype dominance patterns. However, we note that minor differences in correlative strength between H3 and N2 epitope distance are not necessarily biologically relevant and could be attributed to noise in epidemiological or virological data or the limited number of influenza seasons in our study. The fraction of ILI cases in children relative to adults was negatively correlated with N2 epitope distance, consistent with the expectation that cases are more restricted to immunologically naïve children in seasons with low antigenic novelty [7,78]. Regarding epidemic timing, the number of days from epidemic onset to peak (a proxy for epidemic speed) decreased with N2 epitope distance, but other measures of epidemic timing, such as peak week, onset week, and spatiotemporal synchrony across HHS regions, were not significantly correlated with H3 or N2 antigenic change.

The local branching index (LBI) is traditionally used to predict the success of individual clades, with a high LBI value indicating high viral fitness [35,62]. In our epidemiological analysis, low diversity of H3 or N2 LBI values, in the current or prior season, correlated with greater epidemic intensity, higher transmission rates, and shorter seasonal duration. This outcome suggests that low LBI diversity is indicative of a rapid selective sweep by one successful clade and that high LBI diversity is indicative of multiple co-circulating clades with variable seeding times over the course of an epidemic. A caveat is that LBI estimation is more sensitive to sequence sub-sampling schemes than strain-level measures. If an epidemic is very short and intense (e.g., 1-2 months), a phylogenetic tree with our sub-sampling scheme (50 sequences per month) may not incorporate enough sequences to capture the true diversity of LBI values in that season.

Positive associations between H3 antigenic drift and population-level epidemic burden are consistent with
previous observations from theoretical models [25,26,83]. For example, phylodynamic models of
punctuated antigenic evolution have reproduced key features of A(H3N2) phylogenetic patterns and case
dynamics, such as the sequential replacement of antigenic clusters, the limited standing diversity in HA
after a cluster transition, and higher incidence and attack rates in cluster transition years [25,26,83]. Our
results also corroborate empirical analyses of surveillance data [6,27,28,66] and forecasting models of
annual epidemics [29,30] that found direct, quantitative links between HA antigenic novelty and the
number of influenza cases or deaths in a season. Moving beyond the paradigm of antigenic clusters, Wolf
et al., 2010 and Bedford et al., 2014 demonstrated that smaller, year-to-year changes in H3 antigenic drift
also correlate with seasonal severity and incidence [6,27]. A more recent study did not detect an
association between antigenic drift and city-level epidemic size in Australia [31], though the authors used
a binary indicator to signify seasons with major HA antigenic transitions and did not consider smaller,
more gradual changes in antigenicity. While Lam and colleagues did not observe a consistent effect of
antigenic change on epidemic magnitude, they found a negative relationship between the cumulative prior
incidence of an antigenic variant and its probability of successful epidemic initiation in a city [31].

We did not observe a clear relationship between H3 receptor binding site (RBS) distance and epidemic
burden, even though single substitutions at these seven amino acid positions are implicated in major
antigenic transitions [68,84]. The outperformance of the RBS distance metric by a broader set of epitope
sites could be attributed to the tempo of antigenic cluster changes. A(H3N2) viruses are characterized by
both continuous and punctuated antigenic evolution, with transitions between antigenic clusters occurring
every 2 to 8 years [6,26,32,33,36,37,67,68,85]. Counting substitutions at only a few sites may fail to
capture more modest, gradual changes in antigenicity that are on a time scale congruent with annual
outbreaks. Further, a broader set of epitope sites may better capture the epistatic interactions that
underpin antigenic change in HA [86]. Although the 7 RBS sites were responsible for the majority of
antigenic phenotype in Koel et al.'s experimental study [68], their findings do not necessarily contradict
studies that found broader sets of sites associated with antigenic change. Mutations at other epitope sites
may collectively add to the decreased recognition of antibodies or affect viral fitness through alternate
mechanisms (e.g., compensatory or permissive mutations) [26,32,36,62,68,86-88].

A key result from our study is the direct link between NA antigenic drift and A(H3N2) incidence patterns.
Although HA and NA both contribute to antigenicity [20,89] and undergo similar rates of positive selection
[34], we expected antigenic change in HA to exhibit stronger associations with seasonal incidence, given
its immunodominance relative to NA [90]. H3 and N2 epitope distance were both moderately correlated
with epidemic size, peak incidence, and subtype dominance patterns, but, except for subtype dominance,
H3 epitope distance had higher variable importance rankings in random forest models and N2 epitope
distance was not retained after post-hoc model selection of top ranked random forest features. However,
N2 epitope distance but not H3 epitope distance was associated with faster epidemic speed and a greater
fraction of ILI cases in adults relative to children. Antigenic changes in H3 and N2 were independent
across the 22 seasons of our study, consistent with previous research [34,74,76]. Thus, the similar
predictive performance of HA and NA epitope distance for some epidemic metrics does not necessarily
stem from the coevolution of HA and NA.

HI $\log_2$ titer distance was positively correlated with different measures of epidemic impact yet
underperformed in comparison to H3 and N2 epitope distances. This outcome was surprising given that
we expected our method for generating titer distances to produce more realistic estimates of immune
cross-protection between viruses than epitope-based measures. Our computational approach for inferring
HI phenotype dynamically incorporates newer titer measurements and assigns antigenic weight to
phylogenetic branches rather than fixed sequence positions [35,63], while our method for calculating
epitope distance assumes that the contributions of specific sites to antigenic drift are constant through
time, even though beneficial mutations previously observed at these sites are contingent on historical
patterns of viral fitness and host immunity [26,35,62]. HI titer measurements have been more useful than

epitope substitutions in predicting future A(H3N2) viral populations [35] and vaccine effectiveness [91], with the caveat that these targets are more proximate to viral evolution than epidemic dynamics.

HI titer measurements may be more immunologically relevant than epitope-based measures, yet several factors could explain why substitutions at epitope sites outperformed HI titer distances in epidemiological predictions. First, epitope distances may capture properties that affect viral fitness (and in turn outbreak intensity) but are unrelated to immune escape, such as intrinsic transmissibility, ability to replicate, or epistatic interactions. A second set of factors concern methodological issues associated with HI assays. The reference anti-sera for HI assays are routinely produced in ferrets recovering from their first influenza virus infection. Most humans are infected by different influenza virus strains over the course of their lifetimes, and one's immune history influences the specificity of antibodies generated against drifted influenza virus strains [92-95]. Thus, human influenza virus antibodies, especially those of adults, have more heterogeneous specificities than anti-sera from immunologically naïve ferrets [92].

A related methodological issue is that HI assays disproportionately measure anti-HA antibodies that bind near the receptor binding site and, similar to the RBS distance metric, may capture only a partial view of the antigenic change occurring in the HA protein [31,78,96,97]. A recent study of longitudinal serological data found that HI titers are a good correlate of protective immunity for children, while time since infection is a better predictor of protection for adults [97]. This outcome is consistent with the concept of antigenic seniority, in which an individual's first exposure to influenza virus during childhood leaves an immunological "imprint", and exposure to new strains "back boosts" one's antibody response to strains of the same subtype encountered earlier in life [78,98,99]. Ranjeva et al.'s study and others suggest that human influenza virus antibodies shift focus from the HA head to other more conserved epitopes as individuals age [78,96]. Given that HI assays primarily target epitopes adjacent to the RBS, HI assays using ferret or human serological data are not necessarily suitable for detecting the broader immune responses of adults. A third explanation for the underperformance of HI titers concerns measurement error. Recent A(H3N2) viruses have reduced binding efficiency in HI assays, which can skew estimates of immune cross-reactivity between viruses [100]. These combined factors could obfuscate the relationship between the antigenic phenotypes inferred from HI assays and population-level estimates of A(H3N2) incidence.

Novel antigenic variants are expected to have higher infectivity in immune populations, leading to earlier epidemics and more rapid geographic spread [19], but few studies have quantitatively tied antigenic drift to epidemic timing or geographic synchrony. Previous studies of pneumonia and influenza-associated mortality observed greater severity or geographic synchrony in seasons with major antigenic transitions [21,24]. A more recent Australian study of lab-confirmed cases also noted greater spatiotemporal synchrony during seasons in which novel H3 antigenic variants emerged, although their assessment was based on virus typing alone (i.e., influenza A or B) [17]. A subsequent Australian study with finer-resolution data on subtype incidence and variant circulation determined that more synchronous epidemics were not associated with drifted A(H3N2) strains [31], and a US-based analysis of ILI data also failed to detect a relationship between HA antigenic cluster transitions and geographic synchrony [16]. In our study, the earliest epidemics tended to occur in seasons with transitions between H3 antigenic clusters (e.g., the emergence of the FU02 cluster in 2003-2004) or vaccine mismatches (e.g., N2 mismatch in 1999-2000, H3 mismatch in 2014-2015) [32,74,101], but there was not a statistically significant correlation between antigenic change and earlier epidemic onsets or peaks. Regarding epidemic speed, the length of time from epidemic onset to peak decreased with N2 epitope distance but not H3 epitope distance. The relationship between antigenic drift and epidemic timing may be ambiguous because external seeding events or climatic factors, such as temperature and absolute humidity, are more important in driving influenza seasonality and the onsets of winter epidemics [7,10-14,16]. Alternatively, the resolution of our epidemiological surveillance data (HHS regions) may not be granular enough to detect a signature of antigenic drift in epidemic timing, though studies of city-level influenza dynamics were also unable to identify a clear relationship [16,31].

After exploring individual correlations between evolutionary indicators and annual epidemics, we considered the effects of influenza A(H1N1) incidence and B incidence on A(H3N2) virus circulation within a season. We detected strong negative associations between A(H1N1) incidence and A(H3N2) epidemic size, peak incidence, transmissibility, and excess mortality, consistent with previous animal, epidemiological, phylodynamic, and theoretical studies that found evidence for cross-immunity between IAV subtypes [53-55,57,59,102]. For example, individuals recently infected with seasonal influenza A viruses are less likely to become infected during subsequent pandemic waves [52,53,102-104], and the early circulation of one influenza virus type or subtype is associated with a reduced total incidence of the other type/subtypes within a season [31,57]. Due to the shared evolutionary history of their internal genes [79], pre-2009 seasonal A(H1N1) viruses may impact A(H3N2) virus circulation to a greater extent than A(H1N1)pdm09 viruses, which have a unique combination of genes that were not identified in animals or humans prior to 2009 [81,105]. We observed similar relationships between A(H3N2) epidemic metrics and A(H1N1) incidence during pre- and post-2009 pandemic seasons, with slightly stronger correlations observed during the pre-2009 period. However, given the small sample size (12 pre-2009 seasons and 9 post-2009 seasons), we cannot fully answer this question.

In our study, univariate correlations between A(H1N1) and A(H3N2) incidence were more pronounced than those observed between A(H3N2) incidence and evolutionary indicators, and A(H1N1) epidemic size was the highest ranked feature by random forest models predicting epidemic size and peak incidence. Consequently, interference between the two influenza A subtypes may be more impactful than viral evolution in determining the size of annual A(H3N2) outbreaks. Concerning epidemic timing, we did not detect a relationship between A(H3N2) antigenic change and the relative timing of A(H3N2) and A(H1N1) cases; specifically, A(H3N2) incidence did not consistently lead A(H1N1) incidence in seasons with greater H3 or N2 antigenic change. Overall, we did not find any indication that influenza B incidence affects A(H3N2) epidemic burden or timing, which is not unexpected, given that few T and B cell epitopes are shared between the two virus types [106].

Lastly, we used random forest models and multivariable linear regression models to assess the relative importance of viral evolution, prior population immunity, co-circulation of other influenza viruses, and vaccine-related parameters in predicting regional A(H3N2) epidemic dynamics. We chose conditional inference random forest models as our primary method of variable selection because several covariates were collinear, relationships between some predictors and target variables were nonlinear, and our goal was inferential rather than predictive. We performed leave-one-season-out cross-validation to tune each model, but, due to the limited number of seasons in our dataset, we were not able to test predictive performance on an independent test set. With the caveat that models were likely overfit to historical data, random forest models produced accurate predictions of regional epidemic size, peak incidence, and subtype dominance patterns, while predictions of epidemic intensity and transmission rates were less exact. The latter two measures could be more closely tied to climatic factors, the timing of influenza case importations from abroad, or mobility patterns [7,13,14,16] or they may be inherently more difficult to predict because their values are more constrained. Random forest models tended to underpredict epidemic burden in seasons with major antigenic transitions, particularly the SY97 seasons (1998-1999, 1999-2000) and the FU02 season (2003-2004), potentially because antigenic jumps of these magnitudes were infrequent during our 22-season study period. An additional step of post-hoc model selection demonstrated that models with only three covariates could also produce accurate fits to observed epidemiological data.

Our study is subject to several limitations, specifically regarding geographic resolution and data availability. First, our analysis is limited to one country with a temperate climate and its findings concerning interactions between A(H3N2), A(H1N1), and type B viruses may not be applicable to tropical or subtropical countries, which experience sporadic epidemics of all three viruses throughout the year [107]. Second, our measure of population-level influenza incidence is derived from regional CDC outpatient data because those data are publicly available starting with the 1997-1998 season. State level outpatient data are not available until after the 2009 A(H1N1) pandemic, and finer resolution data from

electronic health records are accessible in theory but not in the public domain. Access to ILI cases
aggregated at the state or city level, collected over the course of decades, would increase statistical
power and enable us to add more location-specific variables to our analysis, such as climatic and
environmental factors. A third limitation is that we measured influenza incidence by multiplying the rate of
influenza-like illness by the percentage of tests positive for influenza, which does not completely eliminate
the possibility of capturing the activity of other co-circulating respiratory pathogens [11]. Surveillance data
based on more specific diagnosis codes would ensure the exclusion of patients with non-influenza
respiratory conditions. Fourth, our data on the age distribution of influenza cases were derived from ILI
encounters across four broad age groups and did not include test positivity status, virus type/subtype, or
denominator information. Despite the coarseness of these data, we found statistically significant
correlations in the expected directions between N2 antigenic change and the fraction of cases in children
relative to adults. Lastly, a serological assay exists for NA, but NA titer measurements are not widely
available because the assay is labor-intensive and inter-lab variability is high. Thus, we could not test the
performance of NA antigenic phenotype in predicting epidemic dynamics.

Beginning in early 2020, non-pharmaceutical interventions (NPIs), including lockdowns, school closures,
physical distancing, and masking, were implemented in the United States and globally to slow the spread
of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), the virus responsible for the COVID-
19 pandemic. These mitigation measures disrupted the transmission of seasonal influenza viruses and
other directly-transmitted respiratory viruses throughout 2020 and 2021 [108-113], and population
immunity to influenza is expected to have decreased substantially during this period of low circulation
[114,115]. COVID-19 NPIs relaxed during 2021 and 2022, and co-circulation of A(H3N2) and
A(H1N1)pdm09 viruses in the United States resumed during the 2022-2023 influenza season. Our study
concludes with the 2018-2019 season, and thus it is unclear whether our modeling approach would be
useful in projecting seasonal burden during the post-pandemic period, without an additional component to
account for COVID-19-related perturbations to influenza transmission. Further studies will need to
determine whether ecological interactions between influenza viruses have changed or if the effects of
viral evolution and subtype interference on seasonal outbreaks are different in the post-pandemic period.

In conclusion, relationships between A(H3N2) antigenic drift, epidemic impact, and age dynamics are
moderate, with genetic distances based on broad sets of H3 and N2 epitope sites having greater
predictive power than serology-based antigenic distances for the timeframe analyzed. Influenza
epidemiological patterns are consistent with increased population susceptibility in seasons with high
antigenic novelty, and our study is the first to link NA antigenic drift to epidemic burden, timing, and the
age distribution of cases. It is well established that anti-HA and anti-NA antibodies are independent
correlates of immunity [45,47-49,116-118], and the influenza research community has advocated for NA-
based vaccines [39,119]. The connection between NA drift and seasonal incidence further highlights the
importance of monitoring evolution in both HA and NA to inform vaccine strain selection and epidemic
forecasting efforts. Although antigenic change in both HA and NA was correlated with epidemic dynamics,
ecological interactions between influenza A subtypes appear to be more influential than viral evolution in
determining the intensity of annual A(H3N2) epidemics. The aim of our study was to retrospectively
assess the potential drivers of annual A(H3N2) epidemics, yet we cautiously suggest that one could
project the size or intensity of future epidemics based on sequence data and A(H1N1)pdm09 incidence
alone [27,57].

Methods

Unless otherwise noted, data processing and statistical analyses were performed using R version 4.3.0.

Influenza epidemic timing and burden

Influenza-like illness and virological surveillance data

We obtained weekly epidemiological and virological data for influenza seasons 1997-1998 to 2018-2019,
at the U.S. HHS region level [120]. We defined influenza seasons as calendar week 40 in a given year to
calendar week 20 in the following year, with the exception of the 2008-2009 season, which ended in 2009
week 16 due to the emergence of the A(H1N1)pdm09 virus [57].

We extracted syndromic surveillance data for the ten HHS regions from the U.S. Outpatient Influenza-like
Illness Surveillance Network (ILINet) [120]. ILINet consists of approximately 3,200 sentinel outpatient
healthcare providers throughout the United States that report the total number of consultations for any
reason and the number of consultations for influenza-like illness (ILI) every week. ILI is defined as fever
(temperature of 100°F [37.8°C] or greater) and a cough and/or a sore throat. The indicator is based on
the weekly proportion of outpatient consultations for influenza-like illness and is available weighted or
unweighted by regional population size. The number of ILI encounters by age group are also provided (0-
4, 5-24, 25-64, and ≥65), but these data are not weighted by total encounters or population size.

Data on weekly influenza virus type and subtype circulation were obtained from the US CDC's World
Health Organization (WHO) Collaborating Center for Surveillance, Epidemiology and Control of Influenza
[121]. We estimated the weekly number of respiratory samples testing positive for influenza A(H1N1),
A(H1N1)pdm09, A(H3N2), or B at the HHS region level (see Supplementary Methods for details on data
processing). We defined influenza A subtype dominance in each season based on the proportion of
influenza A virus (IAV) positive samples typed as A(H3N2). We defined seasons as A(H3N2) or A(H1N1)
or A(H1N1)pdm09 dominant when ≥70% of IAV positive samples were typed as one IAV subtype and co-
dominant when one IAV subtype comprised 50-69% of IAV positive samples.

For each HHS region, we estimated weekly incidences of influenza A(H3N2), A(H1N1), and B by
multiplying the percentage of influenza-like illness among outpatient visits, weighted by regional
population, with the percentage of respiratory samples testing positive for a particular type/subtype
[57,77]. We combined pre-2009 seasonal A(H1N1) and A(H1N1)pdm09 viruses as A(H1N1) and the
Victoria and Yamagata lineages of influenza B as influenza B. ILI x percent positive (ILI+) is considered a
robust estimate of influenza activity and has been used in multiple prior modeling studies
[6,18,57,77,122]. We used linear interpolation to estimate missing values for time spans of up to four
consecutive weeks.

The emergence of A(H1N1)pdm09 in 2009 altered influenza testing and reporting patterns. We adjusted
weekly incidences for differences in reporting rates between the pre-2009 pandemic period – defined as
1997 week 40 to 2009 week 17 – and the post-pandemic period – defined as the weeks after 2010 week
33. For each region, we scaled pre-pandemic incidences by the ratio of mean weekly ILI+ (for all influenza
type/subtypes combined) in the post-pandemic period to that of the pre-pandemic period. Incidences for
HHS Region 10 were not adjusted for pre- and post-pandemic reporting because surveillance data for this
region were not available before 2009. To account for differences in reporting rates across HHS regions,
we next scaled each region's type/subtype incidences by its mean weekly ILI+ for the entire study period.
Scaled incidences were used in all downstream analyses of epidemic burden and timing.

Epidemic burden and timing

Epidemic burden: We considered three complementary indicators of epidemic burden, separately for
each influenza type/subtype, HHS region, and season. We defined *peak incidence* as the maximum
weekly scaled incidence and *epidemic size* as the cumulative weekly scaled incidence. We also
estimated *epidemic intensity* based on a method previously developed to study variation in the shape
(i.e., sharpness) of influenza epidemics across US cities [123]. Epidemic intensity was based on the
inverse Shannon entropy of the weekly incidence distribution. Epidemic intensity increases when
incidence is more concentrated in particular weeks and decreases when incidence is more evenly spread
across weeks.

Specifically, we defined the incidence distribution p_{ij} as the fraction of influenza incidence in season j that
occurred during week i in a given region, and epidemic intensity v_j as the inverse of the Shannon entropy
of the incidence distribution:

641
$$v_j = \left(-\sum_i p_{ij}\ln p_{ij}\right)^{-1}$$

Epidemic intensity values were normalized to fall between 0 and 1.

Transmission intensity: For each region, we used the Epidemia R package to model annual A(H3N2)
epidemics and to estimate time-varying (instantaneous) reproduction numbers, effective Rt [124,125](see
Supplementary Methods for model details). Epidemia implements a semi-mechanistic Bayesian approach
using the probabilistic programming language Stan [126].

To generate seasonal indicators of transmission intensity, we extracted posterior draws of daily Rt
estimates for each region and season, calculated the median value for each day, and averaged daily
median values by epidemic week. For each region and season, we averaged Rt estimates from the
weeks spanning epidemic onset to epidemic peak (*initial Rt*) and averaged the two highest Rt estimates
(*maximum Rt*). Initial Rt and maximum Rt produced qualitatively similar results in downstream analyses;
we opted to report results for maximum Rt.

Excess pneumonia and influenza deaths attributable to A(H3N2): To measure the epidemic severity each
season, we obtained estimates of seasonal excess mortality attributable to influenza A(H3N2) from
Hansen et al., 2022 [127]. Excess mortality is a measure of the mortality burden of a given pathogen in
excess of a seasonally adjusted baseline, obtained by regressing weekly deaths from broad disease
categories against indicators of influenza virus circulation. Hansen et al. used pneumonia and influenza
(P&I) excess deaths, which is considered the most specific indicator of influenza burden [128]. Deaths
with a mention of P&I (ICD 10: J00-J18) were aggregated by week and age group (<1, 1-4, 5-49, 50-64,
and ≥65) for 1998-2018. Age-specific generalized linear models were fit to observed weekly P&I death
rates, while accounting for influenza and respiratory syncytial virus (RSV) activity and seasonal and
temporal trends. Hansen et al. estimated the weekly national number of excess A(H3N2)-associated
deaths by subtracting the baseline death rate expected in the absence of A(H3N2) circulation (A(H3N2)
model terms set to zero) from the observed P&I death rate. We summed the number of excess A(H3N2)
deaths per 100,000 people from October to May to obtain seasonal age-specific estimates.

Epidemic onset and peak timing: We estimated the regional onsets of A(H1N1), A(H1N1)pdm09,
A(H3N2), and B epidemics each season by fitting piecewise linear models to subtype-specific incidence
curves from week 30 to the first week of maximum incidence. We did not estimate epidemic onsets for
regions with insufficient signal, which we defined as fewer than three weeks of consecutive incidence
and/or greater than 30% of weeks with missing data in a particular season. The timing of the changepoint
in incidence represents epidemic establishment (i.e., sustained transmission) rather the timing of
influenza introduction or arrival [16]. We were able to estimate A(H3N2) onset timing for most seasons,
except for three A(H1N1) dominant seasons: 2000-2001 (0 regions), 2002-2003 (3 regions), and 2009-
2010 (0 regions). We defined epidemic peak timing as the first week of maximum incidence. To measure
spatiotemporal synchrony, we calculated seasonal variation (standard deviation, s.d.) in regional onset
and peak timing [19,27]. To measure the speed of viral spread, we calculated the number of days
between onset and peak and seasonal duration (the number of weeks with non-zero incidence) for each
region. As a sensitivity analysis, we used wavelets to estimate timing differences between A(H3N2),
A(H1N1), A(H1N1)pdm09, and B epidemics (see Supplementary Methods).

Age patterns: We calculated the seasonal proportion of ILI encounters in each age group (0-4 years, 5-24
years, 25-64 years, and ≥65 years). Data for more narrow age groups are available after 2009, but we
chose these four categories to increase the number of seasons in our analysis.

Influenza vaccination coverage and A(H3N2) vaccine effectiveness

Influenza vaccination coverage and effectiveness vary between years and would be expected to affect
the population impact of seasonal outbreaks, and in turn our epidemiologic indicators. We obtained
seasonal estimates of national vaccination coverage for adults 18-49 years and adults ≥65 years from
studies utilizing vaccination questionnaire data collected by the National Health Interview Survey [129-
135]. We did not consider the effects of vaccination coverage in children, due to our inability to find
published estimates for most influenza seasons in our study.

We obtained seasonal estimates of adjusted A(H3N2) vaccine effectiveness (VE) from 32 observational
studies [136-167]. Most of these studies had case-control test-negative designs (N = 30) and took place
in North America (N = 25) or Europe (N = 6). When possible, we limited VE estimates to those for healthy
adults or general populations. When multiple VE studies were available for a given season, we calculated
mean VE as the weighted average of *m* different VE point estimates:

695
$$\frac{\sum_{i=1}^{m} \delta_{VE_i}^{-1/2} VE_i}{\sum_{i=1}^{m} \delta_{VE_i}^{-1/2}}$$

696 Wherein δ_{VE} denotes the width of the 95% confidence interval (CI) for VE_i [91].

697 The 95% CI for the weighted mean VE was calculated as:

698
$$\frac{1}{m} \sqrt{\sum_{i=1}^{m} \left(\delta_{VE_i}\right)^2}$$

Correlations among epidemic metrics

We used Spearman's correlation coefficients to measure pairwise relationships between A(H3N2)
epidemiological indictors. We adjusted P-values for multiple testing using the Benjamini and Hochberg
method [168].

Indicators of influenza A(H3N2) evolution

We considered multiple indicators of influenza evolution based on genetic and phenotypic (serologic)
data, separately for HA and NA.

HA and NA sequence data

We downloaded all H3 sequences and associated metadata from the Global Initiative on Sharing Avian
Influenza Data (GISAID) EpiFlu database [60]. We focused our analysis on complete H3 sequences that
were sampled between January 1, 1997, and October 1, 2019. We prioritized viruses with corresponding
HI titer measurements provided by the WHO Global Influenza Surveillance and Response System
(GISRS) Collaborating Centers and excluded all egg-passaged viruses and sequences with ambiguous
year, month, and day annotations. To account for variation in sequence availability across global regions,
we subsampled the selected sequences five times to representative sets of 50 viruses per month, with
preferential sampling for North America. Each month 25 viruses (when available) were selected from
North America, with even sampling across nine other global regions (Africa, Europe, China, South Asia,
Japan and Korea, Oceania, South America, Southeast Asia, and West Asia) for the remaining 25 viruses.
To ensure proper topology early in the phylogeny, we included reference strains that had been collected
no earlier than 5 years prior to January 1, 1997. The resultant sets of H3 sequences included 10,088 to
10,090 sequences spanning December 25, 1995 – October 1, 2019.

As with the H3 analysis, we downloaded all N2 sequences and associated metadata from GISAID and selected complete N2 sequences that were sampled between January 1, 1997, and October 1, 2019. We excluded all sequences with ambiguous year, month, and day annotations, forced the inclusion of reference strains collected no earlier than 5 years prior to January 1, 1997, and compiled five replicate subsampled datasets with preferential sampling for North America (9,007 to 9,009 sequences; June 8, 1995 – October 1, 2019).

HA serologic data

Hemagglutination inhibition (HI) measurements from ferret sera were provided by WHO GISRS Collaborating Centers in London, Melbourne, Atlanta, and Tokyo. We converted these raw two-fold dilution measurements to log_2 titer drops normalized by the corresponding log_2 autologous measurements [35,63].

Although a phenotypic assay exists for NA, NA inhibiting antibody titers are not routinely measured for influenza surveillance. Therefore, we could not include a phenotypic marker of NA evolution in our study.

Phylogenetic inference

For each set of H3 and N2 sequences, we aligned sequences with the augur align command [169] and MAFFT v7.407 [170]. We inferred initial phylogenies with IQ-TREE v1.6.10 [171]. To reconstruct time-resolved phylogenies, we applied TreeTime v0.5.6 [172] with the augur refine command [173].

Viral fitness metrics

Following Huddleston et al., 2020 [35], we defined the following fitness metrics for each influenza season:

Antigenic drift: We estimated antigenic drift for each H3 strain using either genetic or serologic data. We implemented three sequence-based metrics based on substitutions at putative epitope sites: 129 sites in HA1 [21,64,66,67,174], 7 sites adjacent to the receptor-binding site (RBS) [68], and 34 sites in the HA stalk [70], hereon *HA epitope distance*, *HA RBS distance*, and *HA stalk footprint distance*. To estimate antigenic drift with hemagglutination inhibition (HI) titer data, hereon *HI log_2 titer distance*, we applied the phylogenetic tree model from Neher et al., 2016 [63] to the H3 phylogeny and available HI data for its sequences. The tree model estimates the antigenic drift per branch in units of log_2 titer change.

To estimate N2 antigenic drift, we implemented two sequence-based metrics that count substitutions at putative epitope sites in the NA head: 223 sites [34] or 53 sites [69], hereon *NA epitope distance.*

Mutational load: To estimate mutational load for each H3 and N2 strain, an inverse proxy of viral fitness [61], we implemented metrics that count substitutions at putative non-epitope sites in HA (N = 200) and NA (N = 246), hereon *HA non-epitope distance* and *NA non-epitope distance*. Mutational load metrics produce higher values for strains that are less fit compared to previously circulating strains.

Clade growth: We estimated the seasonal growth of H3 clades and N2 clades with the local branching index (LBI) [62]. To calculate LBI for each H3 and N2 strain, we applied the LBI heuristic algorithm as originally described by Neher et al., 2014 [62] to H3 and N2 phylogenetic trees, respectively. We set the neighborhood parameter, τ, to 0.4 and only considered viruses sampled between the current season t and the previous season t - 1 as contributing to recent clade growth in the current season t. To estimate the diversity of clade growth rates in each season, we binned LBI values by units of 2 into 10 categories ((0-2),(2-4], (4-6], (6-8], (8-10], (10-12],(12-14], (14-16],(16-18], (18-20]) and estimated the Shannon entropy of LBI categories. Here, the Shannon entropy [175] considers both the richness and relative abundance of viral clades with different growth rates and is calculated as follows:

761
$$H' = - \sum_i p_i \ln p_i$$

wherein p_i is the proportion of LBI values belonging to the ith bin.

Antigenic and genetic distance relative to prior seasons

We estimated genetic and antigenic distances between influenza viruses circulating in consecutive
seasons by calculating the mean distance between viruses circulating in the current season *t* and viruses
circulating during the prior season (*t* – 1 year; one season lag) or two prior seasons ago (*t* – 2 years; two
season lag). Seasonal genetic and antigenic distances are greater when currently circulating strains are
more antigenically distinct from previously circulating strains. We used Spearman's correlation
coefficients to measure pairwise relationships between scaled H3 and N2 evolutionary indictors. We
adjusted P-values for multiple testing using the Benjamini and Hochberg method [168].

Univariate relationships between viral fitness, (sub)type interference and A(H3N2) epidemic
impact

We measured univariate associations between national indicators of A(H3N2) viral fitness and regional
A(H3N2) epidemic parameters – peak incidence, epidemic size, effective Rt, epidemic intensity, subtype
dominance, excess P&I deaths, onset timing, peak timing, spatiotemporal synchrony, the number of
weeks from onset to peak, and seasonal duration. We first measured Spearman correlation coefficients
between pairs of scaled fitness indicators and epidemic metrics using 1000 bootstrap replicates of the
original dataset (1000 samples with replacement).

Next, we fit regression models with different distribution families (Gaussian or Gamma) and link functions
(identity, log, or inverse) to observed data and used Bayesian information criterion (BIC) to select the best
fit model, with lower BIC values indicating a better fit to the data. For subtype dominance, epidemic
intensity, and age-specific proportions of ILI cases, we fit Beta regression models with logit links. For
each epidemic metric, we fit the best-performing regression model to the resampled dataset. To measure
the effects of sub(type) interference on A(H3N2) epidemics, the same approach was applied to measure
the univariate relationships between A(H1N1) or B epidemic size and A(H3N2) peak incidence, epidemic
size, effective Rt, epidemic intensity, and excess mortality. As a sensitivity analysis, we tested univariate
relationships between A(H3N2) epidemic metrics and A(H1N1) epidemic size during pre-2009 seasons
(seasonal A(H1N1) viruses) and post-2009 seasons (A(H1N1)pdm09 viruses) separately.

All predictors were centered and scaled prior to measuring Spearman's correlations or fitting regression
models.
791
792 **Selecting relevant predictors of A(H3N2) epidemic impact**
793
794 Next, we explored multivariable approaches that would shed light on the potential mechanisms driving
795 annual epidemic impact. Considering that we had many predictors and relatively few observations (22
796 seasons x 9-10 HHS regions), several covariates were collinear, and our goal was explicative rather than
797 predictive, we settled on methods that tend to select few covariates.
798
We first used conditional inference random forest models to select relevant predictors of A(H3N2)
epidemic size, peak incidence, effective Rt, epidemic intensity, and subtype dominance (party and caret
R packages) [176-179]. Candidate predictors included viral fitness indicators: genetic and antigenic
distance from previously circulating strains and the Shannon entropy of H3 and N2 LBI values in the
current and prior season; proxies for prior natural immunity: A(H3N2) epidemic size in the prior season,
influenza A(H1N1) epidemic size and B epidemic size in the current and prior seasons, and the dominant
sub(type) in the prior season [12]; and vaccine-related parameters: national adult vaccination coverage in

the current and previous season, A(H3N2) vaccine effectiveness in the current and previous season, and H3 and N2 epitope distances between circulating A(H3N2) viruses in the United States and the A(H3N2) vaccine strain in the same season. We did not conduct variable selection analysis for excess A(H3N2) mortality due to data limitations (one national estimate per season). Metrics related to epidemic timing were also excluded from this analysis because we found weak or non-statistically significant associations with most of the candidate evolutionary predictors in univariate analyses.

We created each forest by generating 3,000 regression trees from 10 repeats of a leave-one-season-out (jackknife) cross-validated sample of the data. Due to the small size of our dataset, evaluating the predictive accuracy of random forest models on a quasi-independent test set produced unstable estimates. Consequently, we included all data in the training set and report root mean squared error (RMSE) and R^2 values from the best tuned model. We used permutation importance (N = 50 permutations) to estimate the relative importance of each predictor in determining model outcomes. Permutation importance is the decrease in prediction accuracy when a single feature (predictor) is randomly permuted, with larger values indicating more important variables. Because our features were collinear, we used conditional permutation importance to compute feature importance scores, rather than the standard marginal procedure [177,178,180,181].

As an alternative method for variable selection, we performed LASSO regression on the same cross-validated dataset and report RMSE and R^2 values from the best tuned model (glmnet and caret R packages)[179,182]. Unlike random forest models, this approach assumes linear relationships between predictors and the target variable. LASSO models (L1 penalty) are more restrictive than ridge models (L2 penalty) and elastic net models (combination of L1 and L2 penalties) and will arbitrarily select one variable from a set of collinear variables.

To further reduce the set of predictors for each epidemic metric, we performed model selection with linear regression models that considered all combinations of the top 10 ranked predictors from conditional inference random forest models. Candidate models were limited to three independent variables, and models were compared using BIC. We did not include HHS region or season as fixed or random effects in these models because these variables either did not improve model fit (region) or caused convergence issues (season).

All predictors were centered and scaled prior to fitting random forest or regression models.

Data availability

Sequence data are available from GISAID using accession ids provided in Supplementary file 1. Source code for phylogenetic analyses, inferred HI titers from serological measurements, and evolutionary fitness measurements are available in the GitHub repository https://github.com/blab/perofsky-ili-antigenicity. The five replicate trees for HA and NA can be found at https://nextstrain.org/groups/blab/ under the keyword "perofsky-ili-antigenicity". Epidemiological data, datasets combining seasonal evolutionary fitness measurements and epidemic metrics, and source code for calculating epidemic metrics and performing statistical analyses are available in the GitHub repository https://github.com/aperofsky/H3N2_Antigenic_Epi. Raw serological measurements are restricted from public distribution by previous data sharing agreements.

Acknowledgements

We thank the Influenza Division at the US Centers for Disease Control and Prevention, the Victorian Infectious Diseases Reference Laboratory at the Australian Peter Doherty Institute for Infection and Immunity, the Influenza Virus Research Center at the Japan National Institute of Infectious Diseases, the Crick Worldwide Influenza Centre at the UK Francis Crick Institute for sharing HI titer data. We gratefully acknowledge the authors, originating and submitting laboratories of the sequences from the GISAID

EpiFlu Database on which this research is based (listed in Appendix 1). We thank members of the Fogarty International Center's Division of International Epidemiology and Population Studies (DIEPS) and the Bedford Lab for useful discussions.

Funding information

ACP, CH, and CV were supported by the in-house research division of the Fogarty International Center, US National Institutes of Health. ACP was supported by the NSF Infectious Disease Evolution Across Scales (IDEAS) Research Collaboration Network. JH was supported by NIH NIAID awards F31 AI140714 and R01 AI165821. The work done at the Crick Worldwide Influenza Centre was supported by the Francis Crick Institute receiving core funding from Cancer Research UK (FC001030), the Medical Research Council (FC001030) and the Wellcome Trust (FC001030). SF, KN, NK, SW and HH were supported by the Ministry of Health, Labour and Welfare, Japan (10110400 and 10111800). SW was supported by the Japan Agency for Medical Research and Development (JP22fk0108118 and JP23fk0108662). The WHO Collaborating Centre for Reference and Research on Influenza is supported by the Australia Government Department of Health and Aged Care. The Melbourne WHO Collaborating Centre for Reference and Research on Influenza is supported by the Australian Government Department of Health. Influenza virus work in the Krammer laboratory was partially supported by the NIAID Centers of Excellence for Influenza Research and Surveillance (CEIRS) contract HHSN272201400008C, NIAID Centers of Excellence for Influenza Research and Response (CEIRR) contract 75N93021C00014 (FK), and NIAID CIVIC contract (75N93019C00051). TB was supported by NIH awards NIGMS R35 GM119774 and NIAID R01 AI127893. TB is an Investigator of the Howard Hughes Medical Institute. Funding sources were not involved in study design, data collection and interpretation, or the decision to submit the work for publication.

Disclaimer

The conclusions of this study do not necessarily represent the views of the National Institutes of Health, the Centers for Disease Control and Prevention, or the US government.

Author contributions

Amanda C Perofsky: Conceptualization, Data curation, Software, Formal analysis, Funding acquisition, Validation, Investigation, Visualization, Methodology, Writing - original draft, Project administration, Writing - review and editing; John Huddleston: Data curation, Software, Formal Analysis, Validation, Investigation, Visualization, Methodology, Writing - review and editing; Chelsea Hansen: Data curation, Software, Formal Analysis, Investigation, Writing – review and editing; John R Barnes, Thomas Rowe, Xiyan Xu, Rebecca Kondor, David E Wentworth, Nicola Lewis, Lynne Whittaker, Burcu Ermetal, Ruth Harvey, Monica Galiano, Rodney Stuart Daniels, John W McCauley, Seiichiro Fujisaki, Kazuya Nakamura, Noriko Kishida, Shinji Watanabe, Hideki Hasegawa, Sheena G Sullivan, Ian Barr, Kanta Subbarao: Resources, Investigation, Methodology, Writing - review and editing; Florian Krammer: Data curation, Resources, Investigation, Funding acquisition, Writing - review and editing; Trevor Bedford: Conceptualization, Resources, Software, Supervision, Methodology, Project administration, Funding acquisition; Cécile Viboud: Conceptualization, Resources, Supervision, Methodology, Project administration, Funding acquisition, Writing - review and editing

Competing interests

The WHO Collaborating Centre for Reference and Research on Influenza in Melbourne has a collaborative research and development agreement (CRADA) with CSL Seqirus for isolation of candidate vaccine viruses in cells and an agreement with IFPMA for isolation of candidate vaccine viruses in eggs. SGS reports honoraria from CSL Seqirus, Moderna, Pfizer, and Evo Health. The Icahn School of Medicine at Mount Sinai has filed patent applications relating to influenza virus vaccines, SARS-CoV-2 serological assays, and SARS-CoV-2 vaccines which list FK as co-inventor. Mount Sinai has spun out companies, Kantaro and Castlevax, to market the SARS-CoV-2 related technologies. FK has consulted for Merck and Pfizer (before 2020), and is currently consulting for Pfizer, Seqirus, 3rd Rock Ventures, GSK and Avimex. The Krammer laboratory is also collaborating with Pfizer on animal models of SARS-

912 CoV-2 and with Dynavax on universal influenza virus vaccines. All other authors declare no competing
913 interests.
914
Supplementary Methods

Influenza virological surveillance data

Data on weekly influenza type and subtype circulation were obtained from the US CDC's World Health
Organization (WHO) Collaborating Center for Surveillance, Epidemiology and Control of Influenza [121].
Approximately 100 public health laboratories and 300 clinical laboratories located throughout the United
States report influenza test results to the US CDC, through either the US WHO Collaborating
Laboratories Systems or the National Respiratory and Enteric Virus Surveillance System (NREVSS).
Clinical laboratories test respiratory specimens for diagnostic purposes whereas public health laboratories
primarily test specimens to characterize influenza virus type, subtype, and lineage circulation. Public
health laboratories often receive samples that have already tested positive for influenza at a clinical
laboratory.

We estimated the weekly number of respiratory samples testing positive for influenza A(H1N1), A(H3N2),
or B at the HHS region level. Beginning in the 2015/2016 season, reports from public health and clinical
laboratories are presented separately in the CDC's weekly influenza updates. From 2015 week 40
onwards, we used clinical laboratory data to estimate the proportion of respiratory samples testing
positive for any influenza type/subtype and the proportion of samples testing positive for influenza A or B.
We used public health laboratory data to estimate the proportion of influenza A isolates typed as A(H3N2)
or A(H1N1)pdm09 in each week. Untyped influenza A-positive isolates were assigned to either A(H3N2)
or A(H1N1) according to their proportions among typed isolates. We combined seasonal and pandemic
A(H1N1) as seasonal A(H1N1) influenza and the Victoria and Yamagata lineages of influenza B as
influenza B. We defined influenza A subtype dominance in each season based on the proportion of
influenza A positive samples typed as A(H1N1) or A(H3N2).

A(H3N2) epidemiological model

Prior to R_t estimation, we computed daily case counts by disaggregating weekly A(H3N2) incidence rates
to daily rates (tempdisagg package) [183] and rounding the resultant values to integers. Observed cases
were modelled as a function of latent infections in the population, assuming a negative binomial
distribution. We assumed an infection ascertainment rate of 0.45 [184], a lognormal-distributed infection-
to-symptom-onset time period with mean 1.4 days and standard deviation 1.5 days [185], and a
lognormal-distributed onset-to-case-observation time period with mean 2 days and standard deviation 1.5
days [186]. Thus, the time distribution for infection-to-case-observation was

945
$$\pi \sim \mathrm{lognormal}(1.4,1.5) + \mathrm{lognormal}(2,1.5)$$

Instead of using the renewal equation to propagate infections, we treated infections as latent parameters
in the model, because the additional variance around infections leads to a posterior distribution that is
easier to sample [125]. For the generation time, we assumed a discretized Weibull distribution with mean
3.6 days and standard deviation 1.6 days [187]. To control for temporal autocorrelation, we modelled R_t
as a daily random walk. We assigned the intercept a normal prior with mean $\log 2$ and variance 0.2, which
gives the initial reproduction number R_0 a prior mean of approximately 2.

Epidemic trajectories for each region and season were fit independently using Stan's Hamiltonian Monte
Carlo sampler [188]. For each model, we ran 4 chains, each for 10,000 iterations (including a burn-in
period of 2,000 iterations that was discarded), producing a total posterior sample size of 32,000. We
verified convergence by confirming that all parameters had sufficiently low R hat values (all R hat < 1.1)
and sufficiently large effective sample sizes (>15% of the total sample size).

Wavelet analysis

We applied a wavelet approach to quantify the relative timing of influenza A(H3N2), A(H1N1), and B epidemics in each HHS region. Incidence time series were square root transformed and normalized and then padded with zeros to reduce edge effects. Wavelet coherence was used to determine the degree of synchrony between A(H3N2) versus A(H1N1) incidence and A(H3N2) versus B incidence within each region at multi-year time scales. Statistical significance was assessed using 10,000 Monte Carlo simulations. Coherence measures time- and frequency-specific associations between two wavelet transforms, with high coherence indicating that two non-stationary signals (time series) are associated at a particular time and frequency [82].

Following methodology developed for influenza and other viruses [19,82,189-191], we used continuous wavelet transformations (Morlet) to calculate the phase of seasonal A(H3N2), A(H1N1), and B epidemics. We reconstructed weekly time series of phase angles using wavelet reconstruction [19,192] and extracted the major one-year seasonal component (period 0.8 to 1.2 years) of the Morlet decomposition of A(H3N2), A(H1N1), and B time series. To estimate the relative timing of A(H3N2) and A(H1N1) incidence or A(H3N2) and B incidence in each region, phase angle differences were calculated as phase in A(H3N2) minus phase in A(H1N1) (or B), with a positive value indicating that A(H1N1) (or B) lags A(H3N2).

References

1. Gerdil, C. (2003) The annual production cycle for influenza vaccine. *Vaccine* 21, 1776-1779. 10.1016/s0264-410x(03)00071-9
2. He, D. *et al.* (2015) Global Spatio-temporal Patterns of Influenza in the Post-pandemic Era. *Sci Rep* 5, 11013. 10.1038/srep11013
3. Wraith, S. *et al.* (2022) Homotypic protection against influenza in a pediatric cohort in Managua, Nicaragua. *Nat Commun* 13, 1190. 10.1038/s41467-022-28858-9
4. Hay, A.J. *et al.* (2001) The evolution of human influenza viruses. *Philos Trans R Soc Lond B Biol Sci* 356, 1861-1870. 10.1098/rstb.2001.0999
5. Ferguson, N.M. *et al.* (2005) Strategies for containing an emerging influenza pandemic in Southeast Asia. *Nature* 437, 209-214. 10.1038/nature04017
6. Bedford, T. *et al.* (2014) Integrating influenza antigenic dynamics with molecular evolution. *Elife* 3, e01914. 10.7554/eLife.01914
7. Bedford, T. *et al.* (2015) Global circulation patterns of seasonal influenza viruses vary with antigenic drift. *Nature* 523, 217-220. 10.1038/nature14460
8. Simonsen, L. (1999) The global impact of influenza on morbidity and mortality. *Vaccine* 17, S3-S10. 10.1016/S0264-410X(99)00099-7
9. Viboud, C. *et al.* (2004) Influenza epidemics in the United States, France, and Australia, 1972-1997. *Emerg Infect Dis* 10, 32-39. 10.3201/eid1001.020705
10. Chattopadhyay, I. *et al.* (2018) Conjunction of factors triggering waves of seasonal influenza. *Elife* 7. 10.7554/eLife.30756
11. Kramer, S.C. and Shaman, J. (2019) Development and validation of influenza forecasting for 64 temperate and tropical countries. *PLoS Comput Biol* 15, e1006742. 10.1371/journal.pcbi.1006742
12. Lee, E.C. *et al.* (2018) Deploying digital health data to optimize influenza surveillance at national and local scales. *PLoS Comput Biol* 14, e1006020. 10.1371/journal.pcbi.1006020
13. Shaman, J. *et al.* (2010) Absolute humidity and the seasonal onset of influenza in the continental United States. *PLoS Biol* 8, e1000316. 10.1371/journal.pbio.1000316
14. Shaman, J. and Kohn, M. (2009) Absolute humidity modulates influenza survival, transmission, and seasonality. *Proc Natl Acad Sci U S A* 106, 3243-3248. 10.1073/pnas.0806852106
15. Bedford, T. *et al.* (2010) Global migration dynamics underlie evolution and persistence of human influenza A (H3N2). *PLoS Pathog* 6, e1000918. 10.1371/journal.ppat.1000918
16. Charu, V. *et al.* (2017) Human mobility and the spatial transmission of influenza in the United States. *PLoS Comput Biol* 13, e1005382. 10.1371/journal.pcbi.1005382

17. Geoghegan, J.L. *et al.* (2018) Continental synchronicity of human influenza virus epidemics
despite climatic variation. *PLoS Pathog* 14, e1006780. 10.1371/journal.ppat.1006780
18. Pei, S. *et al.* (2018) Forecasting the spatial transmission of influenza in the United States. *Proc*
Natl Acad Sci U S A 115, 2752-2757. 10.1073/pnas.1708856115
19. Viboud, C. *et al.* (2006) Synchrony, waves, and spatial hierarchies in the spread of influenza.
Science 312, 447-451. 10.1126/science.1125237
20. Nelson, M.I. and Holmes, E.C. (2007) The evolution of epidemic influenza. *Nat Rev Genet* 8, 196-
205. 10.1038/nrg2053
21. Wiley, D.C. *et al.* (1981) Structural identification of the antibody-binding sites of Hong Kong
influenza haemagglutinin and their involvement in antigenic variation. *Nature* 289, 373-378.
10.1038/289373a0
22. Fiore, A.E. *et al.* (2009) Prevention and control of seasonal influenza with vaccines:
recommendations of the Advisory Committee on Immunization Practices (ACIP), 2009. *MMWR*
Recomm Rep 58, 1-52
23. Boni, M.F. *et al.* (2004) Influenza drift and epidemic size: the race between generating and
escaping immunity. *Theor Popul Biol* 65, 179-191. 10.1016/j.tpb.2003.10.002
24. Greene, S.K. *et al.* (2006) Patterns of influenza-associated mortality among US elderly by
geographic region and virus subtype, 1968-1998. *Am J Epidemiol* 163, 316-326.
10.1093/aje/kwj040
25. Koelle, K. *et al.* (2009) Understanding the dynamics of rapidly evolving pathogens through
modeling the tempo of antigenic change: influenza as a case study. *Epidemics* 1, 129-137.
10.1016/j.epidem.2009.05.003
26. Koelle, K. *et al.* (2006) Epochal evolution shapes the phylodynamics of interpandemic influenza A
(H3N2) in humans. *Science* 314, 1898-1903. 10.1126/science.1132745
27. Wolf, Y.I. *et al.* (2010) Projection of seasonal influenza severity from sequence and serological
data. *PLoS Curr* 2, RRN1200. 10.1371/currents.RRN1200
28. Wu, A. *et al.* (2010) Correlation of influenza virus excess mortality with antigenic variation:
application to rapid estimation of influenza mortality burden. *PLoS Comput Biol* 6.
10.1371/journal.pcbi.1000882
29. Axelsen, J.B. *et al.* (2014) Multiannual forecasting of seasonal influenza dynamics reveals
climatic and evolutionary drivers. *Proc Natl Acad Sci U S A* 111, 9538-9542.
10.1073/pnas.1321656111
30. Du, X. *et al.* (2017) Evolution-informed forecasting of seasonal influenza A (H3N2). *Sci Transl*
Med 9. 10.1126/scitranslmed.aan5325
31. Lam, E.K.S. *et al.* (2020) The impact of climate and antigenic evolution on seasonal influenza
virus epidemics in Australia. *Nat Commun* 11, 2741. 10.1038/s41467-020-16545-6
32. Smith, D.J. *et al.* (2004) Mapping the antigenic and genetic evolution of influenza virus. *Science*
305, 371-376. 10.1126/science.1097211
33. Bedford, T. *et al.* (2011) Strength and tempo of selection revealed in viral gene genealogies. *BMC*
Evol Biol 11, 220. 10.1186/1471-2148-11-220
34. Bhatt, S. *et al.* (2011) The genomic rate of molecular adaptation of the human influenza A virus.
Mol Biol Evol 28, 2443-2451. 10.1093/molbev/msr044
35. Huddleston, J. *et al.* (2020) Integrating genotypes and phenotypes improves long-term forecasts
of seasonal influenza A/H3N2 evolution. *Elife* 9. 10.7554/eLife.60067
36. Shih, A.C. *et al.* (2007) Simultaneous amino acid substitutions at antigenic sites drive influenza A
hemagglutinin evolution. *Proc Natl Acad Sci U S A* 104, 6283-6288. 10.1073/pnas.0701396104
37. Suzuki, Y. (2008) Positive selection operates continuously on hemagglutinin during evolution of
H3N2 human influenza A virus. *Gene* 427, 111-116. 10.1016/j.gene.2008.09.012
38. Chen, Y.Q. *et al.* (2018) Influenza Infection in Humans Induces Broadly Cross-Reactive and
Protective Neuraminidase-Reactive Antibodies. *Cell* 173, 417-429 e410.
10.1016/j.cell.2018.03.030

39. Eichelberger, M.C. *et al.* (2018) Neuraminidase as an influenza vaccine antigen: a low hanging
fruit, ready for picking to improve vaccine effectiveness. *Curr Opin Immunol* 53, 38-44.
10.1016/j.coi.2018.03.025
40. Wohlbold, T.J. *et al.* (2015) Vaccination with adjuvanted recombinant neuraminidase induces
broad heterologous, but not heterosubtypic, cross-protection against influenza virus infection in
mice. *mBio* 6, e02556. 10.1128/mBio.02556-14
41. Brett, I.C. and Johansson, B.E. (2005) Immunization against influenza A virus: comparison of
conventional inactivated, live-attenuated and recombinant baculovirus produced purified
hemagglutinin and neuraminidase vaccines in a murine model system. *Virology* 339, 273-280.
10.1016/j.virol.2005.06.006
42. Couch, R.B. *et al.* (1974) Induction of partial immunity to influenza by a neuraminidase-specific
vaccine. *J Infect Dis* 129, 411-420. 10.1093/infdis/129.4.411
43. Johansson, B.E. *et al.* (1993) Infection-permissive immunization with influenza virus
neuraminidase prevents weight loss in infected mice. *Vaccine* 11, 1037-1039. 10.1016/0264-
410x(93)90130-p
44. Kilbourne, E.D. (1976) Comparative efficacy of neuraminidase-specific and conventional
influenza virus vaccines in induction of antibody to neuraminidase in humans. *J Infect Dis* 134,
384-394. 10.1093/infdis/134.4.384
45. Murphy, B.R. *et al.* (1972) Association of serum anti-neuraminidase antibody with resistance to
influenza in man. *N Engl J Med* 286, 1329-1332. 10.1056/NEJM197206222862502
46. Schulman, J.L. *et al.* (1968) Protective effects of specific immunity to viral neuraminidase on
influenza virus infection of mice. *J Virol* 2, 778-786. 10.1128/JVI.2.8.778-786.1968
47. Couch, R.B. *et al.* (2013) Antibody correlates and predictors of immunity to naturally occurring
influenza in humans and the importance of antibody to the neuraminidase. *J Infect Dis* 207, 974-
981. 10.1093/infdis/jis935
48. Memoli, M.J. *et al.* (2016) Evaluation of Antihemagglutinin and Antineuraminidase Antibodies as
Correlates of Protection in an Influenza A/H1N1 Virus Healthy Human Challenge Model. *mBio* 7,
e00417-00416. 10.1128/mBio.00417-16
49. Monto, A.S. *et al.* (2015) Antibody to Influenza Virus Neuraminidase: An Independent Correlate of
Protection. *J Infect Dis* 212, 1191-1199. 10.1093/infdis/jiv195
50. Grebe, K.M. *et al.* (2008) Heterosubtypic immunity to influenza A virus: where do we stand?
Microbes Infect 10, 1024-1029. 10.1016/j.micinf.2008.07.002
51. Ulmer, J.B. *et al.* (1998) Protective CD4+ and CD8+ T cells against influenza virus induced by
vaccination with nucleoprotein DNA. *J Virol* 72, 5648-5653. 10.1128/JVI.72.7.5648-5653.1998
52. Sridhar, S. *et al.* (2013) Cellular immune correlates of protection against symptomatic pandemic
influenza. *Nat Med* 19, 1305-1312. 10.1038/nm.3350
53. Epstein, S.L. (2006) Prior H1N1 influenza infection and susceptibility of Cleveland Family Study
participants during the H2N2 pandemic of 1957: an experiment of nature. *J Infect Dis* 193, 49-53.
10.1086/498980
54. Sonoguchi, T. *et al.* (1985) Cross-subtype protection in humans during sequential, overlapping,
and/or concurrent epidemics caused by H3N2 and H1N1 influenza viruses. *J Infect Dis* 151, 81-
88. 10.1093/infdis/151.1.81
55. Ferguson, N.M. *et al.* (2003) Ecological and immunological determinants of influenza evolution.
Nature 422, 428-433. 10.1038/nature01509
56. Cowling, B.J. *et al.* (2014) Incidence of influenza virus infections in children in Hong Kong in a 3-
year randomized placebo-controlled vaccine study, 2009-2012. *Clin Infect Dis* 59, 517-524.
10.1093/cid/ciu356
57. Goldstein, E. *et al.* (2011) Predicting the epidemic sizes of influenza A/H1N1, A/H3N2, and B: a
statistical method. *PLoS Med* 8, e1001051. 10.1371/journal.pmed.1001051
58. Steinhoff, M.C. *et al.* (1993) Effect of heterosubtypic immunity on infection with attenuated
influenza A virus vaccines in young children. *J Clin Microbiol* 31, 836-838. 10.1128/jcm.31.4.836-
838.1993

59. Gatti, L. *et al.* (2022) Cross-reactive immunity potentially drives global oscillation and opposed alternation patterns of seasonal influenza A viruses. *Sci Rep* 12, 8883. 10.1038/s41598-022-08233-w

60. Shu, Y. and McCauley, J. (2017) GISAID: Global initiative on sharing all influenza data - from vision to reality. *Euro Surveill* 22. 10.2807/1560-7917.ES.2017.22.13.30494

61. Luksza, M. and Lassig, M. (2014) A predictive fitness model for influenza. *Nature* 507, 57-61. 10.1038/nature13087

62. Neher, R.A. *et al.* (2014) Predicting evolution from the shape of genealogical trees. *Elife* 3. 10.7554/eLife.03568

63. Neher, R.A. *et al.* (2016) Prediction, dynamics, and visualization of antigenic phenotypes of seasonal influenza viruses. *Proc Natl Acad Sci U S A* 113, E1701-1709. 10.1073/pnas.1525578113

64. Bush, R.M. *et al.* (1999) Predicting the evolution of human influenza A. *Science* 286, 1921-1925. 10.1126/science.286.5446.1921

65. Webster, R.G. and Laver, W.G. (1980) Determination of the number of nonoverlapping antigenic areas on Hong Kong (H3N2) influenza virus hemagglutinin with monoclonal antibodies and the selection of variants with potential epidemiological significance. *Virology* 104, 139-148. 10.1016/0042-6822(80)90372-4

66. Wilson, I.A. and Cox, N.J. (1990) Structural basis of immune recognition of influenza virus hemagglutinin. *Annu Rev Immunol* 8, 737-771. 10.1146/annurev.iy.08.040190.003513

67. Wolf, Y.I. *et al.* (2006) Long intervals of stasis punctuated by bursts of positive selection in the seasonal evolution of influenza A virus. *Biol Direct* 1, 34. 10.1186/1745-6150-1-34

68. Koel, B.F. *et al.* (2013) Substitutions near the receptor binding site determine major antigenic change during influenza virus evolution. *Science* 342, 976-979. 10.1126/science.1244730

69. Krammer, F. (2023) Unpublished.

70. Kirkpatrick, E. *et al.* (2018) The influenza virus hemagglutinin head evolves faster than the stalk domain. *Sci Rep* 8, 10432. 10.1038/s41598-018-28706-1

71. Krammer, F. (2019) The human antibody response to influenza A virus infection and vaccination. *Nat Rev Immunol* 19, 383-397. 10.1038/s41577-019-0143-6

72. Margine, I. *et al.* (2013) H3N2 influenza virus infection induces broadly reactive hemagglutinin stalk antibodies in humans and mice. *J Virol* 87, 4728-4737. 10.1128/JVI.03509-12

73. Nachbagauer, R. *et al.* (2016) Age Dependence and Isotype Specificity of Influenza Virus Hemagglutinin Stalk-Reactive Antibodies in Humans. *mBio* 7, e01996-01915. 10.1128/mBio.01996-15

74. Sandbulte, M.R. *et al.* (2011) Discordant antigenic drift of neuraminidase and hemagglutinin in H1N1 and H3N2 influenza viruses. *Proc Natl Acad Sci U S A* 108, 20748-20753. 10.1073/pnas.1113801108

75. Kilbourne, E.D. *et al.* (1990) Independent and disparate evolution in nature of influenza A virus hemagglutinin and neuraminidase glycoproteins. *Proc Natl Acad Sci U S A* 87, 786-790. 10.1073/pnas.87.2.786

76. Schulman, J.L. and Kilbourne, E.D. (1969) Independent variation in nature of hemagglutinin and neuraminidase antigens of influenza virus: distinctiveness of hemagglutinin antigen of Hong Kong-68 virus. *Proc Natl Acad Sci U S A* 63, 326-333. 10.1073/pnas.63.2.326

77. Goldstein, E. *et al.* (2012) Improving the estimation of influenza-related mortality over a seasonal baseline. *Epidemiology* 23, 829-838. 10.1097/EDE.0b013e31826c2dda

78. Gostic, K.M. *et al.* (2019) Childhood immune imprinting to influenza A shapes birth year-specific risk during seasonal H1N1 and H3N2 epidemics. *PLoS Pathog* 15, e1008109. 10.1371/journal.ppat.1008109

79. Webster, R.G. *et al.* (1992) Evolution and ecology of influenza A viruses. *Microbiological Reviews* 56, 152-179. doi:10.1128/mr.56.1.152-179.1992

80. Garten, R.J. *et al.* (2009) Antigenic and Genetic Characteristics of Swine-Origin 2009 A(H1N1) Influenza Viruses Circulating in Humans. *Science* 325, 197-201. doi:10.1126/science.1176225

81. Smith, G.J. *et al.* (2009) Origins and evolutionary genomics of the 2009 swine-origin H1N1
influenza A epidemic. *Nature* 459, 1122-1125. 10.1038/nature08182
82. Johansson, M.A. *et al.* (2009) Multiyear climate variability and dengue--El Nino southern
oscillation, weather, and dengue incidence in Puerto Rico, Mexico, and Thailand: a longitudinal
data analysis. *PLoS Med* 6, e1000168. 10.1371/journal.pmed.1000168
83. Bedford, T. *et al.* (2012) Canalization of the evolutionary trajectory of the human influenza virus.
BMC Biol 10, 38. 10.1186/1741-7007-10-38
84. Petrova, V.N. and Russell, C.A. (2018) The evolution of seasonal influenza viruses. *Nat Rev*
Microbiol 16, 47-60. 10.1038/nrmicro.2017.118
85. Koelle, K. and Rasmussen, D.A. (2015) The effects of a deleterious mutation load on patterns of
influenza A/H3N2's antigenic evolution in humans. *Elife* 4, e07361. 10.7554/eLife.07361
86. Kryazhimskiy, S. *et al.* (2011) Prevalence of epistasis in the evolution of influenza A surface
proteins. *PLoS Genet* 7, e1001301. 10.1371/journal.pgen.1001301
87. Gong, L.I. *et al.* (2013) Stability-mediated epistasis constrains the evolution of an influenza
protein. *Elife* 2, e00631. 10.7554/eLife.00631
88. Myers, J.L. *et al.* (2013) Compensatory hemagglutinin mutations alter antigenic properties of
influenza viruses. *J Virol* 87, 11168-11172. 10.1128/JVI.01414-13
89. Webster, R.G. *et al.* (1982) Molecular mechanisms of variation in influenza viruses. *Nature* 296,
115-121. 10.1038/296115a0
90. Altman, M.O. *et al.* (2015) Lamprey VLRB response to influenza virus supports universal rules of
immunogenicity and antigenicity. *Elife* 4. 10.7554/eLife.07467
91. Ndifon, W. *et al.* (2009) On the use of hemagglutination-inhibition for influenza surveillance:
surveillance data are predictive of influenza vaccine effectiveness. *Vaccine* 27, 2447-2452.
10.1016/j.vaccine.2009.02.047
92. Hensley, S.E. (2014) Challenges of selecting seasonal influenza vaccine strains for humans with
diverse pre-exposure histories. *Curr Opin Virol* 8, 85-89. 10.1016/j.coviro.2014.07.007
93. Lee, J.M. *et al.* (2019) Mapping person-to-person variation in viral mutations that escape
polyclonal serum targeting influenza hemagglutinin. *Elife* 8. 10.7554/eLife.49324
94. Li, Y. *et al.* (2013) Immune history shapes specificity of pandemic H1N1 influenza antibody
responses. *J Exp Med* 210, 1493-1500. 10.1084/jem.20130212
95. Miller, M.S. *et al.* (2013) Neutralizing antibodies against previously encountered influenza virus
strains increase over time: a longitudinal analysis. *Sci Transl Med* 5, 198ra107.
10.1126/scitranslmed.3006637
96. Henry, C. *et al.* (2019) Influenza Virus Vaccination Elicits Poorly Adapted B Cell Responses in
Elderly Individuals. *Cell Host Microbe* 25, 357-366 e356. 10.1016/j.chom.2019.01.002
97. Ranjeva, S. *et al.* (2019) Age-specific differences in the dynamics of protective immunity to
influenza. *Nat Commun* 10, 1660. 10.1038/s41467-019-09652-6
98. Cobey, S. and Hensley, S.E. (2017) Immune history and influenza virus susceptibility. *Curr Opin*
Virol 22, 105-111. 10.1016/j.coviro.2016.12.004
99. Zhang, A. *et al.* (2019) Original Antigenic Sin: How First Exposure Shapes Lifelong Anti-Influenza
Virus Immune Responses. *J Immunol* 202, 335-340. 10.4049/jimmunol.1801149
100. Zost, S.J. *et al.* (2017) Contemporary H3N2 influenza viruses have a glycosylation site that alters
binding of antibodies elicited by egg-adapted vaccine strains. *Proc Natl Acad Sci U S A* 114,
12578-12583. 10.1073/pnas.1712377114
101. Xie, H. *et al.* (2015) H3N2 Mismatch of 2014-15 Northern Hemisphere Influenza Vaccines and
Head-to-head Comparison between Human and Ferret Antisera derived Antigenic Maps. *Sci Rep*
5, 15279. 10.1038/srep15279
102. Cowling, B.J. *et al.* (2010) Protective efficacy of seasonal influenza vaccination against seasonal
and pandemic influenza virus infection during 2009 in Hong Kong. *Clin Infect Dis* 51, 1370-1379.
10.1086/657311
103. Fox, S.J. *et al.* (2017) Seasonality in risk of pandemic influenza emergence. *PLoS Comput Biol*
13, e1005749. 10.1371/journal.pcbi.1005749

104. Laurie, K.L. *et al.* (2015) Interval Between Infections and Viral Hierarchy Are Determinants of
Viral Interference Following Influenza Virus Infection in a Ferret Model. *J Infect Dis* 212, 1701-
1710. 10.1093/infdis/jiv260
105. Garten, R.J. *et al.* (2009) Antigenic and genetic characteristics of swine-origin 2009 A(H1N1)
influenza viruses circulating in humans. *Science* 325, 197-201. 10.1126/science.1176225
106. Terajima, M. *et al.* (2013) Cross-reactive human B cell and T cell epitopes between influenza A
and B viruses. *Virol J* 10, 244. 10.1186/1743-422x-10-244
107. Yang, W. *et al.* (2020) Dynamic interactions of influenza viruses in Hong Kong during 1998-2018.
PLoS Comput Biol 16, e1007989. 10.1371/journal.pcbi.1007989
108. Cowling, B.J. *et al.* (2020) Impact assessment of non-pharmaceutical interventions against
coronavirus disease 2019 and influenza in Hong Kong: an observational study. *Lancet Public
Health* 5, e279-e288. 10.1016/S2468-2667(20)30090-6
109. Huang, Q.S. *et al.* (2021) Impact of the COVID-19 nonpharmaceutical interventions on influenza
and other respiratory viral infections in New Zealand. *Nat Commun* 12, 1001. 10.1038/s41467-
021-21157-9
110. Olsen, S.J. *et al.* (2020) Decreased influenza activity during the COVID-19 pandemic-United
States, Australia, Chile, and South Africa, 2020. *Am J Transplant* 20, 3681-3685.
10.1111/ajt.16381
111. Olsen, S.J. *et al.* (2021) Changes in Influenza and Other Respiratory Virus Activity During the
COVID-19 Pandemic - United States, 2020-2021. *MMWR Morb Mortal Wkly Rep* 70, 1013-1019.
10.15585/mmwr.mm7029a1
112. Qi, Y. *et al.* (2021) Quantifying the Impact of COVID-19 Nonpharmaceutical Interventions on
Influenza Transmission in the United States. *J Infect Dis* 224, 1500-1508. 10.1093/infdis/jiab485
113. Tempia, S. *et al.* (2021) Decline of influenza and respiratory syncytial virus detection in facility-
based surveillance during the COVID-19 pandemic, South Africa, January to October 2020. *Euro
Surveill* 26. 10.2807/1560-7917.ES.2021.26.29.2001600
114. Ali, S.T. *et al.* (2022) Prediction of upcoming global infection burden of influenza seasons after
relaxation of public health and social measures during the COVID-19 pandemic: a modelling
study. *Lancet Glob Health* 10, e1612-e1622. 10.1016/S2214-109X(22)00358-8
115. Baker, R.E. *et al.* (2020) The impact of COVID-19 nonpharmaceutical interventions on the future
dynamics of endemic infections. *Proc Natl Acad Sci U S A* 117, 30547-30553.
10.1073/pnas.2013182117
116. Gaglani, M. *et al.* (2016) Influenza Vaccine Effectiveness Against 2009 Pandemic Influenza
A(H1N1) Virus Differed by Vaccine Type During 2013-2014 in the United States. *J Infect Dis* 213,
1546-1556. 10.1093/infdis/jiv577
117. Gill, P.W. and Murphy, A.M. (1977) Naturally acquired immunity to influenza type A: a further
prospective study. *Med J Aust* 2, 761-765. 10.5694/j.1326-5377.1977.tb99276.x
118. Hope-Simpson, R.E. (1971) Hong Kong influenza variant. *Br Med J* 3, 531.
10.1136/bmj.3.5773.531-b
119. Krammer, F. *et al.* (2018) NAction! How Can Neuraminidase-Based Immunity Contribute to Better
Influenza Virus Vaccines? *mBio* 9. 10.1128/mBio.02332-17
120. Centers for Disease Control and Prevention (2023). FluView Interactive. Centers for Disease
Control and Prevention
121. World Health Organization (2023). FluNet.
122. Pei, S. *et al.* (2021) Optimizing respiratory virus surveillance networks using uncertainty
propagation. *Nat Commun* 12, 222. 10.1038/s41467-020-20399-3
123. Dalziel, B.D. *et al.* (2018) Urbanization and humidity shape the intensity of influenza epidemics in
U.S. cities. *Science* 362, 75-79. 10.1126/science.aat6030
124. Cori, A. *et al.* (2013) A new framework and software to estimate time-varying reproduction
numbers during epidemics. *Am J Epidemiol* 178, 1505-1512. 10.1093/aje/kwt133
125. Scott, J.A. *et al.* (2021) Epidemia: An R package for semi-mechanistic bayesian modelling of
infectious diseases using point processes. *arXiv preprint arXiv:2110.12461*,

126. Carpenter, B. *et al.* (2017) Stan: A Probabilistic Programming Language. *Journal of Statistical*
Software 76, 1 - 32. 10.18637/jss.v076.i01
127. Hansen, C.L. *et al.* (2022) Mortality Associated With Influenza and Respiratory Syncytial Virus in
the US, 1999-2018. *JAMA Netw Open* 5, e220527. 10.1001/jamanetworkopen.2022.0527
128. Simonsen, L. and Viboud, C. (2012) The art of modeling the mortality impact of winter-seasonal
pathogens. *J Infect Dis* 206, 625-627. 10.1093/infdis/jis419
129. Centers for Disease Control and Prevention (2019). Flu Vaccination Coverage, United States,
2018–19 Influenza Season | FluVaxView | Seasonal Influenza (Flu) | CDC.
130. Jang, S.H. and Kang, J. (2021) Factors Associated with Influenza Vaccination Uptake among
U.S. Adults: Focus on Nativity and Race/Ethnicity. *Int J Environ Res Public Health* 18.
10.3390/ijerph18105349
131. Lu, P.J. *et al.* (2019) Seasonal Influenza Vaccination Coverage Trends Among Adult Populations,
U.S., 2010-2016. *Am J Prev Med* 57, 458-469. 10.1016/j.amepre.2019.04.007
132. Lu, P.J. *et al.* (2013) Seasonal influenza vaccination coverage among adult populations in the
United States, 2005-2011. *Am J Epidemiol* 178, 1478-1487. 10.1093/aje/kwt158
133. National Center for Health Statistics (2008). TABLE: Self-reported influenza vaccination coverage
trends 1989-2008 among adults by age group, risk group, race/ethnicity, health-care worker
status, and pregnancy status. National Health Interview Survey (NHIS)
134. Ward, B. *et al.* (2014). Early Release of Selected Estimates Based on Data From the 2014
National Health Interview Survey. In Statistics, N.C.f.H., ed.
135. Ward, B. *et al.* (2016). Early Release of Selected Estimates Based on Data From the 2015
National Health Interview Survey (05/2016). In Statistics, N.C.f.H., ed.
136. Belongia, E.A. *et al.* (2011) Influenza vaccine effectiveness in Wisconsin during the 2007-08
season: comparison of interim and final results. *Vaccine* 29, 6558-6563.
10.1016/j.vaccine.2011.07.002
137. Bridges, C.B. *et al.* (2000) Effectiveness and cost-benefit of influenza vaccination of healthy
working adults: A randomized controlled trial. *JAMA* 284, 1655-1663. 10.1001/jama.284.13.1655
138. Castilla, J. *et al.* (2016) Effectiveness of subunit influenza vaccination in the 2014-2015 season
and residual effect of split vaccination in previous seasons. *Vaccine* 34, 1350-1357.
10.1016/j.vaccine.2016.01.054
139. Flannery, B. *et al.* (2019) Influenza Vaccine Effectiveness in the United States During the 2016-
2017 Season. *Clin Infect Dis* 68, 1798-1806. 10.1093/cid/ciy775
140. Flannery, B. *et al.* (2020) Spread of Antigenically Drifted Influenza A(H3N2) Viruses and Vaccine
Effectiveness in the United States During the 2018-2019 Season. *J Infect Dis* 221, 8-15.
10.1093/infdis/jiz543
141. Flannery, B. *et al.* (2016) Enhanced Genetic Characterization of Influenza A(H3N2) Viruses and
Vaccine Effectiveness by Genetic Group, 2014-2015. *J Infect Dis* 214, 1010-1019.
10.1093/infdis/jiw181
142. Jackson, M.L. *et al.* (2017) Influenza Vaccine Effectiveness in the United States during the 2015-
2016 Season. *N Engl J Med* 377, 534-543. 10.1056/NEJMoa1700153
143. Janjua, N.Z. *et al.* (2012) Estimates of influenza vaccine effectiveness for 2007-2008 from
Canada's sentinel surveillance system: cross-protection against major and minor variants. *J Infect*
Dis 205, 1858-1868. 10.1093/infdis/jis283
144. Kawai, N. *et al.* (2003) A prospective, Internet-based study of the effectiveness and safety of
influenza vaccination in the 2001-2002 influenza season. *Vaccine* 21, 4507-4513. 10.1016/s0264-
410x(03)00508-5
145. Kissling, E. *et al.* (2013) Low and decreasing vaccine effectiveness against influenza A(H3) in
2011/12 among vaccination target groups in Europe: results from the I-MOVE multicentre case-
control study. *Euro Surveill* 18. 10.2807/ese.18.05.20390-en
146. Lester, R.T. *et al.* (2003) Use of, effectiveness of, and attitudes regarding influenza vaccine
among house staff. *Infect Control Hosp Epidemiol* 24, 839-844. 10.1086/502146
147. McLean, H.Q. *et al.* (2014) Impact of repeated vaccination on vaccine effectiveness against
influenza A(H3N2) and B during 8 seasons. *Clin Infect Dis* 59, 1375-1385. 10.1093/cid/ciu680

148. Ohmit, S.E. *et al.* (2014) Influenza vaccine effectiveness in the 2011-2012 season: protection
against each circulating virus and the effect of prior vaccination on estimates. *Clin Infect Dis* 58,
319-327. 10.1093/cid/cit736
149. Pebody, R. *et al.* (2017) End-of-season influenza vaccine effectiveness in adults and children,
United Kingdom, 2016/17. *Euro Surveill* 22. 10.2807/1560-7917.ES.2017.22.44.17-00306
150. Centers for Disease Control and Prevention (2004) Assessment of the effectiveness of the 2003-
04 influenza vaccine among children and adults--Colorado, 2003. *MMWR Morb Mortal Wkly Rep*
53, 707-710
151. Rolfes, M.A. *et al.* (2019) Effects of Influenza Vaccination in the United States During the 2017-
2018 Influenza Season. *Clin Infect Dis* 69, 1845-1853. 10.1093/cid/ciz075
152. Simpson, C.R. *et al.* (2015) Trivalent inactivated seasonal influenza vaccine effectiveness for the
prevention of laboratory-confirmed influenza in a Scottish population 2000 to 2009. *Euro Surveill*
20. 10.2807/1560-7917.es2015.20.8.21043
153. Skowronski, D. *et al.* (2005) Effectiveness of vaccine against medical consultation due to
laboratory-confirmed influenza: results from a sentinel physician pilot project in British Columbia,
2004-2005. *Can Commun Dis Rep* 31, 181-191
154. Skowronski, D.M. *et al.* (2007) Estimating vaccine effectiveness against laboratory-confirmed
influenza using a sentinel physician network: results from the 2005-2006 season of dual A and B
vaccine mismatch in Canada. *Vaccine* 25, 2842-2851. 10.1016/j.vaccine.2006.10.002
155. Skowronski, D.M. *et al.* (2009) Component-specific effectiveness of trivalent influenza vaccine as
monitored through a sentinel surveillance network in Canada, 2006-2007. *J Infect Dis* 199, 168-
179. 10.1086/595862
156. Skowronski, D.M. *et al.* (2010) Association between the 2008-09 seasonal influenza vaccine and
pandemic H1N1 illness during Spring-Summer 2009: four observational studies from Canada.
PLoS Med 7, e1000258. 10.1371/journal.pmed.1000258
157. Skowronski, D.M. *et al.* (2012) A sentinel platform to evaluate influenza vaccine effectiveness and
new variant circulation, Canada 2010-2011 season. *Clin Infect Dis* 55, 332-342.
10.1093/cid/cis431
158. Skowronski, D.M. *et al.* (2014) Low 2012-13 influenza vaccine effectiveness associated with
mutation in the egg-adapted H3N2 vaccine strain not antigenic drift in circulating viruses. *PLoS*
One 9, e92153. 10.1371/journal.pone.0092153
159. Skowronski, D.M. *et al.* (2014) Influenza A/subtype and B/lineage effectiveness estimates for the
2011-2012 trivalent vaccine: cross-season and cross-lineage protection with unchanged vaccine.
J Infect Dis 210, 126-137. 10.1093/infdis/jiu048
160. Skowronski, D.M. *et al.* (2017) Interim estimates of 2016/17 vaccine effectiveness against
influenza A(H3N2), Canada, January 2017. *Euro Surveill* 22. 10.2807/1560-
7917.ES.2017.22.6.30460
161. Skowronski, D.M. *et al.* (2022) Influenza Vaccine Effectiveness by A(H3N2) Phylogenetic
Subcluster and Prior Vaccination History: 2016-2017 and 2017-2018 Epidemics in Canada. *J*
Infect Dis 225, 1387-1398. 10.1093/infdis/jiaa138
162. Skowronski, D.M. *et al.* (2016) A Perfect Storm: Impact of Genomic Variation and Serial
Vaccination on Low Influenza Vaccine Effectiveness During the 2014-2015 Season. *Clin Infect*
Dis 63, 21-32. 10.1093/cid/ciw176
163. Skowronski, D.M. *et al.* (2017) Serial Vaccination and the Antigenic Distance Hypothesis: Effects
on Influenza Vaccine Effectiveness During A(H3N2) Epidemics in Canada, 2010-2011 to 2014-
2015. *J Infect Dis* 215, 1059-1099. 10.1093/infdis/jix074
164. Treanor, J.J. *et al.* (2012) Effectiveness of seasonal influenza vaccines in the United States
during a season with circulation of all three vaccine strains. *Clin Infect Dis* 55, 951-959.
10.1093/cid/cis574
165. Valenciano, M. *et al.* (2018) Exploring the effect of previous inactivated influenza vaccination on
seasonal influenza vaccine effectiveness against medically attended influenza: Results of the
European I-MOVE multicentre test-negative case-control study, 2011/2012-2016/2017. *Influenza*
Other Respir Viruses 12, 567-581. 10.1111/irv.12562

166. van Doorn, E. *et al.* (2017) Influenza vaccine effectiveness estimates in the Dutch population from 2003 to 2014: The test-negative design case-control study with different control groups. *Vaccine* 35, 2831-2839. 10.1016/j.vaccine.2017.04.012

167. Zimmerman, R.K. *et al.* (2016) 2014-2015 Influenza Vaccine Effectiveness in the United States by Vaccine Type. *Clin Infect Dis* 63, 1564-1573. 10.1093/cid/ciw635

168. Benjamini, Y. and Hochberg, Y. (1995) Controlling the False Discovery Rate - a Practical and Powerful Approach to Multiple Testing. *J R Stat Soc B* 57, 289-300. DOI 10.1111/j.2517-6161.1995.tb02031.x

169. Hadfield, J. *et al.* (2018) Nextstrain: real-time tracking of pathogen evolution. *Bioinformatics* 34, 4121-4123. 10.1093/bioinformatics/bty407

170. Katoh, K. *et al.* (2002) MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res* 30, 3059-3066. 10.1093/nar/gkf436

171. Nguyen, L.T. *et al.* (2015) IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* 32, 268-274. 10.1093/molbev/msu300

172. Sagulenko, P. *et al.* (2018) TreeTime: Maximum-likelihood phylodynamic analysis. *Virus Evol* 4, vex042. 10.1093/ve/vex042

173. Huddleston, J. *et al.* (2021) Augur: a bioinformatics toolkit for phylogenetic analyses of human pathogens. *J Open Source Softw* 6. 10.21105/joss.02906

174. Munoz, E.T. and Deem, M.W. (2005) Epitope analysis for influenza vaccine design. *Vaccine* 23, 1144-1148. 10.1016/j.vaccine.2004.08.028

175. Shannon, C.E. (1948) A mathematical theory of communication. *The Bell system technical journal* 27, 379-423

176. Hothorn, T. *et al.* (2006) Survival ensembles. *Biostatistics* 7, 355-373. 10.1093/biostatistics/kxj011

177. Strobl, C. *et al.* (2007) Bias in random forest variable importance measures: illustrations, sources and a solution. *BMC Bioinformatics* 8, 25. 10.1186/1471-2105-8-25

178. Strobl, C. *et al.* (2008) Conditional variable importance for random forests. *BMC Bioinformatics* 9, 307. 10.1186/1471-2105-9-307

179. Kuhn, M. (2008) Building Predictive Models in R Using the caret Package. *Journal of Statistical Software* 28, 1 - 26. 10.18637/jss.v028.i05

180. Altmann, A. *et al.* (2010) Permutation importance: a corrected feature importance measure. *Bioinformatics* 26, 1340-1347. 10.1093/bioinformatics/btq134

181. Debeer, D. and Strobl, C. (2020) Conditional permutation importance revisited. *BMC Bioinformatics* 21, 307. 10.1186/s12859-020-03622-2

182. Friedman, J. *et al.* (2010) Regularization Paths for Generalized Linear Models via Coordinate Descent. *J Stat Softw* 33, 1-22. 10.18637/jss.v033.i01

183. Sax, C. and Steiner, P. (2013) Temporal Disaggregation of Time Series. *R J.* 5, 80

184. Biggerstaff, M. *et al.* (2014) Influenza-like illness, the time to seek healthcare, and influenza antiviral receipt during the 2010-2011 influenza season-United States. *J Infect Dis* 210, 535-544. 10.1093/infdis/jiu224

185. Lessler, J. *et al.* (2009) Incubation periods of acute respiratory viral infections: a systematic review. *Lancet Infect Dis* 9, 291-300. 10.1016/s1473-3099(09)70069-6

186. Russell, K.E. *et al.* (2018) Comparison of outpatient medically attended and community-level influenza-like illness-New York City, 2013-2015. *Influenza Other Respir Viruses* 12, 336-343. 10.1111/irv.12540

187. Cowling, B.J. *et al.* (2009) Estimation of the serial interval of influenza. *Epidemiology* 20, 344-347. 10.1097/EDE.0b013e31819d1092

188. Hoffman, M.D. and Gelman, A. (2014) The No-U-Turn sampler: adaptively setting path lengths in Hamiltonian Monte Carlo. *J. Mach. Learn. Res.* 15, 1593-1623

189. Grenfell, B.T. *et al.* (2001) Travelling waves and spatial hierarchies in measles epidemics. *Nature* 414, 716-723. 10.1038/414716a

190. Liebhold, A. *et al.* (2004) Spatial Synchrony in Population Dynamics. *Annual Review of Ecology, Evolution, and Systematics* 35, 467-490. 10.1146/annurev.ecolsys.34.011802.132516

191. Weinberger, D.M. *et al.* (2012) Influenza epidemics in Iceland over 9 decades: changes in timing and synchrony with the United States and Europe. *Am J Epidemiol* 176, 649-655. 10.1093/aje/kws140

192. Torrence, C. and Compo, G.P. (1998) A Practical Guide to Wavelet Analysis. *Bulletin of the American Meteorological Society* 79, 61-78. https://doi.org/10.1175/1520-0477(1998)079<0061:APGTWA>2.0.CO;2

Figures

1434

Figure 1. Antigenic and genetic evolution of seasonal influenza A(H3N2) viruses, 1997 - 2019. A-B.
Temporal phylogenies of hemagglutinin (H3) and neuraminidase (N2) gene segments. Tip color denotes
the Hamming distance from the root of the tree, based on the number of substitutions at epitope sites in
H3 (N = 129 sites) and N2 (N = 223 sites). "X" marks indicate the phylogenetic positions of US
recommended vaccine strains. **C-D.** Seasonal genetic and antigenic distances are the mean distance
between A(H3N2) viruses circulating in the current season t versus the prior season (t − 1), measured by
C. four sequence-based metrics (HA receptor binding site (RBS), HA stalk footprint, HA epitope, and NA
epitope) and **D.** hemagglutination inhibition (HI) titer measurements. **E.** The Shannon entropy of H3 and
N2 local branching index (LBI) values in each season. Vertical bars in **C, D,** and **E** and are 95%
confidence intervals of seasonal estimates from five bootstrapped phylogenies.

1445
Figure 2. Annual influenza A(H3N2) epidemics in the United States, 1997 - 2019. A. Weekly
incidence of influenza A(H3N2) (red), A(H1N1) (blue), and B (green) averaged across ten HHS regions
(Region 1: Boston; Region 2: New York City; Region 3: Washington, DC; Region 4: Atlanta; Region 5:
Chicago; Region 6: Dallas, Region 7: Kansas City; Region 8: Denver; Region 9: San Francisco; Region
10: Seattle). Time series are 95% confidence intervals of regional incidence estimates. Incidences are the
proportion of influenza-like illness (ILI) visits among all outpatient visits, multiplied by the proportion of
respiratory samples testing positive for each influenza type/subtype. Vertical dashed lines indicate
January 1 of each year. **B.** Intensity of weekly influenza A(H3N2) incidence in ten HHS regions. White
tiles indicate weeks when influenza-like-illness data or virological data were not reported. Weekly time
series for A(H1N1) and B are in Figure S5.

1456

Figure 3. A(H3N2) antigenic drift correlates with larger, more intense annual epidemics. A(H3N2) epidemic size, peak incidence, epidemic intensity, and transmissibility (effective reproduction number, R_t) increase with antigenic drift, measured by **A.** hemagglutinin (H3) epitope distance, and **B.** neuraminidase (N2) epitope distance, and **C.** hemagglutination inhibition (HI) log_2 titer distance. Seasonal antigenic drift is the mean titer distance or epitope distance between viruses circulating in the current season t versus the prior season (t – 1) or two prior seasons (t – 2). Distances are scaled to aid in direct comparison of evolutionary indicators. Point color indicates the dominant influenza A virus (IAV) subtype based on CDC influenza season summary reports (red: A(H3N2), blue: A(H1N1), purple: A(H1N1)pdm09, orange: A(H3N2)/A(H1N1)pdm09 co-dominant), and vertical bands are 95% confidence intervals of regional estimates. Seasonal mean A(H3N2) epidemic metric values were fit as a function of antigenic or genetic distance using LMs (epidemic size, peak incidence), Gaussian GLMs (effective R_t: inverse link), or Beta GLMs (epidemic intensity) with 1000 bootstrap resamples.

Figure 4. The proportion of influenza positive samples typed as A(H3N2) increases with antigenic drift. A-B. Seasonal A(H3N2) subtype dominance increases with H3 and N2 epitope distance. Seasonal epitope distance is the mean epitope distance between viruses circulating in the current season t versus the prior season ($t-1$) or two prior seasons ($t-2$). Distances were scaled to aid in direct comparison of evolutionary indicators. Point color indicates the dominant influenza A virus (IAV) subtype based on CDC influenza season summary reports (red: A(H3N2), blue: A(H1N1), purple: A(H1N1)pdm09, orange: A(H3N2)/A(H1N1)pdm09 co-dominant), and vertical bands are 95% confidence intervals of regional estimates. Seasonal mean A(H3N2) dominance was fit as a function of H3 or N2 epitope distance using Beta GLMs with 1000 bootstrap resamples. **C-D.** Regional patterns of influenza type and subtype incidence during two seasons when A(H3N2) was nationally dominant. **C.** Widespread A(H3N2) dominance during 2003-2004 after the emergence of a novel antigenic cluster, FU02 (A/Fujian/411/2002-like strains). **D.** Spatial heterogeneity in subtype circulation during 2007-2008, a season with low A(H3N2) antigenic novelty relative to the prior season. Pie charts represent the proportion of influenza positive samples typed as A(H3N2) (red), A(H1N1) (blue), or B (green) in each HHS region. Data for Region 10 (purple) were not available for seasons prior to 2009. The sizes of regional pie charts are proportional to the total number of influenza positive samples.

1487

Figure 5. The effects of influenza A(H1N1) and B epidemic size on A(H3N2) epidemic burden. A.
Influenza A(H1N1) epidemic size negatively correlates with A(H3N2) epidemic size, peak incidence,
transmissibility (effective reproduction number, R_t), and epidemic intensity. **B.** Influenza B epidemic size
does not significantly correlate with A(H3N2) epidemic metrics. Point color indicates the dominant
influenza A virus (IAV) subtype based on CDC influenza season summary reports (red: A(H3N2), blue:
A(H1N1), purple: A(H1N1)pdm09, orange: A(H3N2)/A(H1N1)pdm09 co-dominant), and vertical and
horizontal bands are 95% confidence intervals of regional estimates. Seasonal mean A(H3N2) epidemic
metrics were fit as a function of mean A(H1N1) or B epidemic size using Gaussian GLMs (inverse link:
epidemic size, peak incidence; log link: effective R_t) or Beta GLMs (epidemic intensity) with 1000
bootstrap resamples.

Conditional Permutation Importance

Figure 6. Variable importance rankings from conditional inference random forest models predicting A(H3N2) epidemic dynamics. Ranking of variables in predicting regional A(H3N2) **A.** epidemic size, **B.** peak incidence, **C.** effective reproduction number, Rt, **D.** epidemic intensity, and **E.** subtype dominance. Each forest was created by generating 3,000 regression trees from a repeated leave-one-season-out cross-validated sample of the data. Variables are ranked by their conditional permutation importance, with differences in prediction accuracy scaled by the total (null model) error. Black error bars are 95% confidence intervals of conditional permutation scores. Abbreviations: HI titer = hemagglutination inhibition $\log_2$ titer distance, t - 1 = one-season lag, t - 2 = two-season lag, LBI = local branching index, peak = peak incidence, distance to vaccine = epitope distance between currently circulating strains and the recommended vaccine strain, VE = vaccine effectiveness.

1509
Figure 7. Observed versus predicted values of seasonal region-specific A(H3N2) A. epidemic size,
B. peak incidence, C. effective reproduction number, Rt, D. epidemic intensity, and E. subtype
dominance from conditional random forest models. Results are facetted by HHS region and epidemic
metric. Point color and size corresponds to the degree of hemagglutinin (H3) epitope distance in viruses
circulating in season t versus viruses circulating two seasons ago ($t-2$). Large, yellow points indicate
seasons with high antigenic novelty, and small blue points indicate seasons with low antigenic novelty.
Regional Spearman's correlation coefficients and associated P-values are in the top left section of each
facet.

    Tables

1519

1520   **Table 1. Evolutionary indicators of seasonal viral fitness.** Evolutionary indicators are labeled by the
1521   influenza gene for which data are available (hemagglutinin, HA or neuraminidase, NA), the type of data
1522   they are based on, and the component of influenza fitness they represent. Table format is adapted from
1523   Huddleston et al., 2020 [35].
1524

Evolutionary indicator	Influenza gene	Data type	Fitness category	Citations
Mean HI titer $\log_2$ distance from the prior season	HA	Hemagglutinin inhibition assays using ferret sera	Antigenic drift	Huddleston et al., 2020; Neher et al., 2016
Mean epitope distance from the prior season	HA and NA	Sequences	Antigenic drift	Bhatt et al., 2011; Bush et al., 1999; Krammer, unpublished; Webster and Laver, 1980; Wiley et al., 1981; Wilson and Cox, 1990; Wolf et al., 2010
Mean receptor binding site distance from the prior season	HA	Sequences	Antigenic drift	Koel et al., 2013
Mutational load (mean non-epitope distance from the prior season)	HA and NA	Sequences	Functional constraint	Łuksza and Lässig, 2014
Mean stalk "footprint" distance from the prior season	HA	Sequences	Negative control	Kirkpatrick et al., 2018
Mean local branching index	HA and NA	Sequences	Clade growth	Huddleston et al., 2020; Łuksza and Lässig, 2014
Shannon entropy of local branching index	HA and NA	Sequences	Diversity of clade growth rates	Huddleston et al., 2020; Neher et al., 2014

1525
1526

Table 2. Seasonal metrics of A(H3N2) epidemic dynamics. Epidemic metrics are defined and labeled
by which outcome category they represent.

Epidemic Outcome	Definition	Outcome category	Citations
Epidemic size	Cumulative weekly incidence	Burden	
Peak incidence	Maximum weekly incidence	Burden	
Maximum time-varying effective reproduction number, Rt	The number of secondary cases arising from a symptomatic index case, assuming conditions remain the same	Transmissibility	Cori et al., 2013; Scott et al., 2021
Epidemic intensity	Inverse Shannon entropy of the weekly incidence distribution (i.e., the spread of incidence across the season)	Sharpness of the epidemic curve	Dalziel et al., 2018
Subtype dominance	The proportion of influenza positive samples typed as A(H3N2)	Viral activity	
Excess pneumonia and influenza mortality attributable to A(H3N2) virus	Mortality burden in excess of a seasonally adjusted baseline	Severity	Hansen et al., 2022; Simonsen and Viboud, 2012
Onset week	Winter changepoint in incidence	Timing	Charu et al., 2017
Peak week	First week of maximum incidence	Timing	
Spatiotemporal synchrony	Variation (s.d.) in regional onset or peak timing	Speed	Viboud et al., 2006
Onset to peak	Number of days between onset week and peak week	Speed	
Seasonal duration	Number of weeks with non-zero incidence	Speed	

 Table 3. Predictors of seasonal A(H3N2) epidemic burden, transmissibility, intensity, and subtype
1532 **dominance.** Variables retained in the best fit model for each epidemic outcome were determined by BIC.
1533

Outcome	Best Minimal Model[1]	R^2	Adj. R^2	RMSE
Epidemic Size	H3 epitope distance (t-2) + H1 epidemic size + H3 epidemic size (t-1)	0.74	0.69	9.88
Peak Incidence	H3 epitope distance (t-2) + H1 epidemic size + Dominant IAV Subtype (t-1)	0.69	0.63	2.09
Effective Rt	HI titer distance (t-2) + H1 epidemic size + H3 LBI Diversity (t-1)	0.71	0.65	0.1
Epidemic Intensity	HI titer distance (t-2) + N2 distance to vaccine strain + vaccination coverage (t-1)	0.79	0.75	0.07
Subtype Dominance	H3 epitope distance (t-2) + N2 epitope distance (t-1) + Dominant IAV Subtype (t-1)	0.56	0.48	0.2

1534

1535 [1]Candidate models were limited to 3 independent variables and considered all combinations of the top 10
1536 ranked predictors from conditional inference random forest models (Figure 6).

Supplementary Figures

Figure S1. Comparison of seasonal antigenic drift measured by substitutions at hemagglutinin (H3) epitope sites and HI titer measurements, from 1997-1998 to 2018-2019. We used Spearman correlation tests to measure associations between H3 epitope distance and HI titer distance at **A.** one-season lags and **B.** two-season lags. Seasonal antigenic distance is the mean distance between strains circulating in season t and strains circulating in the prior season $t-1$ year (one season lag) or two seasons ago $t-2$ years (two season lag). Seasonal distances are scaled because epitope distance and HI titer distance use different units of measurement. Point labels indicate the current influenza season, and point color denotes the relative timing of influenza seasons, with earlier seasons shaded dark purple (e.g., 1997-1998) and later seasons shaded light yellow (e.g., 2018-2019). H3 epitope distance and HI titer (tree model) distance at two-season lags capture expected "jumps" in antigenic drift during key seasons previously associated with major antigenic transitions [32], such as the SY97 cluster seasons (1997-1998, 1998-1999, 1999-2000) and the FU02 cluster season (2003-2004).

1551

Figure S2. Pairwise correlations between H3 and N2 evolutionary indicators (one season lags). We measured Spearman's correlations between seasonal measures of H3 and N2 evolution, including H3 RBS distance, H3 epitope distance, H3 non-epitope distance, H3 stalk footprint distance, HI titer distance, N2 epitope distance based on 223 or 53 epitope sites, N2 non-epitope distance, mean clade growth of H3 and N2 (local branching index, LBI), and the Shannon entropy of H3 and N2 LBI values. Seasonal distances were estimated as the mean distance between strains circulating in the current season t and those circulating in the prior season ($t - 1$). The Benjamini and Hochberg method was used to adjust P-values for multiple testing. The color of each circle indicates the strength and direction of the association, from dark red (strong positive correlation) to dark blue (strong negative correlation). Stars within circles indicate statistical significance (adjusted $P < 0.05$).

Figure S3. Pairwise correlations between H3 and N2 evolutionary indicators (two season lags). We measured Spearman's correlations between seasonal measures of H3 and N2 evolution, including H3 RBS distance, H3 epitope distance, H3 non-epitope distance, H3 stalk footprint distance, HI titer distance (tree model), N2 epitope distance based on 223 or 53 epitope sites, N2 non-epitope distance, mean clade growth of H3 and N2 (local branching index, LBI), and the Shannon entropy of H3 and N2 LBI values. Seasonal distances were estimated as the mean distance between strains circulating in the current season t and those circulating in the prior season ($t - 1$). The Benjamini and Hochberg method was used to adjust P-values for multiple testing. The color of each circle indicates the strength and direction of the association, from dark red (strong positive correlation) to dark blue (strong negative correlation). Stars within circles indicate statistical significance (adjusted $P < 0.05$).

1573

Figure S4. Comparison of seasonal antigenic drift measured by substitutions at hemagglutinin (H3) and neuraminidase (N2) epitope sites, from 1997-1998 to 2018-2019. We used Spearman correlation tests to measure associations between H3 epitope distance and N2 epitope distance at **A.** one-season lags and **B.** two-season lags. Seasonal epitope distance is the mean distance between strains circulating in season t and strains circulating in the prior season $t-1$ (one season lag) or two seasons ago $t-2$ (two season lag). Point labels indicate the current influenza season, and point color denotes the relative timing of influenza seasons, with earlier seasons shaded dark purple (e.g., 1997-1998) and later seasons shaded light yellow (e.g., 2018-2019). N2 epitope distance at one-season lags captures expected "jumps" in antigenic drift during key seasons previously associated with major antigenic transitions [32], such as the SY97 cluster seasons (1997-1998, 1998-1999, 1999-2000) the FU02 cluster season (2003-2004), and the CA04 cluster season (2004-2005).

Figure S5. Intensity of weekly incidence of A. influenza A(H1N1) and B. influenza B in ten HHS regions, 1997 - 2019. Seasonal and pandemic A(H1N1) were combined as A(H1N1), and the Victoria and Yamagata lineages of influenza B were combined as influenza B. White tiles indicate weeks when either influenza-like-illness cases or virological data were not reported. Data for Region 10 were not available in seasons prior to 2009.

Figure S6. Pairwise correlations between seasonal A(H3N2), A(H1N1), and B epidemic metrics. We
measured Spearman's correlations among indicators of A(H3N2) epidemic timing, including onset week,
peak week, regional variation (s.d.) in onset and peak timing, and the number of days from onset to peak,
indicators of A(H3N2) epidemic magnitude, including epidemic intensity (i.e., the "sharpness" of the
epidemic curve), transmissibility (maximum effective reproduction number, Rt), subtype dominance
patterns, epidemic size, and peak incidence. We also considered relationships between the circulation of
other types/subtypes and A(H3N2) epidemic burden and timing. The Benjamini and Hochberg method
was used to adjust P-values for multiple testing. The color of each circle indicates the strength and
direction of the association, from dark red (strong positive correlation) to dark blue (strong negative
correlation). Stars within circles indicate statistical significance (adjusted P < 0.05).

1603

Figure S7. Univariate correlations between A(H3N2) viral fitness and epidemic impact. Mean Spearman correlation coefficients, 95% confidence intervals of correlation coefficients, and corresponding p-values of bootstrapped (N = 1000) viral fitness indicators (rows) and epidemic metrics (columns). Point color indicates the strength and direction of the association, from dark red (strong positive correlation) to dark blue (strong negative correlation), and stars indicate statistical significance (* P < 0.05, ** P < 0.01, *** P < 0.001). Abbreviations: HI = hemagglutination inhibition, RBS: receptor binding site, t - 1 = one-season lag, t - 2 = two-season lag, LBI = local branching index.

Dominant IAV ● H3 ● H1 ● H1pdm ● H3/H1pdm

Figure S8. Low diversity in the growth rates of circulating A(H3N2) clades is associated with more intense epidemics and higher transmissibility. A(H3N2) effective Rt and epidemic intensity negatively correlate with the diversity of LBI values among circulating A(H3N2) lineages in the current or prior season, measured by the Shannon entropy of **A.** H3 local branching index (LBI) values in the prior season ($t-1$), and **B.** the Shannon entropy of N2 LBI values in the current season t. LBI values are scaled to aid in direct comparisons of H3 and N2 LBI diversity. Point color indicates the dominant influenza A subtype based on CDC influenza season summary reports (red: A(H3N2), blue: A(H1N1), purple: A(H1N1)pdm09, orange: A(H3N2)/A(H1N1)pdm09 co-dominant), and vertical bands are 95% confidence intervals of regional estimates. Mean A(H3N2) epidemic metric values were fit as a function of seasonal LBI diversity using Gaussian GLMs (effective Rt: inverse link) or Beta GLMs (epidemic intensity: logit link) with 1000 bootstrap resamples.

Figure S9. Excess influenza A(H3N2) mortality increases with H3 and N2 antigenic drift, but correlations are not statistically significant. The number of excess influenza deaths attributable to A(H3N2) (per 100,000 people) were estimated from a seasonal regression model fit to weekly pneumonia and influenza-coded deaths [127]. Seasonal epitope distance is the mean distance between strains circulating in season t and those circulating in the prior season ($t-1$) or two seasons ago ($t-2$). Distances are scaled to aid in direct comparison of evolutionary indicators. Point color indicates the dominant influenza A subtype based on CDC influenza season summary reports (red: A(H3N2), blue: A(H1N1), purple: A(H1N1)pdm09, orange: A(H3N2)/A(H1N1)pdm09 co-dominant), and vertical bars are 95% confidence intervals of excess mortality estimates. National excess mortality estimates were fit as a function of seasonal H3 or N2 epitope distance using Gaussian GLMs (log link) with 1000 bootstrap resamples.

1635

Figure S10. Regional patterns of influenza type and subtype incidence from seasons 1997-1998 to 2018-2019. Pie charts represent the proportion of influenza positive samples that were typed as A(H3N2), A(H1N1) or A(H1N1)pdm09, and B in each HHS region. Data for Region 10 (purple) were not available in seasons prior to the 2009 A(H1N1) pandemic.

1640

Figure S11. Univariate correlations between A(H3N2) viral fitness and epidemic timing. Mean Spearman correlation coefficients, 95% confidence intervals of correlation coefficients, and corresponding p-values of bootstrapped (N = 1000) viral fitness indicators (columns) and epidemic timing metrics (rows). Epidemic timing metrics are the week of epidemic onset, regional variation (s.d.) in onset timing, the week of epidemic peak, regional variation (s.d.) in peak timing, the number of days between epidemic onset and peak, and seasonal duration. Color indicates the strength and direction of the association, from dark red (strong positive correlation) to dark blue (strong negative correlation), and stars indicate statistical significance (* P < 0.05, ** P < 0.01, *** P < 0.001). Abbreviations: HI = hemagglutination inhibition, RBS: receptor binding site, t - 1 = one-season lag, t - 2 = two-season lag, LBI = local branching index.

Figure S12. Seasonal duration increases with diversity in clade growth rates of circulating H3 and N2 lineages, measured as the Shannon entropy of local branching index (LBI) values. **A.** H3 LBI diversity and **B.** N2 LBI diversity during the current season positively correlate with seasonal duration. LBI values are scaled to aid in direct comparisons of H3 and N2 LBI diversity. Point color indicates the dominant influenza A subtype based on CDC influenza season summary reports (red: A(H3N2), blue: A(H1N1), purple: A(H1N1)pdm09, orange: A(H3N2)/A(H1N1)pdm09 co-dominant). Mean values of regional season duration were fit as a function of H3 LBI diversity or N2 LBI diversity using Gaussian GLMs (inverse link) with 1000 bootstrap resamples.

1659

Figure S13. Epidemic speed increases with N2 antigenic drift. N2 epitope distance correlates with fewer days from epidemic onset to peak (**A**), while the relationship between H3 epitope distance and epidemic speed is less apparent (**B**). Seasonal epitope distance is the mean distance between strains circulating in season *t* and those circulating in the prior season (*t* − 1) or two seasons ago (*t* − 2). Distances are scaled to aid in direct comparison of evolutionary indicators. Point color indicates the dominant influenza A subtype based on CDC influenza season summary reports (red: A(H3N2), blue: A(H1N1), purple: A(H1N1)pdm09, orange: A(H3N2)/A(H1N1)pdm09 co-dominant). Mean values of regional days from onset to peak were fit as a function of H3 or N2 epitope distance using Gamma GLMs (inverse link) with 1000 bootstrap resamples.

1669

Figure S14. The timing of epidemic onsets and peaks are weakly correlated with H3 and N2 antigenic change. A. Epidemic onsets are earlier in seasons with increased H3 epitope distance ($t-2$), but the correlation is not statistically significant. **B.** Epidemic peaks are earlier in seasons with increased H3 epitope distance ($t-2$) or increased N2 epitope distance ($t-1$), but correlations are not statistically significant. Seasonal epitope distance is the mean distance between strains circulating in season t and those circulating in the prior season ($t-1$) or two seasons ago ($t-2$). Distances are scaled to aid in direct comparison of evolutionary indicators. Point color indicates the dominant influenza A subtype based on CDC influenza season summary reports (red: A(H3N2), blue: A(H1N1), purple: A(H1N1)pdm09, orange: A(H3N2)/A(H1N1)pdm09 co-dominant). Mean values of regional epidemic onsets and peaks were fit as a function of H3 or N2 epitope distance using LMs with 1000 bootstrap resamples.

Figure S15. Univariate correlations between A(H3N2) antigenic change and the age distribution of outpatient influenza-like illness (ILI) cases. Mean Spearman correlation coefficients, 95% confidence intervals of correlation coefficients, and corresponding p-values of bootstrapped (N = 1000) evolutionary indicators (rows) and the proportion of ILI cases in individuals aged < 5 years, 5-24 years, 25-64 years, and ≥ 65 years (columns). Color indicates the strength and direction of the association, from dark red (strong positive correlation) to dark blue (strong negative correlation), and stars indicate statistical significance (* P < 0.05, ** P < 0.01, *** P < 0.001). Abbreviations: HI = hemagglutination inhibition, RBS: receptor binding site, t - 1 = one-season lag, t - 2 = two-season lag.

1689

Figure S16. N2 epitope distance correlates with the age distribution of outpatient influenza-like illness (ILI) cases. Seasonal epitope distance is the mean distance between strains circulating in season t and those circulating in the prior season ($t-1$) or two seasons ago ($t-2$). Distances are scaled to aid in direct comparison of evolutionary indicators. Point color indicates the dominant influenza A subtype based on CDC influenza season summary reports (red: A(H3N2), blue: A(H1N1), purple: A(H1N1)pdm09, orange: A(H3N2)/A(H1N1)pdm09 co-dominant), and vertical bars are 95% confidence intervals of regional age distribution estimates. The fraction of cases in each age group were fit as a function of seasonal H3 or N2 epitope distance using Beta GLMs (logit link) with 1000 bootstrap resamples.

1698
Figure S17. National excess influenza A(H3N2) mortality decreases with A(H1N1) epidemic size
but not B epidemic size. Excess influenza deaths attributable to A(H3N2) (per 100,000 people) were
estimated from a seasonal regression model fit to weekly pneumonia and influenza-coded deaths . Point
color indicates the dominant influenza A subtype based on CDC influenza season summary reports (red:
A(H3N2), blue: A(H1N1), purple: A(H1N1)pdm09, orange: A(H3N2)/A(H1N1)pdm09 co-dominant), and
vertical bands are 95% confidence intervals of model estimates. National excess mortality estimates were
fit as a function of seasonal A(H1N1) or B epidemic size using Gaussian GLMs (log link) with 1000
bootstrap resamples.

1707
Figure S18. The effect of influenza A(H1N1) epidemic size on A(H3N2) epidemic burden during the
entire study period (1997-2019) (top), pre-2009 seasons (middle), and post-2009 seasons (bottom).
Influenza A(H1N1) epidemic size inversely correlates with A(H3N2) epidemic size, peak incidence,
transmissibility (maximum effective reproduction number, Rt), and epidemic intensity. Point color indicates
the dominant influenza A virus (IAV) subtype based on CDC influenza season summary reports (red:
A(H3N2), blue: A(H1N1), purple: A(H1N1)pdm09, orange: A(H3N2)/A(H1N1)pdm09 co-dominant), and
vertical and horizontal bands are 95% confidence intervals of regional estimates. Seasonal mean
A(H3N2) epidemic metrics were fit as a function of mean A(H1N1) epidemic size using Gaussian GLMs
(epidemic size, peak incidence: inverse link; effective Rt: log link) or Beta GLMs (epidemic intensity: logit
link) with 1000 bootstrap resamples.

Figure S19. Wavelet analysis of influenza A and B epidemic timing. A. A(H3N2) incidence precedes A(H1N1) incidence in most seasons. Although A(H1N1) incidence sometimes leads or is in phase with A(H3N2) incidence (negative or zero phase lag), the direction of seasonal phase lags is not clearly associated with A(H1N1) epidemic size. **B.** A(H3N2) incidence leads B incidence (positive phase lag) during each season, irrespective of B epidemic size. Point color indicates the dominant influenza A subtype based on CDC influenza season summary reports (red: A(H3N2), blue: A(H1N1), purple: A(H1N1)pdm09, orange: A(H3N2)/A(H1N1)pdm09 co-dominant), and vertical bars are 95% confidence intervals of regional estimates. To estimate the relative timing of influenza subtype incidences, phase angle differences were calculated as phase in A(H3N2) minus phase in A(H1N1) (or B), with a positive value indicating that A(H1N1) (or B) incidence lags A(H3N2) incidence. To calculate seasonal phase lags, we averaged pairwise phase angle differences from epidemic week 40 to epidemic week 20. Seasonal phase lags were fit as a function of seasonal A(H1N1) or B epidemic size using LMs with 1000 bootstrap resamples.

1732
Figure S20. Variable importance rankings from LASSO models predicting A(H3N2) epidemic
dynamics. Ranking of variables in predicting seasonal A(H3N2) **A.** epidemic size, **B.** peak incidence, **C.**
transmissibility (effective reproduction number, Rt), **D.** epidemic intensity (inverse Shannon entropy), and
E. subtype dominance. Models were tuned using a repeated leave-one-season-out cross-validated
sample of the data. Variables are ranked by their coefficient estimates, with differences in prediction
accuracy scaled by the total (null model) error. Abbreviations: HI titer = hemagglutination inhibition $\log_2$
titer distance, t - 1 = one-season lag, t - 2 = two-season lag, LBI = local branching index, peak = peak
incidence, distance to vaccine = epitope distance between currently circulating strains and the
recommended vaccine strain, VE = vaccine effectiveness.

Figures S21. Relationships between the predictive accuracy of random forest models and H3 epitope distance. Root mean squared errors between observed and model-predicted values were averaged across regions for each season, and results are facetted according to epidemic metric. Point color corresponds to the degree of H3 epitope distance in viruses circulating in season t relative to those circulating two seasons ago ($t-2$), with bright yellow points indicating seasons with greater antigenic novelty. Spearman correlation coefficients and associated P-values are provided in the top left section of each facet.

Figures S22. Relationships between the predictive accuracy of random forest models and N2 epitope distance Root mean squared errors between observed and model-predicted values were averaged across regions for each season, and results are facetted according to epidemic metric. Point color corresponds to the degree of N2 epitope distance in viruses circulating in season t relative to those circulating in the prior season ($t-1$), with bright yellow points indicating seasons with greater antigenic novelty. Spearman correlation coefficients and associated P-values are provided in the top left section of each facet.