Daily variation in blood glucose levels during continuous enteral feeding in patients on the Intensive Care Unit: a retrospective observational study

F.W. Hiemstra[1,2], D.J. Stenvers[3,4,5], A. Kalsbeek[3,6,7], E. de Jonge[1], D.J. van Westerloo[1], L. Kervezee[2]

Author affiliations

[1] Department of Intensive Care, Leiden University Medical Center, Albinusdreef 2, 2333 ZA, Leiden, The Netherlands

[2] Group of Circadian Medicine, Department of Cell and Chemical Biology, Leiden University Medical Center, Albinusdreef 2, 2333 ZA, Leiden, The Netherlands

[3] Department of Endocrinology and Metabolism, Amsterdam UMC location University of Amsterdam, Meibergdreef 9, 1105 AZ Amsterdam, The Netherland

[4] Department of Endocrinology and Metabolism, Amsterdam UMC location Vrije Universiteit, Meibergdreef 9, 1105 AZ Amsterdam,, The Netherlands

[5] Amsterdam Gastroenterology Endocrinology Metabolism, Amsterdam UMC, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands

[6] Netherlands Institute for Neuroscience (NIN), Royal Dutch Academy of Arts and Sciences (KNAW), Meibergdreef 47, 1105 BA Amsterdam, The Netherlands

[7] Laboratory of Endocrinology, Department of Laboratory Medicine, Amsterdam UMC location University of Amsterdam, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands

Correspondence to:

Laura Kervezee, PhD

Group of Circadian Medicine, Department of Cell and Chemical Biology, Leiden University Medical Center

Address: Albinusdreef 2, 2333 ZA, Leiden, The Netherlands

Email: l.kervezee@lumc.nl

Telephone: +31 71 5269764

Abstract

Background

The circadian timing system coordinates daily cycles in physiological functions, including glucose metabolism and insulin sensitivity. Here, the aim was to characterize the 24-hour variation in glucose levels in critically ill patients during continuous enteral feeding after controlling for potential sources of bias.

Methods

Time-stamped clinical data from adult patients who stayed in the Intensive Care Unit (ICU) for at least 4 days and received enteral feeding were extracted from the Medical Information Mart for Intensive Care (MIMIC)-IV database. Linear mixed-effects and XGBoost modelling were used to determine the effect of time of day on blood glucose values.

Findings

In total, 208,153 glucose measurements collected during enteral feeding were available from 6,944 ICU patients. Using linear mixed-effects modelling, time of day had a significant effect on blood glucose levels ($p<0\cdot001$), with a peak of $9\cdot4$ [$9\cdot4–9\cdot5$; estimated marginal means, 95% CI] mmol/L at 10:00 in the morning and a trough of $8\cdot5$ [$8\cdot4–8\cdot5$] mmol/L at 03:00 at night. A similar impact of time of day on glucose levels was found with the XGBoost regression model.

Interpretation

These results revealed marked 24-hour variation in glucose levels in ICU patients even during continuous enteral feeding. This 24-hour pattern persists after adjustment for potential sources of bias, suggesting it may be the result of endogenous biological rhythmicity.

Funding

This work was supported by a VENI grant (2020-09150161910128) from the Netherlands Organization for Health Research and Development (ZonMw) and an institutional project grant from the LUMC.

Keywords

Intensive Care, Critical Illness, Glucose, Enteral Feeding, Daily Variation, Circadian Rhythm, Electronic Health Records

Introduction

The circadian timing system coordinates daily cycles in physiological functions, including glucose metabolism.[1] Local tissue clocks work in concert with the central circadian clock in the suprachiasmatic nucleus to optimally prepare the body for the predictable daily variations in feeding and fasting and rest and activity. As such, physiological processes affecting plasma glucose levels, such as endogenous glucose production, glucose-induced insulin secretion, and local insulin sensitivity all show marked 24-hour rhythms.[2] In healthy humans, this results in a lower increase in glucose levels in response to food intake in the morning as compared to in the evening and night. In contrast, in people with diabetes mellitus, these 24-hour rhythms give rise to the so called dawn phenomenon; a sharp rise in blood glucose levels in the early morning as a result of reduced insulin sensitivity at the time of highest endogenous glucose production in the early morning.[3]

The relevance of these daily changes in glucose regulation in real-world settings is exemplified by various lines of evidence. For example, a large-scale study in which adults not known to have diabetes were randomized to have their blood glucose examined in either the morning or the afternoon.[4] Despite a shorter fasting duration for afternoon blood draws (breakfast was allowed), fasting glucose levels were lower in the afternoon than in the morning. It was estimated that half of the diabetes cases would go undiagnosed in the afternoon group with the current diagnostic criteria, underscoring that time of blood sampling is an important consideration in the diagnosis of diabetes mellitus specifically, and in clinical practice in general.

In hospitalized patients, various outcomes related to glucose regulation have also been reported to vary by time of day. In general, several retrospective studies have found that low glucose levels occur more frequently at night compared to other times of day in adult inpatients, both in those with diabetes mellitus[5-7] and without diabetes mellitus.[8] In addition, various retrospective studies have shown that blood glucose levels vary by time of day in critically ill patients in the Intensive Care Unit (ICU), with higher levels during the day and lower levels at night.[9-11] Indeed, the importance of clock time for the prediction of glucose values is highlighted by the finding that time of day emerged as the third most important feature to forecast glucose values in ICU patients in a machine learning algorithm based on more than 70 clinical features derived from electronic health records.[12]

The origin of this daily variation in glucose control in hospitalized patients is unknown. On the one hand, some authors argued that this effect can be ascribed to 24-hour variation in underlying physiological processes, such as circadian changes in insulin sensitivity.[9] On the other hand, this time-of-day dependent effect has mostly been attributed to daily variations in health care processes rather than physiology, such as daily changes in feeding support or exogenous insulin administration.[10,11]

Moreover, selection bias can play a role, as glucose levels are examined more frequently in patients with poor glucose control, while sampling in more stable patients may be restricted to a routine check during clinical rounds in the morning.[13] Indeed, controlling for uneven sampling by using time-weighted average glucose levels was found to diminish the 24-hour variation in glucose levels in ICU patients.[13] Clearly, identifying and adjusting for sources of bias is an essential step in the analysis of electronic health record data. This is underlined by the finding that the hour of the day a laboratory test is ordered in the hospital can be more predictive for survival than the actual value of the test result.[14]

Thus, it remains unexplored to what extent blood glucose levels show 24-hour variation in clinical settings after controlling for the various sources of bias introduced by daily rhythms in clinical practice. Therefore, the goal of this study was to characterize 24-hour variation in blood glucose levels in critically ill patients during periods of continuous enteral tube feeding and to evaluate the effect of potential sources of bias that may influence these results. To this end, we made use of the Medical Information Mart for Intensive Care-IV (MIMIC-IV) database, containing detailed time-stamped, annotated clinical data of > 50,000 adult ICU patients, providing a wealth of data to study daily patterns in physiology in a real-world critical care setting. As such, this study aims to shed light on the daily dynamics of blood glucose regulation in critically ill patients, facilitating a deeper understanding of their daily physiological patterns.

Methods

Data source

Data were obtained from the MIMIC-IV (version 2.0) database.[15,16] This is a publicly available, single-center critical care database consisting of detailed, time-stamped clinical data from over 50,000 patients ≥ 18 years and admitted to the ICU of the Beth Israel Deaconess Medical Center (Boston, Massachusetts) between 2008 and 2019. Two authors (FWH and LK) obtained access to the database by completing the Collaborative Institutional Training Initiative program and were responsible for data extraction and analysis (certification numbers: 45894318 and 35567467, respectively). The study was reported in accordance with the Strengthening the Reporting of Observational studies in Epidemiology (STROBE) statement[17] and the REporting of studies Conducted using Observational Routinely collected health Data (RECORD) statement.[18] The code of data extraction and analysis is available on GitHub (https://github.com/fwhiemstra/mimic-glucose-24h-variation-in-icu).

Data extraction and pre-processing

Data from the MIMIC-IV database were extracted using structured query language within PostgreSQL (version 14, PostgreSQL Global Development Group) and pre-processing was performed using R

programming language (version 4.2.1). Patients and ICU stays were selected from the database using the following inclusion criteria: 1) length of ICU stay ≥ 4 days; 2) ICU stay was not a readmission; 3) glucose measurements were taken during the ICU stay; and 4) enteral tube feeding was administered during the ICU stay. All glucose measurements that were taken while patients received enteral feeding during their ICU stay were extracted. Both point-of-care capillary tests as well as lab (serum and whole blood) measurements were included. In the case of simultaneous glucose measurements, whole blood measurements were preferred over serum measurements, that were in their turn preferred over point-of-care capillary measurements, following a similar approach as Robles Arévalo et colleagues.[19] Glucose measurements with a value > 55·6 mmol/L (1,000 mg/dL) were excluded from the dataset, as they exceed the limit of accuracy for the laboratory analyzer that was used in the Beth Israel Deaconess Medical Center.[19] For capillary samples, measurements with a value > 27·8 mmol/L (500 mg/dL) were excluded, according to the limit of accuracy for the point-of-care glucometers (Precision Xceed, Abbott Laboratories, Chicago, Illinois, United States). The charting time was used to time-stamp the glucose measurements as this is considered in the MIMIC-IV database to be the closest proxy to the time the data was actually measured. However, in case the 'store time' (the time the measurement was recorded by the care giver) was recorded before the 'chart time' (the time the actual data entry occurred), the 'store time' was used as time-stamp for the glucose measurement, as described previously.[19] Time of day was rounded down to the nearest integer hour on a scale from 0 to 23. Next, covariates were extracted from the MIMIC-IV database and added to the dataset. For each patient, the following variables were extracted: age, sex, diagnosis of diabetes, and in-hospital mortality. Age was categorized into: < 50 years, 50-70 years, and >70 years. These boundaries were chosen based on the data distribution of the studied ICU population. Sample-level variables that vary with the timing of the glucose sampling within an individual ICU stay, were collected and time-matched to each glucose measurement in the dataset (type of mechanical ventilation, administration of insulin, dextrose or glucocorticoids, Richmond Agitation-Sedation Scale (RASS) score, and rate of enteral feeding). To take into account the variation in the duration of action between different types of insulin, time corrections were applied to the start and end time of the insulin administrations before matching the insulin events to the glucose measurements. Start times of insulin administrations were corrected by adding 0.5 hours and end times were extended, depending on their mode of actions, with 2, 4, 10 and 12 hours for rapid-, short-, intermediate and long-action insulin, respectively.[19] A time correction was also used for the administration of glucocorticoids by adding 24 hour to the end time of administration, to account for the persisting effects of glucocorticoids on glucose metabolism. In the subsequent sections of this article, the glucose measurements taken during administration of insulin and glucocorticoids, include those that fall within the time correction windows. Finally, days since ICU admission and time to the next glucose measurement (using all glucose measurements from the MIMIC-IV dataset, including those not collected during enteral feeding) were derived for each glucose

measurement in the data. An overview of the extracted variables is provided in Supplementary Figure S1.

Statistical analysis

Statistical analysis and visualization was performed using R programming language (version 4.2.1). To visualize variation of glucose levels over time, the glucose levels were normalized per patient and plotted against time. Next, a linear mixed-effects model was used to assess the effect of time of day on glucose levels, adjusted for relevant variables (LME4 package, version 1.1.30).[20] Glucose levels (in mmol/L) were used as dependent variable and patient IDs were added as random effect. As fixed effects, variables that have a known, direct effect on glucose levels were included (Supplementary Figure S1). Four different linear mixed-effects models were fitted, with each subsequent model incorporating additional fixed effects to evaluate the effect of adding fixed effects to the model (Supplementary Table S1). In model 1, only an intercept was used. In model 2, patient-level variables were added (age categories, sex, diabetes mellitus diagnosis (yes/no)) as fixed effects, followed by subsequent addition of sample-level variables in model 3 (administration of dextrose, glucocorticoids or insulin, feeding rate at the time of glucose measurement; yes/no). Finally, in model 4, in addition to the patient-level and sample-level variables, time (in hourly bins) was included as fixed effect. The models were fitted using the entire glucose data set, employing the Restricted Maximum Likelihood approach to estimate the variance of random effects, and subsequently subjected to likelihood ratio tests for model comparison. A significance level of 0.05 was considered as statistically significant and was used to select the final model. Estimated marginal means were calculated with the Maximum Likelihood approach and used to visualize the glucose levels over the 24-hour period. Residual plots were created for the final linear mixed-effects model to assess whether the model assumptions of linearity and constant variance were met.

As an alternative approach, the same dataset was analyzed with the decision tree-based machine learning method Extreme Gradient Boosting (XGBoost) to assess the time of day effect using SHapley Additive exPlanations (SHAP) values.[21,22] A detailed description of this method is available in the Supplementary Methods S1: XGBoost analysis using SHAP values.

Next, sensitivity analyses were performed on various subgroups, based on variables that are considered to have an indirect effect on glucose levels or that may potentially bias the time-of-day dependent effect: type of mechanical ventilation, RASS scores (both as proxy for no oral intake), in-hospital mortality, days in ICU (both as proxy for severity of illness), sample type and sampling frequency (to rule out sampling bias) (Supplementary Figure S1). Furthermore, two subgroups were made to distinguish between patients with insulin resistance (defined as those who received an average daily administration

of more than 10 units of insulin) and without insulin resistance (defined as those who did not receive insulin at any point during their ICU stay). For each subgroup, a linear mixed-effects model (using the same variables and settings as in model 4) was fitted. Estimated marginal means were used to visualize the 24-hour patterns for each of the subgroups.

Results

Baseline characteristics

A total of 6,944 ICU stays (52,467 ICU days) were included in the dataset, with a total of 208,153 valid glucose measurements that were taking during enteral feeding (Figure 1). Baseline characteristics of the included patients and characteristics of the included glucose measurements are summarized in Table 1 and Supplementary Table S2, respectively. A representative example of the timing of enteral feeding, glucose measurements, and the administration of dextrose, glucocorticoids, and insulin during a day in the ICU is shown in Figure 2A and a typical pattern of enteral nutritional support across the ICU stay is shown in Figure 2B. Overall, the included glucose measurements were taken throughout the 24-hour period with an increase of the number of measurements at some time points, corresponding to the timing of routine lab rounds (Figure 2C). Time-of-day distributions of dosing rates of enteral feeding and administrations of insulin, dextrose and glucocorticoids are shown in Figure 2D and Supplementary Figure S2, respectively.

Analysis of 24-hour variation in glucose levels during enteral feeding

Without adjustment for potential confounders, normalized glucose levels showed substantial variation over time, with a peak at 10:00h and trough at 03:00h (Supplementary Figure S3). The linear mixed-effects model fits improved significantly each time new variables were added in the subsequent model (Supplementary Table S3). The inclusion of sample-level variables showed the most substantial improvement (model 2 vs model 3: χ^2 (df = 4) = 4133·5, , p <0·001), followed by the inclusion of time of day (χ^2 (df = 23) = 2181·8, p < 0·001). Therefore, model 4, incorporating patient-level, sample-level, and time of day variables, was selected as the final model. Blood glucose levels were significantly affected by time of day (p<0·001), with a peak of 9·4 [9·4 – 9·5; estimated marginal means, 95% CI] mmol/L at 10:00 in the morning and a trough of 8·5 [8·4 – 8·5] mmol/L at 03:00 at night (Table 2; Figure 3A). The effect of diabetes on glucose levels was found to be most pronounced (2·4 [2·3 – 2·5; coefficient, 95% CI], p<0·001), followed by administration of glucocorticoids (1·1 [1·0 – 1·2], p<0·001) (Table 2). Model diagnostics showed no violation of model assumptions (Supplementary Figure S4).

Following selection of the optimal XGBoost model by hyperparameter tuning (optimal parameters: maximum tree depth: 5; Learning rate: 0·05; Number of estimators: 150; Minimum child weight: 5; Fraction of columns to be subsampled in each tree: 0·75; Subsample ratio of training instances: 0·75), a 24-hour pattern was observed in the SHAP values for time of day, with an increase around 10:00h and a decrease between 01:00h and 03:00h (Figure 3B), resembling the 24-hour pattern observed with the linear-mixed effects model (Figure 3A). Time of day was the third most important feature, preceded by diabetes and enteral nutrition rate > 50 mL/h and followed by administration of glucocorticoids (Supplementary Figure S5A-B).

In all subgroups of type of mechanical ventilation, in-hospital mortality, RASS-scores, sample type, days in ICU and sampling frequencies, similar 24-hour patterns were observed, with expected differences in 24-hour baseline levels (Figure 4). Remarkably, the peak-to-trough difference in glucose levels was more than two-fold larger in the subgroup with insulin resistance (10·0 [9·9 – 10·2] versus 11·2 [11·1 – 11·3] mmol/L at 03:00 and 10:00, respectively) compared to the subgroup without insulin resistance (7·0 [6·9 – 7·1] mmol/L at 03:00 and 7·5 [7·4 – 7·6] at 10:00) (Figure 5).

Discussion

Our results indicate that glucose levels in critically ill patients show marked 24-hour variation during continuous enteral feeding. This daily pattern persists when controlling for potential sources of bias and confounding factors, suggesting that the observed 24-hour variation is due to biological processes rather than health care processes, patient characteristics, or treatment effects. These findings were confirmed with an alternative machine learning approach, thereby showing the consistency of the results obtained with the linear mixed-effects model. Remarkably, the effect of time of day on glucose levels, even during continuous nutritional support, is equal or even larger than covariates that are commonly considered in analyses, such as sex and age. This highlights the importance of incorporating time of day in analyses of biological time series data.

We found a decrease of glucose levels during nighttime, followed by a substantial rise of glucose levels in the early morning. Previous studies also observed a nocturnal decrease in glucose levels in critically ill patients, although the timing of the through varied among studies.[9-11] However, those studies included glucose measurements regardless of nutritional status. The daily variation in glucose levels observed in those studies may therefore be driven by an interfering daily pattern in nutrition, particularly when considering oral intake. Since most oral food intake presumably occurs during the daytime and not during the nighttime, any observed 24-hour pattern in glucose levels may be attributable to patterns in food intake rather than endogenous variation in glucose control. A strength of our study is that we only included glucose measurements that were taken while patients received enteral feeding in order to

investigate 24-hour variation in glucose levels under constant nutritional intake. In addition, we verified that the timing of glucose measurements, enteral nutrition, and administration of insulin, dextrose, and glucocorticoids were equally distributed over the 24-hour period.

The subgroup analyses were performed to assess the effect of potential sources of bias that were not or insufficiently available in the MIMIC-IV data. Subgroups were created based on available variables that might indirectly affect glucose levels through other confounders. As oral food intake may not be well documented in electronic health records, we sought to rule out that the observed daily variation in glucose levels is driven by patients that have oral intake – most likely to occur during daytime - in addition to enteral feeding. The sensitivity analyses showing that the 24-hour pattern persists in patients on invasive mechanical ventilation or sedated patients (with RASS score $\leq$ -2), who in general do not have any oral food intake, rules out this possibility. Furthermore, the glucose sampling method used (lab or point-of-care tests) can be influenced by health care processes and acuteness of situation. Lab tests are more likely to be a routine test, while point-of-care tests can be more likely to performed when glucose needs to be closely monitored. Also, severity of illness could affect the glucose levels. Indicators of severity of illness, such as Sequential Organ Failure Assessment (SOFA) scores, were not consistently present in the data. Therefore, in-hospital mortality and days in the ICU were regarded to be associated with illness severity. To assess the effect of sampling bias, time to the next glucose measurements was used a proxy of sampling frequency. For all these potential sources of bias, the expected differences in mean 24-hour levels were observed, but the 24-hour pattern with the morning rise and its timing remained consistent among the subgroups.

Linear mixed-effects models prove to be valuable in the analyses conducted in our study as this approach allows for adjustment of several factors with a known influence on the dependent variable (fixed effects) and accounts for individual variability (random effects). As an addition to the main analyses, the same dataset was analyzed with an XGBoost regression model. XGBoost is a widely adopted machine learning algorithm that uses decision trees within a gradient boosting framework. Although XGBoost is commonly employed for prediction tasks, it is used in our study to assess the effect of a variable of interest (time of day) on an observed parameter (glucose levels), as was also done by Abbaspour and colleagues.[23,24] While linear mixed-effects models assume a linear relationship between fixed effects and outcome variable, XGBoost can capture more intricate patterns and non-linear associations that might exist in the data. Similar results were found when analyzing the data with an XGBoost model as with the linear mixed-effects model, which shows reliability and consistency of our findings. Moreover, this indicates that the use of a machine learning model may be a powerful alternative to traditional linear mixed-effects models in the analysis of time series data.

The 24-hour pattern in glucose levels with low levels at night and an increase in the early morning that we observe in our study is markedly different from what could be expected based on studies on circadian rhythms in glucose control in healthy individuals. In healthy people, glucose excursion are higher when food is consumed during the biological night compared to the day.[25] A study among healthy male participants who received continuous enteral feeding in controlled laboratory conditions revealed that glucose levels are highest during the second half of the night and lowest during the day, i.e., a 24-hour rhythm that is advanced by several hours compared to the pattern observed in our study.[26] However, the sharp rise in blood glucose levels observed in our analysis during the early morning is reminiscent of the dawn phenomenon observed in people with diabetes mellitus. This phenomenon occurs as a result of reduced insulin sensitivity at the time of highest endogenous glucose production.[3] The morning increase that we observe in critically ill patients may reflect increased insulin resistance that is frequently observed in the ICU population.[27] This notion gains further support from the observation that patients without insulin resistance exhibit a less pronounced morning peak, in contrast to patients with insulin resistance. Albeit correlational, our findings add real-world support to the body of literature from pre-clinical studies and studies in healthy human participants that show that glucose metabolism is anything but constant across the 24-hour period.[2]

Although the variation of about 1 mmol/L across 24 hours may not affect glucose monitoring or insulin administration protocols in the ICU, our study does provide insight into the factors that influence glucose levels, and can thereby contribute to the understanding of daily fluctuations of glucose levels in patients during clinical practice at the ICU. The substantial effect of time of day on blood glucose levels observed in our study emphasizes the importance of considering the 24-hour variation in scientific research on glucose regulation in critically ill patients. Relying solely on single measurements or averages derived from unequally distributed measurements may inaccurately reflect the average daily glucose control.[13] Likewise, clinical trials investigating glucose control in critically ill patients may need to consider this daily variation in their sampling protocol and ensuing study endpoints.

One of the strengths of our study is the large number of glucose measurements that was included in the analysis. The availability of a large, rich, clinical dataset enabled the adjustment of multiple confounding factors. Additionally, a heterogeneous group of patients admitted to the ICU was included, thereby making the results more applicable to the mixed ICU population. Nonetheless, several limitations should be considered when interpreting our findings. The retrospective design introduces inherent limitations, as we were constrained to data available in the electronic health records.[28] As a result, it was not possible to control the timing of blood glucose measurements or the ensuing medical decisions. Another limitation is introduced by the variability in the delay and persisting effects after the start and end of administration of the assessed medication on glucose levels. To account for this, we incorporated fixed correction times to the start and end times of insulin and glucocorticoid

administration, but any interindividual variability in the duration or magnitude of their effect was not taken into account. Also, the use of single-center data in this study may introduce bias due to center-specific factors, patient population, or variations in patient management practices. Moreover, it remains to be investigated to what extent patient-level factors that influence 24-hour patterns in glucose levels impact our findings, such as sleep[29-31], neurotrauma[32], or circadian clock function.[33]

In general, our study provides insight into the glucose control during continuous enteral feeding in critically ill patients. An interesting question is to what extent the administration of continuous feeding gives rise to the 24-hour pattern in blood glucose values and is responsible for the high prevalence of insulin resistance that is observed in ICU patients in general. Although this type of nutritional support is common practice in ICUs worldwide, it is known from pre-clinical work and studies in healthy human participants that food intake at the wrong time of day (i.e. during the biological night) leads to unfavorable glucose responses.[2] Indeed, it has been questioned whether continuous feeding is the most optimal feeding schedule from various perspectives (incl. glycemic control, circadian rhythmicity, muscle metabolism, and gastrointestinal function).[34,35] Future studies that compare the effect of continuous feeding to other schedules on various clinical outcomes could additionally investigate the impact on daily glucose variability.

To conclude, our findings demonstrate the presence of a substantial 24-hour pattern in blood glucose levels during continuous enteral feeding in critically ill patients that persists after adjustment for relevant sources of bias. This suggests that the observed 24-hour variation is due to the endogenous biological variation rather than as a result of bias introduced by health care processes, patient characteristics, or treatment effects. In addition, our study provides a framework for addressing these sources of bias when analyzing the effect of time of day on physiological variables available in electronic health records. Our findings contribute to a deeper understanding of daily fluctuations in glucose levels observed at the bedside by nurses and clinicians working at the ICU. Future research should focus on multicenter cohorts and incorporate individual circadian phase and brain states in the analysis to further enhance our understanding of these observations.

Contributors

The study was conceptualized by LK. Data extraction, preprocessing and analysis was performed by FWH and LK. DJvW, EdJ, DJS and AK participated in discussing and interpretation of the results. FWH wrote the first version of the manuscript. All authors have edited and reviewed the manuscript and approved the final version.

Declaration of interest

All authors declare no competing interests.

Acknowledgements

The authors thank Marc D. Ruben for valuable discussions. This work was supported by a VENI grant (2020-09150161910128 to L.K.) from the Netherlands Organization for Health Research and Development (ZonMw) and an institutional project grant from the Leiden University Medical Center (to L.K. and D.J.v.W.). We would like to thank the researchers of the MIT Computational Physiology Laboratory and collaborating researchers for developing, maintaining and publicly sharing of the MIMIC-IV clinical database.

Data Sharing Statement

The MIMIC-IV dataset is publicly available via: https://mimic.mit.edu/. The code used for the curation of the dataset and analyses are available at https://github.com/fwhiemstra/mimic-glucose-24h-variation-in-icu.

References

Kalsbeek A, la Fleur S, Fliers E. Circadian control of glucose metabolism. *Mol Metab*. 2014;3(4):372-83.

Stenvers DJ, Scheer F, Schrauwen P, la Fleur SE, Kalsbeek A. Circadian clocks and insulin resistance. *Nat Rev Endocrinol*. 2019;15(2):75-89.

Peng F, Li X, Xiao F, Zhao R, Sun Z. Circadian clock, diurnal glucose metabolic rhythm, and dawn phenomenon. *Trends in Neurosciences*. 2022;45(6):471-82.

Troisi RJ, Cowie CC, Harris MI. Diurnal variation in fasting plasma glucose: implications for diagnosis of diabetes in patients examined in the afternoon. *JAMA*. 2000;284(24):3157-9.

Rajendran R, Kerry C, Rayman G, MaGIC study group. Temporal patterns of hypoglycaemia and burden of sulfonylurea-related hypoglycaemia in UK hospitals: a retrospective multicentre audit of hospitalised patients with diabetes. *BMJ Open*. 2014;4(7):e005165.

Kerry C, Mitchell S, Sharma S, Scott A, Rayman G. Diurnal temporal patterns of hypoglycaemia in hospitalized people with diabetes may reveal potentially correctable factors. *Diabetic Medicine*. 2013;30(12):1403-6.

Bailon RM, Cook CB, Hovan MJ, Hull BP, Seifert KM, Miller-Cage V, et al. Temporal and Geographic Patterns of Hypoglycemia among Hospitalized Patients with Diabetes Mellitus. *Journal of Diabetes Science and Technology*. 2009;3(2):261-8.

Ulmer BJ, Kara A, Mariash CN. Temporal Occurrences and Recurrence Patterns of Hypoglycemia During Hospitalization. *Endocrine Practice*. 2015;21(5):501-7.

Pidcoke HF, Salinas J, Wanek SM, Concannon M, Loo F, Wirfel KL, et al. Patterns of exogenous insulin requirement reflect insulin sensitivity changes in trauma. *Am J Surg*. 2007;194(6):798-803; discussion

Smith SM, Oveson KE, Strauss W, Raven K, Lefevre MC, Ahmann AJ, et al. Ultradian variation of blood glucose in intensive care unit patients receiving insulin infusions. *Diabetes Care*. 2007;30(10):2503-5.

Egi M, Bellomo R, Stachowski E, French CJ, Hart G, Stow P. Circadian rhythm of blood glucose values in critically ill patients. *Crit Care Med*. 2007;35(2):416-21.

Fitzgerald O, Perez-Concha O, Gallego B, Saxena MK, Rudd L, Metke-Jimenez A, et al. Incorporating real-world evidence into the development of patient blood glucose prediction algorithms for the ICU. *J Am Med Inform Assn*. 2021;28(8):1642-50.

Vogelzang M, van der Horst IC, Zijlstra F, Nijsten MW. Circadian variation of glucose levels: biology or timing of measurements? *Crit Care Med*. 2007;35(7):1800-1; author reply 1-2.

Agniel D, Kohane IS, Weber GM. Biases in electronic health record data due to processes within the healthcare system: retrospective observational study. *BMJ*. 2018;361:k1479.

Johnson A, Bulgarelli L, Pollard T, Horng S, Celi L, Mark R. MIMIC-IV (version 2.0). PhysioNet. 2022.

Goldberger AL, Amaral LA, Glass L, Hausdorff JM, Ivanov PC, Mark RG, et al. PhysioBank, PhysioToolkit, and PhysioNet: components of a new research resource for complex physiologic signals. *Circulation*. 2000;101(23):E215-20.

von Elm E, Altman DG, Egger M, Pocock SJ, Gotzsche PC, Vandenbroucke JP, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet*. 2007;370(9596):1453-7.

Benchimol EI, Smeeth L, Guttmann A, Harron K, Moher D, Petersen I, et al. The REporting of studies Conducted using Observational Routinely-collected health Data (RECORD) statement. *PLoS Med*. 2015;12(10):e1001885.

Robles Arevalo A, Maley JH, Baker L, da Silva Vieira SM, da Costa Sousa JM, Finkelstein S, et al. Data-driven curation process for describing the blood glucose management in the intensive care unit. *Sci Data*. 2021;8(1):80.

Bates D, Machler M, Bolker BM, Walker SC. Fitting Linear Mixed-Effects Models Using lme4. *J Stat Softw*. 2015;67(1):1-48.

Lundberg SM, Lee S-I. A unified approach to interpreting model predictions. Proceedings of the 31st International Conference on Neural Information Processing Systems; Long Beach, California, USA: Curran Associates Inc.; 2017. p. 4768–77.

Chen T, Guestrin C. XGBoost: A Scalable Tree Boosting System. Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining; San Francisco, California, USA: Association for Computing Machinery; 2016. p. 785–94.

Abbaspour S, Aguirre AD, Westover MB, Klerman EB. The Effects of the Timing of Diuretic Administration and Other Covariates on Urine Volume in Hospitalized Patients. *medRxiv*. 2022:2022.12.27.22283947.

Abbaspour S, Robbins GK, Blumenthal KG, Hashimoto D, Hopcia K, Mukerji SS, et al. Identifying Modifiable Predictors of COVID-19 Vaccine Side Effects: A Machine Learning Approach. *Vaccines*. 2022;10(10):1747.

Poggiogalle E, Jamshed H, Peterson CM. Circadian regulation of glucose, lipid, and energy metabolism in humans. *Metabolism*. 2018;84:11-27.

Simon C, Gronfier C, Schlienger JL, Brandenberger G. Circadian and ultradian variations of leptin in normal man under continuous enteral nutrition: relationship to sleep and body temperature. *J Clin Endocrinol Metab*. 1998;83(6):1893-9.

McCowen KC, Malhotra A, Bistrian BR. Stress-induced hyperglycemia. *Crit Care Clin*. 2001;17(1):107-24.

Sauer CM, Chen L-C, Hyland SL, Girbes A, Elbers P, Celi LA. Leveraging electronic health records for data science: common pitfalls and how to avoid them. *The Lancet Digital Health*. 2022;4(12):e893-e8.

DePietro RH, Knutson KL, Spampinato L, Anderson SL, Meltzer DO, Van Cauter E, et al. Association Between Inpatient Sleep Loss and Hyperglycemia of Hospitalization. *Diabetes Care*. 2017;40(2):188-93.

Briancon-Marjollet A, Weiszenstein M, Henri M, Thomas A, Godin-Ribuot D, Polak J. The impact of sleep disorders on glucose metabolism: endocrine and molecular mechanisms. *Diabetol Metab Syndr*. 2015;7:25.

St-Onge MP, Cherta-Murillo A, Darimont C, Mantantzis K, Martin FP, Owen L. The interrelationship between sleep, diet, and glucose metabolism. *Sleep Med Rev*. 2023;69:101788.

Jalloh I, Carpenter KL, Helmy A, Carpenter TA, Menon DK, Hutchinson PJ. Glucose metabolism following human traumatic brain injury: methods of assessment and pathophysiological findings. *Metab Brain Dis*. 2015;30(3):615-32.

Mason IC, Qian J, Adler GK, Scheer F. Impact of circadian disruption on glucose metabolism: implications for type 2 diabetes. *Diabetologia*. 2020;63(3):462-72.

Arabi YM, Casaer MP, Chapman M, Heyland DK, Ichai C, Marik PE, et al. The intensive care medicine research agenda in nutrition and metabolism. *Intensive Care Med*. 2017;43(9):1239-56.

Kouw IWK, Heilbronn LK, van Zanten ARH. Intermittent feeding and circadian rhythm in critical illness. *Curr Opin Crit Care*. 2022;28(4):381-8.

Tables

Table 1: Baseline characteristics of included patients.

Patient characteristics	Patients (n=6,944)
Sex, n (%)	
Male	3,954 (57%)
Female	2,990 (43%)
Age (years), median [IQR]	66 [54-77]
Length of stay (days), median [IQR]	10 [7-16]
Diabetes diagnosis, n (%)	2,132 (31%)
In-hospital mortality, n (%)	1,970 (28%)
Mechanical ventilation during ICU stay, n (%)	
Invasive ventilation	6,232 (90%)
Non-invasive ventilation only	605 (8.7%)
None / No ventilation data available	107 (1.5%)

Table 2: Summary of final linear mixed-effects model.

Variable	Estimates	95% CI	p-value
(Intercept)	7·2	7·0 – 7·3	**<0·001**
Time of day 03:00 [Ref = 00:00]*	-0·4	-0·4 – -0·3	**<0·001**
Time of day 10:00 [Ref = 00:00]*	0·6	0·5 – 0·7	**<0·001**
Diabetes diagnosis [ref = No]	2·4	2·3 – 2·5	**<0·001**
Aged 51-70 y [ref = 18-50 y]	0·5	0·4 – 0·7	**<0·001**
Aged >70 y [ref = 18-50 y]	0·7	0·6 – 0·9	**<0·001**
Male sex [ref = Female]	-0·2	-0·3 – -0·1	**<0·001**
Insulin administration [ref = No]	-0·6	-0·60 – -0·5	**<0·001**
Dextrose administration [ref = No]	0·2	0·2 – 0·3	**<0·001**
Glucocorticoids administration [ref = No]	1·1	1·0 – 1·2	**<0·001**
Feeding rate > 50 mL [Ref = ≤ 50 mL]	0·6	0·6 – 0·7	**<0·001**

*: Time of day was included as a categorical variable with 24 levels of 1-h time bins; only 03:00 (trough) and 10:00 (peak) are shown here due to space constraints. Estimates at all time bins are shown in Figure 3A.

** The start and end time of administration of insulin and glucocorticoids (end times only) were corrected to account for the delaying and/or persisting effect of insulin and glucocorticoids on glucose levels (see Methods for details).

Figures

Figure 1: Inclusion flowchart of ICU stays and glucose measurements.

Figure 2: Distributions of glucose measurements and enteral feeding rates over the 24-hour period and across the ICU stay. (A) Representative example of the timing of enteral feeding, glucose measurements and administrations of dextrose, glucocorticoids and insulin. Black symbols represent glucose measurements taken while the patient received enteral nutritional support (included in the dataset), grey symbols represent glucose measurements taken while the patient did not receive enteral nutritional support (not included in the dataset). (B) Representative examples of enteral nutritional support across the ICU stay, taken from a patient included in the dataset. (C) Distribution of included glucose measurements, and (D) dosing rate of enteral feeding (at the time of glucose measurements) by time of day.

Figure 3: 24-h variation in glucose levels during the entire ICU stay during enteral feeding. (A) Model-predicted glucose levels by time of day (in hourly bins). Data presented as estimated marginal means ± 95% CI derived from the final linear mixed effects model, adjusted for patient-level and sample-level covariates. (B) SHAP values for time of day (as a continuous variable) from the XGBoost model. Each black dot represents an individual glucose measurement. The SHAP value represents shows how much the time of day feature affected the glucose level. A positive SHAP value reflects a positive impact on the predicted glucose level, while negative values reflect negative impacts. Its absolute value represents the magnitude of this effect. A sample of 5000 randomly selected features of the glucose measurements was used in this SHAP analysis.

Figure 4: Subgroup analysis of 24-hour variation in glucose levels. (A) Ventilation mode. (B) Survivor status. (C) Depth of sedation. (D) Days in ICU. (E) Sample type. (F) Time to next glucose sample. For each subgroup, a linear mixed-effects model was fitted. Data presented as estimated marginal means ± 95% CI, adjusted for covariates. Number of patients and measurements per group are shown in Supplementary Table S4.

Figure 5: 24-h variation in glucose levels in patient with and without insulin resistance. Insulin resistance is defined by the administration of insulin during the ICU stay (see Methods for details). Data presented as estimated marginal means ± 95% CI, adjusted for covariates.