

Sensitivity of the African Neuropsychology Battery Memory Subtests and Learning Slopes in Discriminating APOE 4 and Amyloid Pathology in Adult Individuals in the Democratic Republic of Congo

Jean Ikanga, Ph.D.^{1,2}, Sarah D. Patrick, Ph.D.³, Megan Schwinne, M.PH.⁴ Saranya Sundaram Patel, Ph.D.¹, Emmanuel Epenge, M.D.⁵, Guy Gikelekele, M.D.², Nathan Tshengele, M.D.², Immaculee Kavugho, M.A.⁶, Samuel Mampunza, M.D., Ph.D.², Kevin E. Yarasheski, Ph.D.⁷, Charlotte E. Teunissen, Ph.D.⁸, Anthony Stringer, Ph.D.¹, Allan Levey, M.D., Ph.D.⁹, Julio C. Rojas, M.D., Ph.D.¹⁰ Brandon Chan, BA¹⁰ Argentina Lario Lago PhD¹¹ Joel H. Kramer PsyD¹¹ Adam L. Boxer MD PhD¹⁰ Andreas Jeromin, PhD¹¹, Alvaro Alonso, M.D., Ph.D.¹², & Robert J. Spencer, PhD³

¹ Emory University School of Medicine, Department of Rehabilitation Medicine, Atlanta, GA 30322, USA

² University of Kinshasa and Catholic University of Congo, School of Medicine, Kinshasa, Department of Psychiatry, B.P. 7463 Kinshasa I, Democratic Republic of Congo

³ Veteran Affairs Ann Arbor Healthcare System, Ann Arbor, Michigan, USA

⁴ Emory University, School of Medicine, Department of Biomedical Informatics, Atlanta, GA 30322, USA

⁵ University of Kinshasa, Department of neurology, Kinshasa, B.P. 7463 Kinshasa I, Democratic Republic of Congo

⁶ Memory Clinic of Kinshasa, Kinshasa, B.P. 7463 Kinshasa I, Democratic Republic of Congo

⁷ C₂N Diagnostics, C₂N Diagnostics, Saint Louis, MO 63110, USA

⁸ Amsterdam University Medical Centers, Neurochemistry Laboratory, Department of Clinical Chemistry, Amsterdam Neuroscience, Neurodegeneration, Amsterdam University Medical Centers, Vrije Universiteit, 1081 HV Amsterdam, the Netherlands

⁹ Emory University, School of Medicine, Department of Neurology, Atlanta, Georgia 30322, USA

¹⁰ University of San Francisco, Memory and Aging Center, Weill Institute for Neurosciences, Department of Neurology, University of California San Francisco

¹¹ ALZpath, Inc.

¹² Emory University, Rollins School of Public Health, Department of Epidemiology, Georgia, Atlanta, GA, 30307, USA

*Corresponding author. Please send general questions about the study to Dr. Jean Ikanga.

Jean N. Ikanga, Ph.D.
Department of Rehabilitation Medicine
Emory University
1441 Clifton Rd NE
Atlanta, GA 30322, USA
Email: jikanga@emory.edu

Manuscript word count:

ABSTRACT

Background: The current study examined the sensitivity of two memory subtests and their corresponding learning slope metrics derived from the African Neuropsychology Battery (ANB) to detect amyloid pathology and APOE ϵ 4 status in adults from Kinshasa, the Democratic Republic of the Congo.

Methods: 85 participants were classified for the presence of β -amyloid pathology and based on allelic presence of APOE ϵ 4. All participants were screened using CSID and AQ, underwent verbal and visuospatial memory testing from ANB, and provided blood samples for plasma A β ₄₂, A β ₄₀, and APOE proteotype. Pearson correlation, linear and logistic regression were conducted to compare amyloid pathology and APOE ϵ 4 status with derived learning scores, including initial learning, raw learning score, learning over trials, and learning ratio.

Results: Our sample included 35 amyloid positive and 44 amyloid negative individuals as well as 42 without and 39 with APOE ϵ 4. All ROC AUC ranges for the prediction of amyloid pathology based on learning scores were low, ranging between 0.56-0.70 (95% CI ranging from 0.44-0.82). The sensitivity of all the scores ranged between 54.3-88.6, with some learning metrics demonstrating good sensitivity. Regarding APOE ϵ 4 prediction, all AUC values ranged between 0.60-0.69, with all sensitivity measures ranging between 53.8-89.7. There were minimal differences in the AUC values across learning slope metrics, largely due to the lack of ceiling effects in this sample.

Discussion: This study demonstrates that some ANB memory subtests and learning slope metrics can discriminate those that are normal from those with amyloid pathology and those with and without APOE ϵ 4, consistent with findings reported in Western populations.

INTRODUCTION

Problems learning and remembering are common across neurodegenerative conditions and can contribute to declines in everyday functioning¹. In clinical evaluations, learning and memory are frequently evaluated using learning tasks, in which examinees are presented with verbal or visual stimuli over several trials and asked to learn, and then subsequently recall, the information after a delay. The primary test scores include the total learning score (i.e., the total amount of information learned over the repeated trials) and the short- and long-delay memory scores (i.e., the amount of information an examinee recalled after delays). These scores are associated with amyloid and tau biomarkers^{2,3} and brain volume,⁴ aiding in clinically relevant outcomes such as the early detection of Alzheimer’s disease (AD).^{5,6}

Clinicians often use supplementary scores to parse apart the cognitive process test takers use when completing these memory tasks. One of these scores is learning slope, which refers to the rate at which the examinee improves throughout the learning trials. Healthy adults are expected to benefit from repeated exposure to information; however, patients with neurological conditions, such as AD, Korsakoff’s syndrome, or amnesic mild cognitive impairment, commonly demonstrate a lack of improvement from practice or repetition,^{7–10} or more formally known as a “shallow” or “flat” learning slope.

Learning slope calculations can be applied to practically any measure involving repeated administration of to-be-learned stimuli and has been used with a variety of visual and verbal list learning tasks, including the California Verbal Learning Test (CVLT),¹¹ Hopkins Verbal Learning Test-Revised (HVLTR),¹² the Rey Auditory Verbal Learning Test (RAVLT),¹³ Repeatable Battery for the Assessment of Neuropsychological Status (RBANS),^{14,15} and Brief Visual Memory Test-Revised (BVMTR).¹⁶ The most used learning slope metric, the raw

learning slope (RLS), is usually calculated by subtracting the first learning trial score from the final learning trial score. Unfortunately, the interpretability of RLS is constrained by ceiling effects for examinees who perform well on the first trial because there is limited opportunity to learn new information on subsequent trials. For example, an examinee who learns 2 words on a 16-word list can achieve an RLS score of 14, whereas an examinee who learned 10 words on the first trial can obtain a maximum RLS of only 6. Spencer and colleagues addressed the drawbacks to RLS with their proposal of a new method, the learning ratio (LR), which directly addresses the initial learning score.¹⁷ This score mathematically accounts for initial learning trial performance by calculating information learned as a percentage of the amount of information left to be learned.

$$LR = \frac{\text{Final Trial Performance} - \text{First Trial Performance}}{\text{Maximum Score per Trial} - \text{First Trial Performance}}$$

Although the RLS score or some close derivation thereof is usually depicted in test manuals, the LR score has a growing number of normative datasets for several learning and memory tests, including the RAVLT,¹⁸ HVLt-R and BVMT-R,¹⁹ RBANS,²⁰ and African Neuropsychology Battery (ANB),²¹ providing clinicians with anchors for evaluating performance. By eliminating confounds from ceiling effects, the LR may result in stronger associations and predictive ability for clinically relevant outcomes than the RLS, which typically produces weak to null findings when evaluated in the context of demographic characteristics and other performance characteristics.¹⁰ When both scores are directly compared, LR performs equal to or better than RLS for correlations with memory tests,^{3,17,18,22,23} diagnostic discrimination,^{3,18,23} as well as other known risk factors for AD, including hippocampal volume, AD-specific biomarkers, and APOEε4 status.²²⁻²⁴ Thus, the LR score is a relatively new but

promising metric that warrants additional examination, particularly in the context of detecting AD using non-invasive assessment methods.

AD is associated with brain-based changes including the presence of abnormally high amyloid beta plaque deposition and pathologic tau aggregations.²⁵ These physiological changes can be detected years before the development of clinical AD symptoms.²⁶ Tau burden²⁷ and amyloid deposition²⁸ consistently correlate with performance on memory measures. Amyloid beta and tau burden are typically measured using positron emission tomography (PET) scans, cerebrospinal fluid (CSF), or blood plasma. Unfortunately, both PET and CSF are costly, invasive, and inaccessible to many patients. Such practical concerns are most important in communities with limited access to medical and financial resources, such as in the Democratic Republic of Congo (DRC). In contrast, blood plasma is a faster, cheaper, and more accessible alternative.^{29,30} Plasma P-tau181 values are correlated with increased tau PET measurements and have shown promise in discriminating between AD and non-AD degenerative disease, reflecting severity of AD across the AD continuum, and progression to AD.²⁹

Administering paper-and-pencil cognitive tests is cheaper, more portable/accessible, and less invasive than amyloid or tau PET imaging, cerebrospinal fluid (CSF) sampling and analysis, and more recently blood sampling for plasma biomarker analysis. Research on memory test process scores and AD biomarkers indicate that worse LR scores are associated with greater cerebral beta-amyloid deposition (PET) and APOE ϵ 4 positivity,²² though these scores have not yet been evaluated in the context of blood plasma biomarker status for the ANB. Given the practical advantages of the LR score, any indications of subtle cognitive inefficiencies that are associated with AD-specific plasma biomarkers and downstream cognitive changes suggest that this score may be a worthwhile metric for identifying early indications of AD pathology.

This study examined two memory subtests and their corresponding learning slope metrics from the ANB to aid in detecting amyloid pathology and APOE ϵ 4 allele presence.²⁵ We hypothesized that ANB memory and learning slope scores would have adequate to good test receiver operator characteristics (ROC; e.g., area under the curve, specificity, sensitivity) to classify patients into plasma amyloid positive or negative. Based on previous studies,²² we predicted that the LR score would bear a stronger relationship to amyloid pathology and APOE ϵ 4 allele than would the other learning slopes.

METHODS

Study population

Participants were selected from our previous study,²⁹ a matched case-control study to identify risk factors of AD in Sub-Saharan Africa (SSA). Participants were at least 65 years or older, had a family member or close friend to serve as an informant, and fluent in French or Lingala. Participants were excluded if they had history of schizophrenia, neurological, or other or medical conditions potentially affecting the CNS. In the absence of established diagnostic criteria for AD in SSA, we used two screening measures with high sensitivity and specificity for identifying individuals with dementia in Western cohorts, the Alzheimer's Questionnaire (AQ)³¹ and the Community Screening Instrument for Dementia (CSID),³² which evaluate clinical symptoms of dementia. The AQ distinguishes between those with AD from healthy controls.³¹ The CSID Questionnaire has been extensively used in many international and SSA dementia studies.³²

Based on cognitive and functional deficits for Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition, Text Revision (DSM-5-TR) diagnostic criteria,³³ we used Brazzaville cut-offs of CSID, the closest city from Kinshasa, to classify participants.³⁴ Similar to our prior study,³⁵ participants were classified using CSID and AQ scores (see Figure 1), which yielded 4 groups: major neurocognitive disorder/dementia, mild neurocognitive disorder (MND), subjective cognitive impairment, and healthy control (HC), i.e., normal cognition.

A panel consisting of a neurologist, psychiatrist and neuropsychologist reviewed screening tests, clinical interview, and neurological examination of subjects. 56 individuals were confirmed with a diagnosis of dementia and 58 were considered HC. Participants were matched based on

age, education, and gender. Plasma biomarkers were obtained for 85 subjects (75%), resulting in a final sample of 44 dementia and 41 HC. The remaining 29 subjects refused to provide blood samples. Written informed consent was obtained prior to participants' undergoing any study procedures. Participants were financially compensated for their time. The informed consent document and all research procedures were approved by the Ethics Committee/Institutional Review Boards of the University of Kinshasa.




Figure 1. Flow diagram of participant classifications using the CSI-D and the AQ in the current study.

Abbreviations: CSID (Community Screening Instruments for Dementia); AQ (Alzheimer's Questionnaire); MajNCD (major neurocognitive disorder); HC (healthy controls); MND (mild neurocognitive disorder).

Procedure

Participants underwent a comprehensive clinical evaluation, including cognitive testing, self-report questionnaires, and standard psychiatric and neurological evaluations to be diagnosed with dementia or to be considered as HC by an expert panel [neurologist (EE), psychiatrist (GG), and neuropsychologist (JI)]. Subjects were interviewed to obtain demographic, socioeconomic,

and medical history and subsequently administered cognitive testing with ANB subtests.

Afterwards, blood samples were obtained at Medical Center of Kinshasa (CMK) by a phlebotomist. Sample collection protocol and quantification of fluid biomarkers are presented below.

Measures

Plasma biomarkers

Blood samples were drawn in the CMK blood laboratory by venipuncture into dipotassium ethylene diamine tetra acetic acid (K₂ EDTA) tubes. Samples were centrifuged within 15 minutes, and 5 mL of plasma was aliquoted into 0.5 mL polypropylene tubes and stored initially at -20° C for less than a week and then moved to a -80 °C freezer for longer term storage at a CMK laboratory.³⁶ These aliquots were shipped frozen on dry ice to Emory University and APOE isoform-specific peptides were analyzed at C₂N Diagnostics (St. Louis, MO) as described.³⁷ We used LC-MS/MS to detect and identify the APOE isoform-specific peptides (ϵ 2, ϵ 3, ϵ 4), with the purpose being to classify participants into APOE ϵ carriers and non-carriers

Plasma biomarker concentrations were measured using commercially available Neurology 4-PLEX E (A β ₄₀, A β ₄₂), P-tau181 (P-tau181 v2; 1) Quanterix kits for the Simoa HD-X platform (Billerica, MA) at the University of San Francisco. P-tau217 was measured using the proprietary ALZpath P-tau-217 CARE Advantage kit (lot #MAB231122, ALZpath, Inc.) for the Simoa HD-X platform. The instrument operator was blinded to clinical variables. All analytes were measured in duplicate. For A β ₄₀ and A β ₄₂, all samples were measured above the lower limit of quantification (LLOQ) of 1.02 pg/mL and 0.378 pg/mL. The average coefficient of variation (CV) for A β ₄₀ and A β ₄₂ were 6.0% and 6.5%. For P-tau181, all samples were measured above

the kit lower limit of quantification (LLOQ) of 0.085 pg/mL, with an average CV of 11.6%. For P-tau217 the LLOQ was 0.024 pg/mL and the average CV was 19.8%.

In the absence of a gold standard and universal cut-offs available across different assays measuring plasma β -amyloid or tau, we have chosen diagnostics β -amyloid ($A+ = A\beta_{42/40} < 0.056$) cutoff.³⁸ Based on this cutoff, we classified participants into either normal ($A-$) and amyloid pathology ($A+$) groups.

Cognition

Cognitive functioning was evaluated using the learning and memory subtests from the ANB, including the: African List Memory Test (ALMT; verbal learning and memory; long delay free recall correct score) and African Visuospatial Memory Test (AVMT; visuospatial memory; long delay recall correct score). The ANB has been shown to have good psychometric properties in evaluating effects of aging and neurological disease alongside providing culturally and linguistically appropriate neuropsychological measures for SSA countries.³⁵

African List Memory Test (ALMT). The ALMT is a measure of unstructured verbal learning and memory. It consists of 12 words from 4 semantic categories (body parts, means of transportation, animals, and food). The list is read at the rate of one word per 3 seconds across 3 learning trials. Examinees are asked to recall words from the target list immediately following the presentation of the list (raw scores ranging from 0 to 12). A second 12-word interference list is then presented after the third learning trial, followed by the short-delayed recall, and then a long-delay free recall trial approximately 20 minutes later.³⁹

African Visuospatial Memory Test (AVMT). The AVMT is measure of visuospatial learning and memory. This task involves encoding and retaining traditional cultural symbols found in the arts, including woodcarving, textiles, and prints, inherent to many SSA countries.

The examinee is first presented with a page of 4 symbols organized in a 2x2 matrix. The page is displayed for 10 seconds, after which examinees were asked to reproduce (on a blank sheet of paper) as many of the symbols, in their correct location on the page. For each of the 4 individual symbols, scores range from 0 to 5 yielding a total score (accuracy of the drawing and location) from 0 to 20 for each trial.³⁹

Calculation of Initial Learning and Learning Slopes

Initial learning

Initial learning was assessed using Trial 1 score for each measure. A weighted initial learning score was calculated by adjusting the point values for each test by the number of items, with the test awarding the most points per trial - AVMT (20 points) – serving as the standard. Because the highest score of items in ANB memory tests is 20, we made 20 the total for each trial to match the maximum total of 20 in AVMT. The weighted average is adjusted for the number of items for each test (e.g., $1.67 * 12 = 20$). Therefore, ALMT contains 12 possible points that are multiplied by 1.67 so the total approximated the 20 points of AVMT.

We used the following formula:

$$\text{Initial learning} = (1.67 * \text{ALMT Trial 1}) + (\text{AVMT Trial 1})$$

Learning Slope

For learning slope we calculated raw learning score (RLS), LOT (learning over trials) (modified from Morrison et al., 2018), learning ratio (LR) (Spencer et al., 2020), and total LR.

See below for equations.

$$\begin{aligned} \text{RLS} &= \text{Trial 3 performance} - \text{Trial 1 performance} \\ \text{Total RLS} &= \text{AVMT RLS} + (1.67 * \text{ALMT RLS}) \\ \text{LOT} &= \text{Sum of performance on Trials 1 through 3} - (3 * \text{Trial 1 performance}) \end{aligned}$$

$$\begin{aligned}\text{LOT Total} &= \text{Sum of AVMT trials} - (3 * \text{Trial 1 AVMT performance}) + (1.67 * \text{Sum of ALMT trials}) - (5 * \text{Trial 1 ALMT performance}) \\ \text{LR} &= \frac{\text{Trial 3 performance} - \text{Trial 1 performance}}{\text{Total Points Available for a Trial} - \text{Trial 1 performance}} \\ \text{Total LR} &= \frac{\text{AVMT RLS} + (1.67 * \text{ALMT RLS})}{40 - (\text{AVMT Trial 1}) - (1.67 * \text{ALMT Trial 1})}\end{aligned}$$

Statistical Analyses

Statistical analyses were performed using SAS statistical software.³³ Pearson correlations were calculated between individual learning slope performances and ANB memory measures. Linear regression was used to assess between-group differences of various learning slope scores (RLS, LOT, and LR) in which amyloid pathology and APOEε4 status served as predictor variables. These analyses controlled for demographic characteristics (i.e., age, education, gender). We used receiver operating characteristic (ROC) curve analyses to calculate areas under curve (AUCs) to describe diagnostic accuracy of amyloid pathology based on screening tests (CSID, AQ), ANB memory scores, and learning slope metrics (RLS, LOT, LR). We divided the participants based on amyloid positivity or negativity based on our previous cutoffs for amyloid pathology. We used Hosmer and colleagues ROC-AUC categories,⁴⁰ which considered the value of <0.600 as “failure”, values between 0.600 and 0.699 as “poor”, values between 0.700 and 0.799 as “fair”, values between 0.800 and 0.899 as “good”, and values 0.900 or greater as “excellent”. Cutoff scores for screening tests and ANB learning slopes were determined based on optimal sensitivity and specificity for detecting the presence of β-amyloid pathology. We obtained Youden’s J indices (sensitivity + specificity – 100) for each plasma biomarker. We estimated the predicted value of the measure and then the cutoffs. We also calculated the Cohen’s d as follows:

$$\text{Cohen's } d = \frac{M1 - M2}{SD_{pooled}}$$

$$SD_{pooled} = \sqrt{\frac{((n_1 - 1)s_1^2) + ((n_2 - 1)s_2^2)}{n_1 + n_2 - 2}}$$

RESULTS

Demographic, Cognitive, and Clinical Characteristics of Sample

Nearly half of participants had amyloid pathology (44%), and nearly half were APOEε4 positive (49%). Demographic data, neuropsychological scores, and plasma biomarker characteristics stratified by amyloid pathology are presented in Table 1. Groups did not significantly differ between those with and without amyloid pathology with regard to demographics (e.g., age, education, sex). The groups did not differ with respect to cognition after controlling for age, education, and gender. Based on Cohen's d results, AQ total score, delayed recall total, RLS total, and LOT total had at least moderate effect sizes, while the LR total score had a large effect size. Regarding subtest scores, most of the learning slope scores from each of the subtests ranged from small to moderate effect sizes.

Table 1. Demographic, neuropsychological, and behavioral variables stratified by amyloid pathology with the Cohen's d

Variable	Normal (A-) (n=44)	Amyloid Pathology (A+) (n=35)	Cohen's d
Age (years)	71.92 (7.46)	74.82 (7.02)	0.40
Education (years)	9.20 (5.54)	7.03 (5.20)	-0.40
Sex (% female)^a	22 (50.00%)	24 (63.16%)	--
CSID Total	26.21 (7.42)	23.61 (7.61)	-0.35
AQ Total	9.51 (8.45)	14.74 (8.24)	-0.63**
Initial Learning ALMT	4.20 (2.27)	3.71 (2.05)	-0.23
Initial Learning AVMT	3.30 (3.00)	1.83 (2.04)	-0.56*

Initial Learning Total	10.32 (6.19)	8.03 (4.52)	-0.42
Delayed Recall ALMT	3.89 (3.37)	2.63 (3.25)	-0.38
Delayed Recall AVMT	5.36 (4.86)	2.34 (3.28)	-0.71**
Delayed Recall Total	11.85 (9.94)	6.73 (8.23)	-0.56*
RLS ALMT	2.07 (1.66)	1.40 (1.59)	-0.41
RLS AVMT	2.93 (3.55)	1.14 (2.21)	-0.59**
RLS Total	6.39 (4.98)	3.48 (3.56)	-0.66**
LOT ALMT	3.55 (2.68)	2.29 (2.08)	-0.52*
LOT AVMT	5.14 (6.30)	2.60 (4.17)	-0.46*
LOT Total	11.10 (8.05)	6.45 (5.79)	-0.65**
LR ALMT	0.27 (0.24)	0.18 (0.23)	-0.38
LR AVMT	0.19 (0.23)	0.06 (0.13)	-0.68**
LR Total	0.24 (0.21)	0.11 (0.12)	-0.74**

^a Represented as n (%)

* p<.05

**p<.01

Regarding APOE ϵ 4 status, some significant group differences were observed.

Specifically, those without APOE ϵ 4 performed significantly better than those with the ϵ 4 allele across cognitive screening measures and tasks of verbal and visual learning and memory. For the primary test scores, initial learning for visuospatial information had a small effect size, while delayed recall, RLS, and LOT scores for visuospatial information had moderate effect sizes. The learning slope metrics were comparable with initial learning total score having a small effect size alongside mostly moderate effect sizes for RLS, LOT, and delayed recall scores (see Table 2).

Table 2. Demographic, neuropsychological, and behavioral variables stratified by APOE ϵ 4 status with Cohen's D

Variable	APOϵ4- (n=42)	APOϵ4+ (n=39)	Cohen's d
Age (years)	71.85 (7.06)	74.10 (8.20)	0.29
Education (years)	8.21 (5.44)	8.54 (5.29)	0.06
Sex (% female)^a	27 (62.79%)	20 (48.78%)	
CSID Total	27.01 (7.86)	23.03 (6.91)	-0.54*

AQ Total	9.54 (8.45)	14.56 (8.48)	0.59**
Initial Learning ALMT	4.50 (2.06)	3.49 (2.22)	-0.47*
Initial Learning AVMT	3.07 (2.77)	2.23 (2.57)	-0.31
Initial Learning Total	10.59 (5.49)	8.05 (5.48)	-0.46*
Delayed Recall ALMT	4.45 (3.39)	2.18 (3.02)	-0.71**
Delayed Recall AVMT	5.43 (4.52)	2.95 (4.53)	-0.55*
Delayed Recall Total	12.86 (9.71)	6.59 (9.00)	-0.67**
RLS ALMT	2.10 (1.86)	1.44 (1.48)	-0.39
RLS AVMT	3.10 (3.41)	1.44 (3.32)	-0.49*
RLS Total	6.59 (5.00)	3.83 (4.83)	-0.56*
LOT ALMT	3.38 (2.70)	2.46 (2.39)	-0.36
LOT AVMT	5.64 (5.88)	2.54 (5.23)	-0.56*
LOT Total	11.33 (7.83)	6.68 (7.10)	-0.58**
LR ALMT	0.29 (0.27)	0.17 (0.21)	-0.33*
LR AVMT	0.19 (0.21)	0.09 (0.20)	-0.05*
LR Total	0.24 (0.19)	0.13 (0.19)	-0.26*

^a Represented as n (%)

* p<.05

**p<.01

We calculated the sensitivity and specificity of screening and memory subtests to predict amyloid positivity or negativity and the presence of APOEε4 allele (see Table 3). In predicting amyloid pathology, sensitivity of cognitive tests varies between 56.8 (CSID) to 82.9 (Delayed AVMT). The sensitivity of cognitive tests to predict APOEε4 allele was between 60.5 (Aβ_{42/40}) and 74.4 (Delayed AVMT).

Table 3. Sensitivity and Specificity of cognitive tests to predict A+ or APOEε4+ status

Test	CIP Threshold	Aβ _{42/40}			APOEε4		
		High Aβ _{42/40} (A-)	Low Aβ _{42/40} (A+)	Sensitivity/ Specificity	APOEε4 -	APOEε4 +	Sensitivity/ Specificity
CSID Screening	≥25.5 (HC)	25	16	56.8 / 59.5	26	15	62.5 / 65.0
	<25.5 (Dem)	17	21		14	25	
AQ Screening	<13 (HC)	24	13	65.8 / 55.8	25	12	70.7 / 61.0
	≥13 (Dem)	19	25		16	29	
Delayed ALMT	≥5 (HC)	23	13	62.9 / 52.3	26	11	71.8 / 61.9
	<5 (Dem)	21	22		16	28	

Delayed AVMT	≥ 5 (HC)	25	6	82.9 / 56.8	23	10	74.4 / 54.8
	< 5 (Dem)	19	29		19	29	
Aβ_{42/40}	> 0.056 (HC)	-	-	-	26	17	60.5 / 60.5
	≤ 0.056 (Dem)	-	-	-	15	23	

*CIP = cognitive impairment

**p-value < 0.05 means that healthy controls and AD are significantly classified differently between the 2 metrics. Therefore, p-value ≥ 0.05 signifies the groups were classified similarly.

We calculated Pearson correlation coefficients to examine the relationship between total learning slope metrics, initial learning, and delayed recall (See Table 4). Among the learning slope options, LR had the strongest correlations with both initial and delayed recall. The learning slope metrics generally had high correlations with each other.

Table 4: Pearson correlations between total learning slope performances

Variable	Initial Learning Total	RLS Total	LOT Total	LR Total
RLS Total	0.18 (0.056)	-		
LOT Total	0.20 (0.036)	0.94 ($< .0001$)	-	
LR Total	0.39 ($< .0001$)	0.95 ($< .0001$)	0.89 ($< .0001$)	-
Delayed Recall Total	0.82 ($< .0001$)	0.48 ($< .0001$)	0.51 ($< .0001$)	0.63 ($< .0001$)

Analyses of ROC-AUC, Cutoffs, and Sensitivity/Specificity

Tables 5 and 6 display the ROC-AUC values for the screening measures, ANB memory subtests, learning slopes, and total scores when differentiating individuals between groups with amyloid pathology and APOE ϵ 4 status. Our results showed that all AUC/ROC ranges fell between 0.56-0.70 (95% CIs ranging from 0.44-0.82) and the sensitivity varied between 51.4-88.6.

Table 5. Receiver operating characteristic area under the curve, cut scores, and sensitivity/specificity with differentiating amyloid biomarker groups for screening tests, ANB memory subtests, and learning slopes across the total sample

Variable	AUC (95% CI)	Cut Score	Sensitivity/Specificity
CSID Total	0.61 (0.48,0.74)	27.33	64.9 / 54.8

AQ Total	0.68 (0.57,0.80)	6.00	76.3 / 53.5
Delayed Recall ALMT	0.61 (0.49,0.73)	5.00	74.3 / 45.5
Delayed Recall AVMT	0.68 (0.56,0.79)	4.00	77.1 / 63.6
Delayed Recall Total	0.65 (0.53,0.77)	12.02	68.6 / 54.5
Initial Learning ALMT	0.56 (0.44,0.69)	6.00	82.9 / 27.3
Initial Learning AVMT	0.65 (0.53,0.77)	3.00	77.1 / 50.0
Total Initial Learning	0.63 (0.50,0.75)	10.35	77.1 / 43.2
RLS ALMT	0.62 (0.50,0.75)	3.00	80.0 / 43.2
RLS AVMT	0.67 (0.55,0.78)	3.00	77.1 / 50.0
RLS Total	0.70 (0.58,0.82)	2.67	54.3 / 81.8
LOT ALMT	0.64 (0.52,0.76)	4.00	88.6 / 34.1
LOT AVMT	0.62 (0.49,0.74)	6.00	77.1 / 36.4
LOT Total	0.69 (0.57,0.81)	5.37	51.4 / 79.5
LR ALMT	0.62 (0.50,0.75)	0.33	80.0 / 38.6
LR AVMT	0.68 (0.56,0.80)	0.17	85.7 / 45.5
LR Total	0.70 (0.58,0.82)	0.10	60.0 / 72.7

Regarding the APOE ϵ 4, all ROC/AUC values ranged between 0.60-0.70. With regards to total scores, the sensitivity of initial learning, delayed recall, and the learning slopes ranged from 43.6-89.7 (Table 6).

Table 6. Receiver operating characteristic area under the curve, cut scores, and sensitivity/specificity with differentiating APOE ϵ 4+ for cognitive screening tests, ANB memory subtests, and learning slopes across the total sample

Variable	AUC (95% CI)	Cut Score	Sensitivity/ Specificity
CSID Total	0.67 (0.55, 0.80)	27.33	70.0 / 60.0
AQ Total	0.65 (0.53, 0.77)	9.00	70.7 / 61.0
Delayed Recall ALMT	0.68 (0.57, 0.80)	5.00	71.8 / 61.9
Delayed Recall AVMT	0.68 (0.57, 0.80)	5.00	82.1 / 50.0
Delayed Recall Total	0.70 (0.58, 0.81)	10.01	71.8 / 64.3
Initial Learning ALMT	0.64 (0.52, 0.76)	4.00	53.8 / 81.0
Initial Learning AVMT	0.60 (0.48, 0.72)	5.00	89.7 / 31.0
Initial Learning Total	0.66 (0.53, 0.78)	6.68	43.6 / 81.0
RLS ALMT	0.63 (0.51, 0.75)	3.00	82.1 / 47.6
RLS AVMT	0.69 (0.57, 0.80)	2.00	69.2 / 61.9
RLS Total	0.68 (0.56, 0.79)	5.68	74.4 / 50.0
LOT ALMT	0.61 (0.48, 0.73)	5.00	84.6 / 33.3
LOT AVMT	0.69 (0.58, 0.81)	1.00	53.8 / 73.8
LOT Total	0.69 (0.57, 0.81)	4.37	46.2 / 83.3
LR ALMT	0.66 (0.54, 0.78)	0.33	87.2 / 45.2
LR AVMT	0.69 (0.57, 0.81)	0.06	69.2 / 61.9
LR Total	0.69 (0.57, 0.80)	0.11	56.4 / 71.4

DISCUSSION

The presence of amyloid and APOE ϵ 4 allelic presence is each associated with AD, and early detection of these factors can open possibilities for intervention. Performance-based tests are a low cost and technologically portable method for potentially screening for these important biological prognostic factors. These results indicated that memory scores from the ANB have potential utility for detecting both APOE ϵ 4+ status and amyloid deposition. In stratifying by amyloid pathology, most total scores, including learning slope and delayed recall, approached adequate utility as a screening measure aside from initial learning did not distinguish between those with and without amyloid deposition. Most subtests also showed adequate utility, aside from delayed recall ALMT, RLS ALMT, and LR ALMT. A similar pattern of findings emerged when screening for APOE ϵ 4 status. In these analyses, each of the total scores and ANB subtest scores (aside from initial learning AVMT, RLS ALMT, and LOT ALMT), including initial learning, approached the threshold for being an adequate screen.

Research by Hammers and colleagues (2022) has suggested that the LR score has diagnostic advantages over other learning slope metrics such as RLS and LOT scores.¹⁸ In the present study, however, each learning slope score was roughly equivalent as a screening tool, with measures generally resulting in “fair” diagnostic properties and moderate differences in group means. The equivalence observed across measures is likely due to the lack of ceiling effects encountered in the present study compared to prior research that has demonstrated that ceiling effects are readily apparent in RLS scores when LR scores have been shown to be a superior measure over RLS scores. Our analyses revealed that the LR score was significantly correlated with initial learning, though shared a much stronger correlation with the other learning slope metrics that are dependent on raw improvement over the course of the test. The lack of

ceiling effect essentially renders each of the learning slope scores as being interchangeable. In fact, no participant achieved a perfect score on the ALMT or AVMT.

Results also demonstrate that the ANB learning scores show promise in detecting the presence of amyloid and APOE ϵ 4status, but the diagnostic statistics fall just shy of justifying their use in isolation. Although these results do not provide strong support for using these measures independently, additional research is needed to examine whether they may provide useful information when combined with other variables. For instance, data from other tests or biomedical variables could be combined with ANB scores to provide a more refined screening for AD pathology. It is important to note that these results pertain to potentially subtle manifestations and/or risk factors for AD and were not applied to variables associated with more pronounced cognitive pathology, such as significant tau burden or neurodegeneration. We suspect that ceiling effects will be less problematic for advanced disease burden, and, therefore, each of the learning slope scores are likely to function interchangeably in this context as well. Clinically, additional research into understanding these learning metrics and cut-offs is warranted before using them in the SSA.

This study is the first in the SSA to attempt to predict the amyloid pathology and APOE ϵ 4status using cognitive screening measures (CSID and AQ), ANB memory subtests (ALMT and AVMT), and learning slope metrics (initial learning, RLS, LOT, and LR). This study has several limitations. First, this current study has a small sample of participants, which limited the detection of differences that could have been clinically and significantly relevant to discriminate the two groups. Thus, future studies should replicate these findings with larger sample sizes. Second, the screening measures used (CSID and AQ) have not been validated in the SSA/DRC. Third, this study included only subjects with suspected dementia and healthy

controls. Those with cognitive difficulties seen in between the spectrum (e.g., MCI, subjective memory complaints) were excluded, leaving only the extremes of the dementia spectrum. Future studies should include all the 4 groups (healthy controls, MCI, subjective memory complaint and dementia). Additionally, this study analyzed only amyloid pathology and APOE ϵ 4 allele status. Other limitations include lack of normative data for the fluid biomarkers used here in the studied population, relatively small sample size for a biomarker study, and no replication cohort. Future studies should aim to replicate our findings along the AD pathology continuum, tau pathology, neurofilament light (NfL), as well as use other plasma biomarkers (e.g., glial fibrillar acidic protein [GFAP]), as they may provide greater insight into the progression of cerebral amyloid and tau pathology and cognitive decline in SSA populations. A major caveat is that amyloid positive was determined with plasma A β _{42/40} by Simoa, which is far from being the best biomarker for amyloid positivity. The gold standard is amyloid brain PET. Of the plasma biomarkers P-tau₂₁₇ by mass spectrometry or ECL and A β ₄₂ / A β ₄₀ by mass spectrometry seem the best with AUC > 0.90. Plasma P-tau₁₈₁ by Simoa is a reasonably robust alternative. It is not just a marker of tau pathology. It also is an accurate predictor of brain amyloid pathology detected by neuropathology of brain amyloid PET.⁴¹

REFERENCES

1. Salmon DP, Bondi MW. Neuropsychological Assessment of Dementia. *Annu Rev Psychol.* 2009;60:257. doi:10.1146/ANNUREV.PSYCH.57.102904.190024
2. Hammers DB, Atkinson TJ, Dalley BCA, et al. Amyloid positivity using [18F]Flutemetamol-PET and cognitive deficits in nondemented community-dwelling older adults. *Am J Alzheimers Dis Other Demen.* 2017;32(6):320-328. doi:10.1177/1533317517698795/ASSET/IMAGES/LARGE/10.1177_1533317517698795-FIG1.JPEG
3. Hammers DB, Suhrie K, Dixon A, Gradwohl BD, Duff K, Spencer RJ. Validation of HVLt-R, BVMT-R, and RBANS Learning Slope Scores along the Alzheimer's Continuum. *Archives of Clinical Neuropsychology.* 2022;37(1):78-90. doi:10.1093/ARCLIN/ACAB023
4. Bonner-Jackson A, Mahmoud S, Miller J, Banks SJ. Verbal and non-verbal memory and hippocampal volumes in a memory clinic population. *Alzheimers Res Ther.* 2015;7(1):1-10. doi:10.1186/S13195-015-0147-9/TABLES/5
5. Salmon DP. Disorders of memory in Alzheimer's disease. In: Cermak LS, ed. *Handbook of Neuropsychology: Memory and Its Disorders.* Elsevier Science Publishers B.V.; 2000:155-195.
6. Caselli RJ, Langlais BT, Dueck AC, et al. Neuropsychological decline up to 20 years before incident mild cognitive impairment. *Alzheimer's & Dementia.* 2019;16(3):512-523. doi:10.1016/J.JALZ.2019.09.085
7. Machulda MM, Hagen CE, Wiste HJ, et al. Practice effects and longitudinal cognitive change in clinically normal older adults differ by Alzheimer imaging biomarker status. <http://dx.doi.org/101080/1385404620161241303>. 2016;31(1):99-117. doi:10.1080/13854046.2016.1241303
8. Duff K, Hammers DB, Dalley BCA, et al. Short-Term Practice Effects and Amyloid Deposition: Providing Information Above and Beyond Baseline Cognition. *J Prev Alzheimers Dis.* 2017;4(2):87. doi:10.14283/JPAD.2017.9
9. Lim YY, Baker JE, Bruns L, et al. Association of deficits in short-term learning and A β and hippocampal volume in cognitively normal adults. *Neurology.* 2020;95(18):e2577-e2585. doi:10.1212/WNL.00000000000010728
10. Thomas KR, Eppig J, Edmonds EC, et al. Word-list intrusion errors predict progression to mild cognitive impairment. *Neuropsychology.* 2018;32(2):235-245. doi:10.1037/NEU0000413
11. Delis DC, Kramer JH, Kaplan E, Ober BA. *California Verbal Learning Test, Second Edition.* The Psychological Corporation; 2000.
12. Brandt J. The hopkins verbal learning test: Development of a new memory test with six equivalent forms. <https://doi.org/101080/13854049108403297>. 2007;5(2):125-142. doi:10.1080/13854049108403297
13. Schmidt M. *Rey Auditory Verbal Learning Test: A Handbook*. Western Psychological Services; 1996.
14. Randolph C. *Repeatable Battery for the Assessment of Neuropsychological Status (RBANS)*. Psychological Corporation; 1998. Accessed August 15, 2023. <https://portal4care.cdhlh.be/nl/BESTNL/Cognitief%20functioneren/Repeatable%20Battery>

- %20for%20the%20Assessment%20of%20Neuropsychological%20Status%20(RBANS).pdf
15. Randolph C. *RBANS Update: Repeatable Battery for the Assessment of Neuropsychological Status*. PsychCorp; 2012.
 16. Benedict RH. *Brief Visuospatial Memory Test--Revised*. PAR; 1997. Accessed August 15, 2023.
https://scholar.google.com/scholar_lookup?title=The%20brief%20visuospatial%20memory%20test%20-%20revised&publication_year=1997&author=R.H.B.%20Benedict
 17. Spencer RJ, Gradwohl BD, Williams TF, Kordovski VM, Hammers DB. Developing learning slope scores for the repeatable battery for the assessment of neuropsychological status. <https://doi.org/10.1080/2327909520201791870>. Published online 2020:1-7. doi:10.1080/23279095.2020.1791870
 18. Hammers DB, Spencer RJ, Apostolova LG. Validation of and Demographically Adjusted Normative Data for the Learning Ratio Derived from the RAVLT in Robustly Intact Older Adults. *Archives of Clinical Neuropsychology*. 2022;37(5):981-993. doi:10.1093/ARCLIN/ACAC002
 19. Hammers DB, Duff K, Spencer RJ. Demographically-corrected normative data for the HVLTL-R, BVMT-R, and Aggregated Learning Ratio values in a sample of older adults. <https://doi.org/10.1080/1380339520211917523>. 2021;43(3):290-300. doi:10.1080/13803395.2021.1917523
 20. Hammers DB, Duff K, Spencer RJ. Demographically-corrected normative data for the RBANS learning ratio in a sample of older adults. <https://doi.org/10.1080/1385404620211952308>. 2021;36(8):2221-2236. doi:10.1080/13854046.2021.1952308
 21. Ikanga J, Taiwo Z, Lengu K, et al. Performance on the African neuropsychology battery using the learning ratio in a sample of healthy Congolese. *Journal of the International Neuropsychological Society*. 2023;29(5):503-511. doi:10.1017/S135561772200039X
 22. Hammers DB, Suhrie K, Dixon A, et al. Relationship between a novel learning slope metric and Alzheimer's disease biomarkers. <https://doi.org/10.1080/1382558520211919984>. 2021;29(5):799-819. doi:10.1080/13825585.2021.1919984
 23. Hammers DB, Gradwohl BD, Kucera A, Abildskov TJ, Wilde EA, Spencer RJ. Preliminary Validation of the Learning Ratio for the HVLTL-R and BVMT-R in Older Adults. *Cognitive and Behavioral Neurology*. 2021;34(3):170-181. doi:10.1097/WNN.0000000000000277
 24. Hammers DB, Kostadinova R V., Spencer RJ, et al. Sensitivity of memory subtests and learning slopes from the ADAS-Cog to distinguish along the continuum of the NIA-AA Research Framework for Alzheimer's Disease. <https://doi.org/10.1080/138255852022120957>. Published online 2022. doi:10.1080/13825585.2022.2120957
 25. Jack CR, Bennett DA, Blennow K, et al. NIA-AA Research Framework: Toward a biological definition of Alzheimer's disease. *Alzheimer's & Dementia*. 2018;14(4):535-562. doi:10.1016/J.JALZ.2018.02.018
 26. Bateman RJ, Xiong C, Benzinger TLS, et al. Clinical and Biomarker Changes in Dominantly Inherited Alzheimer's Disease. *New England Journal of Medicine*.

- 2012;367(9):795-804.
doi:10.1056/NEJMOA1202753/SUPPL_FILE/NEJMOA1202753_DISCLOSURES.PDF
27. Thomas KR, Weigand AJ, Edwards LC, et al. Tau levels are higher in objective subtle cognitive decline but not subjective memory complaint. *Alzheimers Res Ther.* 2022;14(1). doi:10.1186/s13195-022-01060-1
 28. Thomas KR, Bangen KJ, Weigand AJ, et al. Objective subtle cognitive difficulties predict future amyloid accumulation and neurodegeneration. *Neurology.* 2020;94(4):e397-e406. doi:10.1212/WNL.00000000000008838
 29. Janelidze S, Mattsson N, Palmqvist S, et al. Plasma P-tau181 in Alzheimer's disease: relationship to other biomarkers, differential diagnosis, neuropathology and longitudinal progression to Alzheimer's dementia. *Nature Medicine* 2020 26:3. 2020;26(3):379-386. doi:10.1038/s41591-020-0755-1
 30. Schneider P, Hampel H, Buerger K. Biological Marker Candidates of Alzheimer's Disease in Blood, Plasma, and Serum. *CNS Neurosci Ther.* 2009;15(4):358-374. doi:10.1111/J.1755-5949.2009.00104.X
 31. Malek-Ahmadi M, Sabbagh MN, The Cleo Roberts M. Development and Validation of the Alzheimer's Questionnaire (AQ). *J Nat Sci.* 2015;1(5):e104. Accessed December 18, 2022. /pmc/articles/PMC4423544/
 32. Hall KS, Gao S, Emsley CL, Ogunniyi AO, Morgan O, Hendrie HC. Community screening interview for dementia (CSI 'D'); performance in five disparate study sites. *Int J Geriatr Psychiatry.* 2000;15(6):521-531. doi:10.1002/1099-1166(200006)15:6<521::aid-gps182>3.0.co;2-f
 33. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders.* 5th ed. American Psychiatric Association; 2013.
 34. Guerchet M, M'Belesso P, Mouanga AM, et al. Prevalence of Dementia in Elderly Living in Two Cities of Central Africa: The EDAC Survey. *Dement Geriatr Cogn Disord.* 2010;30(3):261-268. doi:10.1159/000320247
 35. Ikanga J, Reyes A, Kaba D, et al. Prevalence of suspected dementia in a sample of adults living in Kinshasa-Democratic Republic of the Congo. *Alzheimer's & Dementia.* Published online 2023. doi:10.1002/ALZ.13003
 36. Verberk IMW, Misdorp EO, Koelewijn J, et al. Characterization of pre-analytical sample handling effects on a panel of Alzheimer's disease-related blood-based biomarkers: Results from the Standardization of Alzheimer's Blood Biomarkers (SABB) working group. *Alzheimer's & Dementia.* 2022;18(8):1484-1497. doi:10.1002/ALZ.12510
 37. Hu Y, Kirmess KM, Meyer MR, et al. Assessment of a Plasma Amyloid Probability Score to Estimate Amyloid Positron Emission Tomography Findings Among Adults With Cognitive Impairment. *JAMA Netw Open.* 2022;5(4):e228392-e228392. doi:10.1001/JAMANETWORKOPEN.2022.8392
 38. Vergallo A, Mégret L, Lista S, et al. Plasma amyloid β 40/42 ratio predicts cerebral amyloidosis in cognitively normal individuals at risk for Alzheimer's disease. *Alzheimers Dement.* 2019;15(6):764-775. doi:10.1016/J.JALZ.2019.03.009
 39. Ikanga J, Basterfield C, Taiwo Z, et al. The Reliability of the African Neuropsychology Battery in Persons of African Descent. *Archives of Clinical Neuropsychology.* 2022;37(4):839-848. doi:10.1093/ARCLIN/ACAC003
 40. Hosmer Jr DW, Lemeshow S, Sturdivant R. *Applied Logistic Regression.*; 2013. Accessed September 28, 2023.

<https://books.google.com/books?hl=en&lr=&id=64JYAwwAAQBAJ&oi=fnd&pg=PR13&ots=DtkQ2V6moI&sig=n7QQPuwd-TECFjNDYIJPLzmLW3c>

41. Thijssen EH, La Joie R, Wolf A, et al. Diagnostic value of plasma phosphorylated tau181 in Alzheimer's disease and frontotemporal lobar degeneration. *Nat Med*. 2020;26(3):387. doi:10.1038/S41591-020-0762-2