**Unveiling the pathogenesis of non-alcoholic fatty liver disease by**

**decoding biomarkers through integrated single-cell and single-nucleus**

**profiles**

Wenfeng Ma [1,2,3,4], Xin Zhong [1,4], Benqiang Cai [1,4], Mumin Shao [6,7], Xuewen Yu [6,7],

Minling Lv [1,4], Shaomin Xu [1,4], Bolin Zhan [1,4], Qun Li [1,4], Mengqing Ma [1,4], Mikkel

Brejnholt Kjær [5,8], Jinrong Huang [2]*, Yonglun Luo [2,3]*, Henning Grønbæk [5]*, Lin Lin

[2,3]*

1 Department of Liver Disease, Shenzhen Traditional Chinese Medicine Hospital,

Shenzhen, Guangdong 518033, China.

2 Department of Biomedicine, Aarhus University, Aarhus, Denmark.

3 Steno Diabetes Center Aarhus, Aarhus University Hospital, Aarhus, Denmark.

4 Department of Liver Disease, The Fourth Clinical Medical College of Guangzhou

University of Chinese Medicine, Shenzhen, 518033, China.

5 Department of Hepatology and Gastroenterology, Aarhus University Hospital,

Aarhus, Denmark.

6 Department of Pathology, Shenzhen Traditional Chinese Medicine Hospital,

Shenzhen, Guangdong 518033, China.

7 Department of Pathology, The Fourth Clinical Medical College of Guangzhou

University of Chinese Medicine, Shenzhen, 518033, China.

8 Department of Clinical Medicine, Aarhus University, Aarhus, Denmark.

* = corresponding author

## Highlights

• Integrated single-cell and single-nucleus profiles from 82 liver and PBMC

specimens comprising NAFLD patients and healthy controls with increasing

severity were utilized to unveil the NAFLD pathogenesis through decoding

candidate biomarkers of NAFLD.

• In cell-level observations, we decoded 16 up-regulated and 22 down-regulated

secreting genes previously identified as associated with increasing NAFLD severity

in the liver RNA-seq and plasma proteomics data.

• QSOX1, enriched in fibroblasts, and IL1RAP, enriched in hepatocytes, have been

further validated and interpreted in integrated single-cell and single-nucleus

profiles for their potential to predict NAFLD severity.

• The analysis of intercellular crosstalk, focusing on secreted signaling from our

previously identified candidate biomarkers sourced from secreted genes,

highlighted the IL1B-(IL1R1+IL1RAP) pathway between liver monocytes and

hepatocytes/cholangiocytes. This suggests that this pathway might be a potential

reason for the observed downregulation of IL1RAP in NAFLD liver.

## Abstract

**Background and Aims:** Non-alcoholic fatty liver disease (NAFLD) is a heterogenous

liver disease encompassing pathological changes ranging from simple steatosis,

inflammation and fibrosis to cirrhosis. To further unravel NAFLD pathogenesis, we

aimed to decode the candidate NAFLD biomarkers associated with NAFLD severity

using publicly available single-cell RNA sequencing (scRNA-seq) and single-nucleus

RNA sequencing (snRNA-seq) data.

**Methods:** Seurat v5 and anchor-based reciprocal principal components analysis

(RPCA) integration were performed to integrate and analyze the scRNA-seq and

snRNA-seq data of 82 liver and Peripheral Blood Mononuclear Cell (PBMC)

specimens from NAFLD patients and healthy controls to decode the candidate

NAFLD biomarkers generated previously. Using the 'CellChat' R package, we analyzed

ligand-receptor interactions of our candidate biomarkers from secreted genes to

understand their signaling crosstalk and implications in NAFLD's biological processes.

**Results:** We generated a database

(https://dreamapp.biomed.au.dk/NAFLD-scRNA-seq/) to present the NAFLD

pathogenesis by analyzing integrated scRNA-seq and snRNA-seq data. Through

cell-level decoding, we discovered the expression distribution of the candidate

biomarkers associated with NAFLD severity. The analysis of ligand-receptor pairs in

NAFLD liver and PBMC data suggests that the IL1B-(IL1R1+IL1RAP) interaction

between liver monocytes and hepatocytes/cholangiocytes may explain the

correlation between NAFLD severity and IL1RAP down-regulation.

**Conclusions:** We confirmed a strong correlation between liver QSOX1/IL1RAP

concentrations and NAFLD severity at the cellular level. Additionally, our analysis of

comprehensive data unveiled new aspects of NAFLD pathogenesis and intercellular

communication through the use of scRNA and snRNA sequencing data.

(ChiCTR2300073940).

**Lay Summary**

We integrated single-cell RNA sequencing (scRNA-seq) and single-nucleus RNA

sequencing (snRNA-seq) data to unravel non-alcoholic fatty liver disease (NAFLD)

pathogenesis. We decoded candidate biomarkers associated with NAFLD progression,

which were previously screened from RNA sequencing (RNA-seq) data of 625 liver

samples with a novel gene clustering method. A new version of the R package

''Seurat v5' and anchor-based reciprocal principal components analysis (RPCA)

integration were performed to process and integrate scRNA-seq and snRNA-seq data

of 82 liver and Peripheral Blood Mononuclear Cell (PBMC) specimens from NAFLD

patients and healthy controls. The research delved deeper into the cellular

expression patterns of the candidate biomarkers and examined the intercellular

communication of their secreted signaling.

**Graphical abstract**

**Keywords**

## Introduction

Non-alcoholic fatty liver disease (NAFLD) is a multifaceted disease representing a spectrum from Non-alcoholic Fatty Liver (NAFL) of simple steatosis, non-alcoholic steatohepatitis (NASH) to NASH-induced fibrosis and cirrhosis (1, 2). As the global prevalence of NAFLD reached 38.0%, the upward trend in NAFLD and NASH-related liver cancer balanced the declining tendency of virally-caused hepatocellular carcinoma (3-5). Despite the noteworthy advancements in NAFLD research, the lack of FDA-approved treatments underscores the need for a systematic study to understand its pathogenesis (6, 7).

The advent of transcriptomics, specifically single-cell RNA sequencing (scRNA-seq) and single-nucleus RNA sequencing (snRNA-seq) has triggered a paradigm shift in our comprehension of biological processes, cellular diversity, and the intricacies of physiological and disease states (8). These groundbreaking technologies allow us to examine gene expression at the cell level, significantly enhancing our understanding of various aspects of life sciences (9, 10). In 2019, Xiong et al. applied scRNA-seq to reveal the cellular heterogeneity of NAFLD, identifying key cell types and crosstalk during disease progression (11). Tamburini et al. focused on liver lymphatic endothelial cells (LECs) and observed the activation of the IL13 pathway in LECs of NASH patients (12). In 2020, Ramachandran et al. utilizing single-cell transcriptomics, resolved human liver tissue and blood samples of patients with cirrhosis and healthy controls, identifying new pathogenic subpopulations and providing the markers of cell subtypes (13). In 2021, Deczkowska and colleagues discovered an abundance and activation of type 1 conventional dendritic cells (cDC1) in both NAFLD and NASH conditions with scRNA-seq. They further highlighted XCR1+ cDC1 as a significant contributor to liver pathology in NASH (14). Recently, Wang et al. interpreted NASH-associated hepatic stellate cells (HSCs) with snRNA-seq and uncovered an autocrine HSC signaling circuit that emerged in advanced fibrosis (15). These studies underscore the immense potential of transcriptomics application in NAFLD research

for finding molecular signatures and pathways that could guide the development of
diagnosis and therapeutics for NAFLD.

Recently, we demonstrated that Quiescin sulfhydryl oxidase 1 (QSOX1) and
Interleukin-1 receptor accessory protein (IL1RAP) hold potential as indicators of liver
disease severity in NAFLD patients (16). QSOX1, a flavin-linked enzyme that facilitates
disulfide bond formation within the secretory structures of mammalian cells, is
expelled into the extracellular matrix (ECM) and blood as an independent protein
through the standard secretory pathway, encompassing the Golgi apparatus,
secretory vesicles, and the plasma membrane. Furthermore, it is notably amplified
and released from fibroblasts that are actively synthesizing ECM precursors (15-17).
On the other hand, IL1RAP, a member of the IL-1 receptor family, is ubiquitously
present but shows high levels in the liver, placenta, and white blood cells, according
to databases such as NCBI. This protein plays a critical role in inflammation and is
implicated in the initiation and progression of cancer. Studies suggested that both
QSOX1 and IL1RAP hold promise as potential biomarkers of NAFLD or obesity (17,
18).

In 2022, Hao's team unveiled Seurat v5, featuring a 'bridge integration' technique
designed to merge single-cell datasets from diverse molecular domains and various
measurement modalities. Employing concepts from 'dictionary learning' within the
field of representation learning, this method is highly scalable and ideal for handling
expansive datasets of millions of cells. It has wide applicability across different
technologies and molecular modalities, making it valuable for large research
consortia with existing single-cell RNA-seq databases. Additionally, the approach
shows potential in the identification and comprehensive study of rare cellular
populations. Overall, the introduction of bridge integration in Seurat v5 is poised to
make a significant impact on the advancement of single-cell research (19). Compared
to its predecessor (20), Seurat v5 enhanced its capacity for large-scale data
integration and facilitated a more comprehensive analysis of the cellular and nuclear

transcriptomic landscapes. With Seurat v5, we aimed to use publicly available scRNA-seq and snRNA-seq data to decode the candidate NAFLD biomarkers we have previously screened for unraveling NAFLD pathogenesis.

Understanding global cellular interactions necessitates precise representation and analysis of cell-to-cell signaling pathways. "CellChat" serves as a tool, crafted to extract and evaluate intercellular communication networks from sc- and snRNA-seq data (21). It identifies primary signaling pathways for cells, decoding their coordinated actions through network analyses and pattern detection methods. Using manifold learning and quantitative assessments, CellChat classifies signaling routes and differentiates between universal and context-specific pathways across diverse datasets. Consequently, it aids in revealing novel intercellular communications and in building communication blueprints for a range of tissues (22).

## Materials and methods

### Single-cell and Single-nucleus RNA-seq Data Analysis

1. Data Collection

The scRNA-seq and snRNA-seq data from human liver and Peripheral Blood Mononuclear Cell (PBMC) of NAFLD and healthy controls were collected from the NCBI GEO (until June 2023). Only datasets generated with 10x Genomics Single-Cell Reagent Kit were included.

2. Data Alignment

The FASTQ-formatted data were downloaded and converted into feature, barcode and matrix with the human GRCh38 genome using 'cellranger' (cellranger-6.1.2). For snRNA-seq, intron was included for quantification (23). The doublets were removed with the R package 'DoubletFinder' (DoubletFinder-2.0.3) for each sample.

3. Data Normalization and Clustering

179 With the R package 'Seurat V5' (Seurat 4.9.9.9042)
180 (https://satijalab.org/seurat/articles/seurat5_integration.html), the datasets were
181 integrated and processed using the anchor-based reciprocal principal components
182 analysis (RPCA) method. After normalization and batch correction, the scRNA-seq
183 and snRNA-seq data underwent unsupervised clustering.
184
For normalization, the 'LogNormalize' method was utilized with a default scale factor
of 10,000, and subsequent feature identification and data scaling were done by
default. The scaled data was then subjected to Principal Component Analysis (PCA)
with a default setting of computing and storing 50 Principal Components (PCs). To
perform integration in low-dimensional space, we selected the anchor-based RPCA
integration method, which aimed to co-embed shared scRNA-seq and snRNA-seq
library types across batches, yielding a dimensional reduction (integrated.rpca).
192
193 For cell clustering, a shared nearest neighbor (SNN) modularity optimization-based
194 algorithm was employed, with the dimensions of reduction set to 1:15 and the
195 resolution parameter set to 0.2 (liver) and 0.1 (PBMC).
196
4.  Cell Type Annotation and Data Analysis
We annotated liver and PBMC clusters with canonical markers by manually curated
the information to enhance the accuracy of gene annotation. To gain a more nuanced
understanding of transcriptomic variations, we further segmented certain cell types
in additional supplementary files. This enabled us to identify subtype-specific
differences in both healthy and NAFLD conditions. We then compared the
expressions of potential NAFLD biomarkers across multiple clustering methodologies
and depicted the results on distinct UMAP plots.
205
5.  Cell Communication Analysis
The sc- and snRNAseq data of NAFLD liver and PBMC were merged together and
RPCA integrated before normalization. A Seurat object with a new assay of 5,000

cells in each cell type in a tissue was subsetted. The R package 'CellChat' (CellChat

1.6.1) was utilized to analyze the data. We refined the CellChatDB database to

include only interactions involving secreted signaling, as the candidate NAFLD

biomarkers originate from genes that are secreted.

6.  Database Construction

To visualize gene expression variation during the development of NAFLD severity at

the cell level, we generated a NAFLD sc- and snRNA-seq database (NAFLD-DB) using

the tool 'ShinyCell'. (https://github.com/SGDDNB/ShinyCell). This database was

deployed at https://dreamapp.biomed.au.dk/NAFLD-scRNA-seq/.

**Validation of QSOX1 and IL1RAP as Biomarkers in NAFLD at the Liver Cell Level**

1.  Study Design

Our objective was to confirm the *QSOX1* and *IL1RAP* gene expression in cells of

different severity of NAFLD. To address this question, we performed

immunohistochemical staining (IHC) for liver QSOX1/IL1RAP on a group of NAFLD

patients from Shenzhen Traditional Chinese Medicine Hospital, China (SZTCMH).

Details of this cohort have been provided previously (16).

2.  Human Samples and IHC

For IHC, we collected 12 fixed liver tissues from archived histological samples at

SZTCMH between 2014 and 2023, and details were in our earlier study (16).

3.  Statistical Analysis

The significance for all statistical tests was two-sided, with $P < 0.05$. All data analysis

was presented in the plots using R-4.3.1. Statistical analysis in the Violin plots was

performed with Wilcoxon rank sum test to assess differences between unpaired

groups.

## Results

### Alignment of ScRNA-seq and snRNA-seq Data

We incorporated twelve datasets (GSE122083, GSE125188, GSE136103, GSE144430, GSE159977, GSE169446, GSE175793, GSE179886, GSE182159, GSE189539, GSE212837 and GSE217235) into our study. The scRNAseq and snRNA-seq data encompass liver and PBMC samples of both healthy controls and patients with NAFLD (**Supplementary Table 1**). The detailed information regarding the data alignment was presented in the **Supplementary Table 2.**

By employing RPCA integration, we achieved precise comparative analyses across single-cell and single-nucleus sequencing datasets. To address batch effects arising from diverse sequencing technologies and studies involving various experimental batches, donors, and conditions, we conducted integration and normalization, as depicted in **Figure 1A**. After filtering out doublets, the dataset was refined, resulting in a total of 427,787 liver cells (including 214,795 healthy cells and 212,992 NAFLD cells) and 122,238 PBMC cells (including 92,319 healthy cells and 29,919 NAFLD cells) available for subsequent analysis.

### The Effectiveness of Integration and Normalization on Liver Transcriptome Profiles

PCA demonstrated the effectiveness of the RPCA integration and normalization process in eliminating noticeable batch effects of liver transcriptome profiles compared to the unintegrated UMAP, as depicted in **Figure 1B and 1C**. Moreover, sample separation was not primarily driven by health/NAFLD group or library type, as indicated in **Figure 1D and 1E**. Instead, cell type emerged as the primary factor influencing transcriptome profiles, as illustrated in **Figure 1F**.

### Unsupervised Clustering and Annotation of Liver Cells and Nuclei Data

Initially, we categorized all 427,787 liver cells into 12 distinct cell types using gene markers. The identified cell types were annotated using signatures of known

canonical markers (**Figure 2A, B; Supplementary Table 3**).

**Tracing the Cellular Origins of Promising Biomarkers for Liver NAFLD Severity**

In our previous study (16), we pinpointed key genes that are significantly involved in

the advancement of NAFLD. Specifically, we discovered 16 genes, including A2M, C7,

COL6A3, COLEC11, ENPP2, FBLN1, FBN1, FCGBP, IGFBP6, LCN2, LUM, MMP2, PAPLN,

PTGDS, QSOX1, and VWF, that exhibited elevated secretory activity correlating with

the severity of NAFLD. On the other hand, we also identified another set of 22 genes,

namely AZGP1, C1RL, C4BPA, C6, C8B, CFHR3, CNDP1, F2, GC, HP, HPR, IL1RAP, ITIH1,

ITIH2, ITIH4, KLKB1, PON3, SERPINA10, SERPINC1, SERPING1, SMPDL3A, and TTR,

that showed reduced secretory activity, also in relation to the severity of the

condition. These particular genes were not only implicated through RNA-seq data but

were also corroborated by proteomics datasets. The correlation was such that the

expression patterns of these genes seemed to escalate or diminish in tandem with

increasing NAFLD severity. We investigated how these genes contribute to NAFLD

pathogenesis, and why their particular expression patterns are so closely associated

with the progression of NAFLD.

Delving into the intricate cellular milieu, the expression profiles of these potential

biomarkers were meticulously examined across diverse liver cell categories. It was

observed that the HSC/MFB cellular contingent distinctly exhibited pronounced

expression levels of C7, COL6A3, FBN1, and QSOX1 compared to other cellular

phenotypes. Intriguingly, these molecular markers were conspicuously prevalent in

NAFLD-afflicted cohorts over their healthy counterparts (**Figure 2C**). In light of the

overwhelming majority of hepatocytes in hepatic architecture, significant expression

of IL1RAP, ITIH2, TTR, and SERPINA10 was ascertained in these cells, spanning all

identified subtypes. This expression spectrum was notably skewed towards the

healthy clusters versus NAFLD groups, intimating a palpable downregulation at the

holistic hepatic expression matrix (**Figure 2D**).

Cutting-edge developments in medical science have clarified the primary source of myofibroblasts, identifying hepatic stellate cells (HSCs) as their main origin (24, 25). Functioning as lipid storage entities, HSCs share considerable similarity with adipocytes, or fat cells, particularly when the liver is functioning optimally (26). This newfound insight broadens our understanding of liver diseases, including liver fibrosis and cirrhosis, where myofibroblasts play a significant role. Interestingly, the elevated levels of QSOX1, which is enriched in the HSC/MFB, correlate with both the NAS score and fibrosis score, offering a potential biomarker for NAFLD severity.

A more detailed exploration of the HSC/MFB clustering, and annotation using canonical markers has revealed two distinct subsets: hepatic stellate cells and myofibroblasts (**Figure 2E, F; Supplementary Table 3**). Within this context, QSOX1 manifested heightened expression within the NAFLD subset of HSC (**Figure 2G**). In contrast, IL1RAP and CFHR3, bearing the torch for the loftiest expression within hepatocytes amongst all subtypes, evinced diminished expression within the NAFLD demographic compared to its healthier cohort (illustrated in **Figure 2H**).

In a supplementary validation, liver biopsies sourced from patients with varying NAFLD severity were subjected to IHC in our antecedent study (16). Conclusive findings reinforced that QSOX1 resonates with heightened expression specifically in HSC/MFB cells of severe NAFLD patients (**Figure 2I**), while IL1RAP's expression was attenuated in hepatocytes of these patients as seen in **Figure 2J** (16).

**The Effectiveness of Integration and Normalization on PBMC Transcriptome Profiles**
To explore the expression of biomarkers in PBMC between NAFLD patients and healthy controls, we tracked their expression variations using RPCA integrated scRNA-seq data. The results from PCA showed that normalization successfully mitigated discernible batch effects compared to the unintegrated data, as shown in **Figure 3A and B**. The primary factors driving sample differentiation were not the health and NAFLD group, as depicted in **Figure 3C**. Instead, the cell type surfaced as

the primary determinant impacting transcriptome profiles, as illustrated in **Figure 3D**.

**Unsupervised Clustering Identifies Clusters of PBMC Cells**

We first clustered all the 122,238 PBMC cells into 6 cell types, and the identified cell types were annotated using signatures of known canonical markers (**Figure 3E, F; Supplementary Table 3**).

**Tracing the Cellular Origins of Promising Biomarkers for NAFLD severity in PBMC**

We identified 16 genes that are up-regulated (A2M, C7, COL6A3, COLEC11, ENPP2, FBLN1, FBN1, FCGBP, IGFBP6, LCN2, LUM, MMP2, PAPLN, PTGDS, QSOX1, VWF) and 22 that are down-regulated (AZGP1, C1RL, C4BPA, C6, C8B, CFHR3, CNDP1, F2, GC, HP, HPR, IL1RAP, ITIH1, ITIH2, ITIH4, KLKB1, PON3, SERPINA10, SERPINC1, SERPING1, SMPDL3A, TTR) with increasing severity of NAFLD, as evidenced by both RNA-seq and proteomics data. In PBMC, QSOX1 and IL1RAP have the highest expression levels in Natural killer cells (NK cells) across all subtypes. Interestingly, healthy individuals showed higher expression of these genes than those with NAFLD (**Figure 3G, H**). Nonetheless, the baseline expression levels of these genes in PBMC were too faint to support a significant comparison. The low expression within PBMC may not affect their detectability and proportional concentrations in plasma. Even when a gene exhibits varied expression in a distinct cell type or tissue, it's not a given that there will be discernible differences in broader systems like plasma, especially when the initial expression is inherently low (**Figure 3I, J**).

**The intercellular crosstalk in the liver and PBMC of NAFLD patients and healthy controls revealed by integrated sc- and snRNAseq data analysis**

Since our candidate biomarkers were derived from secreted genes, it was essential to delve deeper into the communication mechanisms of these genes. We utilized the 'CellChat' R package to analyze the ligand-receptor interactions, enabling us to better understand the crosstalk of secreted signaling and their potential implications in the biological processes of NAFLD.

The integrated RPCA analysis combined a total of 427,787 liver cells and 122,238 peripheral blood mononuclear cells (PBMCs). The data was then partitioned into subsets comprising 5,000 cells for each cell type within the respective tissues. Subsequent analysis was carried out using the 'CellChat' R package (**Figure 4A**). The examination of ligand-receptor pairs of secreted signaling, which constitutes 61.8% of all categories as depicted in **Figure 4B**, within the transcriptomic data from the liver and PBMCs in the NAFLD group indicates a strong correlation. Specifically, the interaction between IL1B-(IL1R1+IL1RAP) in liver monocytes and hepatocyte/cholangiocyte may provide a convincing rationale for the correlation between NAFLD severity and the down-regulated expression of IL1RAP in the liver, as highlighted in **Figure 4C and F**. In our earlier research (16), we pinpointed potential diagnostic markers from candidate genes by juxtaposing both up-regulated and down-regulated gene clusters against the secretome database from the Human Protein Atlas. Through this cross analysis, we discovered 349 genes that encode secreted proteins: 249 of these genes were up-regulated, while 100 were down-regulated. Of the totally 349 secreted genes, both FGFR2 and JAM3 exhibited up-regulation, while ERBB3 and SDC4 were notably down-regulated. These four genes are cataloged within the ligand-receptor pairs in the CellChat database, as referenced in **Supplementary Table 4**. Yet, when examining the crosstalk analysis derived from the sc- and snRNAseq data of NAFLD patients, only the down-regulated SDC4 (ANGPTL4-SDC4) and ERBB3 (NRG1-ERBB3) emerged as significant receptors in these pairs (see **Supplementary Tables 4 and 5**). The ANGPTL4-SDC4 pairing occurs between liver hepatocytes and hepatocyte/cholangiocyte, as depicted in **Figures 4D and 4G**. Similarly, the NRG1-ERBB3 interaction is evidenced between liver hepatocytes and cholangiocytes, highlighted in **Figures 4E and 4G**. An accompanying violin plot reveals expression patterns of the three crucial down-regulated secreted genes from the crosstalk analysis. Intriguingly, Most of them are predominantly found in hepatocytes, as demonstrated in **Figure 4H**. It may intricate that the significant down regulation of liver gene expression in bulk RNA sequencing may

mostly caused by the gene variation in hepatocyts since it is the biggest population in

the liver. And the down-regulation may caused by the ligands acted on the receptors

in the hepatocytes when the NAFLD severity developed.

**Discussion**

The inherent complexities of NAFLD pathogenesis consistently propel scientists to

extend the boundaries of traditional diagnostic and investigative methodologies,

thus underscoring the escalating importance of genomics technology within this

domain. Our research RPCA integrated publicly available scRNA-seq and snRNA-seq

datasets to identify potential biomarkers, thereby contributing to the enhanced

understanding of the intricate pathogenic processes driving NAFLD (27). Using this

approach, we corroborated the significant correlation between the expression ratio

of QSOX1/IL1RAP and the severity of NAFLD at the cellular level (16). Furthermore,

through an analysis of the integrated large-scale scRNA and snRNA sequencing data

to interpret potential biomarkers, we shed light on previously undiscovered facets of

NAFLD pathogenesis and intercellular crosstalk.

QSOX1 and its less common variant, QSOX2, are unique among sulfhydryl oxidases

because they are specifically found in the Golgi apparatus of animal cells (28).

Studies found elevated expression of QSOX1 in quiescent fibroblasts, and it is also

released into the bloodstream (29, 30). QSOX1 plays a crucial role in the proper

construction of the ECM, particularly elements similar to the basement membrane

(BM) in cell culture. The BM, an ECM layer interfacing bodily cavities or blood vessels

with underlying stromal fibroblasts, is largely composed of elements generated by

fibroblasts (31). The exact reason behind QSOX1's secretion—whether due to

overwhelming the Golgi's retention mechanism or some specialized secretion

pathway—remains unclear. Our findings highlight an increased expression of QSOX1

in fibroblasts within NAFLD-affected liver tissue, suggesting that Golgi function in

fibroblasts could be a relevant focus for NAFLD research (32).

Fibrogenesis is a key driver in NAFLD progression and is chiefly managed by specialized cells called myofibroblasts (33, 34). While these cells are not present in a healthy liver, they can quickly develop following liver damage. Emerging research has established that these myofibroblasts predominantly originate from hepatic stellate cells (HSCs) (24, 25) , which act mainly as lipid storage cells and bear resemblance to fat cells, or adipocytes, under healthy conditions (26).

In our previous article (16), the GO analysis revealed significant enrichment of the up-regulated genes involved in the fibrosis-related process, such as ECM organization. Similar to QSOX1 in terms of expression pattern and location, COL6A3 is associated with liver fibrosis and also plays a role in the ECM organization (35). Increased production of ECM moieties by diverse liver cell types is tightly linked with hepatic fibrosis. Research has demonstrated that adipocytes are affected by the mechanics of their surrounding ECM, such as stiffness and architecture, independent of initial ligand density. In response to a stiffened, modified ECM structure, adipocytes enhance the expression of fibrotic genes and increase ECM deposition. This reaction eventually results in a dysfunctional, fibrotic state (36, 37). This represents a significant research trajectory in the ongoing investigation of NAFLD pathophysiology.

IL1RAP is a crucial co-receptor expressed in numerous tissues and cell types, emphasizing its pivotal role in immune responses, cancers, and infectious diseases (38). The IL1RAP serves as a subunit to the type I IL-1 receptor (IL-1RI), a specific plasma membrane receptor, which facilitates the transmission of numerous biological effects of the pro-inflammatory cytokine IL-1 on an array of target cells (39, 40). Research indicates that one of the isoforms of IL1RAP, known as the soluble isoform of the IL-1 receptor accessory protein (sIL1RAP), has been found to have lower levels in the plasma of obese individuals when compared to their non-obese subjects (17). IL1RAP, enriched in the cytoplasm of liver cells and released into the bloodstream, shows variations in expression levels that correlate with the severity of

NAFLD, as per our study. Nevertheless, its specific role in NAFLD progression remains

an enigma.

CFHR3, displaying similar expression patterns and localization as IL1RAP, also

warrants further investigation for its potential role in diagnosing NAFLD (30). The

complementary DNAs (cDNAs) that encode CFHR3 were extracted from a human

liver cDNA library. These have been located within the 'regulators of complement

activation' (RCA) gene cluster situated on human chromosome 1 (41, 42). CFHR3 has

been linked with some diseases, including hepatocellular carcinoma (HCC), although

the underlying mechanism is still not clearly understood (43). Our research indicates

that CFHR3 expression decreases in correlation with the severity of NAFLD, including

both the NAS and fibrosis scores. This finding opens a new avenue of research into the

pathophysiology of NAFLD.

The IL1 pathway, especially IL1B, has attracted significant attention and has been

extensively researched within the realm of NAFLD (35, 44-47). IL-1β exacerbates liver

steatosis, inflammation, and fibrosis by interacting with the IL-1 receptor, which is

extensively expressed across various liver cell types (44). In 2021, Unamuno et al.

discovered that in the liver of obese patients with NASH, there was a significant

upregulation of mRNA levels for NLRP3 ($P < 0.05$) and its associated proinflammatory

mediator IL1B ($P < 0.01$) (35). In 2018, Pan and colleagues showed that in NASH

patients, IL-1β is primarily secreted by Kupffer cells, which are a type of monocyte in the

liver. This secretion is driven by the activation of the NLRP3 inflammasome, stimulated by

mitochondrial DNA (48). From our studies on cellular interactions within NAFLD, we have

formed and to some extent validated a hypothesis that the IL-1β, when secreted, serves

as a ligand affecting the receptors IL1R1 and IL1RAP in both hepatocytes and

cholangiocytes. This interaction between the ligand and its receptors seems to play a

crucial role in modulating the expression of IL1RAP, which is closely associated with

NAFLD severity. The dynamic interplay between these cellular components provides a

deeper understanding of the disease's underlying mechanisms and offers potential

avenues for NAFLD biomarkers and therapeutic interventions.

The utility of Seurat v5 in processing and integrating scRNA-seq and snRNA-seq data is

apparent in our study. However, we must also acknowledge the inherent limitations of

these techniques (49). Therefore, our findings must be interpreted with caution, and

further confirmatory studies are required to validate the candidate biomarkers and the

NAFLD pathogenesis.

Utilizing the 'CellChat' R package allowed for a streamlined analysis of the data. Given the

vastness of the data, it was segmented into more digestible portions, sketching them into

clusters of 5,000 cells each, ensuring a more organized scrutiny. An outstanding

revelation from this analysis was the significant presence of ligand-receptor pairs of

secreted signaling within the NAFLD group, which constituted a remarkable 61.8% of all

categories. This prevalence underscored their crucial role in the interactions between liver

and PBMC cells. The interaction of IL1B-(IL1R1+IL1RAP) between liver monocytes and

hepatocyte/cholangiocyte was of particular interest, suggesting a plausible explanation for

the detected decline in IL1RAP expression in the liver - a potential indicator of NAFLD

severity. Referring back to our prior research, the crosstalk analysis, built on sc- and

snRNAseq data from NAFLD patients, spotlighted SDC4 (in association with ANGPTL4)

and ERBB3 (aligned with NRG1) as receptors of importance. The relationships between

ANGPTL4-SDC4 and NRG1-ERBB3 with liver hepatocytes and

hepatocyte/cholangiocytes, respectively, reflect their likely significance in the liver's

cellular dynamics. This proposition is bolstered by a violin plot that vividly illustrates the

expression patterns of three crucial down-regulated secreted genes from the crosstalk

analysis, with hepatocytes prominently featuring as their central hub.

Syndecan-4 (SDC4) is a transmembrane protein that is glycosylated and prevalent in

numerous tissues (50). Essential cellular functions such as differentiation, adhesion,

migration, and prognosis are facilitated by SDC4, given its ability to bind to a variety of

external molecules and engage in signal transmission (51-53).

ERBB3, or Receptor tyrosine-protein kinase erbB-3, is a membrane-associated protein

from the EGFR/ERBB family of receptor tyrosine kinases. Research has shown its

involvement in numerous liver conditions, including liver fibrosis (54, 55).

In conclusion, our research sheds light on the intricacies of NAFLD pathogenesis and

highlights the significance of newly identified biomarkers in tracking disease progression.

These insights serve as a foundation for subsequent studies delving into the molecular

intricacies of NAFLD, all aiming towards enhancing disease diagnosis, management, and

the outcomes of patients.

**Abbreviations**

avg_log2FC, log fold-change of the average expression between the two groups; BM,

basement membrane; cDC1, type 1 conventional dendritic cell; cDNAs,

complementary DNAs; ECM, extracellular matrix; F, Fibrosis score; GEO, Gene

Expression Omnibus; GO, Gene Ontology; GRCh37, Genome Reference Consortium

Human Build 37; HCC, hepatocellular carcinoma; HSCs, hepatic stellate cells; IHC,

immunohistochemistry staining; IL-1RI, type I IL-1 receptor; IL1RAP, Interleukin-1

receptor accessory protein; LECs, lymphatic endothelial cells; N, NAS score; NAFL,

Non-alcoholic Fatty Liver; NAFLD, Non-alcoholic fatty liver disease; NAFLD-DB, NAFLD

gene expression database; NAS, NAFLD activity scores; NASH, Non-alcoholic

Steatohepatitis; NK cells, Natural killer cells; PBMC, Peripheral blood mononuclear

cell; PCA, Principal components analysis; PCs, Principal Components; QSOX1,

Quiescin sulfhydryl oxidase 1; RCA, regulators of complement activation; RNA-seq,

RNA sequencing; RPCA, reciprocal principal components analysis; scRNA-seq,

single-cell RNA sequencing; sIL1RAP, the soluble isoform of the IL-1 receptor

accessory protein; SNN, shared nearest neighbor; snRNA-seq, Single-nucleus RNA

sequencing; SZTCMH, Shenzhen Traditional Chinese Medicine Hospital, China; UMAP,

Uniform Manifold Approximation and Projection.

**Financial support**

This research was funded by the Shenzhen Sanming Project of Medicine in Shenzhen, China (grant nos. SZSM201612074).

**Authors' contributions**

WFM and LL designed the study and interpreted the data. The analysis strategy has been developed by LL and WFM. WFM and LL collected, analyzed, and interpreted the data. WFM drafted the manuscript. YLL, HG, and MBK revised the manuscript. WFM and LL performed data analysis and interpretation. BQC, JRH, MLL, XZ, SMX, BLZ, QL, QH, MQM contributed to the validation of the candidate biomarkers. Technical support: LL and JRH. All authors reviewed and approved the final version of the manuscript.

**Data and code availability statement**

The data that support the findings of this study are available from the corresponding author upon reasonable request.

**Conflict of interest**

Henning Grønbæk has received research grants from Abbvie, Intercept, ARLA Food for Health, ADS AIPHIA Development Services AG. Consulting Fees from Ipsen, NOVO, Pfizer. Lecturer for AstraZeneca and EISAI; and on Data Monitoring Committee at CAMURUS AB. All other authors have no conflicts of interest to declare.

**Figure Legend**

**Figure 1. Liver single-cell and nucleus RNA sequencing data integration**
A. Illustration of data processing.
B.Uniform Manifold Approximation and Projection (UMAP) based on the unintegrated data.
C. UMAP based on the anchor-based reciprocal principal components analysis (RPCA) integrated data.
D.UMAP based on the health and NAFLD groups after RPCA integration.
E. UMAP based on the library type after RPCA integration.
F. UMAP based on the input cell type after RPCA integration.

**Figure 2. Integrated liver transcriptome data annotation, analysis and verification**
A. Gene markers of liver cell types.
B. Total 12 distinct cell types were clustered according to the canonical markers.
C. Dot plot of 16 up-regulated secreting marker expressions in cell clusters.
D. Dot plot of 22 down-regulated secreting marker expressions in cell clusters.
E. Gene markers of HSC (hepatic stellate cell)/MFB (myofibroblast) cell types.
F. Total 2 distinct cell types were clustered according to the canonical markers.
G. Violin plot of QSOX1 gene expressions in the hepatic stellate cell and myofibroblast.
H. Violin plot of IL1RAP, CFHR3 gene expressions in the Hepatocytes.
I.Representative immunohistochemical staining (IHC) images of QSOX1 in liver biopsies from mild NAFLD patients (N0-4, F0-2) and severe NAFLD patients (N5-8, F3-4) in our previous research (16).
J. Representative IHC images of IL1RAP in liver biopsies from mild NAFLD patients (N0-4, F0-2) and severe NAFLD patients (N5-8, F3-4) in our previous research (16). Statistical testing was performed using the Wilcoxon rank sum test, with p-values shown in the plot.

**Figure 3. Tracing the Cellular Origins in Liver and PBMC Data Integration**
A. Uniform Manifold Approximation and Projection (UMAP) based on the unintegrated data.Dot plot of 16 up-regulated secreting marker expressions in cell clusters.
B. UMAP based on the anchor-based reciprocal principal components analysis (RPCA) integrated data.
C. UMAP based on the health and NAFLD groups after RPCA integration.
D. UMAP based on the input cell type after RPCA integration.
E. Gene markers of PBMC cell types.
F. Total 6 distinct cell types were clustered according to the canonical markers.
G. Dot plot of 16 up-regulated secreting marker expressions in cell clusters.
H. Dot plot of 22 down-regulated secreting marker expressions in cell clusters.
I. Violin plot of QSOX1 gene expressions in the NK cell.

J. Violin plot of IL1RAP gene expressions in the NK cell.
Statistical testing was performed using the Wilcoxon rank sum test, with p-values shown in the plot.

**Figure 4. The cell communication in the liver and PBMC transcriptomics data**
A. UMAP based on the anchor-based reciprocal principal components analysis (RPCA) integrated liver and PBMC data.
B. Ligand-receptor pairs of secreted signaling, which constitutes 61.8% of all categories.
C. The interaction between IL1B-(IL1R1+IL1RAP) in liver monocytes and hepatocyte/cholangiocyte.
D. The ANGPTL4-SDC4 pairing occurs between liver hepatocytes and hepatocyte/cholangiocyte.
E. The NRG1-ERBB3 interaction is evidenced between liver hepatocytes and cholangiocytes
F. The interaction between IL1B-(IL1R1+IL1RAP) in liver monocytes and hepatocyte/cholangiocyte.
G. The ANGPTL4-SDC4 pairing occurs between liver hepatocytes and hepatocyte/cholangiocyte.
H. Violin plot of IL1B, IL1R1, IL1RAP, ANGPTL4, SDC4, NRG1, ERBB3 gene expressions in the liver and PBMC.

## List of Supplementary Tables

https://drive.google.com/file/d/1Eji-PHKMxil246MClmnh3PZPICD-nlCA/view?usp=sharing

## Supplementary Table 1

The scRNAseq and snRNA-seq data of liver and PBMC samples.

## Supplementary Table 2

The detailed information regarding the data alignment of the scRNAseq and snRNA-seq data.

## Supplementary Table 3

The canonical markers of cell types in the liver and PBMC.

## Supplementary Table 4

The ligand-receptor pairs in the CellChat database.

## Supplementary Table 5

The pathways in the CellChat database.

## Reference

1.    Loomba R, Friedman SL, Shulman GI. Mechanisms and disease consequences of nonalcoholic fatty liver disease. Cell 2021;184:2537-2564.

2.    Powell EE, Wong VW, Rinella M. Non-alcoholic fatty liver disease. Lancet 2021;397:2212-2224.

3.    Vogel A, Meyer T, Sapisochin G, Salem R, Saborowski A. Hepatocellular carcinoma. Lancet 2022;400:1345-1362.

4.    Chen Y, Wang W, Morgan MP, Robson T, Annett S. Obesity, non-alcoholic fatty liver disease and hepatocellular carcinoma: current status and therapeutic targets. Front Endocrinol (Lausanne) 2023;14:1148934.

5.    Younossi ZM, Golabi P, Paik JM, Henry A, Van Dongen C, Henry L. The global epidemiology of nonalcoholic fatty liver disease (NAFLD) and nonalcoholic steatohepatitis (NASH): a systematic review. Hepatology 2023;77:1335-1347.

6.    Khairnar R, Islam MA, Fleishman J, Kumar S. Shedding light on non-alcoholic fatty liver disease: Pathogenesis, molecular mechanisms, models, and emerging therapeutics. Life Sci 2023;312:121185.

7.    Musso G, Saba F, Cassader M, Gambino R. Lipidomics in pathogenesis, progression and treatment of nonalcoholic steatohepatitis (NASH): Recent advances. Prog Lipid Res 2023;91:101238.

8.    Stark R, Grzelak M, Hadfield J. RNA sequencing: the teenage years. Nat Rev Genet 2019;20:631-656.

9.    Andrews TS, Atif J, Liu JC, Perciani CT, Ma XZ, Thoeni C, Slyper M, et al. Single-Cell, Single-Nucleus, and Spatial RNA Sequencing of the Human Liver Identifies Cholangiocyte and Mesenchymal Heterogeneity. Hepatol Commun 2022;6:821-840.

10.    Alvarez M, Benhammou JN, Darci-Maher N, French SW, Han SB, Sinsheimer JS, Agopian VG, et al. Human liver single nucleus and single cell RNA sequencing identify a hepatocellular carcinoma-associated cell-type affecting survival. Genome Med 2022;14:50.

11.    Xiong X, Kuang H, Ansari S, Liu T, Gong J, Wang S, Zhao XY, et al. Landscape of Intercellular Crosstalk in Healthy and NASH Liver Revealed by Single-Cell Secretome Gene Analysis. Mol Cell 2019;75:644-660 e645.

12.    Tamburini BAJ, Finlon JM, Gillen AE, Kriss MS, Riemondy KA, Fu R, Schuyler RP, et al. Chronic Liver Disease in Humans Causes Expansion and Differentiation of Liver Lymphatic Endothelial Cells. Front Immunol 2019;10:1036.

13.    Ramachandran P, Dobie R, Wilson-Kanamori JR, Dora EF, Henderson BEP, Luu NT, Portman JR, et al. Resolving the fibrotic niche of human liver cirrhosis at single-cell level. Nature 2019;575:512-518.

14.    Deczkowska A, David E, Ramadori P, Pfister D, Safran M, Li B, Giladi A, et al. XCR1(+) type 1 conventional dendritic cells drive liver pathology in non-alcoholic steatohepatitis. Nat Med 2021;27:1043-1054.

15.    Wang S, Li K, Pickholz E, Dobie R, Matchett KP, Henderson NC, Carrico C, et al. An autocrine signaling circuit in hepatic stellate cells underlies advanced fibrosis in nonalcoholic steatohepatitis. Sci Transl Med 2023;15:eadd3949.

16.    Ma W, Huang J, Cai B, Shao M, Yu X, Kjær MB, Lv M, et al. A novel gene-screening approach reveals QSOX1/IL1RAP as promising biomarkers for the severity of non-alcoholic fatty liver disease. medRxiv 2023:2023.2007.2026.23293038.

17.   Bozaoglu K, Attard C, Kulkarni H, Cummings N, Diego VP, Carless MA, Shields KA, et al. Plasma
levels of soluble interleukin 1 receptor accessory protein are reduced in obesity. J Clin Endocrinol
Metab 2014;99:3435-3443.

18.   Gao R, Wang J, He X, Wang T, Zhou L, Ren Z, Yang J, et al. Comprehensive analysis of endoplasmic
reticulum-related and secretome gene expression profiles in the progression of non-alcoholic fatty
liver disease. Front Endocrinol (Lausanne) 2022;13:967016.

19.   Hao Y, Stuart T, Kowalski MH, Choudhary S, Hoffman P, Hartman A, Srivastava A, et al. Dictionary
learning for integrative, multimodal and scalable single-cell analysis. Nat Biotechnol 2023.

20.   Hao Y, Hao S, Andersen-Nissen E, Mauck WM, 3rd, Zheng S, Butler A, Lee MJ, et al. Integrated
analysis of multimodal single-cell data. Cell 2021;184:3573-3587 e3529.

21.   Jin S, Guerrero-Juarez CF, Zhang L, Chang I, Ramos R, Kuan CH, Myung P, et al. Inference and
analysis of cell-cell communication using CellChat. Nat Commun 2021;12:1088.

22.   Fang Z, Tian Y, Sui C, Guo Y, Hu X, Lai Y, Liao Z, et al. Single-Cell Transcriptomics of Proliferative
Phase Endometrium: Systems Analysis of Cell-Cell Communication Network Using CellChat. Front Cell
Dev Biol 2022;10:919731.

23.   Slyper M, Porter CBM, Ashenberg O, Waldman J, Drokhlyansky E, Wakiro I, Smillie C, et al. A
single-cell and single-nucleus RNA-Seq toolbox for fresh and frozen human tumors. Nat Med
2020;26:792-802.

24.   Yang W, He H, Wang T, Su N, Zhang F, Jiang K, Zhu J, et al. Single-Cell Transcriptomic Analysis
Reveals a Hepatic Stellate Cell-Activation Roadmap and Myofibroblast Origin During Liver Fibrosis in
Mice. Hepatology 2021;74:2774-2790.

25.   Rosenthal SB, Liu X, Ganguly S, Dhar D, Pasillas MP, Ricciardelli E, Li RZ, et al. Heterogeneity of
HSCs in a Mouse Model of NASH. Hepatology 2021;74:667-685.

26.   She H, Xiong S, Hazra S, Tsukamoto H. Adipogenic transcriptional regulation of hepatic stellate
cells. J Biol Chem 2005;280:4959-4967.

27.   Hundertmark J, Berger H, Tacke F. Single Cell RNA Sequencing in NASH. Methods Mol Biol
2022;2455:181-202.

28.   Reznik N, Fass D. Disulfide bond formation and redox regulation in the Golgi apparatus. FEBS Lett
2022;596:2859-2872.

29.   Coppock DL, Kopman C, Scandalis S, Gilleran S. Preferential gene expression in quiescent human
lung fibroblasts. Cell Growth Differ 1993;4:483-493.

30.   Uhlen M, Karlsson MJ, Hober A, Svensson AS, Scheffel J, Kotol D, Zhong W, et al. The human
secretome. Sci Signal 2019;12.

31.   Grossman I, Alon A, Ilani T, Fass D. An inhibitory antibody blocks the first step in the
dithiol/disulfide relay mechanism of the enzyme QSOX1. J Mol Biol 2013;425:4366-4378.

32.   Ilani T, Reznik N, Yeshaya N, Feldman T, Vilela P, Lansky Z, Javitt G, et al. The disulfide catalyst
QSOX1 maintains the colon mucosal barrier by regulating Golgi glycosyltransferases. EMBO J
2023;42:e111869.

33.   Kisseleva T, Brenner D. Molecular and cellular mechanisms of liver fibrosis and its regression. Nat
Rev Gastroenterol Hepatol 2021;18:151-166.

34.   Kisseleva T. The origin of fibrogenic myofibroblasts in fibrotic liver. Hepatology
2017;65:1039-1043.

35. Unamuno X, Gomez-Ambrosi J, Ramirez B, Rodriguez A, Becerril S, Valenti V, Moncada R, et al.
NLRP3 inflammasome blockade reduces adipose tissue inflammation and extracellular matrix
remodeling. Cell Mol Immunol 2021;18:1045-1057.
36. Kim M, Lee C, Seo DY, Lee H, Horton JD, Park J, Scherer PE. The impact of endotrophin on the
progression of chronic liver disease. Exp Mol Med 2020;52:1766-1776.
37. Di Caprio N, Bellas E. Collagen Stiffness and Architecture Regulate Fibrotic Gene Expression in
Engineered Adipose Tissue. Adv Biosyst 2020;4:e1900286.
38. Frenay J, Bellaye PS, Oudot A, Helbling A, Petitot C, Ferrand C, Collin B, et al. IL-1RAP, a Key
Therapeutic Target in Cancer. Int J Mol Sci 2022;23.
39. Wesche H, Korherr C, Kracht M, Falk W, Resch K, Martin MU. The interleukin-1 receptor accessory
protein (IL-1RAcP) is essential for IL-1-induced activation of interleukin-1 receptor-associated kinase
(IRAK) and stress-activated protein kinases (SAP kinases). J Biol Chem 1997;272:7727-7731.
40. Greenfeder SA, Nunes P, Kwee L, Labow M, Chizzonite RA, Ju G. Molecular cloning and
characterization of a second subunit of the interleukin 1 receptor complex. J Biol Chem
1995;270:13757-13765.
41. Diaz-Guillen MA, Rodriguez de Cordoba S, Heine-Suner D. A radiation hybrid map of complement
factor H and factor H-related genes. Immunogenetics 1999;49:549-552.
42. Jozsi M, Meri S. Factor H-related proteins. Methods Mol Biol 2014;1100:225-236.
43. Wan Z, Li X, Luo X, Wang B, Zhou X, Chen A. The miR-590-3p/CFHR3/STAT3 signaling pathway
promotes cell proliferation and metastasis in hepatocellular carcinoma. Aging (Albany NY)
2022;14:5783-5799.
44. Mirea AM, Tack CJ, Chavakis T, Joosten LAB, Toonen EJM. IL-1 Family Cytokine Pathways
Underlying NAFLD: Towards New Treatment Strategies. Trends Mol Med 2018;24:458-471.
45. Tilg H, Effenberger M, Adolph TE. A role for IL-1 inhibitors in the treatment of non-alcoholic fatty
liver disease (NAFLD)? Expert Opin Investig Drugs 2020;29:103-106.
46. Gehrke N, Hofmann LJ, Straub BK, Ruhle F, Waisman A, Galle PR, Schattenberg JM. Hepatic
interleukin-1 receptor type 1 signalling regulates insulin sensitivity in the early phases of nonalcoholic
fatty liver disease. Clin Transl Med 2022;12:e1048.
47. He Y, Hwang S, Ahmed YA, Feng D, Li N, Ribeiro M, Lafdil F, et al. Immunopathobiology and
therapeutic targets related to cytokines in liver diseases. Cell Mol Immunol 2021;18:18-37.
48. Pan J, Ou Z, Cai C, Li P, Gong J, Ruan XZ, He K. Fatty acid activates NLRP3 inflammasomes in
mouse Kupffer cells through mitochondrial DNA release. Cell Immunol 2018;332:111-120.
49. Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. Integrating single-cell transcriptomic data
across different conditions, technologies, and species. Nat Biotechnol 2018;36:411-420.
50. Tanino Y, Wang X, Nikaido T, Misa K, Sato Y, Togawa R, Kawamata T, et al. Syndecan-4 Inhibits the
Development of Pulmonary Fibrosis by Attenuating TGF-beta Signaling. Int J Mol Sci 2019;20.
51. Becsky D, Gyulai-Nagy S, Balind A, Horvath P, Dux L, Keller-Pinter A. Myoblast Migration and
Directional Persistence Affected by Syndecan-4-Mediated Tiam-1 Expression and Distribution. Int J
Mol Sci 2020;21.
52. Chronopoulos A, Thorpe SD, Cortes E, Lachowski D, Rice AJ, Mykuliak VV, Rog T, et al. Syndecan-4
tunes cell mechanics by activating the kindlin-integrin-RhoA pathway. Nat Mater 2020;19:669-678.
53. Lambert J, Makin K, Akbareian S, Johnson R, Alghamdi AAA, Robinson SD, Edwards DR.
ADAMTS-1 and syndecan-4 intersect in the regulation of cell migration and angiogenesis. J Cell Sci
2020;133.

54.   Scheving LA, Zhang X, Threadgill DW, Russell WE. Hepatocyte ERBB3 and EGFR are required for

maximal CCl4-induced liver fibrosis. Am J Physiol Gastrointest Liver Physiol 2016;311:G807-G816.

55.   Xiong J, Ni J, Chen C, Wang K. miR□148a□3p regulates alcoholic liver fibrosis through targeting

ERBB3. Int J Mol Med 2020;46:1003-1012.

**Healthy control**

**NAFLD spectrum**

**Biomarkers**

18 samples (14 healthy and 4 NAFLD)

**PBMC**

64 samples (32 healthy and 32 NAFLD)

**Liver biopsy**

12 samples (6 mild and 6 severe NAFLD)

**Single cell RNA sequencing**

**IHC**

Created with BioRender.com

**Figure1**

# Figure 2

# Figure 3

# Figure 4