Platelet activation is associated with acute kidney injury after cardiac surgery

Naomi Brown[1,*], Nikol Sullo[1,3*], Nathan Tyson[2], Bryony Eagle-Hemming[1], Florence Y. Lai[1], Sophia Sheikh[1], Kristina Tomkova[1], Lathishia Joel-David[1], Tracy Kumar[1], Hardeep Aujla[1], Mustafa Zakkar[1], Alison H Goodall[1], Gavin J Murphy[1], Marcin J Woźniak[1$]

[1] Department of Cardiovascular Sciences and NIHR Cardiovascular Biomedical Research Unit, University of Leicester, Glenfield Hospital, Leicester, LE3 9QP, UK.

[2] NHS, Glenfield Hospital, Leicester, LE3 9QP, UK.

[*] These authors contributed equally

[3] Current address: Medical School, Swansea University, Swansea, UK

[$] **Address for correspondence:** Dr Marcin J Woźniak, Department of Cardiovascular Sciences, University of Leicester, Clinical Sciences Wing, Glenfield General Hospital, Leicester, LE3 9QP, UK. Tel: +44 116 258 3028, Email: mw299@leicester.ac.uk

Running title: Circulating extracellular vesicles and miR as biomarkers of AKI

Abstract

Background Post-cardiac surgery acute kidney injury (AKI) is common, with high rates of mortality and morbidity. Despite extensive research, there are no reliable biomarkers. We hypothesised that pro-inflammatory changes associated with platelet activation, including circulating extracellular vesicles (EV) and micro-RNA levels, could provide predictive and diagnostic markers for AKI in cardiac surgery patients.

Methods Plasma samples were collected from 95 patients recruited to the MaRACAS study before, immediately after and 6-12, 24 and 48 hours after surgery. EV size distribution and concentrations were measured using NanoSight. EV derivation and platelet and leukocyte activation were measured using flow cytometry. Platelet responsiveness was measured with Multiplate, and circulating soluble biomarkers were measured with a MAGPIX device. Micro-RNA assessment was performed with TaqMan arrays.

Results In the 57% (54/95) of patients who developed AKI post-surgery, the numbers of platelet-derived EVs were higher 24 hours after surgery. TaqMan arrays identified miR-668 downregulated before and miR-92a-1, -920, -518a-3p, -133b and -1262 upregulated after surgery in AKI patients. qRT-PCR confirmed these differences for miR-1262. In addition, in AKI patients, platelets were desensitised to ADP 6-12 hours after surgery, independent of the administration of aspirin or P2Y12 antagonists. AKI patients also had more activated platelets 6-12 hours after surgery, more circulating platelet-granulocyte aggregates before and 6-12 and 24 hours after surgery and higher levels of sICAM1 before and 48 hours after surgery.

Conclusions AKI is associated with increased platelet activation, suggesting that alternative platelet inhibition treatments may be renoprotective. Studies in larger cohorts are required to validate these findings.

Keywords: Surgical Procedure, Cardiovascular, Cell-Derived Microparticles, MicroRNAs, Acute Kidney Injury, Biomarkers

Introduction

Acute kidney injury (AKI) is a common complication of cardiac surgery that is associated with increased morbidity and mortality, [1] prolonged hospital stays and increases healthcare costs. [2] AKI is defined clinically by an acute decrease in glomerular filtration rate (GFR) as determined by changes in serum creatinine (SCr) concentrations and urine output. The mechanisms that underlie AKI are poorly understood, but inflammation and activation of the vascular endothelium and platelets are thought to contribute. [3] No individual biomarker has been established as the gold standard for assessing the risk or diagnosing of AKI in cardiac surgery.

Leukocytes and platelets play important roles in AKI pathophysiology, and their activation could serve as an AKI biomarker. Leukocytes contribute to the inflammatory response observed in AKI by releasing pro-inflammatory cytokines and reactive oxygen species. Platelet activation and aggregation within the renal microvasculature can lead to capillary occlusion, impairing blood flow and worsening tissue injury. [4] In inflammatory states, activated platelets bind monocytes and neutrophils through interaction of P-selectin with P-selectin glycoprotein ligand 1 (PSGL-1). Chemokines and cytokines released by platelets further modulate leukocyte activity, triggering neutrophil extracellular traps formation and enhancing the inflammatory response. [5,6]

Extracellular vesicles (EVs) are small, membrane-bound particles that are secreted by cells into the extracellular space. Two subtypes of EVs are microvesicles (MV) which bud from the plasma membrane and exosomes that are released by exocytosis. EVs are involved in the transfer of proteins and genetic information (mRNA and microRNA) specific to their cellular origin. Hence, changes in the concentration of EVs during pathological processes can both promote pathological changes as well as serve as an indicator for affected tissues or cells. [7]

Micro-RNA (miRNA) are non-coding and highly conserved short RNA oligonucleotides that cells actively secrete or passively release (e.g. dead cells). miRNAs are released within EVs or complexed with proteins (AGO-2 or NPM1 [8]) or high-density lipoproteins. [9,10] They can influence the post-transcriptional expression of target genes, and serve as a means of intracellular communication. [11] Several miRNAs have been associated with the pathogenesis of AKI, [12,13] and studies have shown miRNAs derived from plasma may be used as novel biomarkers for AKI in both adult [14] and paediatric [15] populations.

We hypothesised that pro-inflammatory changes associated with platelet activation, including circulating extracellular vesicles (EV) derived from different cell types and micro-RNA levels, could serve as novel biomarkers of post-surgery AKI. To test this hypothesis, we performed longitudinal measurements from baseline to 48 hours of platelet, leukocyte, and endothelial activation

simultaneously with assessments of EV and miRNA in plasma from adults undergoing cardiac surgery.

Methods

Study design and setting An Observational Case Control Study to Identify the Role of MV and MV Derived Micro-RNA in Post CArdiac Surgery AKI (MaRACAS) was a prospective, single-centre observational study approved by East Midlands - Leicester South Research Ethics Committee (REC reference 13/EM/0383) conducted between 11[th] December 2014 and 30[th] May 2017 in Glenfield General Hospital in Leicester, UK. The protocol was registered at https://clinicaltrials.gov/ct2/show/NCT02315183.

Study cohort included prospectively recruited 95 consecutive adult cardiac surgery patients (> 16 years old) undergoing coronary artery bypass grafting or valve surgery with moderately hypothermic cardiopulmonary bypass (32 – 34 ˚C) blood cardioplegia. All participants were identified as at increased risk for AKI using a modified risk score. [16] Exclusion criteria included: pre-existing inflammatory state (sepsis undergoing treatment, acute kidney injury within 5 days, chronic inflammatory disease or congestive heart failure), emergency or salvage procedure, ejection fraction <30%, critical preoperative state (Kidney Disease: Improving Global Outcomes (KDIGO) Stage 3 AKI or requiring ionotropes, ventilation or intra-aortic balloon pump), or pregnancy.

Blood sampling, storage and bias Blood was collected pre-operatively, on return to the intensive care unit, and 6–12, 24, 48, 72 and 96 hours post-operatively. The total volume collected from each patient was 104.8 ml collected in clotting, EDTA, citrate and hirudin tubes (S-Monovette, Sarstedt, Nuembrecht, Germany). All anticoagulated samples were processed within 2 hours after collection and plasma was frozen at -80 ˚C for subsequent analysis. Samples were identified by a trial acronym and the patient's study ID. Routine measurements, including serum creatinine, were performed in NHS laboratories by personnel who were unaware that the participant was in a trial. For all other analyses, laboratory staff were blinded to the AKI status of patients.

Soluble biomarker analysis Levels of soluble circulating biomarkers, CXCL1, IL-6, -8 and -10 and sICAM1 were measured by multiplex assay on the MAGPIX Analyser (Luminex, Netherlands).

EV analysis was performed in hirudinated plasma samples that had been centrifuged twice at 1,500 × g and stored at –80ºC. Concentration and size distribution were estimated using NanoSight NS500, nanoparticle tracking device (Malvern Instruments, Malvern, United Kingdom) in citrated plasma samples. The flow cytometry analysis of EVs was performed with fluorescently labelled annexin V or antibodies against CD235a, CD41, CD144, CD142, CD14, CD16 CD284 and CD3 (all ThermoFisher). Fluorescently labelled isotype control IgGs were used as controls. Plasma samples (20 μL) were labelled with antibodies at 1:20 dilution in annexin V binding buffer for 25 minutes at room temperature (RT) in a total volume of 100 μL. The samples were analysed by flow cytometry (Cyan ADP; Beckman Coulter, Brea, CA). EV-specific gates (0.1 - 1μm) were determined using

polystyrene particles (Spherotech, USA). Percentage of EVs positive for specific antibodies was adjusted for isotype controls and analysed as described in the statistical analysis section.

MicroRNA analysis RNA from EVs were isolated from circulating citrated plasma using exoRNeasy kit (Qiagen) using the manufacturer's instructions. RNA (100 ng) was reverse transcribed using the Taq-Man Multiplex RT set for TaqMan Array Human MicroRNA Panel v2.0 (Applied Biosystems, Foster City, USA) and Veriti thermal cyclers. Pre-amplified complementary DNA from each pool was analysed using TaqMan Array Human MicroRNA Card A and B v2.0 (Applied Biosystems). Quantitative polymerase chain reaction was carried out on an Applied BioSystems 7900HT thermocycler. The experiments were performed in triplicates. Any assays where the cycling threshold (Ct) value was lower than 20 were removed, and ΔCt (difference in cycling threshold) values were calculated (ΔCt = miRNA Ct−reference miRNA Ct) using reference control U6 snRNA-001973 present in duplicates on each miRNA array. Undetermined assays were assigned maximum cycling value. Data were normalised across arrays using quantile normalisation. Differential expression analysis was performed with *limma* R package. [17]

Selected differentially expressed miRNAs were analysed using rotor-gene (Qiagen) quantitative real-time PCR instruments in samples from all patients using TaqMan Advanced miRNA assays for hsa-miR-17, hsa-miR-223, hsa-miR-486, hsa-miR-668, hsa-miRs-92a-1, hsa-miRs-920, hsa-miRs-518a-3p, hsa-miRs-133b and hsa-miRs-1262 (ThermoFisher) according to the manufacturer's recommendations. The data was standardised against levels of hsa-miR-17 and hsa-miR-486, which showed constant average expression levels across all time-points and patient groups (**sFig. 1**).

Leukocyte and platelets activation were measured by flow cytometry (Cyan ADP, Beckman Coulter, Brea, CA) in citrated blood samples. For leukocyte analysis, red cells in 100 μL of whole blood were lysed with FACS Lysing Solution (BD Biosciences, Oxford, UK) for 10 min at RT. Leukocytes, gated in forward and side scatter, were labelled with FITC-coupled CD64, CD11b, CD16, CD14, and PE-coupled CD163 (Affymetrix, Santa Clara, CA) antibodies at 1:20 dilution for 25 min at room temperature (RT). For platelets 5 μL of whole blood was incubated at RT for 25 min with 1:20 dilutions of either FITC-coupled PAC-1 (BD Biosciences), or CD62P. (Abcam, Cambridge, UK) together with PE-coupled CD41 (Affymetrix) antibodies in phosphate-buffered saline in a total volume of 100 μL. Red blood cells and leukocytes were gated out in forward and side scatter.

Statistical analysis was performed using R Studio Version 4.3.1 [18] and plots prepared with ggplot R package. [19] Data normality was assessed using the Shapiro-Wilk test, with normally distributed variables expressed as mean (± standard deviation [SD]) and non-normally distributed variables expressed as median (interquartile range [IQR]). Analyses were performed between patients with AKI and without AKI. Comparisons of miRNA analysed with PCR-arrays and EV size were adjusted for

multiple comparisons using Benjamini-Hochberg method. Pre-specified outcomes were not adjusted for multiple comparisons. [20-22] Density plots were generated using *density* function, which calculates the distribution of a numeric variable using a kernel density estimates.

Results

Participants

Screening of 230 cardiac surgery patients identified 150 who were eligible for the study. Out of these, fifty-three declined consent and two became ineligible after giving consent. Ninety-five patients completed the study (**Fig. 1A**). Fifty-four of these patients (57%) developed post-cardiac surgery AKI, determined according to the Kidney Disease Improving Global Outcomes (KDIGO) definition, [23] the majority (n=45) exhibiting stage 1, with two patients developing stage 2, and 7 patients developing stage 3 AKI (**Fig. 1A**). Patient baseline demographics and intra-operative characteristics, shown in **Table 1**, were not different between the groups. The mean age was 72.8 (non-AKI) and 70.6 (AKI), with 85% in both groups being male.

The AKI patients had longer post-surgery ventilation times (AKI: 13, 10 -17 hours vs non-AKI 9, 7 - 14 hours, p=0.001) and longer hospital stays (AKI: 10, 7 -13 days vs 8, 6 -9 days, p=0.017). As expected, levels of serum creatinine were significantly higher in AKI patients 6 – 96 hours after surgery (**Fig. 1B, Table 1** and **eTable 1**).

Extracellular vesicles

EV concentrations ranged from 5.46×10^8 – 14.56×10^8 with no overall difference between AKI and non-AKI patients (**Fig. 2A, eTable 1**) or in particle sizes (**Fig. 2B**) at any time point. Derivation analysis included identification of EVs from monocytes (CD14), endothelial cells (CD144 and CD62E), granulocytes (CD16), red cells (CD235a), myeloid cells (CD284), T-cells (CD3) and platelets (CD41). We also assayed EVs positive for annexin V and tissue factor (CD142), which would indicate a pro-thombotic phenotype. The most abundant EVs were CD284-positive (8.77-15.88%) and annexin V-positive (8.198-13.44%), and the least abundant were EVs positive for CD41 (0.05-0.255%). Of these, only platelet-derived EVs (CD41-positive) were expressed at significantly higher levels in patients with AKI 24 hours after surgery (0.095%, 0.05-0.185 vs 0.05%, 0.02-0.18, p=0.028; **Fig. 2C** and **eTable 1**).

MicroRNAs

To identify AKI-related miRNA, 95 samples were pooled into 24 batches at each of three time points (pre, ICU and 24 hours after surgery) and analysed with qRT-PCR arrays for 754 human miRNA species. miR-668 was significantly lower in the AKI patients before surgery, while miRs-92a-1, 920, 518a-3p, 133b, and 1262 were higher in the samples from the AKI patients immediately after surgery (in ICU; **Fig, 2D** and **eTable 2**). Expression of these miRNAs was then verified by qRT-PCR in each of 95 samples at three time-points. Expression of miR-668 and miR-518a-3p were below detection level in most samples. Levels of miR-1262 were significantly higher immediately after surgery (ICU) in AKI patients, which was the result of lower expression levels in the non-AKI group. Levels of miR-133b

were significantly lower in AKI patients before surgery (**Fig. 2E** and **eTable 1**)but not after surgery. Levels of miR-920 and mir-92a-1 were not different in AKI patients.

Platelet response

Aspirin (75mg per day) was given to 61 (64.2%) patients before surgery (32 [61.5%] AKI and 29 [74.4%] non-AKI), which rose to 74 (77.9%) patients after surgery (40 [76.9%] AKI and 34 [87.2%] non-AKI). P2Y12 antagonists (Clopidogrel or Ticagrelor) were stopped five days before surgery. Six patients with AKI (11.5%) and seven patients without AKI received P2Y12 inhibitors after surgery in addition to aspirin. There was no significant difference in anti-platelet administration between AKI groups (**Table 1**).

The platelet response was analysed with a Multiplate aggregometer using ADP, ASPI and TRAP assays. Aspirin was the only drug that could affect platelet response during the study since patients received it daily. As expected, taking aspirin resulted in a lower AUC in the ASPI test, which was, and remained, below the cut-off value for aspirin responsiveness for the majority (78-88%) of aspirinated patients at all time points (**Fig. 3A**). There was no statistical difference in the ASPI responses (**eTable 1**) or in the numbers of aspirin high responders between AKI and non-AKI groups at any time point (**Fig. 3B**).

The range of responses to ADP was broad (**Fig. 3C**) and generally the ADP AUC was reduced following surgery. Interestingly the patients given a P2Y12 antagonist after surgery did not have a significantly lower ADP response than those who were not. The analysis of ADP results showed an overall trend for the AKI patients to have a lower ADP response at every time point, although the statistical difference between the groups was only observed at 6-12 hours after surgery (**Fig. 3D, eTable 1**). The TRAP test, which is insensitive to aspirin, indicated that the platelets showed the highest levels of aggregation 6-12 hours after surgery, which then fell over the subsequent 48 hours to levels that were lower than the responses before surgery (**Fig. 3E**). There was no difference in TRAP response between AKI groups (**eTable 1**).

Thrombo-inflammatory cellular markers

Activation of blood cells implicated in the pathogenesis of AKI was measured using flow cytometry. These included markers of leukocyte activation (CD11b), monocyte activation (CD64 and CD163), and platelet activation (CD41, CD62P and activated GPIIb/IIIa), as well as circulating platelet-monocyte and platelet-granulocyte aggregates (CD14/CD41 and CD16/CD41, respectively).

Significantly higher levels of platelet activation (activated GPIIb/IIIa detected with PAC-1 antibody, **Fig. 4B** and **eTable 1**) were seen in the AKI patients 6-12 hours after surgery. P-selectin expression (CD41/CD62P), a marker of platelet degranulation, was no higher in the AKI patients, and did not change over time, but higher levels of granulocyte-platelet aggregates (a consequence of P-

selectin expression on platelets) were seen in the AKI patients both before surgery and 6-12 and 24 hours afterwards (**Fig. 4B** and **eTable 1**).

Analysis of soluble biomarkers of inflammation included CXCL1, soluble ICAM1 (sICAM1), IL-6, 8 and 10. Out of these, only sICAM1 was altered, and was significantly higher before surgery and 48 hours afterwards in the patients who developed AKI (**Fig. 4C** and **eTable 1**).

Out of all significant pre-operative variables, levels of platelet-granulocyte aggregates (CD16/CD41) correlated positively with levels of sICAM1 and negatively with levels of miR-133b. No significant correlation was observed between postoperative (6-12 hrs) variables.

Sensitivity analysis in aspirin-treated patients

Given that most patients were receiving aspirin, we restricted the analysis to patients on aspirin only. Patients receiving P2Y12 inhibitors were also excluded since they constituted less than 10% of all patients. The results of this analysis were very similar to those of all patients. Patients with AKI had higher levels of serum creatinine 6 – 96 hours after surgery, platelets-derived EV before surgery and at 24 hours post, levels miR-1262 at ICU, platelet-leukocyte aggregates before and 6-12 hours after surgery, and sICAM1 at 48 hours after surgery. Multiplate ADP AUC was lower at 6-12 hours after surgery in patients with AKI. However, we did not observe significant differences for activated GPIIb/IIIa and hsa-miR-133b, platelet-leukocyte aggregates 24 hours after surgery and sICAM1 before surgery (**eTable 3**).

Discussion

Main findings

Longitudinal analysis of cellular activation, inflammatory response, and levels of circulating EV and miRNA has shown a significant elevation of platelet activation, as demonstrated by platelets with activated GPIIb/IIIa and by platelet-granulocyte aggregates in cardiac surgery patients who develop AKI. Levels of platelet-granulocyte aggregates (CD16/CD41) were significantly higher pre-surgery and at 6-12 hours and 24 hours post-surgery in the AKI group. This group also had increased levels of platelets with activated GP IIb/IIIa (PAC-1 positive) and platelet-derived EVs at 6-12 and 24 hours post-surgery, respectively, alongside increased ADP insensitivity 6-12 hours after surgery. Moreover, circulating sICAM1 levels were higher, and levels of plasma miR-133b were lower at baseline in the AKI group. Both of these correlated with levels of platelet-granulocyte aggregates at baseline. miR-1262 was most differentially expressed in the AKI group immediately after surgery and correlated with pre-operative sICAM1 levels.

Clinical significance

The results demonstrate that platelet activation, resulting in platelet leukocyte aggregation are associated with AKI following cardiac surgery. Platelets have been shown to have an important role in experimental AKI, where platelet-granulocyte aggregates facilitate intraglomerular neutrophil recruitment and ROS production. [24] In rodents, platelets guide neutrophils to areas with reduced endothelial integrity like glomeruli, [3] whose glycocalyx is disrupted during AKI. [25] In the current study, sICAM1 and platelet leukocyte aggregates were positively correlated before surgery, suggesting that a pro-inflammatory state could prime the patients' susceptibility to renal injury.

miR-1262 has been associated with mesangial proliferative glomerulonephritis, [26] However, it remains to be tested whether its correlation with sICAM1 is functionally related. There is also little known about the role of miR-133b, which correlated negatively with platelet-granulocyte aggregates. It has been shown to regulate kidney fibrosis, by inhibiting connective tissue growth factor [27] and was previously associated with diabetic nephropathy, where its serum levels increase, [28] but no previous studies have associated miR-133b with either platelets or granulocytes.

Our data suggest that the observed desensitisation of platelets to ADP at 6-12 hours after surgery is not a consequence of the action of P2Y12 antagonists, which were not administered until 24 hours after surgery. It could be explained by the release of ADP in vivo from activated platelets, damaged red cells or other tissues. [29] We also show that aspirin did not affect the observed changes in platelet function in the AKI group. No trial to our knowledge has evaluated the renoprotective effect of anti-platelet agents in cardiac surgery. [16,30] Although, multiple RCTs have compared the effects of different anti-platelet agents on vein graft patency rates, arterial thromboses, and

bleeding post-surgery, [31] none of these trials reported acute kidney injury event rates. Indirect evidence for a renoprotective effect of platelet inhibition comes from small trials of NO donors or prostacyclin [32-34] that have known anti-platelet effects and improve creatinine clearance after cardiac surgery. Together, these results support the reevaluation of existing trials or new trials of novel antiplatelet agents or blockers of P-selectin.

Limitations

As in any other study of AKI biomarkers, the major limitation is the use of serum creatinine to define renal injury. [35] More recent biomarkers, such as TIMP-2 and IGFBP7, have been shown to be useful diagnostic tools for AKI. However, they are not discriminatory in the cardiac surgery cohort. [13] Another limitation is the use of pooled samples for the initial identification of AKI-associated microRNAs. This was dictated by the budgetary limitations, and analysis of all patients, or a selected cohort of patients with well-defined clinical characteristics, could potentially result in the identification of more relevant microRNAs. Sample pooling can also be behind the discrepancies in miR-133b expression between discovery and validation. NanoSight, used to estimate particle concentration, does not discriminate between cell-derived, membrane-bound particles and other particulates in plasma such as lipoproteins. Therefore, we are not able to conclude whether the increases in specific particle sizes are associated with platelet-derived EVs. A final limitation is the small size of the study and high levels of missing data for flow cytometry analysis of platelet and leukocyte activation. We cannot be sure that a larger sample, or one with different eligibility criteria would not provide a different result.

Conclusions

Our analysis does not support EVs and miRNAs as predictive biomarkers of AKI. However, markers of platelet activation have a diagnostic potential and need to be validated in a larger cohort. Moreover, blockers of P-selectin could have a therapeutic potential.

Authors' contributions

Individual contributions to the study were as follows: GJM AHG and MJW designed the study, MJW and NB wrote the manuscript with critical input from GJM and AHG. LJ-D, TK and HA managed the conduct of the study. NS, BE-H, SS and KT collected samples and performed wet lab experiments. NT and MZ extracted clinical data. NB, NS, FYL and MJW performed statistical analyses

Funding

British Heart Foundation (RG/13/6/29947), (CH/12/1/29419) to GJM, MW, TK, and HA; Leicester and Bristol National Institute for Health Research Cardiovascular Biomedical Research Units.

Competing interests

All authors have disclosed that they do not have any potential conflicts of interest.

References

1. Karkouti, K. *et al.* Acute kidney injury after cardiac surgery: focus on modifiable risk factors. *Circulation* **119**, 495–502 (2009).

2. Lagny, M.-G. *et al.* Incidence and outcomes of acute kidney injury after cardiac surgery using either criteria of the RIFLE classification. *BMC Nephrol* **16**, 76 (2015).

3. Jansen, M. P. B., Florquin, S. & Roelofs, J. J. T. H. The role of platelets in acute kidney injury. *Nat Rev Nephrol* **14**, 457–471 (2018).

4. Basile, D. P., Anderson, M. D. & Sutton, T. A. Pathophysiology of acute kidney injury. *Compr Physiol* **2**, 1303–1353 (2012).

5. von Hundelshausen, P. & Weber, C. Platelets as immune cells: bridging inflammation and cardiovascular disease. *Circ Res* **100**, 27–40 (2007).

6. Gomchok, D., Ge, R.-L. & Wuren, T. Platelets in Renal Disease. *International Journal of Molecular Sciences* **24**, 14724 (2023).

7. Oh, S. & Kwon, S.-H. Extracellular Vesicles in Acute Kidney Injury and Clinical Applications. *Int J Mol Sci* **22**, 8913 (2021).

8. Arroyo, J. D. *et al.* Argonaute2 complexes carry a population of circulating microRNAs independent of vesicles in human plasma. *Proc Natl Acad Sci U S A* **108**, 5003–5008 (2011).

9. Vickers, K. C., Palmisano, B. T., Shoucri, B. M., Shamburek, R. D. & Remaley, A. T. MicroRNAs are transported in plasma and delivered to recipient cells by high-density lipoproteins. *Nat Cell Biol* **13**, 423–433 (2011).

10. Wang, K., Zhang, S., Weber, J., Baxter, D. & Galas, D. J. Export of microRNAs and microRNA-protective protein by mammalian cells. *Nucleic Acids Res* **38**, 7248–7259 (2010).

11. Creemers, E. E., Tijsen, A. J. & Pinto, Y. M. Circulating microRNAs: novel biomarkers and extracellular communicators in cardiovascular disease? *Circ Res* **110**, 483–495 (2012).

12. Fan, P. C., Chen, C. C., Chen, Y. C., Chang, Y. S. & Chu, P. H. MicroRNAs in acute kidney injury. *Hum Genomics* **10**, 29 (2016).

13. Miller, D. *et al.* Urinary extracellular vesicles and micro-RNA as markers of acute kidney injury after cardiac surgery. *Sci Rep* **12**, 10402 (2022).

14. Fan, P. C. *et al.* A circulating miRNA signature for early diagnosis of acute kidney injury following acute myocardial infarction. *J Transl Med* **17**, 139 (2019).

15. Sullo, N. *et al.* An Observational Cohort Feasibility Study to Identify Microvesicle and Micro-RNA Biomarkers of Acute Kidney Injury Following Pediatric Cardiac Surgery. *Pediatr Crit Care Med* **19**, 816–830 (2018).

16. Pathak, S. *et al.* Pharmacological interventions for the prevention of renal injury in surgical patients: a systematic literature review and meta-analysis. *Br J Anaesth* **126**, 131–138 (2021).

17. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Research* **43**, e47 (2015).

18. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical   Computing. (2022).

19. Wickham, H. *Ggplot2: Elegant Graphics for Data Analysis*. (Springer, New York, NY, 2009). doi:10.1007/978-0-387-98141-3.

20. Rothman, K. J. No adjustments are needed for multiple comparisons. *Epidemiology* **1**, 43–46 (1990).

21. Saville, D. J. Multiple Comparison Procedures: The Practical Solution. *The American Statistician* **44**, 174–180 (1990).

22. Ridker, P. M. *et al.* C-reactive protein levels and outcomes after statin therapy. *N Engl J Med* **352**, 20–28 (2005).

23. Kidney Disease: Improving Global Outcomes (KDIGO) Acute Kidney Injury Work Group. KDIGO Clinical Practice Guideline for Acute Kidney Injury. *Kidney inter.*, *Suppl.* **2**, 1–138 (2012).

24. Singbartl, K., Forlow, S. B. & Ley, K. Platelet, but not endothelial, P-selectin is critical for neutrophil-mediated acute postischemic renal failure. *FASEB J* **15**, 2337–2344 (2001).

25. Qureshi, S. H., Patel, N. N. & Murphy, G. J. Vascular endothelial cell changes in postcardiac surgery acute kidney injury. *Am J Physiol Renal Physiol* **314**, F726–F735 (2018).

26. Dai, R. *et al.* Differential expression profile of urinary exosomal microRNAs in patients with mesangial proliferative glomerulonephritis. *Aging (Albany NY)* **15**, 866–880 (2023).

27. Cao, D. *et al.* Regulation of connective tissue growth factor expression by miR-133b for the treatment of renal interstitial fibrosis in aged mice with unilateral ureteral obstruction. *Stem Cell Research & Therapy* **12**, 171 (2021).

28. Zhang, X.-L. *et al.* LncRNA CASC2 regulates high glucose-induced proliferation, extracellular matrix accumulation and oxidative stress of human mesangial cells via miR-133b/FOXP1 axis. *Eur Rev Med Pharmacol Sci* **24**, 802–812 (2020).

29. Woulfe, D., Yang, J. & Brass, L. ADP and platelets: the end of the beginning. *J Clin Invest* **107**, 1503–1505 (2001).

30. Abbasciano, R. G. *et al.* Activation of the innate immune response and organ injury after cardiac surgery: a systematic review and meta-analysis of randomised trials and analysis of individual patient data from randomised and non-randomised studies. *Br J Anaesth* **127**, 365–375 (2021).

31. Sandner, S. *et al.* Association of Dual Antiplatelet Therapy With Ticagrelor With Vein Graft Failure After Coronary Artery Bypass Graft Surgery: A Systematic Review and Meta-analysis. *JAMA* **328**, 554–562 (2022).

32. Darcin, O. T. *et al.* Effect of iloprost on renal function in patients undergoing coronary artery bypass grafting: a clinical study. *Ann Thorac Cardiovasc Surg* **19**, 12–17 (2013).

33. Lei, C. *et al.* Nitric Oxide Decreases Acute Kidney Injury and Stage 3 Chronic Kidney Disease after Cardiac Surgery. *Am J Respir Crit Care Med* **198**, 1279–1287 (2018).

34. Kaya, K. *et al.* The effect of sodium nitroprusside infusion on renal function during reperfusion period in patients undergoing coronary artery bypass grafting: a prospective randomized clinical trial. *Eur J Cardiothorac Surg* **31**, 290–297 (2007).

35. Harky, A. *et al.* Acute Kidney Injury Associated with Cardiac Surgery: a Comprehensive Literature Review. *Braz J Cardiovasc Surg* **35**, 211–224 (2020).

Table 1 - Baseline and intraoperative characteristics. Tests between groups were conducted by exact test for categorical variables and t-test or non-parametric Kruskal–Wallis test for continuous variables. Data are presented as n (%) for categorical variables and mean (standard deviation, STD) or median (interquartile range) for continuous variables. Significant values are given in bold. * missing data: smoking staus (n=3), CCS (n=1), LVEF (n=5), left main steam diseased (n=4), Extent of coronary disease (n=4), Antiplatelets drugs (N=4)

	no AKI	AKI	p-value
	(n = 41)	(n = 54)	
Diabetes	20 (48.8%)	31 (57.4%)	0.416
CKD	9 (22.0%)	11 (20.4%)	0.852
Age (years)	70.6 (6.9)	72.8 (7.6)	0.151
Sex - Male	35 (85.4%)	46 (85.2%)	0.980
Caucasian	41 (100.0%)	52 (96.3%)	0.504
BMI	31.7 (5.2)	31.4 (6.1)	0.745
Surgery type			
CABG	19 (46.3%)	19 (35.2%)	0.324
Valve	15 (36.6%)	19 (35.2%)	
both/others	7 (17.1%)	16 (29.6%)	
Medical history			
Myocardial infarction	7 (17.1%)	10 (18.5%)	0.856
CVA/ TIA	5 (12.2%)	8 (14.8%)	0.713
Chronic pulmonary disease	1 (2.4%)	2 (3.7%)	0.999
Previous cardiac surgery	2 (4.9%)	4 (7.4%)	0.696
Smoker			
Current	12 (30.0%)	22 (42.3%)	0.478
Ex-smoker	18 (45.0%)	19 (36.5%)	
Never	10 (25.0%)	11 (21.2%)	
NYHA			
class I	5 (12.2%)	8 (14.8%)	0.491
class II	29 (70.7%)	32 (59.3%)	
class III & IV	7 (17.1%)	14 (25.9%)	
CCS			
asymptomatic	11 (27.5%)	24 (44.4%)	0.110
class I	10 (25.0%)	11 (20.4%)	
class II	16 (40.0%)	19 (35.2%)	
class III	3 (7.5%)	0 (0.0%)	
Left ventricular ejection fraction (LVEF)			
Good(>49%)	24 (63.2%)	35 (67.3%)	0.682
Fair(30-49%)	14 (36.8%)	17 (32.7%)	
Left main stem diseased	4 (10.3%)	9 (17.3%)	0.342
Extent of coronary disease			
Normal	6 (15.4%)	11 (21.2%)	0.287
1VD	11 (28.2%)	9 (17.3%)	

2VD	8 (20.5%)	6 (11.5%)	
3VD	14 (35.9%)	26 (50.0%)	
Anti-platelet drugs*			
Aspirin (pre-operative)	29 (74.4%)	32 (61.5%)	0.288
P2Y12 inhibitors (stopped 5 days before surgery)	5 (12.8)	7 (13.5)	1
Aspirin (1 day after surgery)	34 (87.2%)	40 (76.9%)	0.332
P2Y12 inhibitors (1 days after surgery)	7 (17.9%)	6 (11.5)	0.574
Dual platelet therapy (1 days after surgery)	7 (17.9%)	6 (11.5)	0.574
Serum Troponin I	20 (15 -26)	15 (15 -27)	0.627
Serum creatinine (u mol/L)	88 (73 -103)	85 (73 -105)	0.848
Heart rate (bpm)	68.9 (12.4)	66.4 (11.7)	0.328
MAP (mmHg)	98.7 (11.9)	97.5 (14.4)	0.674
Hb (g/dL)	13.3 (1.7)	12.9 (1.4)	0.246
Hct (%)	39.1 (4.7)	38.3 (4.0)	0.418
WBC (x10^9/L)	7.9 (1.8)	8.1 (1.9)	0.623
Platelets (x10^9/L)	225.6 (76.8)	214.9 (50.9)	0.457
eGFR (ml/min/1.73m2)	82.4 (30.1)	78.2 (22.4)	0.453
Albumin (g/L)	40.5 (2.3)	40.6 (3.6)	0.825
Bilirubin (u mol/L)	13.1 (5.0)	13.1 (6.5)	0.992
AST (IU/L)	23.0 (4.8)	25.1 (7.1)	0.107
ALT (IU/L)	24.1 (11.0)	24.8 (9.9)	0.760
ALP (IU/L)	80.1 (26.5)	83.1 (31.9)	0.634
GGT (IU/L)	24 (18 -34)	32 (19 -58)	0.086
CBP time (min)	93 (82 -109)	106 (82 -139)	0.117
CrossClampTime (min)	55 (46 -77)	65 (47 -97)	0.259
Ventilation time (hr)	9 (7 -14)	13 (10 -17)	**0.001**
ICU stay (hr)	27 (25 -53)	48 (25 -93)	0.080
Hospital LOS (day)	8 (6 -9)	10 (7 -13)	**0.017**

Figure legends

Figure 1 – (**A**) CONSORT diagram. (**B**) Levels of serum creatinine. Asterisks indicate a significant difference between no-AKI and AKI (*p*<0.05).

Figure 2 – (**A**) Total particle concentrations in citrated plasma. (**B**) Mean size distribution of particle concentrations (+/- SD). (**C**) Box plot of EVs positive for platelet marker CD41. (**D**) Bar plot showing the log fold difference in miRNA identified by qRT-PCR arrays between AKI and non-AKI. Only miRNA with adjusted *p*-value<0.05 are shown (see **eTable 2** for details). (**E**) Box plot of validation results for miR-133b and miR-1262. Asterisks indicate a significant difference between AKI and non-AKI groups (p < 0.05). Open and grey boxes in A, C and E represent data from patients with and without AKI, respectively.

Figure 3 – (A) Density plots of the area under the curve (AUC) values acquired with Multiplate ASPI test. The vertical red line indicates the position of a cut-off value for ASPI high responders (AUC = 473). (B) Fraction of ASPI high responders in people with AKI or without. The analysis was performed for aspirin-treated and untreated patients. (C) Density plots of the area under the curve (AUC) values acquired with Multiplate ADP test. (D) Box plot of AUC data acquired using Multiplate ADP assay. Asterisks indicate a significant difference between AKI and non-AKI groups (p < 0.05). Open and grey boxes in B and D represent data from patients with and without AKI, respectively. (E) Density plots of the area under the curve (AUC) values acquired with Multiplate TRAP test.

Figure 4 – (**A**) Box plot of mean expression values for IIb/IIIa detected with PAC-1 antibody in flow cytometry. (**B**) Box plot of a fraction of events positive for CD16 and CD41. (**C**) Box plot of sICAM1 concentrations in citrated plasma acquired with a MAGPIX device. (**D**) Correlations (Spearman) between significant preoperative variables and ICU miR-1262, and between significant variables 6-12 hr after surgery. Blue circles indicate significant positive and red circles negative correlation. Asterisks indicate a significant difference between AKI and non-AKI groups (p < 0.05). Open and grey boxes in A – C represent data from patients with and without AKI, respectively.

Figure 1

Figure 2

Figure 3

Figure 4