Physical activity in adult users of inpatient mental health services: a

scoping review

Garry A. Tew[1]*, Emily Peckham[2], Suzy Ker[3], Jo Smith[3], Philip Hodgson[1,3], Katarzyna K. Machaczek[4],

Matthew Faires[1]

[1] Institute for Health and Care Improvement, York St John University

[2] School of Medical and Health Sciences, Bangor University

[3] Tees Esk and Wear Valleys NHS Foundation Trust

[4] College of Health, Wellbeing and Life Sciences, Sheffield Hallam University

* Corresponding author

E-mail: g.tew@yorksj.ac.uk (GT)

Short title: Physical activity and inpatient mental health services

Abstract

People with severe mental illness engage in considerably less physical activity than those without. They also experience premature mortality of around 10–25 years. A large proportion of these premature deaths are attributed to modifiable behaviours, including physical activity. The inpatient environment provides an opportunity to support people to become more physically active; however, there is limited evidence on which interventions are most successful and what contextual factors affect their delivery. A scoping review was conducted to help understand the extent and type of evidence in this area and identify research gaps. We included studies of physical activity in adult inpatient mental health services published in peer-reviewed journals. Reviews, meta-analyses, and papers focusing on eating disorder populations were excluded. We searched the MEDLINE, CINAHL, PsycINFO, ASSIA and Web of Science databases for relevant studies published in English. We extracted data on study design, participant characteristics, intervention and control conditions, key findings, and research recommendations. We used a descriptive analytical approach and results are presented in tables and figures. Of 25,437 unique records screened, 192 reports from 164 studies were included. Fifty-six studies reported on correlates of physical activity, and 125 studies reported on physical activity interventions. Most intervention studies used a single-group, pre-post design (44%) and included fewer than 100 participants (85%). Ninety percent of interventions delivered physical activity directly to participants, and 51% included group-based sessions. The duration, type, frequency and intensity of sessions varied. Mental health was the most commonly reported outcome (61%), whereas physical activity was rarely an outcome (11%). Overall, there is a modest but growing body of research on physical activity in adult users of inpatient mental health services. More high-quality trials are needed to advance the field, and future research should target neglected intervention types, outcomes, populations and settings.

Keywords

Exercise, Physical activity, Psychiatric inpatient, Severe mental illness, Scoping review

Introduction

People who use mental health services, including people with severe mental illness (SMI), depression, anxiety or stress disorder and people with alcohol and substance use disorders, experience worse health outcomes and a shortened life expectancy when compared to people without these disorders (10-25 years for SMI, 7-20 years for anxiety/stress disorders and alcohol and substance use disorders) [1–4]. Much of this reduced life expectancy is due to modifiable risk factors such as engaging in health risk behaviours, one of which is physical inactivity [5–7]; and it has been reported that between 70 and 75% of people with schizophrenia do not meet public health targets for physical activity [8]. Yet, physical activity interventions have been shown to improve cardiometabolic health outcomes in people with mental illnesses and improve symptoms of depression, cognitive function, feelings of isolation, and quality of life [9–11].

Insufficient physical activity and excessive sedentary behaviour in people receiving care for SMI have been observed in both community and inpatient settings [12,13]. However, the majority of intervention studies have recruited people receiving care in community and outpatient settings, and less research has been done in inpatient settings [14]. While some of the reasons for insufficient physical activity and excessive sedentary behaviour are common across both inpatient and community settings, such as the sedative effects of psychotropic medicine [15] and mental health symptoms (lack of motivation, anhedonia, fatigue and negative body image) [13], there are some unique barriers presented by inpatient settings. For example, lack of appropriate facilities, lack of suitably trained staff to support physical activity, and restrictions on leave for people detained under mental health legislation all contribute to decreased opportunities for physical activity in inpatient settings [16–19]. However, inpatient physical activity programmes provide an opportunity to improve patients' physical and mental health. For this reason, it is important to have an up-to-date understanding of the literature on this topic.

The scope of inpatient physical activity research is broad, including quantitative or qualitative

research such as randomised controlled trials (RCTs) of physical activity interventions or interviews

with service users exploring their perceptions of physical activity. Studies have also explored

correlates of physical activity or been more intervention-focused. Qualitative studies have examined

the views of patients, carers or healthcare professionals, or a particular subpopulation of patients.

Recent reviews regarding inpatient physical activity have focussed on tightly defined inclusion

criteria leading to the inclusion of specific sets of studies, such as RCTs. These include exercise

interventions in mental health inpatient settings [14], physical activity interventions for patients with

SMI in secure forensic settings [20], and studies aiming to increase or evaluate physical activity in

secure settings [21]. A broader review of inpatient physical activity research, that considers all

inpatient settings and quantitative and qualitative evidence would be advantageous to understand

the current state of the literature and inform future research. A scoping review is a suitable way of

achieving this [22]. Understanding and mapping the available evidence on physical activity in

inpatient settings, including qualitative and quantitative research, is important due to their

complementary nature. They answer different questions about physical activity. For example,

quantitative research can shed light on cause-and-effect relationships between various factors

associated with physical activity, while qualitative research can tell us why this is the case. A scoping

review also provides a means to determine whether there is scope and a need for a systematic

review of a particular type or in a specific area of literature, and to identify gaps in the existing

research base that could be filled by future primary research [22]. The overall aim of this scoping

review was therefore to understand the extent and type of evidence regarding physical activity in

adult users of inpatient mental health services. This included study designs used, primary conditions

of participants, types of outcomes assessed, correlates of physical activity explored, intervention

characteristics, outcomes, and research recommendations.

Methods

This scoping review was conducted in accordance with the Joanna Briggs Institute (JBI) methodology for scoping reviews [23] and reported in accordance with the PRISMA Extension for Scoping Reviews (PRISMAScR) [24]. A protocol was prepared in advance and published in the Open Science Framework [25].

Participants/context

We included studies pertaining to adult (≥18 years) users of inpatient mental health services. This included studies where psychiatric inpatients were the participants and studies where other stakeholders were involved (e.g., healthcare professionals giving their views regarding services for psychiatric inpatients). Studies focusing on learning disability populations were included whereas those focusing on eating disorder populations were excluded because of the unique requirement to carefully manage energy input/output in these populations. There were no limits on the country of origin.

Concept

The phenomenon of interest was physical activity. Physical activity has been defined as any bodily movement produced by skeletal muscles and requiring energy expenditure [26]. Exercise is a subset of physical activity that has been defined as any structured and repetitive physical activity that has an objective of improving or maintaining physical fitness [26]. To address the aims of the review we took a broad view of physical activity and included studies that had focused on supervised exercise or promotion of self-managed physical activity.

112 **Types of sources**

113 This scoping review included experimental and quasi-experimental study designs such as RCTs, non-

114 randomised controlled trials, before and after studies and interrupted time-series studies. In

115 addition, analytical observational studies including prospective and retrospective cohort studies,

116 case-control studies and analytical cross-sectional studies were included. This review also considered

117 descriptive observational study designs including case series, individual case reports and descriptive

118 cross-sectional studies for inclusion. Qualitative studies that utilised various methodologies (e.g.,

119 phenomenology, grounded theory, ethnography, action research) were also considered. Literature

120 reviews and meta-analyses were used to identify primary studies, but were excluded from data

121 analysis. Conference abstracts and opinion papers were also excluded.

122

123 **Search strategy**

124 The search strategy targeted peer-reviewed publications. A pilot search of the MEDLINE database

125 was undertaken to identify articles on the topic. The text words contained in the titles and abstracts

126 of relevant articles, and the index terms used to describe them were employed to develop a full

127 search strategy for the MEDLINE, CINAHL, PsycINFO, ASSIA and Web of Science databases (S1 File).

128 The search strategy, including all identified keywords and index terms, was adapted for each

129 included database and/or information source. The database searches were run on October 31, 2022.

130 The reference list of all included sources of evidence, and those of any review articles or meta-

131 analyses, were screened for additional studies. We also conducted forward citation tracking of

132 included studies using Google Scholar. We only included studies that were published in English

133 language from 2007 onwards. The latter relates to amendments in the 1983 Mental Health Act that

134 were made in 2007; prior studies may have employed different clinical practices and patient

135 populations.

Study selection

Following the search, all identified citations were collated and uploaded into Covidence [27], and

duplicates were removed. After a pilot test, the titles and abstracts were screened by pairs of

independent reviewers against the eligibility criteria for the review. Potentially relevant sources

were retrieved in full and assessed in detail against the eligibility criteria by pairs of independent

reviewers. The reasons for excluding sources of evidence in full text that did not meet the eligibility

criteria were documented. Any disagreements that arose between the reviewers at each stage of

the selection process were resolved through discussion or with the aid of additional reviewers.

Data extraction

Data were extracted from papers included in the scoping review by pairs of independent reviewers

using a data extraction form developed by the reviewers in Covidence [27], which was based on the

JBI template extraction tool [28]. The form was piloted on twenty papers to ensure it was fit for

purpose.

The extracted data encompassed specific details about the participants, concept, context, study

methods, key findings relevant to the review questions, and research recommendations. Any

disagreements between the reviewers were resolved through discussion or with the aid of an

additional reviewer. Due to limited timescales and resources, we did not contact any authors to

request missing or additional data.

Data analysis and presentation

A descriptive analytical approach was used to summarise the included studies' contextual, process

and outcome-related data [29,30]. This approach was undertaken to map the key concepts and

available evidence, synthesise existing research findings, and identify research gaps. Physical activity

correlates were categorised based on the Socio-Ecological Model [31–33]. This model was chosen as

it considers the multifaceted and interactive effects of various factors (e.g., interpersonal,

organisational and community), which characterise the delivery of physical activity interventions for

people with SMI. The analysis was reported in accordance with the synthesis without meta-analysis

(SWiM) guideline [34], with data presented in tables and figures where appropriate.

Results

Fig 1 shows the PRISMA flow diagram. The database searches yielded 33,438 records. After
removing duplicates, we screened 25,200 records, from which we reviewed 269 full-text reports,
and finally included 121 reports. Later, we reviewed a further 237 full-text reports that were
identified from review articles or the forward and backward citations of included reports. This
resulted in a further 71 reports being included. Together, 192 reports from 164 studies were
included in the review. The complete reference list of included reports can be found in S2 File.

Fig 1. PRISMA flow diagram.

Descriptive characteristics and methodological information of included articles

Fifty-six (34%) studies reported on correlates of physical activity, 125 (76%) studies reported on
physical activity interventions, and 18 (11%) studies reported on both physical activity correlates and
interventions.

Of the 56 studies on correlates of physical activity, 38 (68%) studies reported on quantitative data
(24 observational studies, 14 interventional studies) and 22 (39%) studies reported on qualitative
data (14 observational studies, 8 interventional studies) (Table 1). Sample sizes varied by study type,
with observational quantitative studies including the largest samples. The majority of studies were
conducted in Europe (62.5%), with most of the remainder originating from Australasia (16%) and
Asia (11%). The three most common settings were psychiatric hospitals (44%), forensic/secure
settings (23%), and 'mixed' settings (18%).

Of the 125 studies on interventions (Table 2), 55 (44%) studies used a single-group, pre-post design
and 48 (38%) studies were RCTs. The majority of studies included fewer than 100 participants (85%)

and were conducted in Europe (49%) and within a psychiatric hospital (46%). The most common

study populations included people with schizophrenia or related psychotic disorders (36%) and a

mixture of diagnoses (21%). Mental health was the most commonly reported outcome (61%),

whereas physical activity was rarely an outcome (11%) (Table 2).

Fourteen (11%) of the intervention studies were qualitative studies. All 14 studies collected

qualitative data from service users who had participated in the intervention; seven studies (50%)

also collected data from healthcare staff, and one study (7%) also collected data from family

members. Most studies (71%) used one-to-one interviews to collect data. Other methods included

observations (n=2), evaluation forms (n=2), document analysis (n=1) and focus groups (n=1).

Table 1. Characteristics of studies of physical activity correlates by type of study*.

	Quantitative		Qualitative		Overall (n=56)
	Observational (n=24)	Interventional (n=14)	Observational (n=14)	Interventional (n=8)	
Sample size	1-100: 9 (37.5%) 101-500: 13 (54%) 500+: 2 (8%)	1-100: 12 (86%) 101-500: 2 (14%) 500+: 0 (0%)	1-15: 6 (43%) 16+: 8 (57%) Missing: 0 (0%)	1-15: 3 (37.5%) 16+: 4 (50%) Missing: 1 (12.5%)	1-100: 38 (68%) 101-500: 15 (27%) 500+: 2 (4%) Missing: 1 (2%)
Continent					
Europe	14 (58%)	11 (79%)	7 (50%)	6 (75%)	35 (62.5%)
North America	1 (4%)	2 (14%)	0 (0%)	0 (0%)	3 (5%)
South America	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Asia	3 (12.5%)	0 (0%)	2 (14%)	1 (12.5%)	6 (11%)
Australasia	4 (17%)	1 (7%)	4 (29%)	1 (12.5%)	9 (16%)
Africa	1 (4%)	0 (0%)	1 (7%)	0 (0%)	2 (4%)
Multiple continents	1 (4%)	0 (0%)	0 (0%)	0 (0%)	1 (2%)
Setting					
Psychiatric hospital	12 (50%)	8 (57%)	3 (21%)	3 (37.5%)	23 (41%)
General hospital	0 (0%)	1 (7%)	1 (7%)	2 (25%)	4 (7%)
Mental health rehabilitation centre	2 (8%)	1 (7%)	0 (0%)	0 (0%)	3 (5%)
Forensic/secure	2 (8%)	3 (21%)	5 (36%)	3 (37.5%)	13 (23%)
Mixed	5 (21%)	1 (7%)	5 (36%)	0 (0%)	10 (18%)
Other	2 (8%)	0 (0%)	0 (0%)	0 (0%)	2 (4%)
Missing	1 (4%)	0 (0%)	0 (0%)	0 (0%)	1 (2%)

199   *Numbers are not exclusive – studies can be counted multiple times, for example, if they include both quantitative and qualitative data.

200   **Table 2. Characteristics of studies of physical activity interventions by type of study*.**

	Randomised controlled trials (n=48)	Other design (n=79)	Overall (n=125)
Sample size	1-100: 41 (85%) 101-500: 7 (15%)	1-100: 66 (83.5%) 101-500: 10 (13%) Missing: 3 (4%)	1-100: 106 (85%) 101-500: 16 (13%) Missing: 3 (2%)
Continent			
Europe	21 (44%)	41 (52%)	61 (49%)
North America	4 (8%)	12 (15%)	15 (12%)
South America	2 (4%)	1 (1%)	3 (2%)
Asia	20 (42%)	12 (15%)	32 (26%)
Australasia	1 (2%)	12 (15%)	13 (10%)
Africa	0 (0%)	1 (1%)	1 (1%)
Multiple continents	0 (0%)	0 (0%)	0 (0%)
Setting			
Psychiatric hospital	25 (52%)	34 (43%)	58 (46%)
General hospital	12 (25%)	18 (23%)	30 (24%)
Forensic/secure	1 (2%)	14 (18%)	15 (12%)
Mental health rehabilitation centre	7 (15%)	10 (13%)	16 (13%)
Mixed	1 (2%)	2 (2.5%)	3 (2%)
Other	2 (4%)	1 (1%)	3 (2%)
Primary condition			
Depression or bipolar disorder	13 (27%)	10 (13%)	22 (18%)
Schizophrenia or related psychotic disorders	21 (44%)	25 (32%)	45 (36%)
Post-Traumatic Stress Disorder	1 (2%)	1 (1%)	2 (2%)
Dementia	2 (4%)	1 (1%)	3 (2%)
Substance misuse	1 (2%)	0 (0%)	1 (1%)
Personality disorder	0 (0%)	1 (1%)	1 (1%)
Learning disability	0 (0%)	1 (1%)	1 (1%)
Mixed	6 (12.5%)	20 (25%)	26 (21%)
Unclear	4 (8%)	20 (25%)	24 (19%)
Type of programme			
Physical activity delivered	38 (79%)	63 (80%)	100 (80%)
Physical activity promoted	6 (12.5%)	3 (4%)	8 (6%)
Mixed	3 (6%)	10 (13%)	13 (10%)
Unclear	1 (2%)	3 (4%)	4 (3%)
Type of physical activity			
Various activities	11 (23%)	21 (27%)	31 (25%)
Structured exercise	15 (31%)	23 (29%)	38 (30%)
Exer-gaming	1 (2%)	4 (5%)	5 (4%)
Sport	2 (4%)	7 (9%)	9 (7%)
Yoga	3 (6%)	8 (10%)	11 (9%)
Unstructured walking/ jogging/ running	11 (23%)	5 (6%)	16 (13%)
Dance	3 (6%)	6 (8%)	8 (6%)
Unclear	2 (4%)	5 (6%)	7 (6%)
Duration of programme			
Variable	0 (0%)	7 (9%)	7 (6%)
Duration of admission	2 (4%)	17 (21.5%)	19 (15%)

Fixed (≤6 weeks)	20 (42%)	5 (6%)	25 (20%)
Fixed (>6 weeks)	23 (48%)	30 (8%)	52 (42%)
Single session	3 (6%)	11 (14%)	13 (10%)
Unclear	0 (0%)	9 (11%)	9 (7%)
Provider			
Study team	15 (31%)	6 (8%)	21 (17%)
Qualified therapist (internal)	16 (33%)	41 (52%)	56 (45%)
Qualified therapist (external)	3 (6%)	10 (13%)	12 (10%)
Member of staff	5 (10%)	9 (11%)	14 (8%)
Member of staff trained by study team	2 (4%)	4 (5%)	6 (8%)
Unclear	8 (17%)	13 (16.5%)	21 (17%)
Who with?			
Individual	6 (12.5%)	8 (10%)	13 (10%)
Group	22 (46%)	43 (54%)	64 (51%)
Mixed	4 (8%)	9 (11%)	13 (10%)
Unclear	16 (33%)	19 (24%)	35 (28%)
Comparator			
No comparator	0 (0%)	53 (67%)	55 (44%)
Usual care	23 (48%)	13 (16.5%)	34 (27%)
Other PA intervention	11 (23%)	4 (5%)	14 (11%)
Other non-PA intervention	7 (15%)	6 (8%)	13 (10%)
Other PA and usual care	3 (6%)	2 (2.5%)	5 (4%)
Other non-PA and usual care	2 (4%)	0 (0%)	2 (2%)
Other PA and other non-PA intervention	1 (2%)	0 (0%)	1 (1%)
Unclear	1 (2%)	1 (1%)	2 (2%)
Outcomes			
PA self-report	3 (6%)	5 (6%)	8 (6%)
PA device-based	1 (2%)	4 (5%)	5 (4%)
PA mixed	1 (2%)	1 (1%)	1 (1%)
Mental health	32 (67%)	45 (57%)	76 (61%)
Physical health	17 (35%)	28 (35%)	45 (36%)
Cognition	11 (23%)	1 (1%)	12 (10%)
Quality of life	10 (21%)	9 (11%)	19 (15%)

*Numbers are not exclusive – studies can be counted multiple times

PA, Physical activity

Correlates of physical activity

Tables 3 and 4 summarise the physical activity correlates examined or highlighted in quantitative

and qualitative studies, respectively. A broad range of individual, interpersonal, environmental and

organisational correlates was reported regarding inpatient physical activity, whereas no wider

societal factors were highlighted. 'Health status' and 'medication side effects' were the most

commonly reported demographic and biological factors in both quantitative (18% and 20%,

respectively) and qualitative (55% and 23%, respectively) studies. 'Self-motivation' was the most

common factor in the 'psychological, cognitive and emotional' category (36% quantitative, 45%

qualitative). Factors were less frequently reported in the 'behavioural' and 'social and cultural'

categories, but 'physical activity enjoyment' (7% quantitative, 5% qualitative) and 'social support'

(25% quantitative, 41% qualitative) were the most common factors, respectively. 'Access to

equipment and facilities' (27% quantitative) and 'environment restrictions' (18% quantitative, 32%

qualitative) were commonly cited physical environment factors. 'Staff capacity' (32% quantitative,

82% qualitative) and 'staff capability' (50% quantitative, 59% qualitative) were among the top-

ranking organisational factors.

Table 5 summarises the themes of research recommendations from the studies of physical activity

correlates. Seventeen (30%) of the 56 studies included research recommendations, which we

grouped according to whether they were related to the research topic, research methods, or patient

involvement.

Table 3. Summary of physical activity correlates examined or highlighted in quantitative studies.

Factor	Number (%) of studies out of 44
Demographic and biological factors	
Medication side-effects	9 (20%)
Health status	8 (18%)
Financial status	4 (9%)
Weight/fatness	3 (7%)
Physical activity status of healthcare staff	3 (7%)
Age	2 (5%)
Education	2 (5%)
Fitness status	1 (2%)
Gender	1 (2%)
Diabetes	1 (2%)
Marital status	1 (2%)
Psychological, cognitive and emotional factors	
Self-motivation	16 (36%)
Psychological health	7 (16%)
Self-efficacy	7 (16%)
Attitudes towards physical activity	6 (14%)
Outcome expectations	6 (14%)
Time availability	5 (11%)
Mental illness stigma	5 (11%)
Physical activity knowledge	3 (7%)
Fear of injury	3 (7%)
Fatigue/Tiredness	3 (7%)
Value of physical health benefits	2 (5%)
Cognitive function	1 (2%)
Mood	1 (2%)
Fear of going outdoors	1 (2%)
Attitudes to staff support	1 (2%)
Self-consciousness	1 (2%)
Behavioural factors	
Physical activity enjoyment	3 (7%)
Alcohol use	2 (5%)
Physical activity duration	1 (2%)
Physical activity intensity	1 (2%)
Physical activity autonomy	1 (2%)
Sleep	1 (2%)
Smoking habits	1 (2%)
Drug use	1 (2%)
Dietary habits	1 (2%)
Social and cultural factors	
Social support from family, friends, or peers	11 (25%)
Staff influence/involvement	5 (11%)
Someone to exercise with	1 (2%)
Acceptance of women exercising	1 (2%)
Acceptance of group exercise	1 (2%)
Religious beliefs	1 (2%)
Physical environment factors	

Access to equipment and facilities	12 (27%)
Environment restrictions	8 (18%)
Safe environment	4 (9%)
Unhealthy environment	2 (5%)
Weather/climate	1 (2%)
Access to green space	1 (2%)
Organisational factors	
Staff capability to support physical activity	22 (50%)
Staff availability/capacity	14 (32%)
Attitudes to/priority for physical activity	10 (23%)
Staff responsibility for physical activity	6 (14%)
Funding	6 (14%)
Opportunities for physical activity	5 (11%)
Staff confidence to support physical activity	2 (5%)
Staff working collaboratively	1 (2%)
Staff awareness of physical health issues	1 (2%)
Hospital size	1 (2%)
Safeguarding procedures	1 (2%)
Awareness of opportunities	1 (2%)
Timing of physical activity sessions	1 (2%)

224 **Table 4. Summary of physical activity correlates examined or highlighted in qualitative studies.**

Factor	Number (%) of studies out of 22
Demographic and biological factors	
Health status	12 (55%)
Medication side-effects	5 (23%)
Psychological, cognitive and emotional factors	
Self-motivation	10 (45%)
Mental illness stigma	7 (32%)
Value of physical health benefits	7 (32%)
Psychological health	3 (14%)
Time availability	2 (9%)
Cognitive function	2 (9%)
Behavioural factors	
Physical activity enjoyment	1 (5%)
Physical activity structure (e.g., intensity/duration)	1 (5%)
Perceived suitability of activities	1 (5%)
Social and cultural factors	
Social support from family, friends or peers	9 (41%)
Acceptance of women exercising	1 (5%)
Acceptance of group exercise	1 (5%)
Physical environment factors	
Environment restrictions	7 (32%)
Safe environment	7 (32%)
Organisational factors	
Staff availability/capacity	18 (82%)
Funding	16 (73%)
Staff capability to support physical activity	13 (59%)
Staff working collaboratively	4 (18%)
Therapeutic relationship with staff	4 (18%)
Links to community groups/services	1 (5%)

226 **Tables 5. Themes of research recommendations from studies of physical activity correlates.**

Theme	Examples of specific research recommendations	Number (%) of articles out of 56
Research topic-related recommendations		
Investigate other groups	"it seems worthwhile to replicate this study in out-patients" (Gerber 2018) "it might be preferable to compare more extreme groups" (Gerber 2018) "subgroups of patients for example, with persons with severe depression" (Ussher 2007)	5 (9%)
Investigate other correlates	"implicit attitudes towards exercise" (Gerber 2018) "consider other comparisons such as by body mass index, perceived physical health or age" (Fraser 2015) "the role of systemic influences on mental health treatment for exercise facilitation" (Ball 2022) "investigation of non-linear relationships between the physical activity and the basic psychological needs is warranted" (Bentzen 2021)	5 (9%)
Investigate other setting	"consider staff perceptions across other healthcare settings" (Kinnafick 2018)	2 (4%)
Research methods-related recommendations		
Longitudinal studies	"the examination of cross-lagged relationships between these variables over time" (Gerber 2018) "further (longitudinal) research to obtain more accurate insights on this topic" (Deenik 2017)	4 (7%)
Intervention studies	"explore the most effective exercise promotion strategies for autonomous exercise engagement for long lasting behaviour change" (Kinnafick 2018) "identify effective exercise interventions and feasible delivery models" (Tetlie 2009)	4 (7%)
Use other data collection methods	"incorporating objective measurements of physical activity and sedentary behaviour" (Seet 2021) "A more structured questionnaire or short interviews with patients might have allowed for the collection of richer descriptions" (Wynaden 2012) "More extensive interviews with the patients and clinical staff and longer period of participant observation" (Bierski 2016)	4 (7%)
Control for additional factors	"controlling more systematically for cognitive capacity" (Gerber 2018) "while controlling for the effects of medication and symptom severity" (Seet 2021)	2 (4%)
Larger studies	"with larger samples of patients" (Ussher 2007) "replicate the findings with larger sample sizes" (Bentzen 2021)	2 (4%)
Explore mechanisms	"regarding the mechanisms to modify such views as a means of increasing such care provision" (Bartlem	1 (2%)

	2018)	
Patient involvement-related recommendations		
Intervention co-produced with patients	"Patient led ideas and co-design should be key principles in programme and environmental design" (Every-Palmer 2018)	1 (2%)

Physical activity interventions

The components of physical activity interventions are summarised in Table 2. Most interventions

delivered physical activity directly to participants (90%) and in group-based sessions (51%). Most

physical activity interventions included structured exercise (30%) such as aerobic and/or resistance

training, physical recreation (26% combined) such as yoga, exer-gaming or walking, or mixed

activities (25%). Fewer studies considered sport- or dance-based interventions (7% and 6%,

respectively). The duration, frequency and intensity of sessions varied (see Table 2). Most

interventions were delivered by health and/or exercise professionals, but few studies assessed

interventions *for* staff. There was also very little evidence on environmental interventions or

interventions that provided participants with physical activity support after discharge from their

inpatient setting.

Table 6 summarises the themes of the findings from the 14 qualitative studies, which we grouped

under the headings of 'perceptions of the intervention' and 'factors influencing intervention delivery

and participation'. Beneficial psychosocial effects of interventions were reported in most studies

(79%), such as participants feeling more relaxed, happier and calmer. Several studies reported on a

range of personal and environmental factors that might influence participation in physical activity.

Examples of personal factors included perceived health benefits, social support, medication side

effects, confinement in a locked facility, and the availability of trained staff. Environmental factors

included physical space and facilities.

Table 7 summarises the themes of research recommendations from the studies of physical activity

interventions. Forty-one (33%) of the 125 studies included research recommendations, which we

grouped according to whether they were related to the research topic or research methods. The

most commonly reported research topic-related recommendation was to investigate other physical

activity interventions (11%), such as different modes of exercise or different behaviour change

strategies. The most commonly reported research methods-related recommendations were longer

follow-up periods (12%) and larger sample sizes (11%).

Table 6. Themes of findings arising from qualitative studies of physical activity interventions.

Theme	Examples of specific findings	Number (%) of articles out of 14
Experiences of the intervention		
Feasibility and acceptability	"Service users gave positive feedback regarding the programme" (Savage 2009) "The intervention was feasible as part of treatment" (Sorenson 2021)	6 (43%)
Physical impacts	"Improvements in physical function" (Wright 2009)	5 (36%)
Psychosocial impacts	"participants experienced interconnectedness and developed a mindful stance" (Schulze 2021) "Gaining a sense of competence" (Wright 2009) "more relaxed, happier and calmer after training sessions" (Savage 2009)	11 (79%)
Factors influencing intervention delivery and participation		
Barriers to participation	"medication side-effects" (Prebble 2011) "the locked environment" (Prebble 2011) "ill health" (Wright 2009) "low confidence" (Wright 2009) "perceived not physically able to undertake some activities" (Wright 2009) "low motivation" (Firth 2017)	6 (43%)
Facilitators of participation	"Enjoyment and some autonomy for service users" (Prebble 2011) "physical, mental and social benefits of participating" (Prebble 2011) "staff and patients exercising together" (Tetlie 2009) "exercise being a mandatory part of treatment" (Tetlie 2009) "instructor who is trained" (Tetlie 2009)	6 (43%)
Factors impeding implementation/delivery	"lack of funding" (Prebble 2011) "staff resistance due to exercise apathy in their own lives" (Prebble 2011) "lack of physical space and facilities" (Prebble 2011)	2 (14%)

 Tables 7. Themes of research recommendations from studies of physical activity interventions.

Theme	Examples of specific research recommendations	Number (%) of articles out of 125
Research topic-related recommendations		
Investigate other interventions/ intervention components	"Future interventions could use motivational interviewing" (Firth 2017) "the effect of different exercise modes" (Gerber 2018) "explore other ways to foster regular physical exercise" (Burge 2017)	14 (11%)
Investigate mechanisms of effect	"consider the impact of symptom severity as a moderator" (Stanton 2016a) "elucidate other ways mindful-yoga impacts patient well-being and treatment" (Spinelli 2020) "explore the mechanisms of improved functional outcomes" (Shimada 2019)	6 (5%)
Investigate other groups	"replicated in cohorts with affective disorders" (Stanton 2016a) "Further studies needed in women" (Areshtanab 2020)	6 (5%)
Investigate optimal exercise strategy	"better identify specific exercise types, sets, repetitions, routines" (Tomasi 2019) "clarifying the type and dose of physical activity" (Ng 2007) "determining the ideal window for exercise intervention" (Furzer 2021)	6 (5%)
Investigate other setting	"it might be better to explore music and exercise interventions in separate study settings to ease the realization of the interventions in a routine clinical setting" (Pitkänen 2019)	1 (1%)
Research methods-related recommendations		
Longer follow-up	"effect of inpatient self-selected exercise on affect and exercise adherence following discharge" (Stanton 2016a) "a longer follow-up period is needed" (Spinelli 2020)	15 (12%)
Larger sample sizes	"Further studies in larger sample sizes are warranted" (Methapatara 2011) "a larger sample size is required" (Shimada 2019)	14 (11%)
Investigate other outcomes	"consider including measures of affect and arousal" (Stanton 2016a) "other domains like cognitive performance, self-esteem, quality of life, stress and body image" (Roy 2018) "variables such as physiological measures should be included" (Shimada 2019)	10 (8%)
Use other data collection methods	"using objective measures of exercise intensity and sleep quality" (Stanton 2016a) "include more comprehensive quantitative measures" (Spinelli 2020) "using an extensive neuropsychological battery" (Shimada 2019)	9 (7%)

Another study including a control group	"using a quiet rest group or an alternate activity group as a comparator" (Stanton 2016a) "a controlled research design" (Spinelli 2020)	8 (6%)
Replications/ confirmatory studies	"More studies to confirm the findings" (Stanton 2016b) "should be replicated in further clinical trial settings" (Thaller 2022)	8 (6%)
Control for confounding variables	"Future studies would need to control for these variables as part of their analysis" (Loh 2015)	2 (2%)

Discussion

Summary of evidence

This scoping review shows that a wide range of research has been published regarding physical

activity and adult users of inpatient mental health services since 2007.

Forty-four quantitative studies explored correlates of physical activity. Organisational factors and

psychological, cognitive and emotional factors were the top two domains cited. Within these, four

out of the top five factors were organisational factors: staff capability, staff capacity, staff attitudes

to physical activity, and access to equipment, while self-motivation was the second most highly cited

correlate of physical activity overall. Similar findings were observed among the 22 qualitative studies

that reported on correlates, with the addition of social support. There was also more emphasis on

stigma, health, value of health and funding for physical activity programmes in the qualitative

studies. Overall, a broad range of correlates have been reported. Given that organisational

correlates were commonly cited, it seems prudent to consider ways to overcome some

organisational barriers in future research and clinical practice.

One hundred and twenty-five interventional studies were identified; only seven of these were large-

scale RCTs with 100 or more participants, and long-term follow-up was rare. We did not identify any

studies that included an economic evaluation, and there were few replications, suggesting that it

would be challenging to evaluate the clinical and cost-effectiveness of physical activity interventions

within the current evidence.

In terms of intervention design, the majority of studies explored an intervention that involved

delivering physical activity (80%) rather than promoting physical activity (6%); 10% used a mixed

model of both delivering and promoting physical activity. Interventions were mainly limited to the

period of inpatient stay, and there was little information on how the person might maintain their

levels of physical activity after discharge. The type of physical activity delivered was generally some

form of structured exercise or physical recreation targeting individuals. However, descriptions of the

interventions were often incomplete, and interventions appeared to lack systematic development.

There was a notable lack of research on environmental interventions and interventions targeting

healthcare professionals. In the current review, and elsewhere [35], we have observed that the

reporting of interventions is often inadequate. Since adequate reporting of interventions is central

to interpreting study findings and translating effective interventions into practice, we recommend

that researchers use relevant reporting guidelines, such as TIDieR [36], when writing study reports.

Research has been conducted in a wide range of countries, including both high-income and low- and

middle-income countries. However, the majority of studies were conducted in European psychiatric

inpatient settings. Given the fact that forensic settings are likely to have a more stable population, in

terms of patients' length of stay, than some other inpatient settings, where patients may stay only a

few days, surprisingly few RCTs have been conducted in forensic settings (one study).

Outcomes collected were mainly mental or physical health-related and the primary outcome was

often not clearly stated. In many studies more than one mental or physical health outcome was

examined, with some of the outcomes being positive and others neutral or negative, leading to

difficulty interpreting the study results. It is important that studies, especially RCTs, clearly state

their primary outcome from the outset to avoid cherry-picking results. Physical activity and health-

related quality of life were rarely included as outcomes; the lack of physical activity outcomes

suggests that increasing levels of physical activity was not the primary objective of many of the

studies.

Strengths and limitations

This review encompasses a wide range of articles and was conducted rigorously and systematically

following a predefined protocol and existing guidance for scoping reviews. It provides a

comprehensive overview of the extent and type of research on physical activity in inpatient mental

health services and identifies important gaps in the literature.

Although a systematic search was conducted, some eligible articles might have been missed. The

review is also limited to English language articles and does not include grey literature. Although a

scoping review was appropriate to meet the objectives of this study, it is subject to limitations that

are typical of this approach. For example, it did not permit a quality assessment of the included

studies. It was also limited in its capacity to conclude what factors affect participation in physical

activity among adult users of inpatient mental health services, what interventions are effective, and

what themes recur in the qualitative literature.

Recommendations for future research

A 2018 meta-review of the evidence on physical activity as a treatment for SMI made several

research recommendations [9], the following of which are relevant for research in relation to the

inpatient context:

- More research is needed to establish pragmatic, scalable methods for delivering physical

activity with the ultimate goal of optimising treatment 'reach'.

- More research is required to ascertain the optimal frequency, intensity, time and type of

physical activity interventions in each inpatient sub-population, acknowledging that these

might vary according to individual needs, preferences and characteristics.

- More research is needed to explore how specific individual, interpersonal, environmental,

and organisational factors influence the uptake and maintenance of physical activity

interventions when they are implemented in clinical practice.

- More research is needed to investigate whether sedentary behaviours can be reduced in

inpatients and tease out the importance of reducing sedentary behaviours in structured

exercise interventions.

- Economic evaluations are required to establish the cost-effectiveness of specific physical

activity interventions.

In addition, we believe that more research is needed to develop and test interventions that span the

transition from inpatient to community settings. For example, it would be useful to study longer-

term outcomes, such as the impact of physical activity programmes on rates of readmission to

inpatient mental health facilities. Finally, this scoping review has identified sets of articles that could

be collated for quantitative or qualitative systematic reviews of physical activity in inpatient mental

health services. Researchers conducting such reviews should consider the quality of the studies,

which was beyond the scope of this review, and to consider the variety of populations represented

in these studies.

Conclusions

In conclusion, this scoping review summarised the extent and type of research on physical activity in

adult users of inpatient mental health services. We identified a modest volume of evidence

regarding correlates of physical activity. A broad range of individual, interpersonal, environmental

and organisational correlates have been reported, but more studies with a longitudinal design are

needed to determine how these and other factors are associated. We also found a large number of

studies reporting on the outcomes of specific physical activity interventions. Most interventions

targeted individuals and involved the delivery of physical activity sessions rather than self-

management interventions. There were very few large-scale RCTs and most studies did not include

physical activity or quality of life outcomes. The findings of this review will help guide further

primary research that is needed to guide clinical practice and policy.

Supporting information

S1 File. Search strategy.

S2 File. Reference list of included reports.

References

1. Correll CU, Solmi M, Veronese N, Bortolato B, Rosson S, Santonastaso P, et al. Prevalence, incidence and mortality from cardiovascular disease in patients with pooled and specific severe mental illness: a large-scale meta-analysis of 3,211,768 patients and 113,383,368 controls. World Psychiatry. 2017;16: 163–180. doi:10.1002/wps.20420

2. Meier SM, Mattheisen M, Mors O, Mortensen PB, Laursen TM, Penninx BW. Increased mortality among people with anxiety disorders: total population study. British Journal of Psychiatry. 2016;209: 216–221. doi:10.1192/bjp.bp.115.171975

3. Archer G, Kuh D, Hotopf M, Stafford M, Richards M. Association Between Lifetime Affective Symptoms and Premature Mortality. JAMA Psychiatry. 2020;77: 806. doi:10.1001/jamapsychiatry.2020.0316

4. Kendler KS, Ohlsson H, Sundquist K, Sundquist J. Drug abuse-associated mortality across the lifespan: a population-based longitudinal cohort and co-relative analysis. Soc Psychiatry Psychiatr Epidemiol. 2017;52: 877–886. doi:10.1007/s00127-017-1398-5

5. Ringen PA, Falk RS, Antonsen B, Faerden A, Mamen A, Rognli EB, et al. Using motivational techniques to reduce cardiometabolic risk factors in long term psychiatric inpatients: a naturalistic interventional study. BMC Psychiatry. 2018;18: 255. doi:10.1186/s12888-018-1832-6

6. Vancampfort D, Firth J, Schuch FB, Rosenbaum S, Mugisha J, Hallgren M, et al. Sedentary behavior and physical activity levels in people with schizophrenia, bipolar disorder and major depressive disorder: a global systematic review and meta-analysis. World Psychiatry. 2017;16: 308–315. doi:10.1002/wps.20458

7. Katzmarzyk PT, Friedenreich C, Shiroma EJ, Lee I-M. Physical inactivity and non-communicable disease burden in low-income, middle-income and high-income countries. Br J Sports Med. 2022;56: 101–106. doi:10.1136/bjsports-2020-103640

8. Scheewe TW, Backx FJG, Takken T, Jörg F, van Strater ACP, Kroes AG, et al. Exercise therapy improves mental and physical health in schizophrenia: a randomised controlled trial. Acta Psychiatr Scand. 2013;127: 464–473. doi:10.1111/acps.12029

9. Stubbs B, Vancampfort D, Hallgren M, Firth J, Veronese N, Solmi M, et al. EPA guidance on physical activity as a treatment for severe mental illness: a meta-review of the evidence and

Position Statement from the European Psychiatric Association (EPA), supported by the
International Organization of Physical Therapists in Mental Health (IOPTMH). European
Psychiatry. 2018;54: 124–144. doi:10.1016/j.eurpsy.2018.07.004

10.   Friedrich B, Mason OJ. Qualitative evaluation of a football intervention for people with
mental health problems in the north east of London. Ment Health Phys Act. 2018;15: 132–
138. doi:10.1016/j.mhpa.2018.10.002

11.   Pearce M, Garcia L, Abbas A, Strain T, Schuch FB, Golubic R, et al. Association Between
Physical Activity and Risk of Depression. JAMA Psychiatry. 2022;79: 550.
doi:10.1001/jamapsychiatry.2022.0609

12.   Kruisdijk F, Deenik J, Tenback D, Tak E, Beekman A-J, van Harten P, et al. Accelerometer-
measured sedentary behaviour and physical activity of inpatients with severe mental illness.
Psychiatry Res. 2017;254: 67–74. doi:10.1016/j.psychres.2017.04.035

13.   Nyboe L, Lund H. Low levels of physical activity in patients with severe mental illness. Nord J
Psychiatry. 2013;67: 43–46. doi:10.3109/08039488.2012.675588

14.   Martland R, Korman N, Firth J, Stubbs B. The efficacy of exercise interventions for all types of
inpatients across mental health settings: A systematic review and meta-analysis of 47 studies.
J Sports Sci. 2023;41: 232–271. doi:10.1080/02640414.2023.2207855

15.   Soundy A, Stubbs B, Probst M, Hemmings L, Vancampfort D. Barriers to and Facilitators of
Physical Activity Among Persons With Schizophrenia: A Survey of Physical Therapists.
Psychiatric Services. 2014;65: 693–696. doi:10.1176/appi.ps.201300276

16.   Long C, Rowell A, Gayton A, Hodgson E, Dolley O. Tackling obesity and its complications in
secure settings. Mental Health Review Journal. 2014;19: 37–46. doi:10.1108/MHRJ-04-2013-
0012

17.   Mateo-Urdiales A, Michael M, Simpson C, Beenstock J. Evaluation of a participatory approach
to improve healthy eating and physical activity in a secure mental health setting. J Public
Ment Health. 2020;19: 301–309. doi:10.1108/JPMH-11-2019-0090

18.   Firth J, Rosenbaum S, Stubbs B, Gorczynski P, Yung AR, Vancampfort D. Motivating factors
and barriers towards exercise in severe mental illness: a systematic review and meta-analysis.
Psychol Med. 2016;46: 2869–2881. doi:10.1017/S0033291716001732

19.   Gilburt H, Rose D, Slade M. The importance of relationships in mental health care: A
qualitative study of service users' experiences of psychiatric hospital admission in the UK.
BMC Health Serv Res. 2008;8: 92. doi:10.1186/1472-6963-8-92

20.   Hassan J, Shannon S, Tully MA, McCartan C, Davidson G, Bunn R, et al. Systematic review of
physical activity interventions assessing physical and mental health outcomes on patients
with severe mental illness (SMI) within secure forensic settings. J Psychiatr Ment Health Nurs.
2022;29: 630–646. doi:10.1111/jpm.12832

21.   Rogers E, Kinnafick F-E, Papathomas A. Physical activity in secure settings: A scoping review of
methods, theory and practise. Ment Health Phys Act. 2019;16: 80–95.
doi:10.1016/j.mhpa.2018.11.004

22.   Munn Z, Peters MDJ, Stern C, Tufanaru C, McArthur A, Aromataris E. Systematic review or
427   scoping review? Guidance for authors when choosing between a systematic or scoping
428   review approach. BMC Med Res Methodol. 2018;18: 143. doi:10.1186/s12874-018-0611-x

429   23.   Peters MDJ, Godfrey C, McInerney P, Munn Z, Tricco AC, Khalil H. Chapter 11: Scoping
430   Reviews (2020 version). . In: JBI Manual for Evidence Synthesis. 2020.

431   24.   Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA Extension for
432   Scoping Reviews (PRISMA-ScR): Checklist and Explanation. Ann Intern Med. 2018;169: 467–
433   473. doi:10.7326/M18-0850

434   25.   Tew G, Peckham E, Ker S, Smith J, Hodgson P, Daniel S. Physical activity in adult users of
435   inpatient mental health services: a scoping review. In: https://osf.io/pj6uw/. 2022.

436   26.   Caspersen CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical fitness:
437   definitions and distinctions for health-related research. Public Health Rep. 1985;100: 126–31.

438   27.   Veritas Health Innovation. Covidence systematic review software. Melbourne, Australia;
439   2022.

440   28.   JBI template source of evidence details, characteristics and results extraction instrument. In:
441   https://jbi-global-wiki.refined.site/space/MANUAL/4687579. 20 Sep 2022.

442   29.   Levac D, Colquhoun H, O'Brien KK. Scoping studies: advancing the methodology.
443   Implementation Science. 2010;5: 69. doi:10.1186/1748-5908-5-69

444   30.   Campbell F, Tricco AC, Munn Z, Pollock D, Saran A, Sutton A, et al. Mapping reviews, scoping
445   reviews, and evidence and gap maps (EGMs): the same but different — the "Big Picture"
446   review family. Syst Rev. 2023;12: 45. doi:10.1186/s13643-023-02178-5

447   31.   Golden SD, Earp JAL. Social Ecological Approaches to Individuals and Their Contexts. Health
448   Education & Behavior. 2012;39: 364–372. doi:10.1177/1090198111418634

449   32.   Bauman AE, Sallis JF, Dzewaltowski DA, Owen N. Toward a better understanding of the
450   influences on physical activity. Am J Prev Med. 2002;23: 5–14. doi:10.1016/S0749-
451   3797(02)00469-5

452   33.   Choi J, Lee M, Lee J, Kang D, Choi J-Y. Correlates associated with participation in physical
453   activity among adults: a systematic review of reviews and update. BMC Public Health.
454   2017;17: 356. doi:10.1186/s12889-017-4255-2

455   34.   Campbell M, McKenzie JE, Sowden A, Katikireddi SV, Brennan SE, Ellis S, et al. Synthesis
456   without meta-analysis (SWiM) in systematic reviews: reporting guideline. BMJ. 2020; l6890.
457   doi:10.1136/bmj.l6890

458   35.   Peckham E, Tew G, Lorimer B, Bailey L, Beeken R, Cooper C, et al. Interventions to increase
459   physical activity and reduce sedentary behaviour in severe mental ill health: How effective
460   are they?' - A systematic review. Ment Health Phys Act. 2023;25: 100547.
461   doi:10.1016/j.mhpa.2023.100547

462   36.   Hoffmann TC, Glasziou PP, Boutron I, Milne R, Perera R, Moher D, et al. Better reporting of
463   interventions: template for intervention description and replication (TIDieR) checklist and
464   guide. BMJ. 2014;348: g1687–g1687. doi:10.1136/bmj.g1687