1

Title

Maternal risk factors contributing to stillbirths in semi-rural Ghana: Insights from

a cross-sectional study.

Author's information

* **Silas Adjei-Gyamfi, MPH, MPhil (Corresponding author)**

Savelugu Municipal Hospital

Ghana Health Service

P. O. Box 45, Savelugu, Northern Region, GHANA

Phone: +233243421318

Email: adjeigyamfis@yahoo.com

ORCID: https://orcid.org/0009-0008-8496-4099

Abigail Asirifi, Dip, BSc

Department of Midwifery

Garden City University College

P. O. Box KS 12775, Kumasi, Ashanti Region, GHANA

Phone: +233545951777

Email: abimaiden80@gmail.com

ORCID: https://orcid.org/0009-0008-8133-4104

Gafaru Mamudu, BSc

Savelugu Municipal Hospital

Ghana Health Service

P. O. Box 45, Savelugu, Northern Region, GHANA

Phone: +233248303028

Email: gafarumamudu@gmail.com

Wisdom Peprah, BSc, MPhil

School of Allied Health Sciences

University for Development Studies

P. O. Box TL 1883, Tamale, Northern Region, GHANA

Phone: +233507093898

Email: pwisdom466@gmail.com

ORCID: https://orcid.org/0000-0003-0263-8158

Clotilda Asobuno, BSc, MPH

Kassena Nankana West District Health Directorate

Ghana Health Service

P. O. Box 21, Paga, Upper East Region, GHANA

Phone: +233245452488

Email: clotildaasobuno@gmail.com

ORCID: https://orcid.org/0009-0008-7471-6462

Constance Siakwan Dapaah, BSc

Ghana Health Service

Atwima Nwabiagya North District Health Directorate

P.O. Box 26, Barekese, Ashanti Region, GHANA

Phone: +233245727275

Email: constancedapaah25@gmail.com

Paul Armah Aryee, MMedSc, MPhil, PhD

School of Allied Health Sciences

University for Development Studies

P. O. Box TL 1883, Tamale, Northern Region, GHANA

Phone: +233200516610

Email: parmah@uds.edu.gh

ORCID: https://orcid.org/0000-0002-9857-1842

Abstract

Background

Stillbirth remains a critical public health challenge in low- and middle-income countries,

impeding progress toward achieving the Sustainable Development Goals. Given the diverse

dynamics of the prevalence and associated factors of stillbirths in various low-resource settings,

such as Ghana, it is essential to have comprehensive knowledge to address and mitigate these

factors that could influence stillbirths. However, there is a paucity of data on maternal factors

contributing to stillbirths in semi-rural areas of Ghana.

Objective

This study aimed to determine the prevalence and associated maternal factors of stillbirths in

semi-rural Ghana.

Design

This is a cross-sectional study that utilised a quantitative approach.

Methods

The study was conducted among 422 mothers who delivered in public health facilities and

reside in the Savelugu Municipality of Northern Ghana. Sociodemographic, obstetric, and

medical-related data were gathered through structured interviews and from record reviews.

Chi-square test of proportions and multiple logistic regression analyses were employed to

identify the independent factors associated with stillbirths at a 95% confidence level.

Results

Stillbirth rate was estimated at 4.0% (95%CI: 2.4%–6.4%). The analysis revealed that anaemia in

the third trimester of pregnancy (aOR:3.15; 95%CI:2.94–6.10; p=0.034), positive sickle cell

status (aOR:2.91; 95%CI:2.50–5.01; p=0.018), and non-utilisation of insecticide-treated bed nets

(ITNs) (aOR:7.75; 95%CI:4.33–8.80; p=0.001) during pregnancy were significantly associated

with increased risk of stillbirths.

Conclusion

The relatively high prevalence of stillbirths in the municipality was influenced by medical factors

(particularly anaemia and positive sickle cell status) and unsafe health practices via the non-use

of ITNs during pregnancy. The management of the Ghana Health Service, alongside program

planners and other stakeholders involved in enhancing perinatal and maternal health, should

focus on implementing effective initiatives such as improved health education on nutrition and

safety practices at the facility and community levels, as well as robust referral systems to

prevent and reduce stillbirths in semi-rural Ghana.

Plain language summary

Stillbirth, which is the death of a baby occuring before or during childbirth, not only remain a public health issue but also cause emotional distress for mothers in most African countries. Despite the importance of understanding the maternal factors of stillbirths, there is limited data available, especially in semi-rural areas of Ghana. A study was therefore undertaken to assess the prevalence and maternal factors associated with stillbirths in semi-rural Ghana. This study involved women who gave birth in public health facilities and live in the Savelugu Municipal area. Researchers collected data through structured interviews and from medical records. The data were analysed using chi-square tests and logistic regression to identify maternal factors linked to stillbirths. The findings showed a stillbirth rate of 4.0%. Key factors associated with stillbirths included maternal anaemia in the third trimester of pregnancy, positive sickle cell status, and not using insecticide-treated bed nets (ITNs) during pregnancy. The study highlights the need for improved health education on nutrition and safe health practices to reduce and/or prevent stillbirths. Priority should be given to managing medical conditions such as anaemia in pregnancy and sickle cell disease, alongside promoting the use of ITNs to safeguard pregnant women against malaria and subsequently reduce the risk of stillbirths.

Keywords

Anaemia,

Risk factors,

Northern Ghana,

Predictors,

Stillbirths,

Stillborns

Number of words

147 Abstract   301 words

148 Main text   3,939 words

Introduction

Stillbirth remains a critical public health challenge in low- and middle-income countries

(LMICs), impeding progress toward achieving the Sustainable Development Goals (1–3). Defined

by the World Health Organisation (WHO), stillbirth is denoted as the birth of a baby with no

signs of life occurring at 28 weeks or more of gestation (3), and it accounts for more than half of

all perinatal and maternal mortalities worldwide (4). Stillbirths, also called stillborns, trigger

maternal psychological trauma which could affect the socio-economic growth and development

in LMICs (1–3).

The aetiology and antecedents of stillbirths are multifactorial and diverse, influenced by

sociological, physiological (medical), and physical factors (1). It is reported that the stillbirth rate

and its determinants are dependent on the geographic and sociodemographic distribution of

the population (3,5). Therefore, the study findings from one setting might not apply to another.

Universally, earlier studies acknowledged that the factors associated with stillbirths included

congenital foetal deformities, foetal distress, preterm births, breech delivery, abruptio

placentae, and male foetus (6–9). In most LMICs, pregnancy-related smoking and alcohol

intake, maternal low socioeconomic status, multiparity, non-use of insecticide-treated bed nets

(ITNs) for malaria prevention during pregnancy, maternal age less than 20 years, unmarried

mothers, employment status, and delayed vaginal delivery were also regarded as some

determinants of stillbirths in other past studies (9–13). In other low-resource settings,

inadequate knowledge about antenatal care (ANC) services, cost, distance and the time needed

to access care are major barriers to the effective uptake of quality ANC and particularly

intrapartum services, which contributes significantly to stillbirths (14,15).

On a global scale, an estimated 2·6 million babies were born with no signs of life in 2015,

which reduced to about two million stillborns (13.9 per 1000 births) in 2019 (5,16). LMICs

particularly, sub-Saharan Africa (SSA) has the highest stillbirth rate globally, estimated at 21.7

per 1000 births, compared to 3.0 for high-income countries (HICs) (16) and accounts for more

than one-third of the global burden of stillbirths (5,16). Furthermore, stillbirth rates vary widely

across regions in the world, from 22·8 per 1000 births in Western and Central Africa to 2·9 in

Western Europe. After West and Central Africa, Eastern and Southern Africa and South Asia

have the second (20.5 per 1000 births) and third-highest (18.2 per 1000 births) stillbirth rates

(16). Estimates of stillbirths in China, South Africa, and Nigeria are found to be 9.0, 35.0 and

40.0 per 1000 births respectively (10,11,14,17). Hug et al. 2021 observed a 2·3% annual decline

in the stillbirth rate globally, with a sluggish decline rate in most Asian and SSA countries (16).

These findings may hinder SSA countries from reaching the Every Newborn Action Plan target of

12 per 1000 births by 2030.

The vast differences in stillbirth rates between HICs and LMICs have led to stillbirth

being equally regarded as a global development metric. Similar to many other SSA nations,

Ghana faces challenges in the routine and adequate monitoring and recording of stillbirths

(5,18). Diverse surveys conducted in Ghana estimate the stillbirth rate to fluctuate between

approximately 14 and 22 per 1000 births (18,19). Also, some population health surveillance

and health facility data from different parts of the country indicate even higher rates. For

instance, the Ghana Maternal Health Survey in 2017 observed that 68% of stillbirths occur in

rural areas compared to urban areas (18). Other findings also revealed 34 stillbirths/1000 births

in Navrongo of the Upper East Region, 59 stillbirths/1000 births in the Ashanti Region and 109

stillbirths/1000 births in a tertiary hospital in the Greater Accra Region (7,20,21), which makes

these high stillbirth occurrences very worrisome in the country. While these longitudinal

studies reported a wide range of health facility, neonatal, paternal and maternal factors, none

have exclusively delved into the broad spectrum of maternal factors that contribute to

stillbirths. Additionally, most of these longitudinal studies were conducted in rural and urban

areas. Hence, a cross-sectional study to provide a snapshot of the current situation in semi-rural

communities is non-existent. Again, to date, little is particularly investigated about maternal

factors contributing to stillbirths in semi-rural Ghana.

On the other hand, epidemiological studies evaluating stillbirths frequently utilise the

concept of perinatal deaths, which encompasses both stillbirths and early neonatal deaths

(5,16). These studies often rely on population health surveillance, community-based oral post-

mortem reports and intricate analytical modelling techniques. However, the utility of these

estimates is constrained by the etiological distinctions between stillbirths and early neonatal

deaths, particularly given that the intrapartum period represents the highest-risk phase for

stillbirths. To achieve the World Health Assembly's national target of a stillbirth prevalence of

12 or less per 1000 births by 2030 in every country through the Every Newborn Action Plan (3),

the maternal factors of stillbirths and their dynamics in different settings or context must be

well understood and appreciated. Although stillbirths are higher in LMICs, including Ghana

(7,20), relevant information on the maternal factors of stillbirths is inadequately and incorrectly

documented in the district and/or regional databank. Hence, adequate evidence of information

on these factors in semi-rural Ghana could trigger useful interventions that may lead to

preventable stillbirths and subsequently contribute to improving maternal and perinatal health.

The purpose of the study is to estimate the prevalence of stillbirths and examine the

maternal factors associated with stillbirths in a semi-rural municipality in the Northern Region

of Ghana. We examined the effect of maternal risk factors on stillbirths by utilising a broad

spectrum of sociodemographic, medical or clinical and obstetric or antenatal characteristics of

the mothers. Given the distinct and unique nature of this study, the findings will be crucial to

inform clinical practice through timely interventions, decision-making, and policy ideation, with

a particular emphasis on maternal factors in the semi-rural Ghana and SSA.

Materials and methods

Study design

This current survey is a cross-sectional study conducted among women who delivered in a semi-rural municipality in the Northern Region of Ghana. The design was employed to determine the prevalence and associated maternal factors of stillbirths. The study followed the STROBE checklist (22,23) of reporting (see Supplementary 1).

Study setting

This study was conducted in Savelugu Municipality of Northern Ghana. Savelugu Municipality is one of the 16 districts/municipalities in the Northern Region of Ghana that shares boundaries with five neighbouring districts, including Gushegu, Karaga, Kumbungu, Nanton, and Tolon. The municipality has five major government (public) health facilities (one hospital and four health centres) that provide comprehensive maternal and newborn services, including free maternal and delivery services (24) to a population of approximately 140,000. In 2019, the coverage of ANC and postnatal care visits was 94% and 98%, respectively (25,26). Due to the increased average monthly delivery count of over 300 deliveries with higher records of unfavourable birth outcomes (25,27,28), this municipality was selected to investigate the factors of stillbirths among women who delivered in government health facilities.

Sample size estimation

The sample size (n) was estimated using Cochran's method with the WHO conservative design

effect (DEF), $n = \frac{Z^2 PQ}{e^2}$ X DEF (29–31). A standard variate (Z) at a 95% confidence interval of 1.96, a

standard error (e) of 5%, a prevalence (P) of stillbirths estimated at 21% in Northern Ghana

(32), Q=1–P and WHO standard conservative DEF size of 1.5 (29,30), was applied to estimate

the preliminary sample size at 384. An additional 10 percent was added to the calculated

sample size to account for fall-outs, incomplete responses and damaged questionnaires

(25,27,33) resulting in a final total sample of approximately 422.

Study population

The study recruited a total of 422 women who visited the five major public facilities in the

Savelugu Municipal for childbirth or delivery purposes from January 1 to June 31, 2020. All

women who lived in the municipality, delivered at the municipal health facilities and had their

health information recorded in the delivery registers were eligible for the study.

Sampling

In the sampling stage, all five major government (public) health facilities were chosen for the

study since all these facilities report on stillbirths every year (34). The sample sizes for each

facility were determined using the probability proportional to size approach within a stratified

random sampling framework (see **Table 1**). The delivery registers from each facility were used

to line-list all eligible postpartum women (who delivered from January 1 to June 31, 2020) to

form a sampling frame. A simple random sampling technique was then employed to select the

required number of respondents from each facility. Specifically, a lottery method without

replacement was utilised to ensure that each participant was selected only once.

The addresses and contact numbers of these selected respondents were recorded to

trace them to their homes for the first part of data collection. Following home-based data

collection, the information of recruited participants were re-traced to the delivery registers to

complete the data collection at the designated health facilities. This approach was utilised to

systematically reduce social desirability bias among the data collectors.

Table 1: Description of proportionate sample size for study health facilities

	Savelugu Municipal Hospital	Savelugu Health Centre	Pong Tamale Health Centre	Moglaa Health Centre	Diare Health Centre	Total
Total half-year facility deliveries in 2020 (X)	1,594	12	103	61	210	1,980
Facility coverage (Y: X/1,980)	80.5%	0.6%	5.2%	3.1%	10.6%	100%
Approximate sample size (Z: Y x 422)	339	3	22	13	45	422

Data collection

Data were collected using a pretested questionnaire by six well-trained data collectors

(midwives and public health officers) from July 16 to September 31, 2020. The data collection

was conducted during the period of COVID-19 pandemic, hence COVI-19 protocols were

followed. The questionnaire was developed based on existing literature, previous studies, and

sociological and biological plausibility (1,5,25,35). We verified the content validity of the

questionnaire by consulting with experts in the field as well as the study authors (see

Supplementary 2). Sociodemographic, obstetric or antenatal and medical-related information

was collected through structured interviews and from record reviews (thus, delivery registers

and maternal and child health record books [MCHRBs]). The data collectors were trained for

one week on study purpose and background, sampling techniques, data collection procedures,

ethical procedures and consent process among others. The data collection procedure was

conducted in two phases. The first phase ran through a period of about two months and was

conducted at the household level after sampling and contacting the eligible study participants.

The second phase of the data collection lasted for about one month and was conducted at the

health facilities level immediately after household-based data collection. This approach was

relevant to reduce social desirability bias among the data collectors. It also mitigated selection

bias among recruited participants by accounting for the non-attendance of health facilities by

mothers who had experienced stillbirths.

Before the commencement of household data collection, the trained data collectors

retrieved the addresses and contact information (telephone numbers) of eligible respondents

and/or their representatives from the delivery registers with the support of the facility

midwives. The data collectors arranged appointments and then visited the homes of the

recruited respondents. Data such as age, marital status, education level, job status, ethnicity

and religious affiliations were collected via structured interviews and were confirmed or

validated in the MCHRBs. Information about first-trimester body mass index (since the foetal

weight in the first trimester is negligible (36), antenatal visits, parity, gravidity, rhesus status,

sickle cell status, tetanus-diphtheria (TD) immunisation status, haemoglobin levels across

pregnancy trimesters, use of pre-pregnancy family planning, and utilisation of ITNs (1,28) were

collected from their MCHRBs at their homes. Haemoglobin electrophoresis was not further

conducted to confirm sickle cell variants by the health facilities. Following completion of the

home-based data collection, information like pre-eclampsia diagnosis, postpartum

haemorrhage, delivery mode, gestational age at birth and birth/delivery outcome prognosis

(stillbirth or livebirth) based on the classification by health professionals (1,28) were

immediately traced and extracted from the delivery registers at the health facilities level.

Study variables

The outcome variable was birth outcome prognosis (stillbirths), categorised into binary

outcomes for stillbirths (coded as 1) or livebirths (coded as 0), whereas exposure variables

included all sociodemographic, obstetric or antenatal, and medical variables of the women,

which were all categorised based on their normal distribution and existing studies (1,5,25,35).

According to the WHO, stillbirths were measured as the birth of a foetus at 28 or more

weeks of gestation with no signs of life, while livebirths were regarded as newborns with signs

of life (3). Haemoglobin levels were dichotomised into anaemia (< 11g/dl) and no anaemia (≥

11g/dl). Gestational age at birth was grouped into < 37 weeks (preterm) and ≥ 37 weeks

(normal-term). Parity was classified as primiparity (one delivery) and multiparity (two or more

deliveries). Gravidity was grouped into primigravida (one pregnancy) and multigravida (two or

more pregnancies) (26–28,33). Standardised WHO categories of pre-pregnancy body mass

index (BMI) of underweight (<18.5 kg/m^2), normal BMI (18.5–24.9 kg/m^2) and overweight

(≥25.0 kg/m^2) were utilised. Sickle cell and rhesus statuses were grouped as positive or

negative, whereas ANC visits were also categorised into ≥ 8 visits and < 8 visits using the new

2016 WHO antenatal model which are also applicable to the Ghanaian health system (25,35).

Statistical analysis

Frequencies and percentages were calculated for the sociodemographic, obstetric or antenatal

and medical variables. Chi-square/Fisher's analysis tests of proportions were employed to

determine the statistical association between birth outcome prognosis (stillbirth) and all

exposure variables at the bivariate level. These analyses were confirmed with univariate logistic

regression. Multicollinearity among the significant exposure variables was then assessed with

variance inflation factor (VIF), where variables with a VIF less than five were selected into

multiple (binary) logistic regression analyses (25,26,28).

Applying simultaneous variable entry, the significant exposure variables with no

multicollinearity issues were forwarded into the multiple logistic regression model and run to

determine significant factors of stillbirths. All analyses were computed using STATA 17.0 (Stata

Corporation, Texas, USA), and a p-value of less than 0.05 was considered significant during

statistical analysis.

                                    Results

358   **Prevalence of stillbirths**

359   The prevalence of stillbirths in the semi-rural municipality of Ghana is presented in Figure 1. All

360   the data from the sampled 422 mothers were used for the analysis. The prevalence of stillbirths

361   in this semi-rural municipality was estimated at 4.0% (95%CI: 2.4% − 6.4%).

362
363   **Figure 1. Prevalence of stillbirths in a semi-rural municipality of Ghana**

Sociodemographic characteristics of study participants

The frequency distribution of sociodemographic characteristics of study participants are

illustrated in Table 2. The average age of the mothers was 27.6 (sd=6.0) years, while nine

percent of all the mothers were teenagers. A large proportion of the participants could not read

or write (no formal education) (43%) and were not employed (64%). Furthermore, more than

two-thirds of the mothers were from the Dagomba ethnic group (81%) and were married (93%).

Finally, Islam was the most practised religion by the study participants (88%).

Table 2: Sociodemographic characteristics of study participants

Variables	Frequency (N)	Proportion (%)
Age category of mother		
< 20 years	38	9.0
20 – 35years	340	80.6
> 35 years	44	10.4
Mean = 27.63 (SD=6.02)		
Marital status of mother		
Married	391	92.7
Not married	31	7.3
Education level of mother		
No education	182	43.1
Primary school	58	13.8
High school	138	32.7
Tertiary	44	10.4
Ethnicity of mother		
Dagomba	340	80.6
Gonja	22	5.2
Frafra/Grusi	34	8.1
Others (Asante, Ewe, Nigeria)	26	6.1
Religion of mother		
Muslim	372	88.2
Christian	50	11.8
Occupation of mother		
Unemployed	269	63.7
Employed	153	36.3
Sex of baby (foetus)		
Male	210	49.8
Female	212	50.2

Obstetric and medical characteristics of study participants

The frequency distribution of medical and obstetric (antenatal) characteristics of respondents

are shown in Table 3. Maternal anaemia was highest in the second (71%), followed by the first

(64%) and third (45%) trimesters of pregnancy. Few of the mothers attended antenatal clinic at

least eight times (28%), ingested sulphadoxine-pyrimethamine at least three times (36%), and

did not use insecticide-treated bed nets (ITNs) during pregnancy (36%). Nearly three-quarters

of the mothers were multiparous (71%) and multigravidae (74%). Additionally, slightly above a

third of the mothers tested positive for sickle cell (38%), whereas almost 90% had positive

blood rhesus status. A small proportion of the mothers had pre-eclampsia/eclampsia (2%),

were diagnosed with antepartum haemorrhage (7%), were immunised against tetanus-

diphtheria (7%), and had glucose-6-phosphate dehydrogenase (G6PD) defect (27%) during the

period of pregnancy.

Table 3: Medical and obstetric characteristics of study participants

Variables	Frequency (N)	Proportion (%)
First-trimester body mass index (BMI)		
Underweight	13	3.1
Normal BMI	279	66.1
Overweight	130	30.8
Number of antenatal care visits		
< 8 visits	304	72.0
≥ 8 visits	118	28.0
Sulphadoxine-pyrimethamine intake		
None	25	5.9
1 – 3 doses	247	58.6
> 3 doses	150	35.5
Pre-eclampsia/Eclampsia		
Yes	9	2.1
No	413	97.9
Antepartum haemorrhage		
Yes	34	8.1
No	388	91.9
Pre-pregnancy family planning use		
Yes	75	17.8
No	347	82.2
Insecticide-treated bed nets use during pregnancy		
Yes	270	64.0
No	152	36.0
Malaria episode during pregnancy		
Yes	59	14.0
No	363	86.0
Gravidity		
Primigravidae	112	26.5
Multigravidae	310	73.5

Parity		
Primiparity	122	28.9
Multiparity	300	71.1
Mode of delivery		
Vaginal delivery	379	89.8
Caesarean section	43	10.2
Gestational age at delivery		
< 37 weeks (preterm)	91	21.6
≥ 37 weeks (normal term)	331	78.4
First-trimester anaemia		
Anaemia	268	63.5
No anaemia	154	36.5
Second-trimester anaemia		
Anaemia	301	71.3
No anaemia	121	28.7
Third-trimester anaemia		
Anaemia	191	45.2
No anaemia	231	54.8
Blood rhesus type		
Positive	379	89.8
Negative	43	10.2
Sickle cell status		
Positive	159	37.8
Negative	263	62.2
G6PD status		
Normal	309	73.2
Complete/partial defect	113	26.8
TD immunization		
Immunized	31	7.4
Not immunized	391	92.6

G6PD: Glucose-6-phosphate dehydrogenase   TD: Tetanus-diphtheria

Bivariate association of stillbirths with background variables

The bivariate association between stillbirths and the background variables are exhibited in

Table 4. Among the sociodemographic, medical and obstetric or antenatal characteristics

(variables) of the study participants, only sulphadoxine-pyrimethamine intake (p=0.012),

insecticide-treated bed net use (p<0.001), parity (p=0.035), third-trimester anaemia status

(p=0.030) and sickle cell status (p=0.003) were significantly associated with birth outcome

prognosis (stillbirths) during bivariable analysis.

Table 4: Bivariate association of stillbirths with background variables

Variables	Bivariate analysis		
	Stillbirths (%)	Livebirths (%)	p-value [a]
Sociodemographic variables			
Age category of mother			
< 20 years	1 (5.9)	37 (9.2)	0.573
20 – 35years	13 (76.5)	327 (80.7)	
> 35 years	3 (17.6)	41 (10.1)	
Mean = 27.63 (SD=6.02)			
Marital status of mother			
Married	15 (88.2)	376 (92.8)	1.000
Not married	2 (11.8)	29 (7.2)	
Education level of mother			
No education	7 (41.2)	175 (43.2)	0.314
Primary school	1 (5.9)	57 (14.1)	
High school	9 (52.9)	129 (31.8)	
Tertiary	0 (0.0)	44 (10.9)	
Ethnicity of mother			
Dagomba	13 (76.5)	327 (80.7)	0.354
Gonja	0 (0.0)	22 (5.4)	
Frafra/Grusi	3 (17.7)	31 (7.7)	
Others (Asante, Ewe, Nigeria)	1 (5.9)	25 (6.2)	
Religion of mother			
Muslim	13 (76.5)	359 (88.6)	0.158
Christian	4 (23.5)	46 (11.4)	
Occupation of mother			
Unemployed	16 (94.1)	253 (62.5)	0.303
Employed	1 (5.9)	152 (37.5)	
Sex of baby (foetus)			
Male	6 (35.3)	204 (50.4)	0.322
Female	11 (64.7)	201 (49.6)	
Medical and obstetric variables			
First-trimester body mass index (BMI)			
Underweight	1 (5.9)	12 (3.2)	0.423
Normal BMI	12 (64.7)	267 (65.9)	
Overweight	5 (29.4)	125 (30.9)	
Number of antenatal care visits			
< 8 visits	14 (82.3)	290 (71.6)	0.333
≥ 8 visits	3 (17.7)	115 (28.4)	
Sulphadoxine-pyrimethamine intake			
None	2 (11.8)	23 (5.7)	**0.012**[*]
1 – 3 doses	12 (70.6)	235 (58.0)	
> 3 doses	3 (17.6)	147 (36.3)	
Pre-eclampsia/Eclampsia			

Yes	2 (11.8)	7 (1.7)	0.768
No	15 (88.2)	398 (98.3)	
Antepartum haemorrhage			
Yes	4 (23.5)	30 (7.4)	0.622
No	13 (76.5)	375 (92.6)	
Pre-pregnancy family planning use			
Yes	3 (17.7)	72 (17.8)	0.989
No	14 (82.3)	333 (82.2)	
Insecticide-treated bed nets use during pregnancy			
Yes	4 (23.5)	266 (65.7)	**< 0.001**[*]
No	13 (76.5)	139 (34.3)	
Malaria episode during pregnancy			
Yes	1 (5.9)	58 (14.3)	0.326
No	16 (94.1)	347 (85.7)	
Gravidity			
Primigravidae	2 (11.8)	110 (27.2)	0.051
Multigravidae	15 (88.2)	295 (72.8)	
Parity			
Primiparity	3 (17.7)	119 (29.4)	**0.035**[*]
Multiparity	14 (82.3)	286 (70.6)	
Mode of delivery			
Vaginal delivery	15 (88.2)	364 (89.9)	0.827
Caesarean section	2 (11.8)	41 (10.1)	
Gestational age at delivery			
< 37 weeks (preterm)	4 (23.5)	87 (21.5)	0.822
≥ 37 weeks (normal term)	13 (76.5)	318 (78.5)	
First-trimester anaemia			
Anaemia	11 (64.7)	257 (63.4)	0.554
No anaemia	6 (35.3)	148 (36.6)	
Second-trimester anaemia			
Anaemia	13 (76.5)	288 (71.1)	0.700
No anaemia	4 (23.5)	117 (28.9)	
Third-trimester anaemia			
Anaemia	13 (76.5)	178 (44.0)	**0.030**[*]
No anaemia	4 (23.5)	227 (55.0)	
Blood rhesus type			
Positive	15 (88.2)	364 (89.9)	0.455
Negative	2 (11.8)	41 (10.1)	
Sickle cell status			
Positive	7 (41.2)	152 (37.5)	**0.003**[*]
Negative	10 (58.8)	253 (62.5)	
G6PD status			
Normal	14 (82.4)	295 (72.8)	0.732
Complete/partial defect	3 (17.6)	110 (27.2)	
TD immunization			
Immunized	3 (17.7)	28 (6.9)	0.121
Not immunized	14 (82.3)	377 (93.1)	

397   [*] *p < 0.05*   [a] *Chi-square/Fisher's exact test*   *G6PD: Glucose-6-phosphate dehydrogenase*   *TD: Tetanus-diphtheria*

Maternal factors associated with stillbirths

The maternal factors associated with stillbirths are presented in Table 5. In the multiple logistic regression analysis, three variables particularly, insecticide-treated bed net use, third-trimester anaemia status and sickle cell status, remained statistically significant in the final model. The model revealed that the odds of mothers with anaemia in the third trimester of pregnancy having stillbirths was 3.15 times (95%CI: 2.94 − 6.10; p=0.034) compared to the non-anaemic mothers. Moreover, mothers with positive sickle cell status had 2.91 (95%CI: 2.50 − 5.01; p=0.018) increased odds of stillbirths compared to their counterparts with negative sickle cell status. Finally, participants who did not sleep under insecticide-treated bed nets throughout pregnancy had 7.75 (95%CI: 4.33 − 8.80; p=0.001) higher odds of stillbirths than those who used it.

Table 5: Multiple logistic regression of maternal factors of stillbirths

Variables	Logistic regression analyses			
	Univariate analysis		Multivariate analysis	
	cOR (95%CI)	p-value	aOR (95%CI)	p-value
Sulphadoxine-pyrimethamine intake				
1 − 3 doses		1.000		1.000
> 3 doses	0.11 (0.15 − 0.88)	0.037[*]	0.17 (0.02 − 1.35)	0.094
None	1.46 (0.31 − 6.82)	0.631	2.68 (0.41 − 7.13)	0.301
Insecticide-treated bed nets use				
Yes		1.000		1.000
No	6.22 (3.99 − 8.40)	0.002[*]	**7.75 (4.33 − 8.80)**	**0.001**[*]
Parity				
Primiparity		1.000		1.000
Multiparity	6.66 (3.87 − 7.70)	0.047[*]	4.64 (0.57 − 7.60)	0.151
Third-trimester anaemia				
No anaemia		1.000		1.000
Anaemia	4.02 (2.29 − 5.12)	0.017[*]	**3.15 (2.94 − 6.10)**	**0.034**[*]
Sickle cell status				

Negative		1.000		1.000
Positive	1.51 (1.10 − 6.41)	0.019[*]	**2.91 (2.50 − 5.01)**	**0.018[*]**
Regression model				
R^2		*0.6030*		
p value		*< 0.001*		

[*] *p < 0.05*   *cOR: crude odds ratio*   *aOR: adjusted odds ratio*   *Bold: significant aOR, p-value, and 95%CI for risk factors*

Discussion

The study assessed the prevalence and contributing factors of stillbirths in a semi-rural locality. The prevalence of stillbirths in the present study was found to be 4.0%, which is in line with a multicenter study in Nigeria [11]. This similarity might be due to the uniformity in the healthcare system and provision of services for pregnant mothers in Ghana and Nigeria. Our study's stillbirth rate was also lower than the estimation in a retrospective cohort study conducted in a tertiary facility in Southern Ghana (10.6%) [20] but higher than the study findings from China (0.9%) and South Africa (3.5%) [10,17]. The discrepancy might be due to variations in the study area and the inclusion and exclusion criteria (methodologies) used in the previous studies. In addition, several unreported cases of stillbirths in the community and/or municipal health facilities could be responsible for this variation. In comparison, China and South Africa have advanced and modernised healthcare systems that offer well-tailored services to pregnant women, thereby contributing to the reduction of stillbirth rates in these countries. Therefore, it could indicate that the care for women during antenatal and labour (intrapartum) might not have received appropriate and standardized obstetric practices and protocols, leading to higher stillbirth incidence in this semi-rural Ghana. On the other side, the lesser prevalence of stillbirths in this study, compared to Southern Ghana, might be due to the deployment of trained traditional birth attendants as agents of facility deliveries and trained

community health nurses to perform the duties of midwives in the Savelugu municipality of

Northern Ghana. This initiative by the Catholic Relief Services (development partner) was

adopted by the Northern Regional Health Directorate of Ghana Health Service to improve

maternal and perinatal health outcomes, which might have contributed to the reduction of the

stillbirth rate in Northern Ghana. Nonetheless, the relatively elevated incidence of stillbirths in

the study area is still alarming as postpartum mothers and their families face psychological

trauma after experiencing stillbirths (11,37) during or following a nine-month journey of

pregnancy.

In this study, maternal anaemia in the third trimester of pregnancy was independently

associated with stillbirths. Whilst a retrospective study in Columbia (38) was in agreement with

the present study, a similar study in Pakistan (39) was not in support. Anaemia is known to be a

significant predictor of malnutrition among pregnant mothers as iron deficiency type is

apparently the most prevalent cause of anaemia during gestation (40). As anaemia causes

inadequate oxygen and nutrient supply to the foetus during the late stage of pregnancy (41,42),

it could lead to a deficient supply of these nutrients and oxygen to the foetus which

subsequently causes hypoxic death of the foetus either in-utero, during childbirth or within a

short period after birth (43). To prevent gestational anaemia and subsequent stillbirths, it is

critical to strengthen public health initiatives and clinical protocols (15). These actions could be

achieved through early screening and diagnosis, prompt patient referral, integrated nutrition

counselling, routine iron-folate supplementation and culturally tailored community-based

education promoting iron-folate concordance and dietary quality (44).

This study also found that mothers with positive sickle cell status had increased odds of

stillbirth delivery. The observed findings align with previous research indicating severe

pregnancy and childbirth complications for both mothers and newborns affected by sickle cell

(45). These complications may stem from the physiological adaptations inherent in the genetic

makeup of pregnant women with sickle cell trait or disease (45,46). Such adaptations can

impact the cardiovascular, haematologic, respiratory, and renal structures in the materno-

foetal system, which can impair the overall health and ability to sustain a healthy pregnancy

(46). Consequently, these physiological challenges may contribute to the reduced likelihood of

delivering a healthy newborn. It therefore becomes imperative for healthcare professionals

especially midwives and medical doctors to recognise this risk and adopt appropriate measures

by providing targeted and special care for pregnant women affected by sickle cell.

Mothers who did not sleep under insecticide-treated bed nets during pregnancy were

more likely to deliver stillbirths. In line with the present findings, mothers' sleep practices,

particularly those who did not sleep under insecticide-treated bed nets during pregnancy, were

prone to stillbirths in a cross-sectional study in Ghana (12) but were contrasting to a meta-

analysis by Flenady and his colleagues (47). Despite the constant supply of free insecticide-

treated bed nets to pregnant mothers for malaria prevention during their first ANC visits in

Ghana (28), it is reported that most mothers do not sleep under the nets due to unfavourable

(hot) weather conditions in Northern Ghana, while others also use the nets for subsistence

farming purposes in their homes (48). The non-use of the bed nets could expose them to

malaria infections and subsequent anaemia in pregnancy, thereby increasing their risk of

stillbirth delivery (49). This underscores the need for expanded and sustained health education

campaigns including outreach activities in both health facilities and community settings to

foster insecticide-treated bed nets utilisation in semi-rural areas of Ghana.

Multiparity was found to be associated with stillbirths in the semi-rural community

during the regression analysis. Evidence from Ethiopia-based comparative studies (50,51)

corroborates the results of the current investigation. Similarly, in a cross-sectional study from

Nigeria, multiparity was associated with stillbirths (11). Most multiparous women perceive

themselves as experienced and often delay or underutilise ANC services during pregnancy,

which may reduce opportunities for early detection of complications associated with stillbirths.

Furthermore, multiparity with short interpregnancy intervals results in nutritional deficiencies

such as iron, folate and vitamin B12, which could lead to foetal growth restrictions and

stillbirths (1,52). While promoting family planning initiatives across communities remains

essential in mitigating multiparity (53,54), prioritising diverse targeted interventions for

multiparous women is equally critical in efforts to prevent the risk of stillbirths in the

municipality.

Interestingly, a large proportion of the study participants who had positive sickle cell

status and did not use insecticide-treated bed nets also made less than eight ANC visits and had

anaemia during each trimester of pregnancy. This implies that fewer ANC visits and low

haemoglobin levels in each trimester during pregnancy could precipitate suboptimal intra-

uterine and gestational environments (41,42) leading to an increased maternal risk for stillbirth

delivery (49,55) in the municipality. Therefore, greater attention should be paid to anaemia in

pregnancy and ANC visits to strengthen maternal and child health outcomes in the country.

Strengths and limitations of the study

The study recorded some limitations. First, the prevalence of stillbirths in the municipality may

be underestimated or overestimated since the study participants were sampled from six-month

cross-sectional retrospective data from health facilities. There could be potential misreporting

(underreporting), misrecording and misclassification of stillbirths and other clinical (health) data

like sickle cell determination by healthcare professionals (27,56,57), which could affect the

estimates of stillbirths and positive sickle cell in the municipality. The data also lacked details on

whether the stillbirths happened in the antepartum or intrapartum stages of delivery. The

absence of this information could imply a considerable blend of antepartum and intrapartum

stillbirths in the dataset. Additionally, the use of retrospective data (such as sickle cell, maternal

weight, multiparity, haemoglobin levels, and gestational age) from records review (MCHRBs

and delivery registers) could be a limitation of the study as a result of possible classification,

measurement and documentation errors (56). It is difficult to ascertain whether the necessary

standard protocols were adequately adhered to before measurement, potentially affecting the

dataset's quality in the analysis. Notwithstanding, the stillbirth estimates and measurements

represent the available information in the municipal health facilities which are used for

decision-making in the municipality.

The study also faced limitations in generalising the maternal factors of stillbirths

because it relied on cross-sectional data. Therefore, the findings of this study could not be used

to draw causal inferences but rather to highlight significant associations of stillbirths to the

scientific community. Furthermore, the reliability of self-reported data may be affected by

recall bias among study participants, posing a methodological limitation in the study.

Ultimately, the study lacked information on certain variables related to the independent

(exposure) variables. Specifically, maternal data on smoking status, alcohol intake and

socioeconomic status were not collected. Despite this, the study provided significant findings

for potential interventions in the municipality and the broader semi-rural Ghana.

On the other hand, the study encountered some notable strengths. At the outset, the

study addresses a significant gap in the maternal and perinatal health literature by

quantitatively examining the association between maternal factors and stillbirths. This directly

connects maternal health factors to perinatal health. The study goes beyond reporting

univariate and bivariate associations by also conducting multivariable analysis to identify

maternal factors of stillbirths, resulting in a more rigorous and robust set of findings.

Conclusion

The prevalence of stillbirths is relatively high in the Savelugu municipality. These stillbirths are

independently influenced by medical factors, particularly anaemia and positive sickle cell

statuses and unsafe health practices such as non-use of ITNs during pregnancy. Timely

identification of these factors among pregnant women (anaemia and sickle cell status) is vital to

prompt early intervention strategies in Ghanaian health facilities. This could be achieved by

strengthening obstetric triage at ANC clinics and labour/delivery rooms and also encouraging

early referrals to higher levels of care. Secondly, due to the frequent misclassification of most

perinatal and neonatal deaths as stillbirths (21), thorough perinatal death audits, which aim to

identify the cause and contributing factors of stillbirths, should be promoted in healthcare

facilities in semi-rural Ghana.

Additionally, the management of Ghana Health Service (especially municipal/district

health directorates), alongside program planners and other stakeholders involved in enhancing

perinatal and maternal health, should also be aware of these maternal factors contributing to

stillbirths. These health authorities should launch and implement effective programs, including

robust referral systems through training and capacity building of health care providers and

community cadres to facilitate early referral and management. Policy efforts should further be

targeted at promoting enhanced health education sessions on undernutrition (anaemia) and

the regular use of ITNs during pregnancy at facility ANC clinics and communities (via community

durbars) in semi-rural Ghana. Finally, we recommend future research to incorporate other

maternal variables, such as socioeconomic status, prenatal smoking and alcohol intake, to

thoroughly examine and provide a more comprehensive understanding of the maternal factors

contributing to stillbirths in both the municipality and the wider Northern Region.

List of abbreviations

ANC: Antenatal care

BMI: Body mass index

G6PD: Glucose-6-phosphate dehydrogenase

ITNs: Insecticide-treated bed nets

MCHRBs: Maternal and child health record books

STATA: Statistics and data

TD: Tetanus-diphtheria

VIF: Variance inflation factor

WHO: World Health Organization

566   **Declarations**

Ethics approval and consent to participate

This study was conducted according to the guidelines laid down in the Declaration of Helsinki,

and all procedures involving research study participants were approved by the Navrongo Health

Research Centre Institutional Review Board with approval number NHRCIRB373. Permission

was also granted by the Ghana Health Service at the municipal directorate and health facilities

level. Written informed consent/assent was obtained from all study participants and/or legal

representatives and also from parents/guardians of participants under 18 years of age. All

participants were able to understand, read and write about consent to participate.

Consent for publication

Not applicable

Acknowledgements

580 The authors are thankful to all research assistants, study participants, and volunteers who
581 contributed in diverse ways to the successful implementation of the study. We are also grateful
582 to Elvis Brown Ayaala, Abdulai Sulemana, Allor Mohammed, and Julien Beweleyir for their
583 supportive supervision and data collection exercise during this study.

Funding statement

586 There was no funding support

Conflict of interest statement

589 The authors declared that they have no conflict of interest.

Availability of data and materials

592 The datasets collected, generated, and/or analysed during the present study are available from
593 the corresponding author upon reasonable request. Data sharing not applicable to this article
594 as no datasets were generated or analysed during the current study.

Authors' contributions

SAG, AA, MG, CSD and PAA designed and conceptualised the study. SAG, AA, MG, WP, CA and

CSD carried out the data collection while SAG, WP, CA and PAA analysed and interpreted the

data. All authors contributed to drafting the initial manuscript, reviewing and approving the

final manuscript.

Authors' acronyms

Silas Adjei-Gyamfi (SAG); Abigail Asirifi (AA); Mamudu Gafaru (MG); Wisdom Peprah (WP);

Clotilda Asobuno (CA); Constance Siakwan Dapaah (CSD); Paul Armah Aryee (PAA)

References

1. Lawn JE, Blencowe H, Waiswa P, Amouzou A, Mathers C, Hogan D, et al. Stillbirths: Rates,

risk factors, and acceleration towards 2030. Vol. 387, The Lancet. Lancet Publishing

Group; 2016. p. 587–603.

2. Bawah AT, Ngala RA, Alidu H, Seini MM, Wumbee JDK, Yeboah FA. Gestational diabetes

mellitus and obstetric outcomes in a Ghanaian community. Pan Afr Med J. 2019;32.

3. WHO. Every newborn: An action plan to end Preventable deaths: Executive summary

Geneva: World Health Organization. WHO, UNICEF. 2014;12 p.

4. Harding EBM, Harrington LT, Lockwood CM, Duncan Brown JR, Hall AC, Brown MA, et al.

Perinatal and neonatal mortality. Br Med J. 2006;281(6254):1567.

5. Blencowe H, Cousens S, Jassir FB, Say L, Chou D, Mathers C, et al. National, regional, and

worldwide estimates of stillbirth rates in 2015, with trends from 2000: A systematic

analysis. Lancet Glob Heal. 2016;4(2):e98–108.

6. Lawn JE, Yakoob M, Haws RA, Soomro T, Darmstadt GL, Bhutta ZA. 3.2 million stillbirths:

Epidemiology and overview of the evidence review. Vol. 9, BMC Pregnancy and

Childbirth. BioMed Central; 2009. p. 1–17.

7.   Dassah ET, Odoi AT, Opoku BK. Stillbirths and very low Apgar scores among vaginal births

in a tertiary hospital in Ghana: A retrospective cross-sectional analysis. BMC Pregnancy

Childbirth. 2014;14(1).

8.   Njoku C., Emechebe C., Eyong E., Ukaga J., Anachuna K. Prevalence and risk factors for

stillbirths in a tertiary hospital in Niger Delta area of Nigeria: A ten year review. Int J Med

Biomed Res. 2016;5(3):106–13.

9.   Mukherjee A, Di Stefano L, Blencowe H, Mee P. Determinants of stillbirths in sub-Saharan

Africa: A systematic review. BJOG An Int J Obstet Gynaecol. 2024;131(2):140–50.

10.   Zhu J, Liang J, Mu Y, Li X, Guo S, Scherpbier R, et al. Sociodemographic and obstetric

characteristics of stillbirths in China: A census of nearly 4 million health facility births

between 2012 and 2014. Lancet Glob Heal. 2016;4(2):e109–18.

11.   Okonofua FE, Ntoimo LFC, Ogu R, Galadanci H, Mohammed G, Adetoye D, et al.

Prevalence and determinants of stillbirth in Nigerian referral hospitals: A multicentre

study. BMC Pregnancy Childbirth. 2019;19(1).

12.   Owusu JT, Anderson FJ, Coleman J, Oppong S, Seffah JD, Aikins A, et al. Association of

maternal sleep practices with pre-eclampsia, low birth weight, and stillbirth among

Ghanaian women. Int J Gynecol Obstet. 2013;121(3):261–5.

http://doi.wiley.com/10.1016/j.ijgo.2013.01.013

13.   Odendaal H, Dukes KA, Elliott AJ, Willinger M, Sullivan LM, Tripp T, et al. Association of
prenatal exposure to maternal drinking and smoking with the risk of stillbirth. JAMA
Netw Open. 2021;4(8):E2121726.

14.   Afulani PA. Determinants of stillbirths in Ghana: Does quality of antenatal care matter?
BMC Pregnancy Childbirth. 2016;16(1):1–17. http://dx.doi.org/10.1186/s12884-016-
0925-9

15.   Bhutta ZA, Darmstadt GL, Haws RA, Yakoob MY, Lawn JE. Delivering interventions to
reduce the global burden of stillbirths: Improving service supply and community demand.
BMC Pregnancy Childbirth. 2009;9(1):1–37. http://dx.doi.org/10.1186/1471-2393-9-S1-
S7

16.   Hug L, You D, Blencowe H, Mishra A, Wang Z, Fix MJ, et al. Global, regional, and national
estimates and trends in stillbirths from 2000 to 2019: A systematic assessment. Lancet.
2021;398(10302):772–85. http://dx.doi.org/10.1016/S0140-6736(21)01112-0

17.   Lavin T, Pattinson RC, Nedkoff L, Gebhardt S, Preen DB. Stillbirth risk across pregnancy by
size for gestational age in Western Cape Province, South Africa: Application of the
fetuses-at-risk approach using perinatal audit data. South African Med J.
2019;109(12):927–33.

18.   Ghana Statistical Service (GSS) Ghana Health Service (GHS) and ICF. 2017 Ghana
Maternal Health Survey (GMHS) Research Policy Briefs. GSS, GHS, and ICF. 2019.
https://statsghana.gov.gh/gssmain/fileUpload/pressrelease/PB14.pdf

19.   Ghana Statistical Service (GSS), Ghana Health Service (GHS), and Macro International.

Ghana Maternal Health Survey 2007. GSS, GHS, and Macro International. Calverton,

Maryland, USA; 2009.

20.   Adu-Bonsaffoh K, Gyamfi-Bannerman C, Oppong SA, Seffah JD. Determinants and

outcomes of preterm births at a tertiary hospital in Ghana. Vol. 79, Placenta. W.B.

Saunders Ltd; 2019. p. 62–7.

21.   Nonterah, A. Engelbert ., Isaiah AA, Edmund WK, Kagura J, Tamimu M, Ayamba YE,

Nonterah WE, et al. Trends and risk factors associated with stillbirths : A case study of

the Navrongo War Memorial Hospital in Northern Ghana. PLoS One. 2020;15(2):1–13.

https://doi.org/10.1371/journal.%0D%0Apone.0229013

22.   Cuschieri S. The STROBE guidelines. Saudi J Anaesth. 2019;13(5):S31–4.

23.   Vandenbroucke JP, Von Elm E, Altman DG, Gøtzsche PC, Mulrow CD, Pocock SJ, et al.

Strengthening the Reporting of Observational Studies in Epidemiology (STROBE):

Explanation and elaboration. PLoS Med. 2007;4(10):1628–54.

24.   Ameyaw EK, Kofinti RE, Appiah F. National health insurance subscription and maternal

healthcare utilisation across mothers' wealth status in Ghana. Health Econ Rev.

2017;7(1):16.https://healtheconomicsreview.biomedcentral.com/articles/10.1186/s1356

1-017-0152-8

25.   Adjei-Gyamfi S, Musah B, Asirifi A, Hammond J, Aryee PA, Miho S, et al. Maternal risk

factors for low birthweight and macrosomia: A cross-sectional study in Northern Region,

695  Ghana. J Heal Popul Nutr. 2023;42(1):1–16. https://doi.org/10.1186/s41043-023-00431-0

696 26. Adjei-Gyamfi S, Asirifi A, Peprah W, Abbey DA, Hamenoo KW, Zakaria MS, et al. Anaemia

697  at 36 weeks of pregnancy: Prevalence and determinants among antenatal women

698  attending peri-urban facilities in a developing country, Ghana. PLOS Glob Public Heal.

699  2024;4(9):e0003631. https://doi.org/10.1371/journal.pgph.0003631

700 27. Adjei-Gyamfi S, Zakaria MS, Asirifi A, Issahaku S, Ibrahim MA, Aryee PA. Maternal

701  anaemia and polycythaemia during pregnancy and risk of inappropriate birth weight for

702  gestational age babies: A retrospective cohort study in the northern belt of Ghana. BMJ

703  Open. 2024;14:e082298. https://doi.org/10.1136/bmjopen-2023-082298

704 28. Adjei-Gyamfi S, Asirifi A, Aiga H. Prevalence and associated risk factors of preterm and

705  post-term births in Northern Ghana: A retrospective study in Savelugu Municipality. J

706  Pediatr Perinatol Child Heal. 2023;07(04):235–48.

707  https://doi.org/10.26502/jppch.74050169

708 29. WHO Web Annex 2. Template protocol for surveys to estimate the prevalence of

709  biomarkers of infection with the hepatitis viruses.In: Consolidated strategic information

710  guidelines for viral hepatitis planning and tracking progress towards elimination. Geneva;

711  2019.

712 30. Adjei-Gyamfi S, Asirifi A, Asobuno C, Korang FK. Knowledge and occupational practices of

713  beauticians and barbers in the transmission of viral hepatitis: A mixed-methods study in

714  Volta Region of Ghana. PLoS One. 2025;20(1):1–24.

715  http://dx.doi.org/10.1371/journal.pone.0306961

31. Cochran WG. Samling Techniques. John Wiley & Sons, Inc. 1977. 448 p.

32. Engmann C, Walega P, Aborigo RA, Adongo P, Moyer CA, Lavasani L, et al. Stillbirths and

early neonatal mortality in rural Northern Ghana. Trop Med Int Heal. 2012;17(3):272–82.

33. Asobuno C, Adjei-Gyamfi S, Aabebe FG, Hammond J, Taikeophithoun C, Amuna NN, et al.

Risk factors for anaemia among pregnant women: A cross-sectional study in Upper East

Region,   Ghana.   PLoS   One.   2024;19(11):e0301654.

https://doi.org/10.1371/journal.pone.0301654

34. GHS. DHIMS-2: District Information Management Systems version 2. Ghana Health

Service. 2020. https://dhims.chimgh.org/dhims

35. Ghana Statistical Service (GSS), Ghana Health Service (GHS) & ICF International. Ghana

Demographic and Health Survey 2014. Rockville, Maryland, USA; 2015.

36. Rasmussen KM, Catalano PM, Yaktine AL. New guidelines for weight gain during

pregnancy: What obstetrician/gynecologists should know. Curr Opin Obs Gynecol.

2009;21(6):521–6.

37. Vogel JP, Souza JP, Mori R, Morisaki N, Lumbiganon P, Laopaiboon M, et al. Maternal

complications and perinatal mortality: Findings of the World Health Organization

multicountry survey on maternal and newborn health. BJOG. 2014;121 Suppl 1(Suppl

1):76–88.

38. Smith C, Teng F, Branch E, Chu S, Joseph KS. Maternal and perinatal morbidity and

mortality associated with anemia in pregnancy. Obstet Gynecol. 2019;134(6):1234–44.

39. Mahmood T, Rehman AU, Tserenpil G, Siddiqui F, Ahmed M, Siraj F, et al. The association

between iron-deficiency anemia and adverse pregnancy outcomes: A retrospective

report from Pakistan. Cureus. 2019;11(10).

40. Garzon S, Cacciato PM, Certelli C, Salvaggio C, Magliarditi M, Rizzo G. Iron deficiency

anemia in pregnancy: Novel approaches for an old problem. Oman Med J.

2020;35(5):e166.

41. Barker DJP. Intrauterine programming of adult disease. Vol. 1, Molecular Medicine

Today. Elsevier Current Trends; 1995. p. 418–23.

42. Wu G, Imhoff-Kunsch B, Girard AW. Biological mechanisms for nutritional regulation of

maternal health and fetal development. Vol. 26, Paediatric and Perinatal Epidemiology.

John Wiley & Sons, Ltd; 2012. p. 4–26.

43. Parks S, Hoffman MK, Goudar SS, Patel A, Saleem S, Ali SA, et al. Maternal anaemia and

maternal, fetal, and neonatal outcomes in a prospective cohort study in India and

Pakistan. BJOG An Int J Obstet Gynaecol. 2019;126(6):737–43.

44. WHO. Global Nutrition Targets 2025: Low birth weight policy brief

(WHO/NMH/NHD/14.5). Geneva; 2014.

45. Smith-Whitley K. Complications in pregnant women with sickle cell disease. Hematology.

2019;2019(1):359–66. https://doi.org/10.1182/hematology.2019000039

46. Jain D, Atmapoojya P, Colah R, Lodha P. Sickle cell disease and pregnancy. Mediterr J

Hematol Infect Dis. 2019;11(1):e2019040. http://dx.doi.org/10.4084/MJHID.2019.040

47.   Flenady V, Koopmans L, Middleton P, Frøen JF, Smith GC, Gibbons K, et al. Major risk

factors for stillbirth in high-income countries: A systematic review and meta-analysis.

Lancet. 2011;377(9774):1331–40.

48.   Savelugu Municipal Health Directorate (Ghana Health Service). Annual Performance

Report 2020: Savelugu Municipal Health Directorate. Savelugu, Northern Region; 2021.

49.   Moore KA, Simpson JA, Scoullar MJL, McGready R, Fowkes FJI. Quantification of the

association between malaria in pregnancy and stillbirth: A systematic review and meta-

analysis. Lancet Glob Heal. 2017 Nov 1;5(11):e1101–12.

50.   Tadese M, Tessema SD, Taye BT. Adverse perinatal outcomes among grand multiparous

and low multiparous women and its associated factors in North Shewa zone public

hospitals: The role of parity. Int J Gen Med. 2021;14:6539–48.

51.   Yimer NB, Tenaw Z, Gedefaw A. Pregnancy outcomes in grand multiparous women: Does

parity matter? A comparative study. Ethiop J Reprod Heal. 2020;12(1):35–45.

52.   Pattinson R, Kerber K, Buchmann E, Friberg IK, Belizan M, Lansky S, et al. Stillbirths: How

can health systems deliver for mothers and babies? Lancet. 2011;377(9777):1610–23.

https://www.sciencedirect.com/science/article/pii/S0140673610623069

53.   Dibaba D, Charkos TG. Challenges in reducing grand multiparity rates in Ethiopia: An

analysis of 2019 EDHS data using a multilevel model approach. Contracept Reprod Med.

2025;10(1). https://doi.org/10.1186/s40834-024-00328-1

54.   Solanke BL, Banjo OO, Oyinloye BO, Asa SS. Maternal grand multiparity and intention to

use modern contraceptives in Nigeria. BMC Public Health. 2018;18(1):1–15. https://link-

springer-com.ezproxy.aut.ac.nz/articles/10.1186/s12889-018-6130-1

55. Idung A, Jombo H, Udonwa NE, Iyanam VE, Idung AU, Jombo HE. Anaemia in pregnancy

at booking: Prevalence and risk factors among antenatal attendees in a Southern Nigeria.

Asian J Med Heal. 2019;15(1):48805.

56. Libretti A, Corsini C, Remorgida V. What's new on female genital mutilation/cutting?

Recent findings about urogynecologic complications, psychological issues, and obstetric

outcomes. Minerva Obstet Gynecol.;77(1):56–61.

https://pubmed.ncbi.nlm.nih.gov/37539674/

57. Libretti A, Bianco G, Corsini C, Remorgida V. Female genital mutilation/cutting: Going

beyond urogynecologic complications and obstetric outcomes. Arch Gynecol Obstet.

2023;308(4):1067–74. https://doi.org/10.1007/s00404-023-06929-6