1

Title

Maternal risk factors contributing to stillbirths in semi-rural Ghana: Insights from

a cross-sectional study.

Author's information

* **Silas Adjei-Gyamfi, MPH, MPhil (Corresponding author)**

 Savelugu Municipal Hospital

 Ghana Health Service

 P. O. Box 45, Savelugu, Northern Region, GHANA

 Phone: +233243421318

 Email: adjeigyamfis@yahoo.com

 ORCID: https://orcid.org/0009-0008-8496-4099

Abigail Asirifi, Dip, BSc

 Department of Midwifery

 Garden City University College

 P. O. Box KS 12775, Kumasi, Ashanti Region, GHANA

 Phone: +233545951777

 Email: abimaiden80@gmail.com

 ORCID: https://orcid.org/0009-0008-8133-4104

Gafaru Mamudu, BSc

25   Savelugu Municipal Hospital

26   Ghana Health Service

27   P. O. Box 45, Savelugu, Northern Region, GHANA

28   Phone: +233248303028

29   Email: gafarumamudu@gmail.com

31   Wisdom Peprah, BSc, MPhil

School of Allied Health Sciences

University for Development Studies

P. O. Box TL 1883, Tamale, Northern Region, GHANA

Phone: +233507093898

Email: pwisdom466@gmail.com

ORCID: https://orcid.org/0000-0003-0263-8158

39   Clotilda Asobuno, BSc, MPH

Kassena Nankana West District Health Directorate

Ghana Health Service

P. O. Box 21, Paga, Upper East Region, GHANA

Phone: +233245452488

Email: clotildaasobuno@gmail.com

ORCID: https://orcid.org/0009-0008-7471-6462

47   Constance Siakwan Dapaah, BSc

48   Ghana Health Service

49   Atwima Nwabiagya North District Health Directorate

50   P.O. Box 26, Barekese, Ashanti Region, GHANA

51   Phone: +233245727275

52   Email: constancedapaah25@gmail.com

Paul Armah Aryee, MMedSc, MPhil, PhD

School of Allied Health Sciences

University for Development Studies

P. O. Box TL 1883, Tamale, Northern Region, GHANA

Phone: +233200516610

Email: parmah@uds.edu.gh

Abstract

Background

Stillbirths do not only trigger maternal psychological trauma but also affect socioeconomic

growth in low- and middle-income countries. Given the diverse dynamics of the prevalence and

associated factors of stillbirths in various low-resource settings, such as Ghana, it is essential to

have comprehensive knowledge to address and mitigate these factors that could influence

stillbirths. However, there is data paucity on maternal factors contributing to stillbirths in semi-

rural areas of Ghana.

Objective

This study aimed to determine the prevalence and associated maternal factors of stillbirths in

semi-rural Ghana.

Design

This is a cross-sectional study that utilised a quantitative approach.

Methods

The study was conducted among 422 mothers who delivered in public health facilities and

reside in a semi-rural municipality in Northern Ghana. Sociodemographic, obstetric, and

medical-related data were gathered through structured interviews and from record reviews.

Chi-square test of proportions and multiple logistic regression analyses were employed to

identify the independent factors associated with stillbirths at a 95% confidence level.

Results

Stillbirth rate was estimated at 4.0% (95%CI: 2.4%–6.4%). The analysis revealed that anaemia in

the third trimester of pregnancy (aOR:3.15; 95%CI:2.94–10.6; p=0.034), positive sickle cell

status (aOR:2.91; 95%CI:1.50–7.01; p=0.018), and non-utilisation of insecticide-treated bed nets

(ITNs) (aOR:7.75; 95%CI:2.33–9.8; p=0.001) during pregnancy were significantly associated with

increased risk of stillbirths.

Conclusion

The relatively high prevalence of stillbirths in the municipality was influenced by medical factors

(particularly anaemia and positive sickle cell status) and unsafe health practices via the non-use

of ITNs during pregnancy. The management of the Ghana Health Service, alongside program

planners and other stakeholders involved in enhancing perinatal and maternal health, should

focus on implementing effective initiatives such as improved health education on nutrition and

safety practices at the facility and community levels, as well as robust referral systems to

prevent and reduce stillbirths in semi-rural Ghana.

113   **Plain language summary**

Stillbirths, which are the death of a baby before or during delivery, not only cause emotional

distress for mothers but also affect the national growth in most countries. Despite the

importance of understanding the maternal factors of stillbirths, there is limited data available,

especially in semi-rural areas of Ghana. A study was therefore undertaken to assess the

prevalence and maternal factors associated with stillbirths in semi-rural Ghana. This study

involved mothers who gave birth in public health facilities and live in the same semi-rural area.

Researchers collected data through structured interviews and from medical records. The data

was analysed using chi-square tests and logistic regression to identify maternal factors linked to

stillbirths. The findings showed a stillbirth rate of 4.0%. Key factors associated with stillbirths

included maternal anaemia in the third trimester of pregnancy, positive sickle cell status, and

not using insecticide-treated bed nets (ITNs) during pregnancy. The study highlights the need

for improved health education on nutrition and safe health practices to reduce and/or prevent

stillbirths. Efforts should focus on addressing medical conditions like anaemia and sickle cell

disease and promoting the use of ITNs to protect pregnant women from malaria and

subsequent stillbirths.

Keywords

Anaemia,

Determinants,

Risk factors,

Northern Ghana,

Predictors,

Stillbirths,

Stillborns

Number of words

Abstract   301 words

Main text   3,896 words

Introduction

Stillbirths, also called stillborns, which usually occur before or during childbirth, not only trigger maternal psychological trauma but also contribute to perinatal and maternal mortalities, which affect the socio-economic growth and development in low- and middle-income countries (LMICs) (1–3). Stillbirth is denoted as the birth of a foetus with no signs of life occurring at 28 weeks or more of gestation (3), and it accounts for more than half of all perinatal mortalities worldwide (4).

On a global scale, an estimated 2·6 million babies were born with no signs of life in 2015, which reduced to about two million stillborns (13.9 per 1000 births) in 2019 (5,6). Sub-Saharan Africa (SSA) has the highest stillbirth rate globally, estimated at 21.7 per 1000 births, compared to 3.0 for high-income countries (HICs) (6) and accounts for more than one-third of the global burden of stillbirths (5,6). Furthermore, stillbirth rates vary widely across regions in the world, from 22·8 per 1000 births in Western and Central Africa to 2·9 in Western Europe. After West and Central Africa, Eastern and Southern Africa and South Asia have the second (20.5 per 1000 births) and third-highest (18.2 per 1000 births) stillbirth rates (6). Estimates of stillbirths in China, Ghana, South Africa, and Nigeria are found to be 9.0, 15.0, 35.0 and 40.0 per 1000 births respectively (7–10). Hug et al. 2021 observed a 2·3% annual decline in the stillbirth rate globally, with a sluggish decline rate in most Asian and SSA countries (6). This finding may hinder SSA countries from reaching the Every Newborn Action Plan target of 12 per 1000 births by 2030.

172   The aetiology and antecedents of stillbirths are multifaceted and are affected by
173 sociological, physiological (medical), and physical factors (1). It is reported that the stillbirth rate
174 and its determinants are dependent on the geographic and sociodemographic distribution of
175 the population (3,5). Therefore, the study findings from one setting might not apply to another.
176 Universally, earlier studies acknowledged that the factors associated with stillbirths included
177 congenital foetal deformities, foetal distress, preterm births, breech delivery, abruptio
178 placentae, and male foetus (11–13). Non-use of insecticide-treated bed nets (ITNs) for malaria
179 prevention during pregnancy, maternal age less than 20 years, unmarried mothers,
180 employment status, multiparity, and delayed vaginal delivery were also regarded as some
181 determinants of stillbirths in other past studies (7,9,14). In low-resource settings, inadequate
182 knowledge about antenatal care (ANC) services, cost, distance and the time needed to access
183 care are major barriers to the effective uptake of quality ANC and particularly intrapartum
184 services, which contributes significantly to stillbirths (10,15).

185   The vast differences in stillbirth rates between HICs and LMICs have led to stillbirth
186 being equally regarded as a global development metric. Similar to many other SSA nations,
187 Ghana faces challenges in the routine and adequate monitoring and recording of stillbirths
188 (5,16). Diverse surveys conducted in Ghana estimate the stillbirth rate to fluctuate between
189 approximately 14 and 22 per 1000 births (16,17). Also, some population health surveillance
190 and health facility data from different parts of the country indicate even higher rates. For
191 instance, the Ghana Maternal Health Survey in 2017 observed that 68% of stillbirths occur in
192 rural areas compared to urban areas (16). Other findings also revealed 34 stillbirths/1000 births
193 in Navrongo of the Upper East Region, 59 stillbirths/1000 births in the Ashanti Region and 109

stillbirths/1000 births in a tertiary hospital in the Greater Accra Region (12,18,19), which makes

these high stillbirth occurrences very worrisome in the country. While these longitudinal

studies reported a wide range of health facility, neonatal, paternal and maternal factors, none

have exclusively delved into the broad spectrum of maternal factors that contribute to

stillbirths. Additionally, most of these longitudinal studies were conducted in rural and urban

areas. Hence, a cross-sectional study to provide a snapshot of the current situation in semi-rural

communities is non-existent. Again, to date, little is particularly investigated about maternal

factors contributing to stillbirths in semi-rural Ghana.

On the other hand, epidemiological studies evaluating stillbirths frequently utilise the

concept of perinatal deaths, which encompasses both stillbirths and early neonatal deaths

(5,6). These studies often rely on population health surveillance, community-based oral post-

mortem reports and intricate analytical modelling techniques. However, the utility of these

estimates is constrained by the etiological distinctions between stillbirths and early neonatal

deaths, particularly given that the intrapartum period represents the highest-risk phase for

stillbirths. To achieve the World Health Assembly's national target of a stillbirth prevalence of

12 or less per 1000 births by 2030 in every country through the Every Newborn Action Plan (3),

the maternal factors of stillbirths and their dynamics in different settings must be well

understood and appreciated. Although stillbirths are higher in LMICs, including Ghana (12,18),

relevant information on the maternal factors of stillbirths is inadequately and incorrectly

documented in the district and/or regional databank. Hence, adequate evidence of information

on these factors in semi-rural Ghana could trigger useful interventions that may lead to

preventable stillbirths and subsequently contribute to improving maternal and perinatal health.

The purpose of the study is to estimate the prevalence of stillbirths and examine the

maternal factors associated with stillbirths in a semi-rural municipality in the Northern Region

of Ghana. We examined the effect of maternal risk factors on stillbirths by utilising a broad

spectrum of sociodemographic, medical or clinical and obstetric or antenatal characteristics of

the mothers. Given the distinct and unique nature of this study, the findings will be crucial to

inform clinical practice through timely interventions, decision-making, and policy ideation, with

a particular emphasis on maternal factors in the semi-rural Ghana and SSA.

Materials and methods

Study design

This current survey is a cross-sectional study conducted among women who delivered in a

semi-rural municipality in the Northern Region of Ghana. The design was employed to

determine the prevalence and associated maternal factors of stillbirths. The study followed the

STROBE checklist of reporting (see Supplementary 1).

Study setting

This study was conducted in Savelugu Municipality of Northern Ghana. Savelugu Municipality is

one of the 16 districts/municipalities in the Northern Region of Ghana that shares boundaries

with five neighbouring districts, including Gushegu, Karaga, Kumbungu, Nanton, and Tolon. The

municipality has five major government (public) health facilities (one hospital and four health

centres) that provide comprehensive maternal and newborn services, including free maternal

and delivery services (20) to a population of approximately 130,000. In 2019, the coverage of

ANC and postnatal care visits was 94% and 98%, respectively (21,22). Due to the increased

average monthly delivery count of over 300 deliveries with higher records of unfavourable birth

outcomes (21,23,24), this municipality was selected to investigate the factors of stillbirths

among women who delivered in government health facilities.

Sample size estimation

The sample size (n) was estimated using Cochran's method with the World Health Organization

(WHO) conservative design effect (DEF), $n = \frac{Z^2 PQ}{e^2}$ X DEF (25–27). A standard variate (Z) at a 95%

confidence interval of 1.96, a standard error (e) of 5%, a prevalence (P) of stillbirths estimated

at 21% in northern Ghana (28), Q=1–P and WHO standard conservative DEF size of 1.5 (26,27),

was applied to estimate the preliminary sample size at 384. An additional 10 percent was added

to the calculated sample size to account for fall-outs, incomplete responses and damaged

questionnaires (21,23,29) resulting in a final total sample of approximately 422.

Study population

The study recruited a total of 422 women who visited the five major public facilities in the

Savelugu Municipal for childbirth or delivery purposes from January 1 to June 31, 2020. All

women who lived in the municipality, delivered at the municipal health facilities and had their

health information recorded in the delivery registers were eligible for the study.

Sampling

In the sampling stage, all five major government (public) health facilities were chosen for the

study since all these facilities report on stillbirths every year (30). The sample sizes for each

facility were determined using the probability proportional to size technique. The delivery

registers from each facility were used to line-list all eligible postpartum women (who delivered

from January 1 to June 31, 2020) to form a sampling frame. A simple random sampling method

was then used to select the required number of respondents from each facility. The addresses

and contact numbers of these selected respondents were recorded to trace them to their

homes.

Data collection

Data was collected using a pretested questionnaire by six well-trained data collectors (midwives

and public health officers) from July 16 to September 31, 2020. The questionnaire was

developed based on existing literature, previous studies, and sociological and biological

plausibility (1,5,21,31). We verified the content validity of the questionnaire by consulting with

experts in the field as well as the study authors. Sociodemographic, obstetric or antenatal and

medical-related information was collected through structured interviews and from record

reviews (thus, delivery registers and maternal and child health record books [MCHRBs]). The

data collectors were trained for one week. The data collection procedure was in two phases.

The first phase was at the household level, while the second was at the health facilities level

after sampling and contacting the eligible study participants.

Before the commencement of household data collection, the trained data collectors

retrieved the addresses and contact information (telephone numbers) of respondents and/or

their representatives from the delivery register with the support of the facility midwives. The

data collectors arranged appointments and then visited the homes of the recruited

respondents. Data not excluding age, marital status, education level, job status, ethnicity and

religious affiliations were collected via structured interviews and were validated in the

MCHRBs. Information about first-trimester body mass index (since the foetal weight in the first

trimester is negligible (32)), antenatal visits, parity, gravidity, rhesus status, sickle cell status,

tetanus-diphtheria (TD) immunisation status, haemoglobin levels across pregnancy trimesters,

use of pre-pregnancy family planning, and utilisation of ITNs (1,24) were collected from their

MCHRBs at their homes. Afterwards, information like pre-eclampsia diagnosis, postpartum

haemorrhage, delivery mode, gestational age at birth and birth/delivery outcome prognosis

(stillbirth or livebirth) based on the classification by health professionals (1,24) were traced and

extracted from the delivery registers at the health facilities level.

Study variables

The outcome variable was birth outcome prognosis (stillbirths), categorised into binary

outcomes for stillbirths (coded as 1) or livebirths (coded as 0), whereas exposure variables

included all sociodemographic, obstetric or antenatal, and medical variables of the women,

which were all categorised based on their normal distribution and existing studies (1,5,21,31).

According to the WHO, stillbirths were measured as the birth of a foetus at 28 or more

weeks of gestation with no signs of life, while livebirths were regarded as newborns with signs

of life (3). Haemoglobin levels were dichotomised into anaemia (< 11g/dl) and no anaemia (≥

11g/dl). Gestational age at birth was grouped into < 37 weeks (preterm) and ≥ 37 weeks

(normal-term). Parity was classified as primiparity (one delivery) and multiparity (two or more

deliveries). Gravidity was grouped into primigravida (one pregnancy) and multigravida (two or

more pregnancies) (22–24,29). Standardised WHO categories of pre-pregnancy body mass

index (BMI) of underweight (<18.5 kg/m^2), normal BMI (18.5–24.9 kg/m^2) and overweight

(≥25.0 kg/m^2) were utilised. Rhesus and sickle cell statuses were grouped as positive or

negative, whereas ANC visits were also categorised into ≥ 8 visits and < 8 visits using the new

2016 WHO antenatal model (21,31).

Statistical analysis

Frequencies and percentages were calculated for the sociodemographic, obstetric or antenatal

and medical variables. Chi-square/Fisher's analysis tests of proportions were employed to

determine the statistical association between birth outcome prognosis (stillbirth) and all

exposure variables at the bivariate level. These analyses were confirmed with univariate logistic

regression. Multicollinearity among the significant exposure variables was then assessed with

variance inflation factor (VIF), where variables with a VIF less than five were selected into

multiple (binary) logistic regression analyses (21,22,24).

Applying simultaneous variable entry, the significant exposure variables with no

multicollinearity issues were forwarded into the multiple logistic regression model and run to

determine significant factors of stillbirths. All analyses were computed using STATA 17.0 (Stata

Corporation, Texas, USA), and a p-value of less than 0.05 was considered significant during

statistical analysis.

Results

340 **Prevalence of stillbirths**

341 Figure 1 presents the prevalence of stillbirths in a semi-rural municipality of Ghana. All the data

342 from the sampled 422 mothers were used for the analysis. The prevalence of stillbirths in this

343 semi-rural municipality was estimated at 4.0% (95%CI: 2.4% – 6.4%).

344
Figure 1. Prevalence of stillbirths in a semi-rural municipality of Ghana

Sociodemographic characteristics of study participants

Table 1 illustrates the frequency and bivariate distribution of sociodemographic characteristics

of study participants. The average age of the mothers was 27.6 (sd=6.0) years, while teenage

mothers were estimated at 9%. A large proportion of the participants could not read or write

(no formal education) (43%), were not employed (64%), and were married (93%). Furthermore,

more than two-thirds of the mothers were from the Dagomba ethnic group (81%). Finally, Islam

was the most practised religion by the study participants (88%).

Obstetric and medical characteristics of study participants

Table 2 shows the frequency and bivariate distribution of medical and obstetric or antenatal

characteristics of respondents. Maternal anaemia was highest in the second (71%), followed by

the first (64%) and third (45%) trimesters of pregnancy. Few of the mothers attended antenatal

clinic at least eight times (28%), ingested sulphadoxine-pyrimethamine at least three times

(36%), and did not use insecticide-treated bed nets (ITNs) during pregnancy (36%). Nearly

three-quarters of the mothers were multiparous (71%) and multigravidae (74%). Additionally,

slightly above a third of the mothers tested positive for sickle cell (38%), whereas almost 90%

had positive blood rhesus status. A small proportion of the mothers had pre-

eclampsia/eclampsia (2%), were diagnosed with antepartum haemorrhage (7%), were

immunised against tetanus-diphtheria (7%), and had glucose-6-phosphate dehydrogenase

(G6PD) defect (27%) during the period of pregnancy.

371 **Table 1: Frequency and bivariate distribution of sociodemographic variables with stillbirths**

Variables	Frequency	Bivariate analysis		
	N (%)	Stillbirths (%)	Livebirths (%)	p-value [a]
Age category of mother				
< 20 years	38 (9.0)	1 (5.9)	37 (9.2)	0.573
20 – 35years	340 (80.6)	13 (76.5)	327 (80.7)	
> 35 years	44 (10.4)	3 (17.6)	41 (10.1)	
Mean = 27.63 (SD=6.02)				
Marital status of mother				
Married	391 (92.7)	15 (88.2)	376 (92.8)	1.000
Not married	31 (7.3)	2 (11.8)	29 (7.2)	
Education level of mother				
No education	182 (43.1)	7 (41.2)	175 (43.2)	0.314
Primary school	58 (13.8)	1 (5.9)	57 (14.1)	
High school	138 (32.7)	9 (52.9)	129 (31.8)	
Tertiary	44 (10.4)	0 (0.0)	44 (10.9)	
Ethnicity of mother				
Dagomba	340 (80.6)	13 (76.5)	327 (80.7)	0.354
Gonja	22 (5.2)	0 (0.0)	22 (5.4)	
Frafra/Grusi	34 (8.1)	3 (17.7)	31 (7.7)	
Others (Asante, Ewe, Nigeria)	26 (6.1)	1 (5.9)	25 (6.2)	
Religion of mother				
Muslim	372 (88.2)	13 (76.5)	359 (88.6)	0.158
Christian	50 (11.8)	4 (23.5)	46 (11.4)	
Occupation of mother				
Unemployed	269 (63.7)	16 (94.1)	253 (62.5)	0.303
Employed	153 (36.3)	1 (5.9)	152 (37.5)	
Sex of baby (foetus)				
Male	210 (49.8)	6 (35.3)	204 (50.4)	0.322
Female	212 (50.2)	11 (64.7)	201 (49.6)	

372 [*] $p < 0.05$ [a] *Chi-square/Fisher's exact test*

381 **Table 2: Frequency and bivariate distribution of medical and obstetric variables with**
382 **stillbirths**

Variables	Frequency	Bivariate analysis		
	N (%)	Stillbirths (%)	Livebirths (%)	p-value [a]
First-trimester body mass index (BMI)				
Underweight	13 (3.1)	1 (5.9)	12 (3.2)	0.423
Normal BMI	279 (66.1)	12 (64.7)	267 (65.9)	
Overweight	130 (30.8)	5 (29.4)	125 (30.9)	
Number of antenatal care visits				
< 8 visits	304 (72.0)	14 (82.3)	290 (71.6)	0.333
≥ 8 visits	118 (28.0)	3 (17.7)	115 (28.4)	
Sulphadoxine-pyrimethamine intake				
None	25 (5.9)	2 (11.8)	23 (5.7)	**0.012**[*]
1 – 3 doses	247 (58.6)	12 (70.6)	235 (58.0)	
> 3 doses	150 (35.5)	3 (17.6)	147 (36.3)	
Pre-eclampsia/Eclampsia				
Yes	9 (2.1)	2 (11.8)	7 (1.7)	0.768
No	413 (98.9)	15 (88.2)	398 (98.3)	
Antepartum haemorrhage				
Yes	34 (8.1)	4 (23.5)	30 (7.4)	0.622
No	388 (91.9)	13 (76.5)	375 (92.6)	
Pre-pregnancy family planning use				
Yes	75 (17.8)	3 (17.7)	72 (17.8)	0.989
No	347 (82.2)	14 (82.3)	333 (82.2)	
Insecticide-treated bed nets use during pregnancy				
Yes	270 (64.0)	4 (23.5)	266 (65.7)	**< 0.001**[*]
No	152 (36.0)	13 (76.5)	139 (34.3)	
Malaria episode during pregnancy				
Yes	59 (14.0)	1 (5.9)	58 (14.3)	0.326
No	363 (86.0)	16 (94.1)	347 (85.7)	
Gravidity				
Primigravidae	112 (26.5)	2 (11.8)	110 (27.2)	0.051
Multigravidae	310 (73.5)	15 (88.2)	295 (72.8)	
Parity				
Primiparity	122 (28.9)	3 (17.7)	119 (29.4)	**0.035**[*]
Multiparity	300 (71.1)	14 (82.3)	286 (70.6)	
Mode of delivery				
Vaginal delivery	379 (89.8)	15 (88.2)	364 (89.9)	0.827
Caesarean section	43 (10.2)	2 (11.8)	41 (10.1)	
Gestational age at delivery				
< 37 weeks (preterm)	91 (21.6)	4 (23.5)	87 (21.5)	0.822
≥ 37 weeks (normal term)	331 (78.4)	13 (76.5)	318 (78.5)	
First-trimester anaemia				
Anaemia	268 (63.5)	11 (64.7)	257 (63.4)	0.554

No anaemia	154 (35.5)	6 (35.3)	148 (36.6)	
Second-trimester anaemia				
Anaemia	301 (71.3)	13 (76.5)	288 (71.1)	0.700
No anaemia	121 (28.7)	4 (23.5)	117 (28.9)	
Third-trimester anaemia				
Anaemia	191 (45.2)	13 (76.5)	178 (44.0)	**0.030**[*]
No anaemia	231 (54.8)	4 (23.5)	227 (55.0)	
Blood rhesus type				
Positive	379 (89.8)	15 (88.2)	364 (89.9)	0.455
Negative	43 (10.2)	2 (11.8)	41 (10.1)	
Sickle cell status				
Positive	159 (37.8)	7 (41.2)	152 (37.5)	**0.003**[*]
Negative	263 (62.2)	10 (58.8)	253 (62.5)	
G6PD status				
Normal	309 (73.2)	14 (82.4)	295 (72.8)	0.732
Complete/partial defect	113 (26.8)	3 (17.6)	110 (27.2)	
TD immunization				
Immunized	31 (7.4)	3 (17.7)	28 (6.9)	0.121
Not immunized	391 (92.6)	14 (82.3)	377 (93.1)	

[*] $p < 0.05$ [a] *Chi-square/Fisher's exact test*   *G6PD:Glucose-6-phosphate dehydrogenase TD:Tetanus-diphtheria*

Bivariate association of stillbirths with background variables

Among the sociodemographic, medical and obstetric or antenatal characteristics (variables) of the study participants, only sulphadoxine-pyrimethamine intake (p=0.012), insecticide-treated bed net use (p<0.001), parity (p=0.035), third-trimester anaemia status (p=0.030) and sickle cell status (p=0.003) were significantly associated with birth outcome prognosis (stillbirths) during bivariate analysis as shown in Tables 1 and 2.

Maternal factors associated with stillbirths

In the multiple logistic regression analysis, as exhibited in Table 3, three variables namely insecticide-treated bed net use, third-trimester anaemia status and sickle cell status, remained statistically significant in the final model. The model revealed that the odds of mothers with

anaemia in the third trimester of pregnancy having stillbirths was 3.15 times (95%CI: 2.94 –

10.6; p=0.034) compared to the non-anaemic mothers. Moreover, mothers with positive sickle

cell status had 2.91 (95%CI: 1.50 – 7.01; p=0.018) increased odds of stillbirths compared to their

counterparts with negative sickle cell status. Finally, participants who did not sleep under

insecticide-treated bed nets throughout pregnancy had 7.75 (95%CI: 2.33 – 9.80; p=0.001)

higher odds of stillbirths than those who used it.

Table 3: Multiple logistic regression of maternal factors of stillbirths

Variables	Logistic regression analyses			
	Univariate analysis		Multivariate analysis	
	cOR (95%CI)	p-value	aOR (95%CI)	p-value
Sulphadoxine-pyrimethamine intake				
1 – 3 doses		1.000		1.000
> 3 doses	0.11 (0.15 – 0.88)	0.037[*]	0.17 (0.02 – 1.35)	0.094
None	1.46 (0.31 – 6.82)	0.631	2.68 (0.41 – 17.3)	0.301
Insecticide-treated bed nets use				
Yes		1.000		1.000
No	6.22 (3.99 – 9.40)	0.002[*]	**7.75 (2.33 – 9.80)**	**0.001[*]**
Parity				
Primiparity		1.000		1.000
Multiparity	6.66 (3.87 – 9.70)	0.047[*]	4.64 (0.57 – 7.60)	0.151
Third-trimester anaemia				
No anaemia		1.000		1.000
Anaemia	4.02 (2.29 – 12.5)	0.017[*]	**3.15 (2.94 – 10.6)**	**0.034[*]**
Sickle cell status				
Negative		1.000		1.000
Positive	1.51 (1.10 – 7.41)	0.019[*]	**2.91 (1.50 – 7.01)**	**0.018[*]**
Regression model				
R^2		0.6030		
p value		< 0.001		

405 [*] *p < 0.05   cOR: crude odds ratio   aOR: adjusted odds ratio   Bold: significant aOR, p-value, and 95%CI*
406 *for risk factors*

Discussion

The global occurrence of stillbirth as a critical public health issue not only increases the rate of

perinatal and neonatal mortalities but also affects the mental health of mothers and their

families (2,3). In poor economic communities like the Ghanaian context, most stillbirths are

generally higher and usually go unrecorded (16,17).

The prevalence of stillbirths in the present study was found to be 4.0%, which is in line

with a multicenter study in Nigeria (9). This similarity might be due to the uniformity in the

healthcare system and provision of services for pregnant mothers in Ghana and Nigeria. Our

study's stillbirth rate was also lower than the estimation in a retrospective cohort study

conducted in a tertiary facility in Southern Ghana (10.6%) (18) but higher than the study

findings from China (0.9%) and South Africa (3.5%) (7,8). The discrepancy might be due to

variations in the study area and the inclusion and exclusion criteria (methodologies) used in the

previous studies. In addition, several unreported cases of stillbirths in the community and/or

municipal health facilities could be responsible for this variation. In comparison, China and

South Africa have advanced and modernised healthcare systems that offer well-tailored

services to pregnant women, thereby contributing to the reduction of stillbirth rates in these

countries. Therefore, it could indicate that the care for women during antenatal and labour

(intrapartum) might not have received appropriate and standardized obstetric practices and

protocols, leading to higher stillbirth incidence in this semi-rural Ghana. On the other side, the

lesser prevalence of stillbirths in this study, compared to Southern Ghana, might be due to the

deployment of trained traditional birth attendants as agents of facility deliveries and trained

community health nurses to perform the duties of midwives in the Savelugu municipality. This

initiative by the Catholic Relief Services (development partner) was adopted by the Northern

Regional Health Directorate of Ghana Health Service to improve maternal and perinatal health

outcomes, which might have contributed to the reduction of the stillbirth rate in Northern

Ghana. Nonetheless, the relatively elevated incidence of stillbirths in the study area is still

alarming as postpartum mothers and their families face psychological trauma after

experiencing stillbirths during or following a nine-month journey of pregnancy (9,33).

In this study, maternal anaemia in the third trimester of pregnancy was independently

associated with stillbirths. Whilst a retrospective study in Columbia (34) was in agreement with

the present study, a similar study in Pakistan (35) was not in support. Anaemia is known to be a

significant predictor of malnutrition among pregnant mothers as iron deficiency type is

apparently the most prevalent cause of anaemia during gestation (36). As anaemia causes

inadequate oxygen and nutrient supply to the foetus during the late stage of pregnancy (37,38),

it could lead to a deficient supply of these nutrients and oxygen to the foetus which

subsequently causes hypoxic death of the foetus either in-utero, during childbirth or within a

short period after birth (39).

This study also found that mothers with positive sickle cell status had increased odds of

stillbirth delivery. The observed findings align with previous research indicating severe

pregnancy and childbirth complications for both mothers and newborns affected by sickle cell

(40). These complications may stem from the physiological adaptations inherent in the genetic

makeup of pregnant women with sickle cell trait or disease (40,41). Such adaptations can

impact the cardiovascular, haematologic, respiratory, and renal structures in the materno-

foetal system, which can impair the overall health and ability to sustain a healthy pregnancy

(41). Consequently, these physiological challenges may contribute to the reduced likelihood of

delivering a healthy newborn.

Finally, mothers who did not sleep under insecticide-treated bed nets during pregnancy

were more likely to deliver stillbirths. In line with the present findings, mothers' sleep practices

including those who did not sleep under insecticide-treated bed nets during pregnancy, were

prone to stillbirths in a cross-sectional study in Ghana (14) but were contrasting to a meta-

analysis by Flenady and his colleagues (42). Despite the constant supply of free insecticide-

treated bed nets to pregnant mothers for malaria prevention during their first ANC visits in

Ghana (24), it is reported that most mothers do not sleep under the nets due to unfavourable

weather conditions (hot weather) in Northern Ghana, while others also use the nets for

subsistence farming purposes in their homes (43). The non-use of the bed nets could expose

them to malaria infections and subsequent anaemia in pregnancy, thereby increasing their risk

of stillbirth delivery (44).

Interestingly, a large proportion of the study participants who had positive sickle cell

status and did not use insecticide-treated bed nets also made less than eight ANC visits and had

anaemia during each trimester of pregnancy. This implies that fewer ANC visits and low

haemoglobin levels in each trimester during pregnancy could precipitate unfavourable intra-

uterine and gestational conditions (37,38) leading to an increased maternal risk for stillbirth

delivery (44,45) in the municipality. Therefore, greater attention should be paid to anaemia in

pregnancy and ANC visits to strengthen maternal and child health outcomes in the country.

Strengths and limitations of the study

The study recorded some limitations. First, the prevalence of stillbirths in the municipality may

be underestimated or overestimated since the study participants were sampled from six-month

cross-sectional retrospective data from health facilities. There could be misreporting and

misrecording of stillbirths by healthcare professionals, which could affect the estimates of

stillbirths in the municipality. The data lacked details on whether the stillbirth happened in the

antepartum or intrapartum stages of delivery. The absence of this information could imply a

considerable blend of antepartum and intrapartum stillbirths in the dataset. Additionally, the

use of retrospective data (including maternal weight, haemoglobin levels, and gestational age)

from records review (MCHRBs and delivery registers) could be a limitation of the study as a

result of possible measurement and documentation errors. It is difficult to ascertain whether

the necessary standard protocols were adequately adhered to before measurement, potentially

affecting the dataset's quality in the analysis. Notwithstanding, the stillbirth estimates and

measurements represent the available information in the municipal health facilities which are

used for decision-making in the municipality.

The study also faced limitations in generalising the maternal factors of stillbirths

because it relied on cross-sectional data. Therefore, the findings of this study could not be used

to draw causal inferences but rather to highlight significant associations of stillbirths to the

scientific community. Ultimately, the study lacked information on certain variables related to

the independent (exposure) variables. Specifically, maternal data on smoking status, alcohol

intake and socioeconomic status were not collected. Despite this, the study provided significant

findings for potential interventions in the municipality and the broader semi-rural Ghana. On

the other hand, the study encountered some notable strengths. At the outset, the study

addresses a significant gap in the maternal and perinatal health literature by quantitatively

examining the association between maternal factors and stillbirths. This directly connects

maternal health factors to perinatal health. The study goes beyond reporting univariate and

bivariate associations by also conducting multivariable analysis to identify maternal factors of

stillbirths, resulting in a more rigorous and robust set of findings.

Conclusion

The prevalence of stillbirths is relatively high in this semi-rural municipality. These stillbirths are

independently influenced by medical factors, particularly anaemia and positive sickle cell

statuses and unsafe health practices such as non-use of ITNs during pregnancy. Timely

identification of these factors among pregnant women (anaemia and sickle cell status) is vital to

prompt early intervention strategies in Ghanaian health facilities. This could be achieved by

strengthening obstetric triage at ANC clinics and labour/delivery rooms and also encouraging

early referrals to higher levels of care. Secondly, due to the frequent misclassification of most

perinatal and neonatal deaths as stillbirths (19), thorough perinatal death audits, which aim to

identify the cause and contributing factors of stillbirths, should be promoted in healthcare

facilities in semi-rural Ghana.

Additionally, the management of Ghana Health Service (especially municipal/district

health directorates), alongside program planners and other stakeholders involved in enhancing

perinatal and maternal health, should also be aware of these maternal factors contributing to

stillbirths. These health implementers should launch effective programs such as robust referral

systems through training and capacity building of health care providers and community cadres

to facilitate early referral and management. Policy efforts should further be targeted at

promoting enhanced health education sessions on undernutrition (anaemia) and the regular

use of ITNs during pregnancy at facility ANC clinics and communities (via community durbars) in

semi-rural Ghana. Finally, we recommend future research to incorporate other maternal

variables, such as socioeconomic status, prenatal smoking and alcohol intake, to thoroughly

examine and provide a more comprehensive understanding of the maternal factors

contributing to stillbirths in both the municipality and the wider Northern Region.

List of abbreviations

ANC: Antenatal care

BMI: Body mass index

G6PD: Glucose-6-phosphate dehydrogenase

ITNs: Insecticide-treated bed nets

MCHRBs: Maternal and child health record books

STATA: Statistics and data

TD: Tetanus-diphtheria

VIF: Variance inflation factor

WHO: World Health Organization

Declarations

Ethics approval and consent to participate

This study was conducted according to the guidelines laid down in the Declaration of Helsinki,

and all procedures involving research study participants were approved by the Navrongo Health

Research Centre Institutional Review Board with approval number NHRCIRB373. Permission

was also granted by the Ghana Health Service at the municipal directorate and health facilities

level. Written informed consent/assent was obtained from all study participants and/or legal

representatives and also from parents/guardians of participants under 18 years of age.

Consent For Publication

Not applicable

Availability of data and materials

The datasets collected, generated, and/or analyzed during the present study are available from

the corresponding author upon reasonable request. Data sharing not applicable to this article

as no datasets were generated or analyzed during the current study.

Competing interest

The authors declared that they have no competing interests.

Funding statement

There was no funding support

Authors' contributions

SAG, AA, MG, CSD and PAA designed and conceptualized the study. SAG, AA, MG, WP, CA and

CSD carried out the data collection while SAG, WP, CA and PAA analyzed and interpreted the

data. All authors contributed to drafting the initial manuscript, reviewing and approving the

final manuscript.

Authors' acronyms

Silas Adjei-Gyamfi (SAG); Abigail Asirifi (AA); Mamudu Gafaru (MG); Wisdom Peprah (WP);

Clotilda Asobuno (CA); Constance Siakwan Dapaah (CSD); Paul Armah Aryee (PAA)

Acknowledgements

The authors are thankful to all research assistants, study participants, and volunteers who

contributed in diverse ways to the successful implementation of the study. We are also grateful

References

594   1.   Lawn JE, Blencowe H, Waiswa P, Amouzou A, Mathers C, Hogan D, et al. Stillbirths: Rates,

595   risk factors, and acceleration towards 2030. Vol. 387, The Lancet. Lancet Publishing

596   Group; 2016. p. 587–603.

597   2.   Bawah AT, Ngala RA, Alidu H, Seini MM, Wumbee JDK, Yeboah FA. Gestational diabetes

598   mellitus and obstetric outcomes in a Ghanaian community. Pan Afr Med J. 2019;32.

599   3.   WHO. Every newborn: An action plan to end Preventable deaths: Executive summary

600   Geneva: World Health Organization. Who, Unicef. 2014;12 p.

601   4.   Harding EBM, Harrington LT, Lockwood CM, Duncan Brown JR, Hall AC, Brown MA, et al.

602   Perinatal and neonatal mortality. Br Med J. 2006;281(6254):1567.

603   5.   Blencowe H, Cousens S, Jassir FB, Say L, Chou D, Mathers C, et al. National, regional, and

604   worldwide estimates of stillbirth rates in 2015, with trends from 2000: A systematic

605   analysis. Lancet Glob Heal. 2016 Feb 1;4(2):e98–108.

606   6.   Hug L, You D, Blencowe H, Mishra A, Wang Z, Fix MJ, et al. Global, regional, and national

607   estimates and trends in stillbirths from 2000 to 2019: A systematic assessment. Lancet.

608   2021;398(10302):772–85. http://dx.doi.org/10.1016/S0140-6736(21)01112-0

7.   Zhu J, Liang J, Mu Y, Li X, Guo S, Scherpbier R, et al. Sociodemographic and obstetric

characteristics of stillbirths in China: A census of nearly 4 million health facility births

between 2012 and 2014. Lancet Glob Heal. 2016 Feb 1;4(2):e109–18.

8.   Lavin T, Pattinson RC, Nedkoff L, Gebhardt S, Preen DB. Stillbirth risk across pregnancy by

size for gestational age in Western Cape Province, South Africa: Application of the

fetuses-at-risk approach using perinatal audit data. South African Med J.

2019;109(12):927–33.

9.   Okonofua FE, Ntoimo LFC, Ogu R, Galadanci H, Mohammed G, Adetoye D, et al.

Prevalence and determinants of stillbirth in Nigerian referral hospitals: A multicentre

study. BMC Pregnancy Childbirth. 2019 Dec 30;19(1).

10.   Afulani PA. Determinants of stillbirths in Ghana: Does quality of antenatal care matter?

BMC Pregnancy Childbirth [Internet]. 2016;16(1):1–17.

http://dx.doi.org/10.1186/s12884-016-0925-9

11.   Lawn JE, Yakoob M, Haws RA, Soomro T, Darmstadt GL, Bhutta ZA. 3.2 million stillbirths:

Epidemiology and overview of the evidence review. Vol. 9, BMC Pregnancy and

Childbirth. BioMed Central; 2009. p. 1–17.

12.   Dassah ET, Odoi AT, Opoku BK. Stillbirths and very low Apgar scores among vaginal births

in a tertiary hospital in Ghana: A retrospective cross-sectional analysis. BMC Pregnancy

Childbirth. 2014 Aug 28;14(1).

13.   Njoku C., Emechebe C., Eyong E., Ukaga J., Anachuna K. Prevalence and risk factors for

stillbirths in a tertiary hospital in Niger Delta area of Nigeria: a ten year review. Int J Med

Biomed Res. 2016 Oct 3;5(3):106–13.

14.   Owusu JT, Anderson FJ, Coleman J, Oppong S, Seffah JD, Aikins A, et al. Association of

maternal sleep practices with pre-eclampsia, low birth weight, and stillbirth among

Ghanaian women. Int J Gynecol Obstet. 2013;121(3):261–5.

http://doi.wiley.com/10.1016/j.ijgo.2013.01.013

15.   Bhutta ZA, Darmstadt GL, Haws RA, Yakoob MY, Lawn JE. Delivering interventions to

reduce the global burden of stillbirths: Improving service supply and community demand.

BMC Pregnancy Childbirth 2009;9(1):1–37. 10.1186/1471-2393-9-S1-S7

16.   Ghana Statistical Service (GSS) Ghana Health Service (GHS) and ICF. 2017 Ghana

Maternal Health Survey (GMHS) Research Policy Briefs [Internet]. GSS, GHS, and ICF.

2019. https://statsghana.gov.gh/gssmain/fileUpload/pressrelease/PB14.pdf

17.   Ghana Statistical Service (GSS), Ghana Health Service (GHS) and MI. Ghana maternal

health survey 2007. GSS, GHS, and Macro International. Calverton, Maryland, USA; 2009.

18.   Adu-Bonsaffoh K, Gyamfi-Bannerman C, Oppong SA, Seffah JD. Determinants and

outcomes of preterm births at a tertiary hospital in Ghana. Vol. 79, Placenta. W.B.

Saunders Ltd; 2019. p. 62–7.

19.   Nonterah, A. Engelbert ., Isaiah AA, Edmund WK, Kagura J, Tamimu M, Ayamba YE,

Nonterah WE, et al. Trends and risk factors associated with stillbirths: A case study of

the Navrongo War Memorial Hospital in Northern Ghana. PLoS One. 2020;15(2):1–13.

https://doi.org/10.1371/journal.%0D%0Apone.0229013

20.   Ameyaw EK, Kofinti RE, Appiah F. National health insurance subscription and maternal

651   healthcare utilisation across mothers' wealth status in Ghana. Health Econ Rev.

652   2017;7(1):16.https://healtheconomicsreview.biomedcentral.com/articles/10.1186/s1356

653   1-017-0152-8

654   21.   Adjei-Gyamfi S, Musah B, Asirifi A, Hammond J, Aryee PA, Miho S, et al. Maternal risk

655   factors for low birthweight and macrosomia: A cross-sectional study in Northern Region,

656   Ghana. J Heal Popul Nutr. 2023;42(1):1–16. https://doi.org/10.1186/s41043-023-00431-0

657   22.   Adjei-Gyamfi S, Asirifi A, Peprah W, Abbey DA, Hamenoo KW, Zakaria MS, et al. Anaemia

658   at 36 weeks of pregnancy: Prevalence and determinants among antenatal women

659   attending peri-urban facilities in a developing country, Ghana. PLOS Glob Public Heal.

660   2024;4(9):e0003631. Available from: https://doi.org/10.1371/journal.pgph.0003631

661   23.   Adjei-Gyamfi S, Zakaria MS, Asirifi A, Issahaku S, Ibrahim MA, Aryee PA. Maternal

662   anaemia and polycythaemia during pregnancy and risk of inappropriate birth weight for

663   gestational age babies: A retrospective cohort study in the northern belt of Ghana. BMJ

664   Open. 2024;14:e082298. https://doi.org/10.1136/bmjopen-2023-082298

665   24.   Adjei-Gyamfi S, Asirifi A, Aiga H. Prevalence and associated risk factors of preterm and

666   post-term births in Northern Ghana: A retrospective study in Savelugu Municipality. J

667   Pediatr Perinatol Child Heal. 2023;07(04):235–48.

668   https://doi.org/10.26502/jppch.74050169

669   25.   Cochran WG. Samling Techniques. John Wiley & Sons, Inc. 1977. 448 p.

670   26.   WHO Web Annex 2. Template protocol for surveys to estimate the prevalence of

671   biomarkers of infection with the hepatitis viruses. In: Consolidated strategic information

672   guidelines for viral hepatitis planning and tracking progress towards elimination. Geneva;

673   2019.

674   27.   Adjei-Gyamfi S, Asirifi A, Asobuno C, Korang FK. Knowledge and occupational practices of

675   beauticians and barbers in the transmission of viral hepatitis: A mixed-methods study in

676   Volta   Region   of   Ghana.   PLoS   One.   2025;20(1):1–24.

677   http://dx.doi.org/10.1371/journal.pone.0306961

678   28.   Engmann C, Walega P, Aborigo RA, Adongo P, Moyer CA, Lavasani L, et al. Stillbirths and

679   early neonatal mortality in rural Northern Ghana. Trop Med Int Heal. 2012;17(3):272–82.

680   29.   Asobuno C, Adjei-Gyamfi S, Aabebe FG, Hammond J, Taikeophithoun C, Amuna NN, et al.

681   Risk factors for anaemia among pregnant women: A cross-sectional study in Upper East

682   Region,   Ghana.   PLoS   One.   2024;19(11):e0301654.

683   https://doi.org/10.1371/journal.pone.0301654

684   30.   GHS. DHIMS-2: District Information Management Systems version 2 [Internet]. Ghana

685   Health Service. 2020. Available from: https://dhims.chimgh.org/dhims

686   31.   Ghana Statistical Service (GSS), Ghana Health Service (GHS) & II. Ghana Demographic and

687   Health Survey 2014. Rockville, Maryland, USA; 2015.

688   32.   Rasmussen KM, Catalano PM, Yaktine AL. New guidelines for weight gain during

689   pregnancy: what obstetrician/gynecologists should know. Curr Opin Obs Gynecol.

2009;21(6):521–6.

33.   Vogel JP, Souza JP, Mori R, Morisaki N, Lumbiganon P, Laopaiboon M, et al. Maternal

complications and perinatal mortality: findings of the World Health Organization

Multicountry Survey on Maternal and Newborn Health. BJOG. 2014;121 Suppl 1(Suppl

1):76–88.

34.   Smith C, Teng F, Branch E, Chu S, Joseph KS. Maternal and Perinatal Morbidity and

Mortality Associated with Anemia in Pregnancy. Obstet Gynecol. 2019 Dec

1;134(6):1234–44.

35.   Mahmood T, Rehman AU, Tserenpil G, Siddiqui F, Ahmed M, Siraj F, et al. The Association

between Iron-deficiency Anemia and Adverse Pregnancy Outcomes: A Retrospective

Report from Pakistan. Cureus. 2019 Oct 7;11(10).

36.   Garzon S, Cacciato PM, Certelli C, Salvaggio C, Magliarditi M, Rizzo G. Iron Deficiency

Anemia in Pregnancy: Novel Approaches for an Old Problem. Oman Med J. 2020

Sep;35(5):e166.

37.   Barker DJP. Intrauterine programming of adult disease. Vol. 1, Molecular Medicine

Today. Elsevier Current Trends; 1995. p. 418–23.

38.   Wu G, Imhoff-Kunsch B, Girard AW. Biological mechanisms for nutritional regulation of

maternal health and fetal development. Vol. 26, Paediatric and Perinatal Epidemiology.

John Wiley & Sons, Ltd; 2012. p. 4–26.

39.   Parks S, Hoffman MK, Goudar SS, Patel A, Saleem S, Ali SA, et al. Maternal anaemia and

maternal, fetal, and neonatal outcomes in a prospective cohort study in India and

Pakistan. BJOG An Int J Obstet Gynaecol. 2019 May 1;126(6):737–43.

40. Smith-Whitley K. Complications in pregnant women with sickle cell disease. Hematology.

2019;2019(1):359–66. Available from: https://doi.org/10.1182/hematology.2019000039

41. Jain D, Atmapoojya P, Colah R, Lodha P. Sickle cell disease and pregnancy. Mediterr J

Hematol Infect Dis 2019;11(1):e2019040. http://dx.doi.org/10.4084/MJHID.2019.040

42. Flenady V, Koopmans L, Middleton P, Frøen JF, Smith GC, Gibbons K, et al. Major risk

factors for stillbirth in high-income countries: a systematic review and meta-analysis.

Lancet. 2011 Apr 16;377(9774):1331–40.

43. Savelugu Municipal Health Directorate (Ghana Health Service). Annual Performance

Report 2020: Savelugu Municipal Health Directorate. Savelugu, Northern Region; 2021.

44. Moore KA, Simpson JA, Scoullar MJL, McGready R, Fowkes FJI. Quantification of the

association between malaria in pregnancy and stillbirth: a systematic review and meta-

analysis. Lancet Glob Heal. 2017 Nov 1;5(11):e1101–12.

45. Idung A, Jombo H, Udonwa NE, Iyanam VE, Idung AU, Jombo HE. Anaemia in Pregnancy

at Booking: Prevalence and Risk Factors among Antenatal Attendees in a Southern

Nigeria General Hospital Article in Asian Journal of Medicine and Health · Asian J Med

Heal. 2019;15(1):48805.