1

Title

Maternal risk factors for stillbirths in a semi-rural municipality in Ghana: Findings

from a retrospective observational study

Author's information

* **Silas Adjei-Gyamfi, MPH, MPhil (Corresponding author)**

    Savelugu Municipal Hospital

    Ghana Health Service

    P. O. Box 45, Savelugu, Northern Region, GHANA

    Phone: +233243421318

    Email: adjeigyamfis@yahoo.com

Abigail Asirifi, Dip, BSc

    Department of Midwifery

    Garden City University College

    P. O. Box KS 12775, Kumasi, Ashanti Region, GHANA

    Phone: +233545951777

    Email: abimaiden80@gmail.com

Gafaru Mamudu, BSc

    Savelugu Municipal Hospital

    Ghana Health Service

P. O. Box 45, Savelugu, Northern Region, GHANA

Phone: +233248303028

Email: gafarumamudu@gmail.com

Wisdom Peprah, BSc, MPhil

School of Allied Health Sciences

University for Development Studies

P. O. Box TL 1883, Tamale, Northern Region, GHANA

Phone: +233507093898

Email: pwisdom466@gmail.com

Clotilda Asobuno, BSc, MPH

Kassena Nankana West District Health Directorate

Ghana Health Service

P. O. Box 21, Paga, Upper East Region, GHANA

Phone: +233245452488

Email: clotildaasobuno@gmail.com

Constance Siakwan Dapaah, BSc

Ghana Health Service

Atwima Nwabiagya North District Health Directorate

P.O. Box 26, Barekese, Ashanti Region, GHANA

Phone: +233245727275

Email: constancedapaah25@gmail.com

Paul Armah Aryee, MMedSc, MPhil, PhD

School of Allied Health Sciences

University for Development Studies

P. O. Box TL 1883, Tamale, Northern Region, GHANA

Phone: +233 200 516610

Email: parmah@uds.edu.gh

Abstract

Background

Stillbirths do not only trigger maternal psychological trauma but also affect the socio-economic

growth in developing countries. Since the prevalence and predictors of stillbirths and their

dynamics are diverse in different low-resource settings like Ghana, it requires adequate

knowledge to mitigate these factors for reducing perinatal deaths. Moreover, data are scarce

on the maternal factors of stillbirth in Ghana. This study, therefore, aimed to determine the

prevalence and associated factors of stillbirths in a semi-rural municipality in Northern Ghana.

Methods

A retrospective observational study was conducted among 422 mothers who delivered in public

health facilities and reside in a semi-rural municipality of Northern Ghana. Sociodemographic,

obstetric, and medical-related data were collected through semi-structured interviews and

from records review. We used binary logistic regression to identify the independent factors of

stillbirths at a confidence level of 95%.

Results

Stillbirth rate was estimated at 4.0% (95%CI: 2.4%–6.4%). The analysis revealed that anaemia in

the third trimester of pregnancy (aOR:3.15; 95%CI:2.94–10.6; p=0.034), positive sickle cell

status (aOR:2.91; 95%CI:1.50–7.01; p=0.018), and non-use of insecticide-treated bed nets

during pregnancy were significantly associated with increased risk of stillbirths (aOR:7.75;

95%CI:2.33–9.8; p=0.001).

Conclusion

The relatively high stillbirth prevalence in the municipality was affected by medical and lifestyle

factors. Management of the municipal health directorate (Ghana Health Service), program

planners, and other stakeholders who are engaged in improving perinatal and maternal health

should target and implement effective programs on anaemia and lifestyle modifications for the

prevention and reduction of stillbirths.

Keywords

Anaemia,

Determinants,

Risk factors,

Northern Ghana,

Predictors,

Stillbirths

Number of words

Abstract   247 words

Main text   2,729 words

Introduction

Stillbirths, also called stillborns, during delivery or childbirth do not only trigger maternal psychological trauma but also contribute to perinatal and maternal mortalities, which affect the socio-economic growth and development in low and middle-income countries (1–3). Stillbirth is denoted as the birth of a foetus with no signs of life occurring at 28 weeks or more of gestation (3), and it accounts for more than half of all perinatal mortalities worldwide (4).

In 2015, an estimated 2·6 million neonates were born with no signs of life globally, which indicates an approximately 19.0% reduced rate since 2000 with a sluggish reduction among Sahelian Africans. About 98.0% of all stillbirths happen in developing countries, with 77.0% occurring among African and Asian pregnancies, especially those from rural settings (5). In China (6), South Africa (7), and Nigeria (8), the prevalence of stillbirths was found to be 0.9%, 3.5%, and 4.0% respectively. Among the Ghanaian population, higher stillbirth rates of 5.9% and 10.6% were respectively recorded in 2013 and 2018 at two popular tertiary hospitals (9,10) which makes these high stillbirth occurrences very worrisome in the country.

The aetiology and antecedents of stillbirths are multifaceted and are affected by sociological, physiological (medical), and physical factors (1). It is reported that the stillbirth rate and its determinants are dependent on the geographic and sociodemographic characteristics of the population (3,5). Therefore, the study findings from one setting might not apply to another. Universally, earlier studies acknowledged that the factors associated with stillbirths include congenital foetal deformities, foetal distress (11), preterm births, breech delivery (10), abruptio placentae, and male foetus (12). Non-use of insecticide-treated bed nets for malaria prevention

during pregnancy (13), maternal age less than 20 years, unmarried mothers (6), employment

status, multiparity, and delayed vaginal delivery (8) were also regarded as the determinants of

stillbirths in other past studies.

To achieve the World Health Assembly's national target of a stillbirth prevalence of 10

or less per 1000 deliveries by 2035 in every country through the Every Newborn Action Plan (3),

the determinants or predictors of stillbirths and their dynamics in different settings must be

well understood and appreciated. Although stillbirths are higher in developing countries,

including Ghana (9,10), relevant information on the predictors or factors of stillbirths is

inadequately documented, readily unavailable and/or scarce in the district, regional, or national

databank. Hence, adequate awareness and proper documentation of the maternal medical,

sociodemographic and obstetric factors of stillbirths in low-resource settings like Ghana could

trigger useful interventions that lead to preventable perinatal mortalities, including stillbirths

and subsequently contribute to improving maternal and child health. This study, therefore,

sought to determine the maternal factors associated with stillbirths in a semi-rural municipality

in the Northern Region of Ghana.

Materials and methods

Study design

This current survey is a retrospective observational study conducted among women who

delivered in a semi-rural municipality in the Northern Region of Ghana. This design was

employed to identify the associated factors of stillbirths.

Study settings

This study was conducted in Savelugu Municipality of Northern Ghana. Savelugu Municipality is

one of the 16 districts/municipalities in the Northern Region of Ghana that shares boundaries

with five neighbouring districts, including Gushegu, Karaga, Kumbungu, Nanton, and Tolon. The

municipality has five major government (public) health facilities (one hospital and four health

centres) that provide comprehensive maternal and newborn services, including free maternal

and delivery services (14) to a population of approximately 130,000. In 2019, the coverage of

antenatal and postnatal visits was 94% and 98%, respectively (15,16). Due to the increased

average monthly delivery count of over 300 deliveries with higher records of unfavourable birth

outcomes (15,17,18), this municipality was selected to investigate the predictors of stillbirths

among women who delivered in government health facilities.

Sample size and sampling procedure

The sample size (n) was estimated using Cochran's method (19), $n = \dfrac{Z^2 PQ}{e^2}$. We used the normal

standard variate (Z) at a 95% confidence interval of 1.96, a standard error (e) of 5%, a

population proportion (P) of 50% since published data are scarce on stillbirths in the region and

Q=1–P, to calculate the preliminary sample size of 384. To account for fall-outs, incomplete

responses and damaged questionnaires, an additional 10 percent was added to the calculated

sample size (15,17,20), resulting in a final total of approximately 422.

In the sampling stage, all five major government (public) health facilities were chosen

for the study since all these facilities report on stillbirths every year (21). The sample sizes for

each facility were determined using the probability proportional to size technique. The delivery

registers from each facility were used to line-list all eligible postpartum women (who delivered

from January 1 to June 31, 2020) to form a sampling frame. A simple random sampling method

was then used to select the required number of respondents from each facility. The addresses

and contact numbers of these selected respondents were recorded to trace them to their

homes.

Study population

The study recruited a total of 422 women who visited the five major public facilities in the

Savelugu Municipal for childbirth or delivery purposes from January 1 to June 31, 2020. All

women who lived in the municipality, delivered at the municipal health facilities and had their

health information recorded in the delivery registers were eligible for the study.

Data collection

Data was collected using a pretested questionnaire by six well-trained data collectors (midwives

and public health officers) from July 16 to September 31, 2020. The questionnaire was

developed based on existing literature, previous studies, and sociological and biological

plausibility (1,5,15,22). The questionnaire is attached as a supplementary material (Suppl 1).

We verified the content validity of the questionnaire by consulting with experts in the field as

well as the study authors. Sociodemographic, obstetric and medical-related information was

collected through semi-structured interviews and from record reviews (delivery registers and

maternal and child health record books [MCHRBs]). The data collectors were trained for one

week. The data collection procedure was in two phases. The first phase was at the household

level, while the second was at the health facilities level after sampling and contacting the

eligible study participants.

Before the commencement of household data collection, the trained data collectors

retrieved the addresses and contact information (telephone numbers) of respondents and/or

their representatives from the delivery register with the support of the facility midwives. The

data collectors arranged appointments and then visited the homes of the recruited

respondents. Data not excluding age, marital status, education level, job status, ethnicity and

religious affiliations were collected via structured interviews and were validated in the

MCHRBs, while information about first-trimester body mass index (since foetal weight at the

first trimester is negligible (23)), antenatal visits, parity, gravidity, rhesus status, sickle cell

status, tetanus-diphtheria (TD) immunization status, haemoglobin levels across pregnancy

trimesters, use of pre-pregnancy family planning, and insecticide-treated bed nets (1,18) were

collected from their MCHRBs at their homes. Afterwards, information like pre-eclampsia

diagnosis, postpartum haemorrhage, delivery mode, birth/delivery outcome prognosis, and

gestational age at birth (1,18) were traced and extracted from the delivery registers at the

health facilities level.

Study variables

The outcome variable was birth outcome prognosis, categorized into binary outcomes for

stillbirths (coded as 1) or livebirths (coded as 0) while exposure variables included all selected

sociodemographic, obstetric, and medical variables which were categorical.

According to the World Health Organization (WHO), stillbirths were measured as the

birth of a foetus at 28 or more weeks of gestation with no signs of life, while livebirths were

regarded as newborns with signs of life (3). Haemoglobin levels were dichotomized into

anaemia (< 11g/dl), and no anaemia (≥ 11g/dl), while gestational age at birth was grouped into

< 37 weeks (preterm) and ≥ 37 weeks (normal-term). Parity was classified as primiparity (one

delivery) and multiparity (two or more deliveries), while gravidity was grouped into

primigravida (one pregnancy) and multigravida (two or more pregnancies) (16–

18,20). Standardized categories of pre-pregnancy body mass index (BMI) of underweight (<18.5

kg/m^2), normal BMI (18.5–24.9 kg/m^2) and overweight (≥25.0 kg/m^2) were used. Rhesus and

sickle cell statuses were grouped as positive or negative, while antenatal visits were also

grouped into ≥ 8 visits and < 8 visits using the new 2016 WHO antenatal model (15,22).

Statistical analysis

Frequencies and percentages were calculated for sociodemographic, medical and obstetric

characteristics. Chi-square/Fisher's analysis test of proportions was employed to determine the

statistical association between birth outcome prognosis (stillbirth) and all exposure

variables. Multicollinearity among the significant exposure variables was then assessed with

variance inflation factor (VIF), where variables with a VIF less than five were selected into

multivariate logistic regression analyses (15,16,18).

Applying simultaneous variable entry, the significant exposure variables with no

multicollinearity were forwarded into the multivariate binary logistic regression model and run

to determine significant factors of stillbirths. All analyses were computed using STATA 17.0

(Stata Corporation, Texas, USA), and a p-value of less than 0.05 was considered significant

during statistical analysis.

<h1 style="text-align:center">Results</h1>

Prevalence of stillbirths in a semi-rural municipality of Ghana

All the data from the sampled 422 mothers were used for the analysis. The prevalence of stillbirths in this semi-rural municipality was estimated at 4.0% (95%CI: 2.4% − 6.4%), as seen in Figure 1.

277
Figure 1. Prevalence of stillbirths in a semi-rural municipality of Ghana

Sociodemographic, medical and obstetric characteristics of study participants

The average age of the mothers was 27.6 (sd=6.0) years, while teenage mothers were

estimated at 9%. A large proportion of the participants could not read nor write (no formal

education) (43%), were unemployed (64%), and were married (92%) (Table 1). Anaemia was

highest in the second (71%), followed by the first (64%) and third (45%) trimesters of

pregnancy. Few of the mothers had delivered once (28%), attended antenatal clinic at least

eight times (28%), ingested sulphadoxine-pyrimethamine at least three times (35%), and did

not use insecticide-treated bed nets during pregnancy (36%) (Table 2).

Table 1: Association of sociodemographic characteristics with stillbirths in a semi-rural
municipality of Ghana

Characteristics	Total (%)	Stillbirths (%)	Livebirths (%)	p-value [a]
Age category of mother				
< 20 years	38 (9.0)	1 (5.9)	37 (9.2)	0.573
20 – 35years	340 (80.6)	13 (76.5)	327 (80.7)	
> 35 years	44 (10.4)	3 (17.6)	41 (10.1)	
Mean = 27.63 (SD=6.02)				
Marital status of mother				
Married	389 (92.2)	1 (5.9)	29 (7.2)	1.000
Not married	33 (7.8)	16 (94.1)	376 (92.8)	
Education level of mother				
No education	182 (43.1)	7 (41.2)	175 (43.2)	0.314
Primary school	58 (13.8)	1 (5.9)	57 (14.1)	
High school	138 (32.7)	9 (52.9)	129 (31.8)	
Tertiary	44 (10.4)	0 (0.0)	44 (10.9)	
Ethnicity of mother				
Dagomba	340 (80.6)	13 (76.5)	327 (80.7)	0.354
Gonja	22 (5.2)	0 (0.0)	22 (5.4)	
Frafra/Grusi	34 (8.1)	3 (17.7)	31 (7.7)	
Others (Asante, Ewe, Nigeria)	26 (6.1)	1 (5.9)	25 (6.2)	
Religion of mother				
Muslim	372 (88.2)	13 (76.5)	359 (88.6)	0.158
Christian	50 (11.8)	4 (23.5)	46 (11.4)	
Occupation of mother				
Unemployed	269 (63.7)	16 (94.1)	253 (62.5)	0.303
Employed	153 (36.3)	1 (5.9)	152 (37.5)	
Sex of baby (foetus)				
Male	210 (49.8)	6 (35.3)	204 (50.4)	0.322
Female	212 (50.2)	11 (64.7)	(49.6)	

290 [*] p < 0.05   [a] Chi-square/Fisher's exact test

293 **Table 2: Association of medical and obstetric characteristics with stillbirths in a semi-rural**
294 **municipality of Ghana**

Characteristics	Total (%)	Stillbirths (%)	Livebirths (%)	p-value [a]
First-trimester body mass index (BMI)				
Underweight	13 (3.1)	1 (5.9)	13 (3.2)	0.423
Normal BMI	279 (66.1)	12 (64.7)	267 (65.9)	
Overweight	130 (30.8)	5 (29.4)	125 (30.9)	
Number of antenatal visits				
< 8 visits	304 (72.0)	14 (82.3)	290 (71.6)	0.333
≥ 8 visits	118 (28.0)	3 (17.7)	115 (28.4)	
Sulphadoxine-pyrimethamine intake				
None	25 (5.9)	2 (11.8)	23 (5.7)	**0.012***
1 – 3 doses	249 (59.0)	14 (82.4)	235 (58.0)	
> 3 doses	148 (35.1)	1 (5.9)	147 (36.3)	
Pre-eclampsia/Eclampsia				
Yes	7 (1.7)	15 (88.2)	7 (1.7)	0.768
No	415 (98.3)	2 (11.8)	398 (98.3)	
Antepartum haemorrhage				
Yes	30 (7.1)	4 (23.5)	30 (7.4)	0.622
No	392 (92.9)	13 (76.5)	375 (92.6)	
Pre-pregnancy family planning use				
Yes	75 (17.8)	3 (17.7)	72 (17.8)	0.989
No	347 (82.2)	14 (82.3)	333 (82.2)	
Insecticide-treated bed nets use during pregnancy				
Yes	270 (64.0)	4 (23.5)	266 (65.7)	**< 0.001***
No	152 (36.0)	13 (76.5)	139 (34.3)	
Malaria episode during pregnancy				
Yes	59 (14.0)	1 (5.9)	58 (14.3)	0.326
No	363 (86.0)	16 (94.1)	347 (85.7)	
Gravidity				
Primigravidae	111 (26.3)	1 (5.9)	110 (27.2)	0.051
Multigravidae	311 (73.7)	16 (94.1)	295 (72.8)	
Parity				
Primiparity	120 (28.4)	1 (5.9)	119 (29.4)	**0.035***
Multiparity	302 (71.6)	16 (94.1)	286 (70.6)	
Mode of delivery				
Vaginal delivery	379 (89.8)	15 (88.2)	364 (89.9)	0.827
Caesarean section	43 (10.2)	2 (11.8)	41 (10.1)	
Gestational age at delivery				
< 37 weeks (preterm)	91 (21.6)	4 (23.5)	87 (21.5)	0.822
≥ 37 weeks (normal term)	331 (78.4)	13 (76.5)	318 (78.5)	
First-trimester anaemia				
Anaemia	268 (63.5)	11 (64.7)	257 (63.4)	0.554
No anaemia	154 (35.5)	6 (35.3)	148 (36.6)	
Second-trimester anaemia				
Anaemia	301 (71.3)	13 (76.5)	288 (71.1)	0.700
No anaemia	121 (28.7)	4 (23.5)	117 (28.9)	

Third-trimester anaemia				
Anaemia	191 (45.2)	13 (76.5)	178 (44.0)	**0.030**[*]
No anaemia	231 (54.8)	4 (23.5)	227 (55.0)	
Blood rhesus type				
Positive	379 (89.8)	15 (88.2)	364 (89.9)	0.455
Negative	43 (10.2)	2 (11.8)	41 (10.1)	
Sickle cell status				
Positive	153 (36.3)	7 (41.2)	152 (37.5)	**0.003**[*]
Negative	269 (63.7)	10 (58.8)	253 (62.5)	
G6PD status				
Normal	311 (73.7)	14 (82.4)	295 (72.8)	0.732
Complete/partial defect	111 (26.3)	3 (17.6)	110 (27.2)	
TD immunization				
Immunized	31 (7.4)	3 (17.7)	28 (6.9)	0.121
Not immunized	391 (92.6)	14 (82.3)	377 (93.1)	

[*] $p < 0.05$ [a] *Chi-square/Fisher's exact test*

Associated factors of stillbirths in a semi-rural municipality of Ghana

Among the selected characteristics of the study participants, sulphadoxine-pyrimethamine intake (p=0.012), insecticide-treated bed nets use (p<0.001), parity (p=0.035), third-trimester anaemia (p=0.030) and sickle cell status (p=0.003) were significantly associated with birth outcome prognosis (stillbirths) during bivariate analysis (Tables 1 and 2).

In the multivariate logistic regression analysis, insecticide-treated bed net use, third-trimester anaemia and sickle cell status remained significant (Table 3). The study revealed that the odds of women with anaemia in the third trimester of pregnancy having stillbirths was 3.15 times (95%CI: 2.94 – 10.6; p=0.034) compared to the non-anaemic women. Moreover, women with positive sickle cell status had 2.91 (95%CI: 1.50 – 7.01; p=0.018) increased odds of stillbirths compared to their counterparts with negative sickle cell status. Finally, participants who did not sleep under insecticide-treated bed nets throughout pregnancy had 7.75 (95%CI: 2.33 – 9.8; p=0.001) higher odds of stillbirths than those who used it.

Table 3: Logistic regression of associated factors of stillbirths in a semi-rural municipality of Ghana

Characteristics	cOR (95%CI)	p-value	aOR (95%CI)	p-value
Sulphadoxine-pyrimethamine intake				
1 – 3 doses		1.000		1.000
> 3 doses	0.11 (0.15 – 0.88)	0.037[*]	0.17 (0.02 – 1.35)	0.094
None	1.46 (0.31 – 6.82)	0.631	2.68 (0.41 – 17.3)	0.301
Insecticide-treated bed nets use				
Yes		1.000		1.000
No	6.22 (3.99 – 9.4)	0.002[*]	7.75 (2.33 – 9.8)	**0.001[*]**
Parity				
Primiparity		1.000		1.000
Multiparity	6.66 (3.87 – 9.7)	0.047[*]	4.64 (0.57 – 7.6)	0.151
Third-trimester anaemia				
No anaemia		1.000		1.000
Anaemia	4.02 (2.29 – 12.5)	0.017[*]	3.15 (2.94 – 10.6)	**0.034[*]**
Sickle cell status				
Negative		1.000		1.000
Positive	1.51 (1.10 – 7.41)	0.019[*]	2.91 (1.50 – 7.01)	**0.018[*]**
Regression model				
R^2	*0.6030*			
p value	*< 0.001*			

314 [*]*p < 0.05*   *cOR: crude odds ratio*   *aOR: adjusted odds ratio*

Discussion

The global occurrence of stillbirth as a critical public health issue does not only increase the rate of perinatal mortalities but also affects the mental health of mothers and their families (2,3). In poor economic communities like the Ghanaian context, most stillbirths are generally higher and usually go unrecorded (24). The prevalence of stillbirths in the present study was found to be 4.0%, which is in line with a multicenter study in Nigeria (8). This similarity might be due to the uniformity in the healthcare system and provision of services for pregnant mothers in Ghana and Nigeria.

Our study's stillbirth rate was also lower than the estimation (10.6%) found in a retrospective cohort study conducted in a tertiary facility in Southern Ghana (9), but higher than the findings (0.9%) from a Chinese report (6). The discrepancy might be due to variations in the study area and the inclusion and exclusion criteria (methodologies) used. Several unreported cases of stillbirths in the community and/or municipal health facilities could be responsible for this variation. Moreover, the lesser prevalence of stillbirths in this study as compared to Southern Ghana might be due to the deployment of trained traditional birth attendants as agents of facility deliveries, and trained community health nurses to perform the duties of midwives in the Savelugu municipality. This was an initiative of the Catholic Relief Services (development partner) which was adopted by the Northern Regional Health Directorate of Ghana Health Service to improve maternal and child health outcomes.

Nonetheless, the relatively high prevalence of stillbirths in the study area is worrying as

postpartum mothers and their families suffer psychological trauma after experiencing stillbirths

throughout the nine-month journey of pregnancy (8,25). It could also indicate that the care for

women during pregnancy, labour, and birth might not have received appropriate and

standardized obstetric practices and protocols, leading to stillbirth.

In this study, maternal anaemia in the third trimester of pregnancy was independently

associated with stillbirths. Whilst a retrospective study in Columbia (26) was in agreement with

the present study, a similar study in Pakistan (27) was not in support. Anaemia is known to be a

significant predictor of malnutrition among pregnant mothers as iron deficiency type is

apparently the most prevalent cause of anaemia during gestation (28). As anaemia causes

inadequate oxygen and nutrient supply to the foetus during the late stage of pregnancy (29,30),

it could lead to a deficient supply of these nutrients to the foetus which subsequently causes

the death of the foetus either in-utero or within a short period after birth (31).

This study also found that women with sickle cell positive have increased odds of

stillbirth delivery. The observed findings align with previous research indicating severe

pregnancy and childbirth complications for both mothers and newborns affected by sickle cell

(32). These complications may stem from the physiological adaptations inherent in the genetic

makeup of pregnant women with sickle cell trait or disease (32,33). Such adaptations can

impact the cardiovascular, haematologic, respiratory, and renal structures in the materno-

foetal system, which can impair the overall health and ability to sustain a healthy pregnancy

(33). Consequently, these physiological challenges may contribute to the reduced likelihood of

delivering a healthy, live-born newborn.

Finally, mothers who did not sleep under insecticide-treated bed nets during pregnancy

were more likely to deliver stillbirths. In line with the present findings, mothers' sleep practices

including those who did not sleep under insecticide-treated bed nets during pregnancy, were

exposed to stillbirths in a cross-sectional study in Ghana (13) but were contrasting to a meta-

analysis by Flenady and his colleagues (34). Despite the constant supply of free insecticide-

treated bed nets to pregnant mothers for malaria prevention during their first antenatal care

(ANC) visits in Ghana (18), it is reported that most mothers do not sleep under the nets due to

unfavourable weather conditions (hot weather) in Northern Ghana, while others also use the

nets for subsistence farming purposes in their homes (35). The non-use of the bed nets could

expose them to malaria infections and subsequent anaemia in pregnancy, thereby increasing

their risk of stillbirth delivery (36).

Interestingly, a large proportion of the study participants who had positive sickle cell

status and did not use insecticide-treated bed nets also made less than eight ANC visits and had

anaemia during each trimester of pregnancy. This implies that fewer ANC visits and low

haemoglobin levels in each trimester during pregnancy could precipitate unfavourable intra-

uterine and gestational conditions (29,30) leading to an increased maternal risk for stillbirth

delivery (36,37) in the municipality. Therefore, greater attention should be paid to anaemia in

pregnancy and antenatal visits to strengthen maternal and child health outcomes in the

country.

Limitations of the study

The prevalence of stillbirths in the municipality may be underestimated or overestimated since

the study participants were sampled from a six-month retrospective data. We must admit that

the use of retrospective data (like maternal weight, haemoglobin levels, and gestational age)

from records review (MCHRBs and delivery registers) could be a limitation of the study as a

result of possible measurement and documentation errors. This study had limitations in the

generalizability of the factors of stillbirths due to the employment of a cross-section of data in

the year 2020. Notwithstanding, the study indicated very important and evidence-based

findings for immediate interventions in the municipality and the broader Northern Region of

Ghana.

Conclusion

The prevalence of stillbirths was relatively high in the municipality. These stillbirths were

independently influenced by medical and lifestyle factors. Timely identification of these

determinants is vital to prompt intervention strategies through strengthening obstetric triage

and indirectly mitigating the higher risk of maternal psycho-emotional problems in the

municipality. Management of Ghana Health Service (via the municipal health directorate),

program planners, and other stakeholders who are engaged in improving perinatal and

maternal health should also be aware of these factors or predictors in order to develop and

launch effective programs for reducing stillbirths in the municipality and the entire Northern

Region of Ghana.

Regarding the global incidence of stillbirths, the World Health Assembly recommends a

national target of stillbirth prevalence of 10 or less per 1000 deliveries by 2035 in developing

countries like Ghana (3). To achieve this goal, education on nutrition and lifestyle, as well as

early detection and swift management of gestational anaemia are very necessary to prevent

stillbirths and consequently reduce perinatal mortalities in underserved settings like Savelugu

municipality of Ghana and parts of sub-Saharan Africa.

List of abbreviations

ANC: Antenatal care

BMI: Body mass index

MCHRBs: Maternal and child health record books

STATA: Statistics and data

TD: Tetanus-diphtheria

VIF: Variance inflation factor

WHO: World Health Organization

Declarations

Ethics approval and consent to participate

This study was conducted according to the guidelines laid down in the Declaration of Helsinki,

and all procedures involving research study participants were approved by the Navrongo Health

Research Centre Institutional Review Board with approval number NHRCIRB373. Permission

was also granted by the Ghana Health Service at the municipal and health facilities level.

Written informed consent/assent was obtained from all study participants and/or legal

representatives and also from parents/guardians of participants under 18 years of age.

Consent For Publication

Not applicable

Availability of data and materials

The datasets collected, generated, and/or analyzed during the present study are available from

the corresponding author on reasonable request.

Competing interest

The authors declared that they have no competing interest

Funding statement

There was no funding support

Authors' contributions

SAG, AA, MG, CSD and PAA designed and conceptualized the study. SAG, AA, MG, WP, CA and

CSD carried out the data collection while SAG, WP, CA and PAA analyzed and interpreted the

data. All authors contributed to drafting the initial manuscript, reviewing and approving the

final manuscript.

Authors' acronyms

Silas Adjei-Gyamfi (SAG); Abigail Asirifi (AA); Mamudu Gafaru (MG); Wisdom Peprah (WP);

Clotilda Asobuno (CA); Constance Siakwan Dapaah (CSD); Paul Armah Aryee (PAA)

Acknowledgements

The authors are thankful to all research assistants, study participants, and volunteers who

contributed in diverse ways to the successful implementation of the study. We are also grateful

to Mr Elvis Brown Ayaala, Mr Abdulai Sulemana, Mr Allor Mohammed, and Mr Julien Beweleyir

for their supportive supervision and data collection exercise during this study.

References

1.   Lawn JE, Blencowe H, Waiswa P, Amouzou A, Mathers C, Hogan D, et al. Stillbirths: Rates,

risk factors, and acceleration towards 2030. Vol. 387, The Lancet. Lancet Publishing

Group; 2016. p. 587–603.

2.   Bawah AT, Ngala RA, Alidu H, Seini MM, Wumbee JDK, Yeboah FA. Gestational diabetes

mellitus and obstetric outcomes in a Ghanaian community. Pan Afr Med J. 2019;32.

3.   WHO. Every newborn: An action plan to end Preventable deaths: Executive summary

Geneva: World Health Organization. WHO, UNICEF. 2014;12 p.

4.   Harding EBM, Harrington LT, Lockwood CM, Duncan Brown JR, Hall AC, Brown MA, et al.

Perinatal and neonatal mortality. Br Med J. 2006;281(6254):1567.

5.   Blencowe H, Cousens S, Jassir FB, Say L, Chou D, Mathers C, et al. National, regional, and

worldwide estimates of stillbirth rates in 2015, with trends from 2000: A systematic

analysis. Lancet Glob Heal. 2016 Feb 1;4(2):e98–108.

6.   Zhu J, Liang J, Mu Y, Li X, Guo S, Scherpbier R, et al. Sociodemographic and obstetric

characteristics of stillbirths in China: A census of nearly 4 million health facility births

between 2012 and 2014. Lancet Glob Heal. 2016 Feb 1;4(2):e109–18.

7.   Lavin T, Pattinson RC, Nedkoff L, Gebhardt S, Preen DB. Stillbirth risk across pregnancy by

size for gestational age in Western Cape Province, South Africa: Application of the

fetuses-at-risk approach using perinatal audit data. South African Med J.

2019;109(12):927–33.

8.   Okonofua FE, Ntoimo LFC, Ogu R, Galadanci H, Mohammed G, Adetoye D, et al.

Prevalence and determinants of stillbirth in Nigerian referral hospitals: A multicentre

study. BMC Pregnancy Childbirth. 2019 Dec 30;19(1).

9.   Adu-Bonsaffoh K, Gyamfi-Bannerman C, Oppong SA, Seffah JD. Determinants and

outcomes of preterm births at a tertiary hospital in Ghana. Vol. 79, Placenta. W.B.

Saunders Ltd; 2019. p. 62–7.

10.   Dassah ET, Odoi AT, Opoku BK. Stillbirths and very low Apgar scores among vaginal births

in a tertiary hospital in Ghana: A retrospective cross-sectional analysis. BMC Pregnancy

Childbirth. 2014 Aug 28;14(1).

11.   Lawn JE, Yakoob M, Haws RA, Soomro T, Darmstadt GL, Bhutta ZA. 3.2 million stillbirths:

Epidemiology and overview of the evidence review. Vol. 9, BMC Pregnancy and

Childbirth. BioMed Central; 2009. p. 1–17.

12.   Njoku C., Emechebe C., Eyong E., Ukaga J., Anachuna K. Prevalence and risk factors for

stillbirths in a tertiary hospital in Niger Delta area of Nigeria: a ten year review. Int J Med

Biomed Res. 2016 Oct 3;5(3):106–13.

13.   Owusu JT, Anderson FJ, Coleman J, Oppong S, Seffah JD, Aikins A, et al. Association of

maternal sleep practices with pre-eclampsia, low birth weight, and stillbirth among

Ghanaian   women.   Int   J   Gynecol   Obstet.   2013;121(3):261–5.

http://doi.wiley.com/10.1016/j.ijgo.2013.01.013

14.   Ameyaw EK, Kofinti RE, Appiah F. National health insurance subscription and maternal

healthcare utilisation across mothers' wealth status in Ghana. Health Econ Rev.

2017;7(1):16. https://doi.org/10.1186/s13561-017-0152-8

15.   Adjei-Gyamfi S, Musah B, Asirifi A, Hammond J, Aryee PA, Miho S, et al. Maternal risk

factors for low birthweight and macrosomia: A cross-sectional study in Northern Region,

Ghana. J Heal Popul Nutr. 2023;42(1):1–16. https://doi.org/10.1186/s41043-023-00431-0

16.   Adjei-Gyamfi S, Asirifi A, Peprah W, Abbey DA, Hamenoo KW, Zakaria MS, et al. Anaemia

at 36 weeks of pregnancy: Prevalence and determinants among antenatal women

attending peri-urban facilities in a developing country, Ghana. PLOS Glob Public Heal.

2024;4(9):e0003631. https://doi.org/10.1371/journal.pgph.0003631

17.   Adjei-Gyamfi S, Zakaria MS, Asirifi A, Issahaku S, Ibrahim MA, Aryee PA. Maternal

anaemia and polycythaemia during pregnancy and risk of inappropriate birth weight for

gestational age babies: A retrospective cohort study in the northern belt of Ghana. BMJ

Open. 2024;14:e082298. https://doi.org/10.1136/bmjopen-2023-082298

18.   Adjei-Gyamfi S, Asirifi A, Aiga H. Prevalence and associated risk factors of preterm and

post-term births in Northern Ghana: A retrospective study in Savelugu Municipality. J

Pediatr   Perinatol   Child   Heal.   2023;07(04):235–48.

https://doi.org/10.26502/jppch.74050169

19.   Cochran WG. Samling Techniques. John Wiley & Sons, Inc. 1977. 448 p.

20.   Asobuno C, Adjei-Gyamfi S, Aabebe FG, Hammond J, Taikeophithoun C, Amuna NN, et al.

523   Risk factors for anaemia among pregnant women: A cross-sectional study in Upper East

524   Region,   Ghana.   PLoS   One.   2024;19(11):e0301654.

525   https://doi.org/10.1371/journal.pone.0301654

526   21.   Ghana Health Service. DHIMS-2: District Information Management Systems version 2.

527   GHS. 2020. https://dhims.chimgh.org/dhims

528   22.   Ghana Statistical Service (GSS), Ghana Health Service (GHS) & II. Ghana Demographic and

529   Health Survey 2014. Rockville, Maryland, USA; 2015.

530   23.   Rasmussen KM, Catalano PM, Yaktine AL. New guidelines for weight gain during

531   pregnancy: what obstetrician/gynecologists should know. Curr Opin Obs Gynecol.

532   2009;21(6):521–6.

533   24.   Okeudo C, Ezem B, Ojiyi E. Stillbirth rate in a teaching hospital in South-eastern Nigeria: A

534   silent tragedy. Ann Med Health Sci Res. 2012;2(2):176.

535   25.   Vogel JP, Souza JP, Mori R, Morisaki N, Lumbiganon P, Laopaiboon M, et al. Maternal

536   complications and perinatal mortality: findings of the World Health Organization

537   Multicountry Survey on Maternal and Newborn Health. BJOG. 2014;121 Suppl 1(Suppl

538   1):76–88.

539   26.   Smith C, Teng F, Branch E, Chu S, Joseph KS. Maternal and Perinatal Morbidity and

540   Mortality Associated with Anemia in Pregnancy. Obstet Gynecol. 2019 Dec

1;134(6):1234–44.

27.   Mahmood T, Rehman AU, Tserenpil G, Siddiqui F, Ahmed M, Siraj F, et al. The Association

between Iron-deficiency Anemia and Adverse Pregnancy Outcomes: A Retrospective

Report from Pakistan. Cureus. 2019 Oct 7;11(10).

28.   Garzon S, Cacciato PM, Certelli C, Salvaggio C, Magliarditi M, Rizzo G. Iron Deficiency

Anemia in Pregnancy: Novel Approaches for an Old Problem. Oman Med J. 2020

Sep;35(5):e166.

29.   Barker DJP. Intrauterine programming of adult disease. Vol. 1, Molecular Medicine

Today. Elsevier Current Trends; 1995. p. 418–23.

30.   Wu G, Imhoff-Kunsch B, Girard AW. Biological mechanisms for nutritional regulation of

maternal health and fetal development. Vol. 26, Paediatric and Perinatal Epidemiology.

John Wiley & Sons, Ltd; 2012. p. 4–26.

31.   Parks S, Hoffman MK, Goudar SS, Patel A, Saleem S, Ali SA, et al. Maternal anaemia and

maternal, fetal, and neonatal outcomes in a prospective cohort study in India and

Pakistan. BJOG An Int J Obstet Gynaecol. 2019 May 1;126(6):737–43.

32.   Smith-Whitley K. Complications in pregnant women with sickle cell disease. Hematology.

2019;2019(1):359–66. https://doi.org/10.1182/hematology.2019000039

33.   Jain D, Atmapoojya P, Colah R, Lodha P. Sickle cell disease and pregnancy. Mediterr J

Hematol Infect Dis. 2019;11(1):e2019040. http://dx.doi.org/10.4084/MJHID.2019.040

34.   Flenady V, Koopmans L, Middleton P, Frøen JF, Smith GC, Gibbons K, et al. Major risk

factors for stillbirth in high-income countries: a systematic review and meta-analysis.

Lancet. 2011 Apr 16;377(9774):1331–40.

35.   Savelugu Municipal Health Directorate (Ghana Health Service). Annual Performance

Report 2020: Savelugu Municipal Health Directorate. Savelugu, Northern Region; 2021.

36.   Moore KA, Simpson JA, Scoullar MJL, McGready R, Fowkes FJI. Quantification of the

association between malaria in pregnancy and stillbirth: a systematic review and meta-

analysis. Lancet Glob Heal. 2017 Nov 1;5(11):e1101–12.

37.   Idung A, Jombo H, Udonwa NE, Iyanam VE, Idung AU, Jombo HE. Anaemia in Pregnancy

at Booking: Prevalence and Risk Factors among Antenatal Attendees in a Southern

Nigeria General Hospital Article in Asian Journal of Medicine and Health · May. Asian J

Med Heal. 2019;15(1):48805.