1 **Genome-wide DNA methylation, imprinting, and gene expression in human placentas**
2 **derived from Assisted Reproductive Technology**
3

4 Auvinen P*[1], Vehviläinen J*[1], Rämö K[1], Laukkanen I[1], Marjonen-Lindblad H[1], Wallén E[1],
5 Söderström-Anttila V[2], Kahila H[3], Hydén-Granskog C[3], Tuuri T[3], Tiitinen A[3], Kaminen-Ahola
6 N[1]
7
AFFILIATIONS
1. Environmental Epigenetics Laboratory, Department of Medical and Clinical Genetics,
Medicum, University of Helsinki, Helsinki, Finland
2. The Family Federation of Finland, Fertility Clinic and University of Helsinki, Helsinki,
Finland
3. Department of Obstetrics and Gynecology, Helsinki University Hospital and University of
Helsinki, Helsinki, Finland
15
16 Corresponding author:
17 Nina Kaminen-Ahola, nina.kaminen@helsinki.fi
18
19 *Equal contribution

20
21 **ABSTRACT**
22

Assisted reproductive technology (ART) has been associated with increased risk for growth
disturbance and imprinting disorders, but the molecular mechanisms and whether they are a
result of the ART procedures or the underlying subfertility are unknown. Here we performed
genome-wide DNA methylation analysis by EPIC Illumina microarrays and gene expression
analysis by mRNA sequencing for a total of 80 ART and 77 control placentas, including
separate procedure- and sex-specific analyses. ART-associated changes enriched in the
pathways of hormonal regulation, insulin resistance, neuronal development, and
vascularization. Observed changes in the number of stromal cells as well
as *TRIM28* and *NOTCH3* expressions in ART placentas indicated impaired angiogenesis and
growth. The enrichment of DNA methylation changes in the imprinted regions and alterations
in *TRIM28, ZFP57*, and *NLRP5* suggested defective stabilization of the imprinting.

Furthermore, downregulated expression of imprinted endocrine signaling molecule *DLK1*,

associated with both ART and subfertility, provides a potential mechanism for the metabolic

and phenotypic features associated with ART.

INTRODUCTION

Approximately one in six couples experience infertility worldwide and the usage of Assisted

Reproductive Technology (ART) has increased over the last decades. To date over ten million

children have been conceived through these technologies[1]. The main procedure of ART is *in*

vitro fertilization (IVF), which may also include intracytoplasmic sperm injection (ICSI), in

which a single sperm cell is injected into the oocyte cytoplasm. In addition to the freshly

transferred embryos (FRESH), also embryo cryopreservation and frozen embryo transfer

(FET) can be a part of IVF or ICSI procedures. Less invasive *in vivo* fertility treatments, such

as intrauterine insemination (IUI) and hormone treatments to induce ovulation, are also widely

used.

Although children conceived by using the treatments are generally healthy, ART has been

associated with increased risks for adverse obstetric and perinatal outcomes such as preterm

birth (PTB), low birth weight (LBW), birth defects, and placental anomalies[2,3].

Additionally, long-term effects such as rapid postnatal growth, female early onset puberty as

well as cardiovascular and metabolic disorders have been associated with the ART phenotype[4-

6]. An increased risk for certain imprinting disorders[7,8] and a tendency towards higher risk of

neurodevelopmental disorders has furthermore been reported[9]. There are method-associated

risks, and it has been reported that FRESH derived children have a higher risk for PTB, LBW,

and for being small for gestational age (SGA) compared to natural conception. A higher risk

of pre-eclampsia and a higher risk for being large for gestational age (LGA) has been noted in

pregnancies after FET compared to FRESH[10]. However, it is unclear whether these

associations are the result of ART technology *per se* or the underlying subfertility.

The mechanisms by which ART or infertility increases the risk for adverse outcomes are

unknown. ART has been suggested to affect fetal development through epigenetic

modifications as the procedures take place during extensive epigenetic reprogramming in the

gametogenesis and early embryogenesis. Indeed, ART-associated DNA methylation (DNAm)

profiles have been reported in several epigenome-wide association studies (EWASs) of human

blood[11-18] and placenta[19-22]. Increasing evidence suggest that the placenta is particularly

susceptible to epigenetic changes caused by ART and/or infertility[23-25] and alterations have

been found especially in imprinted genes[19-22] and also in epigenetically silenced repetitive

elements (REs)[26-28]. However, the results of EWASs are not consistent, ART procedure-

specific EWASs as well as genome-wide gene expression studies for placentas are scarce, and

studies controlled for placental cell type heterogeneity are lacking.

Here, we explored ART-associated genome-wide DNAm by microarrays (Illumina's Infinium

MethylationEPIC) and gene expression by 3'mRNA sequencing (mRNA-seq) of full-term

placental samples from ART and naturally conceived singleton newborns (Table 1). The effect

of cell type heterogeneity was excluded by adjusting the DNAm analyses by placental cell

types. Furthermore, ART-associated phenotypic features were examined. The procedure-

specific changes were investigated by analyzing the ART samples in subgroups: newborns

derived from IVF or ICSI methods as well as from FRESH or FET procedures. Moreover, to

separate the effects of decreased fertility from ART procedures on gene expression, we

analyzed placental samples from pregnancies of couples who had went through IUI as well as

from subfertile (SF) couples who were about to start or already started ART treatment but got

pregnant spontaneously.

Males and female fetuses have been shown to differ in terms of placental DNAm[29,30] as well

as in their ability to adapt to adverse intrauterine environments[31,32]. In the ART studies, it has

been shown that males have a higher risk to be LGA compared to females[33-35], but still only a

few sex-specific EWASs have been performed so far[12,18,22]. In the current study, we performed

sex-specific analyses by examining placentas of male and female newborns separately,

including sex chromosomes.

RESULTS

Participant characteristics and placental phenotypes

General characteristics of 80 ART and 77 naturally conceived control newborns as well as their

mothers were compared. All ART and subgroup-specific information of the phenotype analysis

can be found in the Supplementary Table 1. In general, the phenotypes did not differ

significantly between ART and control newborns. A total of five SGA (6.3%) (IVF male, two

IVF females and two ICSI females, all FRESH) and five LGA (6.3%) (IVF-FRESH, ICSI-

FRESH, and ICSI-FET males as well as ICSI-FRESH and ICSI-FET females) newborns were
in the ART group, and two SGA newborns (2.6%, male and female) in controls. The gestational
age (GA) of ART newborns was significantly shorter, particularly in females (ART: $n = 44$,
control: $n = 35$) ($P = 0.003$, $P = 0.004$, respectively), and this was driven by IVF and FRESH
newborns ($n = 50$, $n = 42$, respectively). In the ART group, one (1.3%) IVF newborn was
preterm (gestational week 36+5). According to the Finnish growth charts[36], the head
circumference (HC) SD (z-score) of ICSI female newborns was significantly smaller compared
to control females ($n = 18$, $n = 35$, respectively) ($P = 0.018$). Furthermore, IUI and SF newborns
as well as their mothers were compared ($n = 12$, $n = 11$, respectively). The GA of SF newborns
was significantly shorter compared to controls ($P = 0.005$) and one female LGA newborn was
observed in this group. The mothers of ART newborns were significantly older and they gained
less weight during pregnancy compared to the mothers of control newborns ($P < 0.0001$, $P =$
0.002, respectively). Also, the mothers of IUI and SF newborns were significantly older, and
the body mass index (BMI) of the IUI mothers was higher compared to controls ($P = 0.016$, P
$= 0.017$, $P = 0.006$, respectively).

When the placentas of ART and control newborns were compared, no significant difference in
weights was observed (Supplementary Table 1). However, in ART subgroups, FET placentas
were significantly heavier compared to FRESH ($n = 38$, $n = 50$, respectively). The DNAm
profiles of placental samples and cell type fragment analysis were utilized to determine
potential differences between ART and control placentas in five major placental cell types:
syncytiotrophoblasts, trophoblasts, stromal cells, Hofbauer cells, and endothelial cells
(Supplementary Fig. 1). When comparing the fragments between ART ($n = 80$) and control (n
$= 77$) samples, we observed significantly lower number of stromal cells in the ART placentas
($P = 0.002$) (Supplementary Fig. 2) as well as ICSI and FRESH subgroup placentas compared
to controls ($n = 30$, $n = 42$, $n = 77$, respectively) ($P = 0.02$ and $P = 0.03$, respectively). Sex-
specific analyses revealed significantly lower number of stromal cells in female ART, ICSI,
and FRESH placentas compared to control females ($n = 44$, $n = 18$, $n = 28$, $n = 35$, respectively)
($P = 0.003$, $P = 0.001$, $P = 0.01$, respectively) (Supplementary Fig. 2). Furthermore, to exclude
the variation caused by different ART procedures, we compared FRESH and FET placentas
derived only from the IVF to the controls. Significantly more endothelial cells were observed
in the IVF-FRESH placentas compared to the IVF-FET placentas ($n = 25$, $n = 25$, respectively)
($P = 0.01$).

ART-associated genome-wide DNAm in placenta

To investigate genome-wide average DNAm level (GWAM) in the ART and control placentas ($n = 80$, $n = 77$, respectively), we used the DNAm level of all 746,966 probes in the microarrays (general characteristics in Supplementary Table 1). Furthermore, we examined ART-associated DNAm in REs by comparing the predicted mean DNAm level of CpGs in Alu, endogenous retrovirus (EVR), and long interspersed nuclear element (LINE1) sequences in ART and control placentas. However, we did not observe significant ART-associated alterations in GWAM levels either overall or at any genomic locations relative to gene or CpG island, or in REs (Supplementary Fig. 3 and 4, respectively).

ART-associated genome-wide DNAm was explored by using linear regression model adjusted for cell type, sex, as well as maternal pre-pregnancy BMI and age as covariates. The analysis resulted in 6728 significantly differentially methylated CpG sites (3524 hypo- and 3204 hypermethylated) with FDR < 0.05 (Fig. 1a, Supplementary Table 2). To separate the most prominent changes and to minimize false positive hits, we focused on the CpG sites with DNAm difference of ≥ 5% between ART and control placentas, which are termed as differentially methylated positions (DMPs). There were 164 DMPs associating with 126 genes in ART placentas (FDR < 0.05, $\Delta\beta \leq -0.05$ and $\Delta\beta \geq 0.05$), of which 99 were hypo- and 65 hypermethylated. Further, we tested for differentially methylated regions (DMRs) defined as a region with maximal allowed genomic distance of 1000 bp containing three or more CpGs with $P < 0.05$ according to Fisher's combined probability test. A total of 787 DMRs associating with 687 genes in ART placentas were revealed (Supplementary Table 3).

To get a comprehensive picture on the biological processes (BPs) in which ART-associated DMRs cluster, we performed pathway analyses (Supplementary Table 4). Gene ontology (GO) enrichment analysis of ART-associated DMRs revealed BPs involved in gliogenesis and hormone secretion (*FDR* < 0.05), specifically insulin secretion ($P < 0.05$). When only DMRs located in the regulatory regions (200 and 1500 bp from transcription starting sites (TTSs)) were analyzed, also inflammatory response and cytokine production brought forth ($P < 0.05$). According to KEGG enrichment analysis, ART-associated DMRs linked to terms such as insulin resistance, Type I and II diabetes mellitus (T1D and T2D), growth hormone synthesis,

secretion, and action, as well as GnRH secretion, estrogen signaling pathway, and TNF
signaling pathway ($P < 0.05$) (Fig. 1b).

Furthermore, placental DMRs in the current study associated with *CHRNE, DUSP22,*
NECAB3, RASL11B, SNCB, and *FSCN2*, as well as both DMRs and DMPs with *PCDHGB4,*
APC2, and *MSX1,* which all had DNAm changes in the same direction as in the blood samples
of ART derived individuals in previous EWASs[14,16,17]. Moreover, *FDFT1, FGF5, MYO7A,*
TRIM72, and *C4orf51* have also been observed in a previous study, in which ART placentas
were compared to placentas from IUI and SF pregnancies[21].

ART-associated genome-wide gene expression in placenta
To study genome-wide ART-associated gene expression, we performed mRNA-seq for ART
and control placentas ($n = 59$, $n = 39$, respectively) (general characteristics in Supplementary
Table 5). When the mRNA-seq model was adjusted by sex, we observed 71 significantly
differentially expressed genes (DEGs) (FDR < 0.05) of which 53 were downregulated and 18
upregulated (Fig. 1c, Supplementary Table 6). One of the most prominently downregulated
gene was *delta like non-canonical Notch ligand 1 (DLK1)* (Fig. 1d). This paternally expressed
imprinted gene is essential for embryonic development and growth, and its malfunction has
been connected to metabolic abnormalities, such as obesity, T2D, and hyperlipidemia in
previous human studies[37]. Other interesting downregulated DEGs were *tripartite motif-*
containing protein 28 (*TRIM28*) and *notch receptor 3* (*NOTCH3*) (Fig. 1d). *TRIM28* is a
mediator of epigenetic modifications and its haplosufficiency in mice has been linked to
testicular degeneration, age-dependent infertility[38], and bi-stable epigenetic obesity[39].
Furthermore, *TRIM28* is essential in decidualization and implantation[40] and is involved in the
regulation of female obesity[41]. *NOTCH3,* which is expressed mainly in vascular smooth muscle
and pericytes, is needed for the development of fully functional arteries in mouse[42], and
recently it has been reported that *NOTCH3* signaling controls human trophoblast stem cell
expansion and proliferation[43]. Interestingly, also ART-associated DMPs and DMRs were
observed in its family member *NOTCH1*, which mediates uterine stromal differentiation and is
essential for decidualization and implantation in mouse[44].

When all the placentas were analyzed, moderate correlations were seen between *TRIM28* and
DLK1 as well as between *TRIM28* and *NOTCH3* expressions ($r = 0.357$***, $n = 118$, and $r =$
0.358***, $n = 118$, respectively) (Supplementary Table 7). However, when the control and

ART placentas were analyzed separately, correlation between *TRIM28* and *NOTCH3* was only

seen in the control ($r = 0.472$*, $n = 39$) but not in the ART placentas. Interestingly, although

there is no evidence of direct interaction between *DLK1* and *NOTCH3*, correlations between

the expressions of the genes in all, control, ART, and SF placentas were observed ($r =$

0.518***, $n = 118$; $r = 0.419$**, $n = 39$; $r = 0.427$**, $n = 59$; $r = 0.733$*, $n = 10$, respectively).

According to the GO:BP enrichment analysis, ART-associated gene expression is linked

predominantly to vasculogenesis including endothelial cell development (*PECAM1, S1PR3,*

COL18A, and *ROBO4),* renal vasculature development (*PECAM1, NOTCH3,* and *CD34*), and

foam cell differentiation (*AGTR1, ABCA1,* and *CETP*) (FDR-corrected *q*-value < 0.05)

(Fig. 1e, Supplementary Table 8). Interestingly, foam cells are a sign for vascular changes in

the placenta, the so-called acute atherosis, which has been observed frequently in non-

transformed spiral arteries in pregnancies associated with pre-eclampsia, SGA, fetal death,

spontaneous PTB, and preterm premature rupture of membranes[45].

The separate effects of ART and subfertility

To separate the effects of ART methods and subfertility, we compared gene expression of

placentas derived from IUI and SF couples to controls ($n = 10$, $n = 10$, $n = 39$, respectively)

(Supplementary Table 9). One DEG, upregulated *TSIX*, a non-coding RNA gene and an

antisense to *X Inactive Specific Transcript* (*XIST)*, was observed in IUI and three

downregulated DEGs (*ATG9B, SRRM2,* and *INSR*) in SF placentas. When both IUI and SF

placentas were compared to controls, five downregulated DEGs (*DLK1, VIM, IGFBP5, TBX2,*

and *TRPC6)* were detected. Notably, decreased counts of *DLK1* were also observed in SF

placentas (Fig. 1d), suggesting that downregulation of *DLK1* associates with subfertility. By

adding IUI and SF samples to the controls we were able to separate the potential effects of

hormonal treatments and subfertility from the effects of *in vitro* culture of ART procedures.

Interestingly, only 13 DEGs associated with *in vitro* culture (Fig. 1f, Supplementary Table 9).

ART-associated changes in IVF and ICSI placentas

When IVF and ICSI placentas were compared separately to controls, 140 IVF-associated DMPs

linking to 98 genes (Fig. 2a, Supplementary Table 10) and 439 DMRs linking to 417 genes

were detected (Supplementary Table 11) ($n = 50$, $n = 30$, $n = 77$, respectively) (general

characteristics in Supplementary Table 1). The ICSI placentas associated with 60 DMPs

linking to 47 genes (Fig. 2b, Supplementary Table 12) as well as 63 DMRs linking to 48 genes

(Supplementary Table 13). Only five DMPs and one gene (*TCN1*) as well as 17 DMR-
associated genes were common between the IVF and ICSI placentas and when the IVF
placentas were compared to ICSI, none DMPs or DMRs were detected (Fig. 2c, Supplementary
Table 14). Although the IVF-associated DMRs and the pathways in which they clustered were
similar to ART result (Supplementary Table 15), this separate comparison revealed ICSI-
associated DMRs which linked to genes *COLGALT1, DGKZ, PAX8, PAX8-AS1,* and *PDE4B*
(Fig. 2c, Supplementary Table 14).

mRNA-seq analysis for the IVF ($n = 38$) and control ($n = 39$) placentas showed 14 DEGs (10
down- and 4 upregulated), whereas a higher number, 46 genes (34 down- and 12 upregulated),
was observed in the ICSI placentas ($n = 21$) (Fig. 2d,f, Supplementary Tables 14 and 16). *DLK1*
was the only common DEG between IVF and ICSI placentas. Interestingly, the upregulated
genes *ZFY, RPS4Y1, PCDH11Y,* and *DDX3Y* in the ICSI placentas locate on chromosome Y
and the expressions of them all correlated significantly with each other in ART male placentas,
but not in the controls (Supplementary Table 7). Since male infertility was a common reason
(44.4%) for the infertility among the analyzed ICSI male samples (Supplementary Table 17)
the upregulated genes *DDX3Y*[46], *ZFY*[47], and *PCDH11Y*[48], which all have been associated with
male infertility in previous studies, are notable. Furthermore, the ICSI-associated DEGs
MAP1B, HBA1, EPHX1, HBB, and *HBG2* enriched in response-to-toxic-substance pathway are
also interesting (FDR-corrected q-value < 0.05), considering that environmental exposures
have been associated with male infertility in previous studies (Supplementary Table 18).
Epoxide hydrolases, such as *EPHX1*, are involved in detoxifying and excreting the
environmental chemicals which are associated with decreased semen quality and male
infertility[49].

ART-associated changes in FRESH and FET placentas
Nearly equal amount of DMPs associated with the FRESH and FET placentas when they were
compared separately to the controls ($n = 42, n = 38, n = 77$, respectively): 99 FRESH-associated
DMPs were annotated to 97 genes and 93 FET DMPs to 59 genes (Fig. 3a-c, Supplementary
Tables 19−21). Furthermore, 253 FRESH- and 100 FET-associated DMRs were observed
(FDR < 0.05) (Fig. 3c, Supplementary Tables 21−23). GO enrichment analysis of FRESH
DMRs revealed BPs involved in pathways such as cell adhesion via plasma membrane
adhesion molecules (FDR < 0.05) and regulation of cytoskeleton organization ($P < 0.05$),
whereas FET DMRs clustered to the regulation of astrocyte differentiation and ion

transmembrane transporter activity, as well as peptidyl-arginine ADP-ribosylation ($P < 0.05$)
(Supplementary Tables 24 and 25, respectively). Interestingly, FRESH DMRs in KEGG
enrichment analysis clustered in T2D and insulin resistance, which did not emerge in the
analysis for FET DMRs.

mRNA-seq analysis for the FRESH, FET, and control ($n = 30$, $n = 29$, $n = 39$, respectively)
placentas showed 41 FRESH-associated DEGs (35 down- and six upregulated) and only six
FET DEGs (five down- and one upregulated) (Fig 3d-f, Supplementary Tables 21 and 26).
Downregulated *DLK1* came up from both FRESH and FET placentas, but *NOTCH3* and
TRIM28 were significantly downregulated only in FRESH placentas (Fig. 3d-f). The only
upregulated FET-associated DEG was *dermatan sulfate epimerase* (*DSE*), which
overexpression has been observed in several cancers and which associates with active
angiogenesis, invasion, and proliferation[50].

Finally, we compared FRESH and FET placentas derived only from the IVF procedure to the
controls ($n = 25$, $n = 25$, $n = 77$, respectively). A total of 34 DMPs and 25 DMRs were
associating with IVF-FRESH placentas, whereas 29 DMPs and 30 DMRs associated with IVF-
FET (Fig. 3g,h, Supplementary Tables 27 and 28, respectively). In both FRESH and IVF-
FRESH placentas, the most interesting DMPs located in the regulatory regions of *ZFP57 zinc*
finger protein (*ZFP57*) (cg13773306) and *NLR family pyrin domain containing 5* (*NLRP5*)
(cg14217229), which have been associated with multilocus imprinting disturbance[51] (Fig. 3i).
Interestingly, *ZFP57,* which also had a DMR in ART, IVF, FRESH, and IVF-FRESH
placentas, is a cofactor of *TRIM28*, and they both are required for the germline imprinting[52,53].
The DNAm level of *ZFP57* DMP (cg13773306) correlated with *TRIM28* expression in ART,
ICSI and FET placentas, ($r = 0.33*$, $n = 59$; $r = 0.6**$, $n = 21$; $r = 0.41*$, $n = 38$, respectively),
with *DLK1* expression in FET placentas ($r = 0.4*$, n = 38), and with *NOTCH3* expression in
ICSI placentas ($r = 0.53*$, $n = 30$), supporting associations between DNAm and expression of
these genes (Supplementary Table 29). Furthermore, the DNAm level of *NRLP5* DMP
(cg14217229) correlated with *NOTCH3* expression in ART, ICSI, and FRESH placentas ($r =$
$0.34**$, $n = 59$; $r = 0.54*$, $n = 21$; $r = 0.44*$, $n = 30$, respectively). There were only one DEG
in IVF-FRESH placentas, downregulated *CBX6* (log fold change = -0.97, FDR = 0.01), and no
DEGs in the IVF-FET placentas were observed ($n = 19$, $n = 19$, $n = 39$, respectively).

Sex-specific differences in ART-derived placentas

To identify sex-specific effects of ART, we performed DNAm analyses including sex
chromosomes for the placentas of male and female newborns separately. The GWAM overall
level was higher in male placentas compared to females, which is consistent with a previous
study[29] (Supplementary Fig. 3). A higher GWAM overall level was observed in ICSI male
placentas compared to control males, and significantly higher compared to IVF males ($n = 12$,
$n = 42$, $n = 24$, respectively) ($P = 0.04$). Furthermore, ICSI male placentas had significantly
higher GWAM level at genomic locations 5'UTR and S_Shore compared to control males as
well as at 5'UTR, S_Shelf and S_Shore compared to IVF males ($P = 0.03$, $P = 0.03$, $P < 0.05$,
$P = 0.01$, $P = 0.03$, respectively). By contrast to the males, differences in GWAM overall level
were not observed between control, IVF, and ICSI female placentas ($n = 35$, $n = 26$, $n = 18$,
respectively). However, the ICSI female placentas had significantly lower GWAM at S_Shelf
compared to IVF females ($P = 0.007$).
315
316   Moreover, a trend of higher GWAM in both male and female FET placentas compared to
317   FRESH and controls was observed (Supplementary Fig. 3). In FET male placentas the GWAM
318   level was significantly higher at N_Shelf compared to FRESH males, and in FET female
319   placentas significantly lower at 1stExon compared to FRESH females ($n = 22$, $n = 14$, $n = 16$,
320     $n = 28$, respectively) ($P = 0.03$ and $P = 0.04$, respectively).
321
322   Sex-specific examination of Alu, LINE1, and EVR REs revealed significantly lower Alu
323   DNAm in ICSI male placentas ($n = 12$) compared to IVF males ($n = 24$) (Supplementary Fig.
324   4). Furthermore, ICSI female placentas ($n = 18$) had significantly lower LINE1 DNAm level
325   compared to control females ($n = 35$) ($P = 0.03$ and $P = 0.04$, respectively).
326
In ART male placentas, there were only one hypermethylated DMP compared to control males
(ART: $n = 36$, control: $n = 42$), whereas 20 DMPs and eight DMRs were observed in ART
females (ART: $n = 44$, control: $n = 35$) (Supplementary Table 30). Interestingly, mRNA-seq
for male placentas (ART: $n = 29$, control: $n = 19$) revealed 21 DEGs (18 down, 3 up) (FDR <
0.05) clustering in pathways of endothelial and epithelial cell development in GO:BP
enrichment analysis, and none DEGs in the ART females (ART: $n = 30$, control: $n = 20$)
(Supplementary Table 31). In IVF male placentas five DEGs (downregulated *PECAM1,*
SPARCL1, DLK1, and *S1PR3* and upregulated *LYPLA1*) were observed compared to control
males, and in IVF female placentas 11 DMPs and four DMRs were observed compared to
control females ($n = 20$, $n = 19$, $n = 26$, $n = 35$ respectively). Despite of the small sample size

in sex-specific analysis of ICSI samples, 16 DEGs were observed in ICSI male placentas (ICSI:
$n = 9$, control: $n = 19$), including five upregulated genes, *DDX3Y, EIF1AY, ZFY, RPS4Y1*, and
PCDH11Y, locating all on chromosome Y. In female placentas (ICSI: $n = 12$, control $n = 20$),
six DEGs were observed including downregulated *CASP2* and upregulated *TFPI2, CSH1,*
CGA, PSG2, and *PSG3*.
342
When FRESH and FET placentas were analyzed sex-specifically, only one downregulated
DEG (*SPARCL),* was observed in FRESH male placentas (FRESH: $n = 11$, control: $n = 19$)
and 21 DMPs and 2 DMRs in the FRESH female placentas (FRESH: $n = 28$, control: $n = 35$)
(Supplementary Table 30 and 31). In FET male placentas, one DMR (*HTT-AS* 3 CpG) (FET:
$n = 22$, control: $n = 42$), and 11 DEGs clustering in GO:BP pathway term protein refolding
(*HSPA13, HSPA5*), were observed (FET: $n = 18$, control: $n = 19$).
349
Associations between placental gene expression and phenotypes
Potential correlations between *TRIM28*, *NOTCH3*, and *DLK1*, and phenotypes of newborns
and their mothers were calculated (Supplementary Table 29). When all samples were
examined, there was a weak correlation between placental *NOTCH3* expression and birth
weight (BW) (SD) ($r = 0.272$**, $n = 118$), and it was driven by male samples in which the
correlations between *NOTCH3* and BW as well as *NOTCH3* and birth length (BL) (SD) were
observed ($r = 0.414$**, $n = 58$, and $r = 0.369$**, $n = 58$, respectively). Furthermore, *NOTCH3*
expression correlated with the BW and BL (SD) of controls ($r = 0.41$*, $n = 39$, and $r = 0.386$*,
$n = 39$, respectively), with BW and BL (SD) and placental weight of ICSI ($r = 0.441$*, $r =$
0.503*, and $r = 0.463$*, $n = 21$, respectively), as well as with BW and BL (SD) of FET
newborns ($r = 0.419$**, $n = 29$; $r = 0.441$*, $n = 29$, respectively). The correlation between
NOTCH3 and BL among the ICSI samples was driven by male newborns.
362
In all male samples there was a negative correlation between *TRIM28* expression and maternal
age ($r = -0.339$, $P = 0.009$, $n = 58$) and in all the female samples *TRIM28* correlated weakly
with GA ($r = 0.257$*, $n = 60$). Furthermore, *TRIM28* expression correlated negatively with
maternal weight gain during pregnancy in the control and ART placentas ($r = -0.418$*, $n = 37$;
$r = -0.295$*, $n = 54$, respectively). In the ART placentas it was driven by ICSI, specifically by
ICSI female placentas ($r = -0.596$**, $n = 19$; $r = -0.655$*, $n = 11$, respectively). Also, placental

TRIM28 expression correlated negatively with maternal weight gain and positively with

maternal BMI in the FRESH placentas ($r = -0.388^*$, $n = 26$; $r = 0.363^*$, $n = 30$, respectively).

Placental *DLK1* expression correlated with ART female BL (SD) ($r = 0.394^*$, $n = 30$) driven

by ICSI females ($r = 0.676^*$, $n = 12$), and negatively with maternal weight gain during the

pregnancy in IVF females ($r = -0.618^*$, $n = 15$).

ART- and subfertility-associated placental imprinting

Finally, we focused on genomic imprinting since imprinting disorders have been associated

with ART in several previous studies[54]. Imprinted genes are a group of parent-of-origin

monoallelically expressed genes, which are maintained by correctly methylated imprinting

control regions (ICRs) established in the germline. A total of 19 ART- and/or ART subgroup-

associated DMRs on the microarrays are annotated to the imprinted genes[55] of which 13 were

hypomethylated (ART: $n = 80$, IVF: $n = 50$, ICSI: $n = 30$, FRESH: $n = 42$, FET: $n = 38$, control:

$n = 77$) (Supplementary Table 32). Six of these (*BLCAP, KCNQ1, KCNQ1OT1, NNAT, PEG3,*

and *ZIM2*) have been associated with ART also in previous EWASs[12,16,21,56] and were found to

be hypomethylated as in the current study. To focus specifically on ICRs, we compared 826

CpGs on ICRs compiled by Ochoa and collagues[57] between ART and control placentas and

observed significant differences in 115 sites (90 hypo- and 25 hypermethylated) ($P < 0.05$)

(Supplementary Table 32). After multiple testing correction, 16 CpG sites remained significant

(Bonferroni adjusted $P < 0.05$). Furthermore, by using genome-wide DNAm data we examined

if the ART-associated DNAm changes were particularly enriched in the ICRs. There were 15

ART-associated CpGs (13 hypo- and 2 hypermethylated) locating in the ICRs (FDR < 0.05)

and according to the hypergeometric distribution test, imprinted regions were significantly

enriched with DNAm changes ($P = 0.009$).

In the current study, we observed ART-associated hypomethylation in *ZFP57* as well as

prominent downregulation of paternally expressed *DLK1,* which locates in a paternally

imprinted *DLK1-DIO3* locus. Since *ZFP57* binds to and stabilizes paternally methylated ICR

in *DLK1-DIO3* locus in mouse[58] and *TRIM28* is a cofactor of *ZFP57*[52,53], we examined ICR

DNAm at this locus of ART (IVF: $n =$ four, ICSI: $n =$ four), SF ($n =$ four), and control ($n =$ six)

placentas by traditional bisulphite sequencing (Fig. 4a). The analyzed ICR sequence is next to

the repeated sequence motifs, which contribute to the DNAm of the ICR[58]. Maternal and

paternal alleles were distinguished by two single nucleotide polymorphisms (SNPs)
(rs1884539 and rs75998174) when possible. Interestingly, although we did not see highly
methylated paternal and demethylated maternal allele in this specific ICR sequence, we
observed significantly decreased total DNAm in the ICR of SF placentas compared to control
(CpG1: $P = 0.003$, CpG2: $P = 0.007$) and ART placentas (CpG1: $P = 0.002$, CpG7: $P = 0.047$)
(Fig. 4b, Supplementary Fig. 5). However, we did not observe significantly decreased DNAm
in the ART placentas compared to controls, which indicated that the ICR hypomethylation is
not the only mechanism which affects the complex *DLK1* regulation[59,60].

DISCUSSION

In this study, we performed the first genome-wide DNAm and gene expression analyses of
human placentas from ART pregnancies as far as we are aware. Those pregnancies are not a
homogeneous group, but differ from each other due to the ART procedures, causes of
underlying infertility, and sex of the child. By combining the omics data to the phenotypic
information and conducting method- and sex-specific analyses, it was possible to perceive the
associations between molecular alterations and phenotypic features. There were only a few
common ART procedure-associated alterations in the IVF and ICSI placentas, suggesting a
strong impact of the underlying sub/infertility on the DNAm and gene expression rather than
the effects of the *in vitro* culture. Furthermore, we confirmed previous findings about the
procedure-specific effect of FET and observed heavier FET placentas compared to the FRESH
placentas. Our observation about the lower amount of gene expression changes in the FET
placentas compared to FRESH was consistent with this. Although freezing might be a
challenging environmental insult for the embryo, it enables to avoid the hormonal disturbances
caused by ovarian stimulation in FRESH. Nearly all (82%) of the FET pregnancies in our study
have been started by natural cycle, which could explain the differences between FRESH and
FET placentas.

In some studies, the more invasive ICSI procedure has been associated with a higher risk of
birth defects[61], which is concerning due to the increased use of this method. In the current
study, more changes in gene expression were observed in the ICSI placentas compared to IVF.
Male factor infertility was a prominent reason for ICSI in this study, which could explain the
observed alterations. This can be seen in a small number of common ART-associated genes in

IVF and ICSI placentas as well as in many male infertility-associated genes on the Y
chromosome in the ICSI male placentas. Indeed, in addition to information about the placenta
itself, placenta could offer a "molecular window" to early development[62-64], even to the germ
cells. The origin of these early alterations and their potential to provide future biomarkers of
infertility should be studied.

Furthermore, in addition to several sex-specific ART-associated alterations, we observed
considerable differences between the ICSI males and females. The placental GWAM of ICSI
females were more hypomethylated compared to ICSI males, and ICSI female newborns were
smaller at birth – even significantly smaller HCs compared to controls were observed. Sex-
specific differences in ICSI have been observed also in previous studies. A significantly higher
number of trophectoderm cells in ICSI male embryos was detected compared to their female
counterparts, which was not observed when IVF male and female embryos were compared[65].
Also previously, ICSI female newborns had significantly smaller BWs (SD) as well as HCs
(SD) at the ages of 18 and 36 months (the information about the HC at birth was lacking)
compared to controls[66]. The number of sex-specific studies so far is surprisingly small and
more studies with larger sample sizes are needed to verify this discovery as well as to find out
whether this is caused by the ICSI method or by the underlying male infertility.

In addition to confirming previous ART-associated alterations observed in blood and placental
samples, we found new candidate genes *DLK1*, *TRIM28*, and *NOTCH3* for ART-associated
phenotypes. By performing cell type fragment analysis, we observed a significantly lower
number of stromal cells in the ART placentas compared to controls. Interestingly, it has been
suggested that differences in placental mesenchymal stromal cells could lead to impaired
vascular development and consequently to restricted growth[67,68]. This is supported by our gene
expression analysis, in which the most prominent ART-associated changes were linked to
vasculogenesis, and where the downregulation of *NOTCH3* and *TRIM28*, both involved in the
regulation of angiogenesis through the VEGFR-DLL4-NOTCH signaling circuit[69], was
observed. Furthermore, *NOTCH3* expression correlated with BW (SD) of control newborns,
with BW (SD), BL (SD), and placental weight of ICSI male newborns, as well as BW (SD)
and BL (SD) of FET newborns. Since *NOTCH3* and *TRIM28* are not significantly
downregulated in placentas of FET newborns − whose size is near to the control newborns −
the downregulation of *NOTCH3* could explain the higher risk of LWB and SGA associated
with FRESH. Interestingly, among the control and FET groups the maternal hormonal function

is not affected: in most cases the embryo was transferred in a natural cycle without the
hormonal disturbances caused by ovarian stimulation. The role of hormones brought forth also
in the enrichment analysis, where ART-associated DMRs linked to terms such as growth
hormone synthesis, secretion, and action, as well as GnRH secretion, and estrogen signaling
pathway. Recently, it has been shown that *TRIM28* modulates estrogen receptor α and
progesterone receptor signaling, which regulate endometrial cell proliferation, decidualization,
implantation, and fetal development[40]. Considering all these observations, the functional links
between *NOTCH3, TRIM28*, and the ART-associated cardiovascular phenotype should be
studied. Also, the separate effects of hormonal ovarian stimulation in ART protocols and
subfertility-associated impaired hormonal function on early development need to be clarified
in the future.
480
Along with *TRIM28* and *NOTCH3*, one of our most interesting discoveries in the ART
placentas concerned the prominent downregulation of imprinted *DLK1* as well as the pathways
of insulin signaling, insulin resistance, T1D and T2D, which came forth in the DNAm analyses.
The developmentally essential *DLK1* is a part of evolutionary conserved Delta-Notch pathway,
and it inhibits adipogenesis by controlling the cell fate of adipocyte progenitors[70,71]. According
to animal studies, mice without paternally expressed *DLK1* exhibit growth retardation and
obesity while *DLK1* overexpression generate decreased fat mass, diet-induced obesity
resistance, and reduced insulin signalling[72,73]. In human, mutations in *DLK1* gene have been
reported as a cause of central precocious puberty associated with obesity and metabolic
syndrome with undetectable DLK1 serum levels[74]. Furthermore, in Temple syndrome, where
DLK1 expression is downregulated due to the maternal uniparental disomy of the imprinted
DLK1 locus, phenotypic features such as prenatal growth failure, short postnatal stature, female
early onset puberty, and truncal obesity have been observed[75]. Since increased risks for SGA,
rapid postnatal growth, and female early onset puberty[6] have all been associated with ART
children, *DLK1* is a plausible candidate gene for the phenotypic effects associated with ART.
496
In addition to the downregulation of *DLK1* expression, we observed significant changes in
ZFP57, TRIM28, and *NLRP5,* which all are needed for the stability of DNAm in the imprinted
regions. We observed enrichment of decreased DNAm in several ICRs – common binding sites
of *ZFP57* and *TRIM28* – suggesting that the machinery, which normally stabilizes the DNAm
of ICRs globally could be impaired. This could explain the high variability in ART-associated
imprinted genes observed in previous studies. Previously, mutations or dysregulation of *ZFP57*

have been associated with imprinting disorders such as transient neonatal diabetes mellitus[76],

which incidence has been shown to be higher among ART newborns[77]. Furthermore, mutations

in *NLRP5* were responsible for early embryonic arrest in infertility patients[78]. More specific

examination of *DLK1-DIO3* locus showed decreased DNAm of the ICR in SF placentas. This

suggests that the instability of DNAm at this locus associates with subfertility, which is

consistent with the decreased *DLK1* expression in the ART, SF, and IUI placentas observed in

the current study, and with the previously reported associations between subfertility and

decreased placental *DLK1* expression[19]. However, *TRIM28* can regulate the expression of

imprinted *DLK1* in several ways[59,60] and therefore the interaction of these two genes should be

examined by functional studies. Also, the effect of ageing on *TRIM28* expression and its

associations with both *DLK1* and *NOTCH3* should be studied profoundly, due to the previous

association between *TRIM28* and age-dependent infertility in mouse[38] as well as our observed

negative correlation between placental *TRIM28* expression and maternal age.

These findings are also interesting in the perspective of fetal programming. According to the

programming hypothesis, adverse conditions such as prenatal malnutrition in the critical

developmental period may permanently program physiological processes. If there is plenty of

food available postnatally, this programming may prove inappropriate and cause metabolic

alterations leading to adult diseases such as obesity, T2D, metabolic syndrome, and

cardiovascular disease[79,80]. Previously, Cleaton and colleagues showed in mouse that both *Dlk1*

expression in maternal tissues and circulating DLK1 derived from the fetus are necessary for

appropriate maternal metabolic adaptations to pregnancy by allowing the switch to fatty acid

use when resources are limited during fasting[81]. Also, they reported that the circulating DLK1

level predicts embryonic mass in mouse and lower DLK1 levels were strongly associated with

high-resistance patterns of flow in the umbilical artery and slow abdominal circumference

growth velocity in a human cohort. Furthermore, DLK1 level enables to differentiate healthy

SGA from pathologically small infants. Our observation about the downregulated placental

DLK1 and *TRIM28* as well as their negative correlation with maternal weight gain during the

pregnancy suggest that they did not effectively inhibit maternal adipogenesis, resulting in

increased maternal weight and impaired maternal metabolic adaptations to the pregnancy. This

negative correlation was seen particularly in our IVF and ICSI female as well as FRESH

placentas, in the subgroups where the relatively smaller placentas and newborns were derived

from. The switch to fatty acid use in maternal tissues during fasting could be impaired,

potentially causing restricted fetal growth and a higher risk for adult disorders in the future.

However, in the ART-associated phenotype the increased risk for LBW, SGA, rapid postnatal
growth, and metabolic disorders would be a consequence of downregulation of *DLK1* caused
by subfertility associated disorder, not a consequence of poor maternal nutrition as in the
example of fetal programming. Notable, significant associations between subfertility and LBW
have been observed in previous studies[82].

In summary, the observed molecular changes in genome-wide DNAm and gene expression in
the ART placentas were relatively subtle, which is a finding in line with the fact that the
majority of the ART conceived children are as healthy as newborns from natural conceptions.
Decreased expression of *TRIM28*, *DLK1,* and *NOTCH3* as well as ART-associated DNAm in
ZFP57 and *NLRP5* bring forth potential mechanisms for the metabolic and phenotypic features
including growth disturbance and imprinting disorders, which have been associated with ART.
Considering that the observed changes associated also with subfertility, they offer a precious
insight to the molecular background of infertility.

METHODS

Epigenetics of ART (epiART) Cohort
Finnish, Caucasian couples applied to fertilization treatment in the Fertility clinic of the Family
Federation of Finland or the Reproductive Medicine Unit of Helsinki University Central
Hospital, Finland were recruited to this study and the samples were collected during the years
2015–2021. The epiART cohort includes 80 newborns fertilized using ART (IVF *n* = 50 and
ICSI *n* = 30) of which 42 using FRESH and 38 using FET embryos. Moreover, the cohort
includes newborns from pregnancies of couples who had went through IUI as well as from SF
couples who were about to start or already started ART treatment but got pregnant
spontaneously. More detailed information about the number of samples and newborns and
maternal characteristics is presented in Table 1 and Supplementary Table 1. Parental infertility
diagnoses of ART, IUI, and SF samples were categorized into male or female factor
infertilities, infertility caused by multiple causes, and unexplained infertility (Supplementary
Fig. 6 and Supplementary Table 17). Four cases in the ART group (5%) (IVF 4% and ICSI
3.3%) and one in the IUI group (8.3%) used medication for chronic somatic disease
hypothyroidism. Two ART cases (2.5%), both in IVF group (4%), had a psychiatric disorder
diagnosis and the other one used medication for it (anxiety disorder). One in the IUI group
(8.3%) had medication for a bipolar disorder. Pregnancy related disorders such as gestational

diabetes mellitus occurred in seven ART cases (8.8%) (IVF 12%, and ICSI 33.3%), two IUI

(16.7%) and two SF (18.2%), hepatogestosis in three ART cases (3.8%) (IVF 4%, ICSI 3.3%)

and pregnancy related hypertension in one IUI (8.3%) and one SF (9.1%). A total of 16 ART

(20 %) (IVF 22% and ICSI 17%), four IUI (33%), and three SF (27%) deliveries were cesarean

sections (CSs). The naturally conceived controls ($n = 77$) were full term neonates born from

uncomplicated pregnancies of healthy Finnish, Caucasian mothers collected during the years

2013–2015 in Helsinki University Hospital, Finland. Five (6.5%) of the deliveries were CSs.

Differences in newborn birth measures and maternal characteristics between study groups were

calculated, depending on the data distribution, by parametric two-tailed Student's *t*-test or

nonparametric Mann-Whitney U test in the case of two groups, and by parametric One-Way

ANOVA followed by Bonferroni post hoc or nonparametric Kruskal-Wallis test followed by

pairwise comparisons in the case of multiple groups. Differences in BW, BL, and HC of the

newborns were calculated using both anthropometric measures and the SDs of measures based

on Finnish growth charts[36], in which GA at birth, twinning, parity, and gender were considered

when calculating the SDs (z-scores) of birth measures. SGA and LGA are characterized by

weight and/or length at least 2 SD scores below or above the mean.

Sample collection and preparation

Placental samples of newborns were collected immediately after delivery. When this was not

possible, the placenta was stored in the fridge for a maximum of 12 h and only DNA was

extracted for further analyses. The placental biopsies (1 cm^3) were collected from the fetal side

of the placenta within a radius of 2–4 cm from the umbilical cord, rinsed in cold 1× PBS, fixed

in RNAlater® (Thermo Fisher Scientific), and stored at −80 °C. Placental genomic DNA was

extracted from one to four (3.5 on average) pieces of placental tissue samples using commercial

QIAamp Fast DNA Tissue or AllPrep DNA/RNA/miRNA Universal Kits (Qiagen) or standard

phenol-chloroform protocol. RNA was extracted from the same placental pieces as DNA (2.8

pieces on average) by AllPrep DNA/RNA/miRNA Universal Kit. RNA quality was assessed

using an Agilent 2100 Bioanalyzer (Agilent Technologies, Inc.), which was provided by the

Biomedicum Functional Genomics Unit (FuGU) at the Helsinki Institute of Life Science and

Biocenter Finland at the University of Helsinki.

DNAm microarrays

DNAm measurements and data processing. Genomic DNA (1000 ng) from available placental samples (sample information in Supplementary Table 1) was sodium bisulfite-converted using the Zymo EZ DNAm™ kit (Zymo Research), and genome-wide DNAm was assessed with Infinium Methylation EPIC BeadChip Kit (Illumina) following a standard protocol at the Institute for Molecular Medicine Finland. The raw DNAm dataset was pre-processed, quality controlled, and filtered using ChAMP R package[83] with default settings: the detection *P*-value cutoff was set at $P = 0.01$, and probe bead count > 3 in at least 95% of samples. All probes and samples passed these quality control thresholds and were included in the subsequent steps. Type-I and Type-II probes were normalized using the *adjustedDasen* method[84] in wateRmelon R package[85]. Potential effects caused by technical factors and biological covariates were studied from singular value decomposition plots. The data was corrected for the effects of array, slide, batch, DNA extraction method, and placental type (stored in a fridge before sampling or not) by the Empirical Bayes method using the R package ComBat[86]. After ComBat adjustment, probes located in sex chromosomes and probes binding to polymorphic and off-target sites were filtered. Moreover, probes in Finnish-specific SNPs, which overlap with any known SNPs with global minor allele frequency were removed. Population-specific masking was obtained from Zhou et al.[87].

Subsequently, a total of 746,966 probes were retained for further downstream analysis. Probes in sex chromosomes (16,945 probes for males and 16,651 probes for females) were included in sex-specific analyses. Annotation information was merged to corresponding probes from IlluminaHumanMethylationEPICanno.ilm10b4.hg19 R package[88], which is based on the file "MethylationEPIC_v-1-0_B4.csv" from Illumina[89]. Probe genomic locations relative to gene and CpG island were annotated based on University of California, Santa Cruz (UCSC) database. If the location information was missing, probe was marked as "unknown." In the case of multiple location entries, group "others" was used. Otherwise, the following abbreviations were used: TSS1500: 1500 bp upstream of transcription start site, TSS200: 200 bp upstream of TSS, UTR: untranslated region, N_shelf: north shelf, N_shore: north shore, S_shore: south shore, S_shelf: south shelf. Probes were annotated to genes based on, in addition to UCSC, also GENCODE Basic V12 and Comprehensive V12 databases.

DMP analysis. Genome-wide differential DNAm analysis by using M values was performed by R package Limma[90], and the linear model was adjusted to consider biological covariates newborn sex and maternal age and pre-pregnancy BMI. Planet R package[91] was used to count

placental cell-type fractions by CIBERSORT method with unfiltered data and used as an

adjusting factor in the model. β values were used for visualization and interpretation of the

results. DMPs were considered as significant when DNAm difference was greater than 5% ($\Delta\beta$

≤ -0.05 and $\Delta\beta \geq 0.05$) and FDR-corrected *P*-value smaller than 0.05. Benjamini-Hochberg

procedure was used to control for FDR. To calculate the differences in placental cell type

composition between the study groups, depending on the data distribution, parametric or

nonparametric two-tailed *t*-test in the case of two groups, and parametric One-Way ANOVA

followed by Bonferroni *post hoc* or nonparametric Kruskal-Wallis test followed by Wilcoxon

Rank Sum Exact test in the case of multiple groups were used.

DMR and Pathway analysis. DMRcate R package[92] was used to analyze DMRs. The method

uses minimum description length for detecting region boundaries in DMR identification.

DMRcate was adjusted to determine probes (≥ 3) in a region with maximal allowed genomic

distance of 1000 bp having FDR < 0.05. DMRs with Fisher's combined probability test *P* <

0.05 were considered significant. DMRs were annotated to imprinted genes based on

Geneimprint database[55]. Pathway analysis was performed for significant DMRs by *goregion*

function in missMethyl R package[93] and GO:BP and KEGG knowledgebases were used as a

source. When the GO terms were not significant after FDR correction, the terms with the

nominal *P*-value < 0.05 were reported.

GWAM analysis. β values of all probes in the array (normalized, ComBat adjusted, and filtered

data) were used to calculate placental GWAM[94] levels sample-wise. Differences between the

study groups were calculated by parametric or nonparametric two-tailed *t*-test depending on

the data distribution in the case of two groups. In the case of multiple groups, One-Way

ANOVA followed by Tukey's honestly significant difference (HSD) test was used.

RE DNAm analysis. Processed DNAm data (M values) was used to predict DNAm in Alu,

EVR, and LINE1 elements sample-wise using Random Forest-based algorithm implemented

by REMP R package[95]. Less reliable predicted results were trimmed according to quality score

threshold 1.7 and missing rate 0.2 (20%). Differences between the study groups were calculated

by parametric or nonparametric two-tailed *t*-test depending on the data distribution in the case

of two groups. In the case of multiple groups, One-Way ANOVA followed by Tukey's HSD

test was used.

ICR analysis. Enrichment of ART-associated CpGs at ICRs was measured with

hypergeometric distribution analysis. In total of 826 CpG sites located at ICRs[57] were identified

in our genome-wide DNAm data, of which 15 CpGs were significantly differentially

methylated (FDR < 0.05). Enrichment of these ICR-associated CpGs were compared to all

CpGs in the EPIC array with *phyper* function in R. All CpGs at the ICRs were compered

between ART and control placentas by two-tailed Student's *t*-test. Bonferroni adjustment was

used for multiple comparison.

Traditional bisulfite sequencing

A total of eight ART (four IVF, four ICSI, two/sex), four SF (two/sex), and six control placental

samples (three/sex) were subjected to bisulfite sequencing. Samples were chosen based on

mRNA-seq counts, control samples having the highest and ART and SF samples the lowest

counts. Paternal and maternal alleles were distinguished according to rs1884539(A/G) and

rs75998174(A/G) polymorphisms at the *DLK1-DIO3* ICR. Primers were designed to

incorporate the polymorphisms using Bisulfite Primer Seeker tool (Zymo Research)[96] (forward

5'GGTTTATAGTTGTTTATGGTTTGTTAAT   and   reverse

5'CTCCAACAAAAATTCCTTAAACTAAATT). The 451 bp amplicon (chr14:101,277,375-

101,277,826) covered 12 CpG sites at the ICR. Two separate bisulfite conversions were

performed for 500 ng of genomic DNA using EZ DNA Methylation™ kit (Zymo Research)

and pooled afterwards. To avoid possible PCR bias, three independent 20 μl PCR reactions

(HotStar PCR kit, Qiagen) were performed per sample using annealing temperature 56 °C.

PCR reactions were gel isolated, and the three reactions of each sample were pooled and

purified using NucleoSpin Gel and PCR Clean-up Kit (Macherey-Nagel). The purified PCR

fragments were ligated into pGEM®-T Easy Vector (Promega) and cloned by standard protocol.

The recombinant-DNA clones were purified using NucleoSpin® Plasmid EasyPure kit

(Macherey-Nagel) according to manufacturer's instructions. The sequences were analyzed by

BIQ Analyzer[97] excluding the clones with lower than 90% conversion rate from the dataset. A

total of eight to 30 clones of each individual were successfully sequenced (Supplementary Fig.

5). DNAm differences in CpG sites between study groups (control, ART, SF) were calculated

by One-Way ANOVA followed by Bonferroni post hoc.

mRNA-seq analysis

Differential expression analysis. Drop-seq pipeline[98] was used to construct the mRNA-seq

count table for available placental RNA samples (sample information in Supplementary Table

5) provided by FuGU. A total of 34,989 transcripts were identified for downstream analysis.

Principal component analysis implemented in DESeq2[99] was used to identify batch effects, and

ComBat-seq[100] was used to adjust separate mRNA-seq batches. Differential expression

analysis was performed by DESeq2 R package, with model adjusting for newborn sex. Genes

were considered as differentially expressed when FDR-corrected *P*-value was < 0.05.

Benjamini-Hochberg procedure was used to control for FDR.

Pathway analysis. enrichgo function in R package clusterProfiler version 4.0[101] was used to

perform gene-set enrichment analysis for significant DEGs. The GO knowledgebase was used

as the source for identifying significantly enriched BP terms (FDR-corrected *q*-value < 0.05).

Benjamini-Hochberg procedure was used to control for FDR.

Correlation analysis

The within sample gene expression correlations were calculated using ComBat-seq-adjusted

and Transcripts Per Million-normalized mRNA-seq counts and the correlations between gene

expression, *ZFP57* (cg13773306), *NLRP5* (cg14217229) DNAm, newborn phenotype, and

maternal characteristics using ComBat-seq-adjusted and DESeq2-normalized mRNA-seq

counts by Spearman rank correlation.

Statistical analysis

All statistical analyses were conducted in R versions 4.2.3 and 4.3.1[102], IBM SPSS Statistics

for Windows, version 29.0 (IBM Corp.), or GraphPad Prism 9 software (GraphPad Software,

Inc.). All data are expressed as the mean with ±SD for a normal distribution of variables.

Statistical analyses were performed as described in the relevant method sections and in the

figure legends.

ABBREVIATIONS

ART: Assisted reproductive technology

BL: Birth length

BMI: Body mass index

BP: Biological process

BW: Birth weight

CS: Cesarean section

DEG: Differentially expressed gene

DLK1: Delta like non-canonical Notch ligand 1

DMP: Differentially methylated position

DMR: Differentially methylated region

DNAm: DNA methylation

DSE: Dermatan sulfate epimerase

EVR: Endogenous retrovirus

EWAS: Eepigenome-wide association study

FET: Frozen embryo transfer

FRESH: Fresh embryo transfer

FuGU: Biomedicum Functional Genomics Unit

GA: Gestational age

GO: Gene ontology

GWAM: Genome-wide average DNAm level

HC: Head circumference

HSD: Honestly significant difference

ICR: Imprinting control region

ICSI: Intracytoplasmic sperm injection

IUI: Intrauterine insemination

IVF: In vitro fertilization

LBW: Low birth weight

LGA: Large for gestational age

LINE1: Long interspersed nuclear element

mRNA-seq: 3'mRNA sequencing

N_shelf: north shelf

N_shore: north shore

NLRP5: NLR family pyrin domain containing 5

NOTHC3: Notch receptor 3

PTB: Preterm birth

RE: Repetitive element

S_shore: south shore

S_shelf: south shelf

SF: Subfertile

SGA: Small for gestational age

SNP: Single nucleotide polymorphism

T1D: Type I diabetes mellitus

T2D: Type II diabetes mellitus

TRIM28: Tripartite motif-containing protein 28

TSS: Transcription starting site

TSS200: 200 bp upstream of TSS

TSS1500: 1500 bp upstream of transcription start site

UCSC: University of California, Santa Cruz

UTR: Untranslated region

XIST: X Inactive Specific Transcript

ZFP57: ZFP57 zinc finger protein

Funding

This project was supported by the Academy of Finland (332212) and University of Helsinki

(Early Career Investigator Funding, Faculty of Medicine) (N.K-A.), Finnish Cultural

Foundation (00190186, 00200185, 00212573, and 00222347) (P.A.), and Research funds from

Helsinki University Hospital (A.T., T.T).

Contributions

P.A., J.V., V.S-A., C.H-G., T.T., A.T. and N.K-A. contributed to the study design. H.K., V.S-

A., and A.T. recruited the study participants. P.A., J.V., K.R., E.W., H.M., C.H. and N.K-A.

contributed to the sample collection and processing. P.A., K.R., I.L. and N.K-A. contributed to

the laboratory experiments. P.A., J.V., K.R., I.L. and N.K-A. contributed to the data analysis.

P.A., J.V. and N.K-A. drafted the manuscript. All authors contributed to the revision of the

manuscript. All authors gave final approval of the version to be published.

Supplementary material

Supplementary Figures 1–6

Supplementary Tables 1–32

Ethics approval and consent to participate

Informed consent was obtained from all participants and the study was approved by the Ethics

Committee of Helsinki University Central Hospital (386/13/03/03/2012, 285/13/03/03/2013).

Competing interests

The authors declare no competing interests.

Availability of data

The datasets supporting the conclusions of the current study are included within the article and

its additional files. Due to the sensitive nature of the patient data used, the data sets are not and

cannot be made publicly available.

Acknowledgements

We gratefully thank all families that participated in this study. We would also like to

acknowledge Teija Karkkulainen, Riikka Vass, and Ira Larsen for their contribution to the

recruitment of the participants as well as Samuli Auvinen for his assistance with bisulphite

sequencing visualization.

REFERENCES

1.   European Ivf Monitoring Consortium, f.t.E.S.o.H.R. et al. ART in Europe, 2018: results

generated from European registries by ESHRE. *Hum Reprod Open* **2022**, hoac022

(2022).

2.   Qin, J.B. et al. Worldwide prevalence of adverse pregnancy outcomes among singleton

pregnancies after in vitro fertilization/intracytoplasmic sperm injection: a systematic

review and meta-analysis. *Arch Gynecol Obstet* **295**, 285-301 (2017).

3.   Vermey, B.G. et al. Are singleton pregnancies after assisted reproduction technology

(ART) associated with a higher risk of placental anomalies compared with non-ART

singleton pregnancies? A systematic review and meta-analysis. *BJOG* **126**, 209-218

(2019).

4.   Ceelen, M., van Weissenbruch, M.M., Vermeiden, J.P.W., van Leeuwen, F.E. & de

Waal, H.A.D.V. Cardiometabolic differences in children born after in vitro fertilization:

Follow-up study. *J Clin Endocr Metab* **93**, 1682-1688 (2008).

5.   Guo, X.Y. et al. Cardiovascular and metabolic profiles of offspring conceived by
838   assisted reproductive technologies: a systematic review and meta-analysis. *Fertility and*
839   *Sterility* **107**, 622-631.e5. (2017).

840   6.   Klemetti, R. et al. Puberty disorders among ART-conceived singletons: a Nordic
841   register study from the CoNARTaS group. *Hum Reprod* **37**, 2402-2411 (2022).

842   7.   Lazaraviciute, G., Kauser, M., Bhattacharya, S., Haggarty, P. & Bhattacharya, S. A
843   systematic review and meta-analysis of DNA methylation levels and imprinting
844   disorders in children conceived by IVF/ICSI compared with children conceived
845   spontaneously. *Hum Reprod Update* **20**, 840-852 (2014).

846   8.   Henningsen, A.A. et al. Imprinting disorders in children born after ART: a Nordic study
847   from the CoNARTaS group. *Hum Reprod* **35**, 1178-1184 (2020).

848   9.   Rönö, K. et al. The neurodevelopmental morbidity of children born after assisted
849   reproductive technology: a Nordic register study from the Committee of Nordic
850   Assisted Reproductive Technology and Safety group. *Fertility and Sterility* **117**, 1026-
851   1037 (2022).

852   10.   Maheshwari, A. et al. Is frozen embryo transfer better for mothers and babies? Can
853   cumulative meta-analysis provide a definitive answer? *Human Reproduction Update*
854   **24**, 35-58 (2018).

855   11.   Melamed, N., Choufani, S., Wilkins-Haug, L.E., Koren, G. & Weksberg, R.
856   Comparison of genome-wide and gene-specific DNA methylation between ART and
857   naturally conceived pregnancies. *Epigenetics-Us* **10**, 474-483 (2015).

858   12.   Estill, M.S. et al. Assisted reproductive technology alters deoxyribonucleic acid
859   methylation profiles in bloodspots of newborn infants. *Fertility and Sterility* **106**, 629-
860   639.e10. (2016).

861   13.   El Hajj, N. et al. DNA methylation signatures in cord blood of ICSI children. *Hum*
862   *Reprod* **32**, 1761-1769 (2017).

863   14.   Novakovic, B. et al. Assisted reproductive technologies are associated with limited
864   epigenetic variation at birth that largely resolves by adulthood. *Nat Commun* **10**, 3922
865   (2019).

866   15.   Tobi, E.W. et al. DNA methylation differences at birth after conception through ART.
867   *Hum Reprod* **36**, 248-259 (2021).

868   16.   Yeung, E.H. et al. Conception by fertility treatment and offspring deoxyribonucleic
869   acid methylation. *Fertility and Sterility* **116**, 493-504 (2021).

17.   Håberg, S.E. et al. DNA methylation in newborns conceived by assisted reproductive
technology. *Nat Commun* **13**, 1896 (2022).

18.   Rahimi, S. et al. Capturing sex-specific and hypofertility-linked effects of assisted
reproductive technologies on the cord blood DNA methylome. *Clin Epigenetics* **15**, 82
(2023).

19.   Litzky, J.F. et al. Placental imprinting variation associated with assisted reproductive
technologies and subfertility. *Epigenetics-Us* **12**, 653-661 (2017).

20.   Xu, N. et al. Comparison of Genome-Wide and Gene-Specific DNA Methylation
Profiling in First-Trimester Chorionic Villi From Pregnancies Conceived With
Infertility Treatments. *Reprod Sci* **24**, 996-1004 (2017).

21.   Choufani, S. et al. Impact of assisted reproduction, infertility, sex and paternal factors
on the placental DNA methylome. *Hum Mol Genet* **28**, 372-385 (2019).

22.   Mani, S. et al. Embryo cryopreservation leads to sex-specific DNA methylation
perturbations in both human and mouse placentas. *Hum Mol Genet* **31**, 3855-3872
(2022).

23.   Mann, M.R.W. et al. Selective loss of imprinting in the placenta following
preimplantation development in culture. *Development* **131**, 3727-3735 (2004).

24.   de Waal, E. et al. The cumulative effect of assisted reproduction procedures on
placental development and epigenetic perturbations in a mouse model. *Hum Mol Genet*
24, 6975-6985 (2015).

25.   Rahimi, S. et al. Moderate maternal folic acid supplementation ameliorates adverse
embryonic and epigenetic outcomes associated with assisted reproduction in a mouse
model. *Hum Reprod* **34**, 851-862 (2019).

26.   Ghosh, J., Coutifaris, C., Sapienza, C. & Mainigi, M. Global DNA methylation levels
are altered by modifiable clinical manipulations in assisted reproductive technologies.
Clin Epigenetics **9**, 14 (2017).

27.   Choux, C. et al. The epigenetic control of transposable elements and imprinted genes
in newborns is affected by the mode of conception: ART versus spontaneous
conception without underlying infertility. *Hum Reprod* **33**, 331-340 (2018).

28.   Barberet, J. et al. Do frozen embryo transfers modify the epigenetic control of imprinted
genes and transposable elements in newborns compared with fresh embryo transfers
and natural conceptions? *Fertility and Sterility* **116**, 1468-1480 (2021).

29.   Gong, S. et al. Genome-wide oxidative bisulfite sequencing identifies sex-specific
methylation differences in the human placenta. *Epigenetics-Us* **13**, 228-239 (2018).

30. Andrews, S.V., Yang, I.J., Froehlich, K., Oskotsky, T. & Sirota, M. Large-scale
placenta DNA methylation integrated analysis reveals fetal sex-specific differentially
methylated CpG sites and regions. *Sci Rep-Uk* **12**, 9396 (2022).

31. Sandman, C.A., Glynn, L.M. & Davis, E.P. Is there a viability-vulnerability tradeoff?
Sex differences in fetal programming. *J Psychosom Res* **75**, 327-335 (2013).

32. Tarrade, A., Panchenko, P., Junien, C. & Gabory, A. Placental contribution to
nutritional programming of health and diseases: epigenetics and sexual dimorphism. *J*
Exp Biol **218**, 50-58 (2015).

33. Kaartinen, N.M. et al. Male gender explains increased birthweight in children born after
transfer of blastocysts. *Hum Reprod* **30**, 2312-2320 (2015).

34. Litzky, J.F. et al. Effect of frozen/thawed embryo transfer on birthweight, macrosomia,
and low birthweight rates in US singleton infants. *Am J Obstet Gynecol* **218**, 433.e1-
433.e10. (2018).

35. Coetzee, K., Ozgur, K., Bulut, H., Berkkanoglu, M. & Humaidan, P. Large-for-
gestational age is male-gender dependent in artificial frozen embryo transfers cycles: a
cohort study of 1295 singleton live births. *Reprod Biomed Online* **40**, 134-141 (2020).

36. Sankilampi, U., Hannila, M.L., Saari, A., Gissler, M. & Dunkel, L. New population-
based references for birth weight, length, and head circumference in singletons and
twins from 23 to 43 gestation weeks. *Ann Med* **45**, 446-454 (2013).

37. Gomes, L.G. et al. DLK1 Is a Novel Link Between Reproduction and Metabolism. *J*
Clin Endocr Metab **104**, 2112-2120 (2019).

38. Tan, J.H.L., Wollmann, H., van Pelt, A.M.M., Kaldis, P. & Messerschmidt, D.M.
Infertility-Causing Haploinsufficiency Reveals TRIM28/KAP1 Requirement in
Spermatogonia. *Stem Cell Rep* **14**, 818-827 (2020).

39. Dalgaard, K. et al. Trim28 Haploinsufficiency Triggers Bi-stable Epigenetic Obesity.
Cell **164**, 353-364 (2016).

40. Li, R. et al. TRIM28 modulates nuclear receptor signaling to regulate uterine function.
Nat Commun **14**, 4605 (2023).

41. Bond, S.T. et al. Deletion of Trim28 in committed adipocytes promotes obesity but
preserves glucose tolerance. *Nat Commun* **12**, 74 (2021).

42. Domenga, V. et al. Notch3 is required for arterial identity and maturation of vascular
smooth muscle cells. *Gene Dev* **18**, 2730-2735 (2004).

43. Dietrich, B.V., K; Lackner, A. I.; Meinhardt, G; Koo, B-K.; Pollheimer, J.; Haider, S;
Knöfler, M. NOTCH3 signalling controls human trophoblast stem cell expansion and
differentiation. *bioRxiv* **2023.07.03.547490**(2023).

44. Afshar, Y. et al. Notch1 mediates uterine stromal differentiation and is critical for
complete decidualization in the mouse. *Faseb J* **26**, 282-294 (2012).

45. Kim, J.Y. & Kim, Y.M. Acute Atherosis of the Uterine Spiral Arteries:
Clinicopathologic Implications. *J Pathol Transl Med* **49**, 462-471 (2015).

46. Dicke, A.K. et al. DDX3Y is likely the key spermatogenic factor in the AZFa region
that contributes to human non-obstructive azoospermia. *Commun Biol* **6**, 350 (2023).

47. Holmlund, H., Yamauchi, Y., Ruthig, V.A., Cocquet, J. & Ward, M.A. Return of the
forgotten hero: the role of Y chromosome-encoded Zfy in male reproduction. *Mol Hum*
Reprod **29**, gaad025 (2023).

48. Anilkumar, T.R. et al. Expression of protocadherin 11Yb (PCDH11Yb) in seminal
germ cells is correlated with fertility status in men. *Reprod Fert Develop* **29**, 2100-2111
(2017).

49. Qin, Y.F. et al. Genetic variants in epoxide hydrolases modify the risk of
oligozoospermia and asthenospermia in Han-Chinese population. *Gene* **510**, 171-174
(2012).

50. Thelin, M.A. et al. Biological functions of iduronic acid in chondroitin/dermatan
sulfate. *Febs J* **280**, 2431-2446 (2013).

51. Eggermann, T. et al. Trans-acting genetic variants causing multilocus imprinting
disturbance (MLID): common mechanisms and consequences. *Clin Epigenetics* **14**, 41
(2022).

52. Li, X.J. et al. A Maternal-Zygotic Effect Gene, Zfp57, Maintains Both Maternal and
Paternal Imprints. *Dev Cell* **15**, 547-557 (2008).

53. Quenneville, S. et al. In Embryonic Stem Cells, ZFP57/KAP1 Recognize a Methylated
Hexanucleotide to Affect Chromatin and DNA Methylation of Imprinting Control
Regions. *Mol Cell* **44**, 361-372 (2011).

54. Cortessis, V.K. et al. Comprehensive meta-analysis reveals association between
multiple imprinting disorders and conception by assisted reproductive technology. *J*
Assist Reprod Gen **35**, 943-952 (2018).

55. Jirtle, R.L. Geneimprint. Imprinted Genes: by Species.
https://www.geneimprint.com/site/genes-by-species. (2023).

56. Barberet, J. et al. DNA methylation profiles after ART during human lifespan: a
systematic review and meta-analysis. *Human Reproduction Update* **28**, 629-655
(2022).

57. Ochoa, E. et al. ImprintSeq, a novel tool to interrogate DNA methylation at human
imprinted regions and diagnose multilocus imprinting disturbance. *Genet Med* **24**, 463-
474 (2022).

58. Hara, S., Terao, M., Tsuji-Hosokawa, A., Ogawa, Y. & Takada, S. Humanization of a
tandem repeat in IG-DMR causes stochastic restoration of paternal imprinting at mouse
Dlk1-Dio3 domain. *Hum Mol Genet* **30**, 564-574 (2021).

59. Alexander, K.A., Wang, X., Shibata, M., Clark, A.G. & Garcia-Garcia, M.J. TRIM28
Controls Genomic Imprinting through Distinct Mechanisms during and after Early
Genome-wide Reprogramming. *Cell Rep* **13**, 1194-1205 (2015).

60. Lu, H.P. et al. TRIM28 Regulates Dlk1 Expression in Adipogenesis. *Int J Mol Sci* **21**,
7245 (2020).

61. Davies, M.J. et al. Reproductive Technologies and the Risk of Birth Defects. *New Engl*
J Med **366**, 1803-1813 (2012).

62. Auvinen, P. et al. Chromatin modifier developmental pluripotency associated factor 4
(DPPA4) is a candidate gene for alcohol-induced developmental disorders. *Bmc Med*
20, 495 (2022).

63. Laufer, B.I. et al. Placenta and fetal brain share a neurodevelopmental disorder DNA
methylation profile in a mouse model of prenatal PCB exposure. *Cell Rep* **38**, 110442
(2022).

64. Zhu, Y.H. et al. Placental methylome reveals a 22q13.33 brain regulatory gene locus
associated with autism. *Genome Biol* **23**, 46 (2022).

65. Dumoulin, J.C. et al. Growth rate of human preimplantation embryos is sex dependent
after ICSI but not after IVF. *Hum Reprod* **20**, 484-91 (2005).

66. Kai, C.M. et al. Serum insulin-like growth factor-I (IGF-I) and growth in children born
after assisted reproduction. *J Clin Endocr Metab* **91**, 4352-4360 (2006).

67. Umapathy, A. et al. Mesenchymal Stem/Stromal Cells from the Placentae of Growth
Restricted Pregnancies Are Poor Stimulators of Angiogenesis. *Stem Cell Rev Rep* **16**,
557-568 (2020).

68. Boss, A.L., Chamley, L.W., Brooks, A.E.S. & James, J.L. Differences in human
placental mesenchymal stromal cells may impair vascular function in FGR.
Reproduction **162**, 319-330 (2021).

69.   Wang, Y.F. et al. TRIM28 regulates sprouting angiogenesis through VEGFR-DLL4-
1004   Notch signaling circuit. *Faseb J* **34**, 14710-14724 (2020).

1005   70.   Mitterberger, M.C. et al. DLK1(PREF1) is a negative regulator of adipogenesis in
1006   CD105(+)/CD90(+)/CD34(+)/CD31(-)/FABP4(-) adipose-derived stromal cells from
1007   subcutaneous abdominal fat pats of adult women. *Stem Cell Res* **9**, 35-48 (2012).

1008   71.   Vietor, I. et al. The negative adipogenesis regulator Dlk1 is transcriptionally regulated
1009   by Ifrd1 (TIS7) and translationally by its orthologue Ifrd2 (SKMc15). *Elife* **12**, e88350
1010   (2023).

1011   72.   Moon, Y.S. et al. Mice lacking paternally expressed Pref-1/Dlk1 display growth
1012   retardation and accelerated adiposity. *Mol Cell Biol* **22**, 5585-5592 (2002).

1013   73.   Villena, J.A. et al. Resistance to High-Fat Diet-Induced Obesity but Exacerbated
1014   Insulin Resistance in Mice Overexpressing Preadipocyte Factor-1 (Pref-1) A New
1015   Model of Partial Lipodystrophy. *Diabetes* **57**, 3258-3266 (2008).

1016   74.   Dauber, A. et al. Paternally Inherited DLK1 Deletion Associated With Familial Central
1017   Precocious Puberty. *J Clin Endocr Metab* **102**, 1557-1567 (2017).

1018   75.   Ioannides, Y., Lokulo-Sodipe, K., Mackay, D.J.G., Davies, J.H. & Temple, I.K. Temple
1019   syndrome: improving the recognition of an underdiagnosed chromosome 14 imprinting
1020   disorder: an analysis of 51 published cases. *J Med Genet* **51**, 495-501 (2014).

1021   76.   Mackay, D.J.G. et al. Hypomethylation of multiple imprinted loci in individuals with
1022   transient neonatal diabetes is associated with mutations in ZFP57. *Nat Genet* **40**, 949-
1023   951 (2008).

1024   77.   Fauque, P. et al. Reproductive technologies, female infertility, and the risk of
1025   imprinting-related disorders. *Clin Epigenetics* **12**, 191 (2020).

1026   78.   Mu, J. et al. Mutations in NLRP2 and NLRP5 cause female infertility characterised by
1027   early embryonic arrest. *J Med Genet* **56**, 471-480 (2019).

1028   79.   Barker, D.J., Winter, P.D., Osmond, C., Margetts, B. & Simmonds, S.J. Weight in
1029   infancy and death from ischaemic heart disease. *Lancet* **2**, 577-580 (1989).

1030   80.   Barker, D.J.P. The developmental origins of chronic adult disease. *Acta Paediatr* **93**,
1031   26-33 (2004).

1032   81.   Cleaton, M.A.M. et al. Fetus-derived DLK1 is required for maternal metabolic
1033   adaptations to pregnancy and is associated with fetal growth restriction. *Nat Genet* **48**,
1034   1473-1480 (2016).

82. Declercq, E. et al. Perinatal outcomes associated with assisted reproductive technology:
the Massachusetts Outcomes Study of Assisted Reproductive Technologies
(MOSART). *Fertil Steril* **103**, 888-895 (2015).

83. Tian, Y. et al. ChAMP: updated methylation analysis pipeline for Illumina BeadChips.
Bioinformatics **33**, 3982-3984 (2017).

84. Wang, Y.C. et al. InterpolatedXY: a two-step strategy to normalize DNA methylation
microarray data avoiding sex bias. *Bioinformatics* **38**, 3950-3957 (2022).

85. Pidsley, R. et al. A data-driven approach to preprocessing Illumina 450K methylation
array data. *Bmc Genomics* **14**, 293 (2013).

86. Johnson, W.E., Li, C. & Rabinovic, A. Adjusting batch effects in microarray expression
data using empirical Bayes methods. *Biostatistics* **8**, 118-127 (2007).

87. Zhou, W.D., Laird, P.W. & Shen, H. Comprehensive characterization, annotation and
innovative use of Infinium DNA methylation BeadChip probes. *Nucleic Acids Res* **45**,
e22 (2017).

88. Hansen, K. IlluminaHumanMethylationEPICanno.ilm10b4.hg19: Annotation for
Illumina's EPIC methylation arrays. R package version 0.6.0. (2017).

89. Illumina. https://www.illumina.com. (2023).

90. Ritchie, M.E. et al. limma powers differential expression analyses for RNA-sequencing
and microarray studies. *Nucleic Acids Res* **43**, e47 (2015).

91. Yuan, V. et al. Cell-specific characterization of the placental methylome. *Bmc*
Genomics **22**, 6 (2021).

92. Kolde, R., Martens, K., Lokk, K., Laur, S. & Vilo, J. seqlm: an MDL based method for
identifying differentially methylated regions in high density methylation array data.
Bioinformatics **32**, 2604-2610 (2016).

93. Phipson, B., Maksimovic, J. & Oshlack, A. missMethyl: an R package for analyzing
data from Illumina's HumanMethylation450 platform. *Bioinformatics* **32**, 286-288
(2016).

94. Li, S. et al. Genome-wide average DNA methylation is determined in utero. *Int J*
Epidemiol **47**, 908-916 (2018).

95. Zheng, Y.A. et al. Prediction of genome-wide DNA methylation in repetitive elements.
Nucleic Acids Res **45**, 8697-8711 (2017).

96. Zymo Research. Bisulfite Primer Seeker. https://zymoresearch.eu/pages/bisulfite-
primer-seeker. (2023).

97.   Bock, C. et al. BiQ analyzer: visualization and quality control for DNA methylation

1069   data from bisulfite sequencing. *Bioinformatics* **21**, 4067-4068 (2005).

1070   98.   Macosko, E.Z. et al. Highly Parallel Genome-wide Expression Profiling of Individual

1071   Cells Using Nanoliter Droplets. *Cell* **161**, 1202-1214 (2015).

1072   99.   Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and

1073   dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

1074   100.   Zhang, Y.Q., Parmigiani, G. & Johnson, W.E. ComBat-seq: batch effect adjustment for

1075   RNA-seq count data. *Nar Genom Bioinform* **2**, lqaa078 (2020).

1076   101.   Wu, T.Z. et al. clusterProfiler 4.0: A universal enrichment tool for interpreting omics

1077   data. *Innovation-Amsterdam* **2**, 100141 (2021).

1078   102.   Team, R.C. R: a language and environment for statistical computing. Vienna: R

1079   Foundation for Statistical Computing. (2023).

1080

Table 1. Samples sizes of each study group in phenotype, genome-wide DNAm and mRNA-seq analyses. Study groups are presented in bold with sample sizes in each analysis including the number of samples in the sex-specific analyses. The ART subgroups are shown on a white background. The subgroups or sex-specific samples which are not included in the analyses are written in light gray. IUI and SF groups (indicated with asterisks) are not included in genome-wide DNAm analysis. ART: assisted reproductive technology, IVF: *in vitro* fertilization, ICSI: intracytoplasmic sperm injection, FRESH: fresh embryo transfer, FET: frozen embryo transfer, IUI: insemination, and SF: subfertile.

Group	Phenotypes and Genome-wide DNAm N (male/female)	Genome-wide mRNA-seq N (male/female)
Control	77 (42/35)	39 (19/20)
ART	80 (36/44)	59 (29/30)
IVF	50 (24/26)	38 (20/18)
IVF-FRESH	25 (10/15)	19 (8/11)
IVF-FET	25 (14/11)	19 (12/7)
ICSI	30 (12/18)	21 (9/12)
ICSI-FRESH	17 (4/13)	11 (3/8)
ICSI-FET	13 (8/5)	10 (6/4)
FRESH	42 (14/28)	30 (11/19)
FET	38 (22/16)	29 (18/11)
IUI	12* (8/4)	10 (6/4)
SF	11* (5/6)	10 (4/6)

Fig. 1. ART-associated differential DNAm and ART-, IUI-, SF-, and *in vitro* culture-associated differential gene expression. a Volcano plot showing the distribution of associations between CpG sites and ART. DMPs with $\Delta\beta \leq -0.065$ and $\Delta\beta \geq 0.065$ are labeled based on UCSC RefGene Name for visualization. **b** Significantly enriched terms identified in KEGG enrichment analysis for DMRs ($P < 0.05$). The 30 most significant pathways are shown. **c** Volcano plot showing the distribution of associations between mRNA expression and ART. The 28 most significant DEGs are labeled for visualization including only confirmed genes. **d** Normalized mRNA-seq counts of *TRIM28, DLK1*, and *NOTCH3* genes in control, ART, IVF, ICSI, FRESH, FET, IUI, and SF placental samples. Data

presented as mean ±SD. *FDR < 0.05 in genome-wide mRNA-seq analysis compared to controls. **e** Significantly enriched terms identified in GO:BP enrichment analysis for ART-associated DEGs (FDR-corrected *q*-value < 0.05). All significant pathways are shown. **f** Volcano plot showing the distribution of associations between mRNA expression and *in vitro* culture. All significant DEGs are labeled for visualization including only confirmed genes. In volcano plots, horizontal line marks FDR 0.05 and vertical line marks effect size ±0.05. Control *n* = 77 and ART *n* = 80 in DNAm and control *n* = 39, ART *n* = 59, IVF *n* = 38, ICSI *n* = 21, IUI *n* = 10, and SF *n* = 10 in mRNA-seq analyses.

Fig. 2. IVF- and ICSI-associated differential DNAm and gene expression. a Volcano plot showing the distribution of associations between CpG sites and IVF. DMPs with $\Delta\beta \leq -0.065$ and $\Delta\beta \geq 0.065$ are labeled based on UCSC RefGene Name for visualization. **b** Volcano plot showing the distribution of associations between CpG sites and ICSI. DMPs with $\Delta\beta \leq -0.055$ and $\Delta\beta \geq 0.055$ are labeled based on UCSC RefGene Name for visualization. **c** Venn diagram showing the number of ART-associated DMPs as well as DMR-associated genes, which are in common between IVF and ICSI

placentas. **d** Volcano plot showing the distribution of associations between mRNA expression and IVF. All significant DEGs are labeled for visualization including only confirmed genes. **e** Volcano plot showing the distribution of associations between mRNA expression and ICSI. The 28 most significant DEGs are labeled for visualization. **f** Venn diagram showing the number of ART-associated DEGs, which are in common between IVF and ICSI placentas. In volcano plots, horizontal line marks FDR 0.05 and vertical line marks effect size ±0.05. Control $n = 77$, IVF $n = 50$, and ICSI $n = 30$ in DNAm and control $n = 39$, IVF $n = 38$, and ICSI $n = 21$ in mRNA-seq analyses.

Fig. 3. FRESH- and FET-associated differential DNAm and gene expression. a Volcano plot showing the distribution of associations between CpG sites and FRESH. DMPs with Δβ ≤ −0.057 and Δβ ≥ 0.057 are labeled based on UCSC RefGene Name for visualization. **b** Volcano plot showing the distribution of associations between CpG sites and FET. DMPs with Δβ ≤ −0.058 and Δβ ≥ 0.058 are labeled based on UCSC RefGene Name for visualization. **c** Venn diagram showing the number of ART-associated DMPs as well as DMR-associated genes, which are in common between FRESH and

FET placentas. **d** Volcano plot showing the distribution of associations between mRNA expression and FRESH. The 26 most significant DEGs are labeled for visualization including only confirmed genes. **e** Volcano plot showing the distribution of associations between mRNA expression and FET. All significant DEGs are labeled for visualization including only confirmed genes. **f** Venn diagram showing the number of ART-associated DEGs, which are in common between FRESH and FET placentas. **g** Volcano plot showing the distribution of associations between CpG sites and IVF-FRESH. All DMPs are labeled for visualization. **h** Volcano plot showing the distribution of associations between CpG sites and IVF-FET. All DMPs are labeled for visualization. **i** Normalized DNAm levels of DMPs in *ZFP57* (cg13773306) and *NLRP5* (cg14217229) in placentas of control, ART, and ART subgroups. *FDR < 0.05 in genome-wide DNAm analyses compared to controls. In volcano plots, horizontal line marks FDR 0.05 and vertical line marks effect size ±0.05. Control n = 77, ART n = 80, IVF n = 50, ICSI n = 30, FRESH n = 42, FET n = 38, IVF-FRESH n = 25, and IVF-FET n = 25 in DNAm and control n = 39, FRESH n = 30, and FET n = 29 in mRNA-seq analyses.

Fig. 4. ART- and SF-associated DNAm profiles at *DLK1-DIO3* ICR. a Schematic figure about *DLK1* and *DLK1-DIO3* ICR. The chromosomal region chr14:101,277,232-101,277,826 at the ICR, containing repeated sequence with nine underlined motifs (chr14:101,277,232-101,277,411) and PCR amplified sequence (chr14:101,277,375-101,277,826), is presented. Primers spanning the amplicon are highlighted in gray and the covered 12 CpG sites and the SNPs (rs1884539A/G and rs75998174A/G) used to distinguish paternal and maternal alleles are bolded. **b** DNAm levels at 12 CpG sites in *DLK1-DIO3* ICR in control ($n = 6$), ART ($n = 8$), IVF ($n = 4$), ICSI ($n = 4$), and SF ($n = 4$) placentas. Differences in DNAm at the CpG sites were calculated between control, ART, and SF placentas. Data presented as mean ±SD. $*P < 0.05$ and $**P < 0.01$, One-Way ANOVA followed by Bonferroni post hoc.