1
2
3
4
Stroke Severity and Cognition in the REGARDS-CARES Study

Jason A. Blake[1], D. Leann Long[2], Amy J. Knight[3], Burel R. Goodin[4], Michael Crowe[1], Suzanne
E. Judd[2], David L. Roth[5], & Olivio J. Clay[1,6]

[1]Department of Psychology, University of Alabama at Birmingham

[2]Department of Biostatistics, University of Alabama at Birmingham

[3]Department of Neurology, University of Alabama at Birmingham

[4]Department of Anesthesiology, Washington University in St. Louis

[5]Center on Aging and Health, Johns Hopkins School of Medicine

[6]Alzheimer's Disease Research Center, University of Alabama at Birmingham

Abstract Word Count: 245
Manuscript Word Count: 5,999
19

Corresponding Author:

Jason A. Blake, MA
Department of Psychology, University of Alabama at Birmingham
Campbell Hall 415
1300 University Boulevard
Birmingham, AL 35233
Email: jablake@uab.edu
Phone: (215)939-3744
ORCID: 0000-0002-4575-2420

Conflict of Interest: For work unrelated to this manuscript, D. Leann Long received
investigator-initiated research support from Amgen, Inc. Otherwise, the authors declare no
conflicts of interest.

33   **ABSTRACT**

Objective: Cognitive impairment after stroke is common, present in 30-50% of survivors. Stroke

severity, indicated by both volume and location, is the most consequential predictor of cognitive

impairment, with severe strokes predicting higher chances of cognitive impairment. The current

investigation examines the associations of two stroke severity ratings and a caregiver-report of

post-stroke functioning with longitudinal cognitive outcomes. **Methods:** The analysis was

conducted on 157 caregivers and stroke survivor dyads who participated in the Caring for Adults

Recovering from the Effects of Stroke (CARES) project, an ancillary study of the REasons for

Geographic and Racial Differences in Stroke (REGARDS) national cohort study. Glasgow Coma

Scale (GCS) and modified Rankin Scale (mRS) collected at hospitalization discharge were

included as two primary predictors of cognitive impairment. The number of caregiver-reported

problems and impairments at nine months following stroke were included as a third predictor.

Cognition was assessed using a biennial telephone battery, incorporating multiple cognitive

assessments and sections from the Montreal Cognitive Assessment. Longitudinal cognitive

scores were analyzed up to five years post-stroke, controlling for baseline (pre-stroke) cognitive

scores and demographic variables of each stroke survivor collected at CARES baseline. **Results:**

Separate mixed models showed significant main effects of GCS (b=0.3280, p=0.0009), mRS

(b=-0.2119, p=0.0002), and caregiver-reported impairments (b=-0.0671, p<0.0001) on

longitudinal cognitive scores. In a combined model including all three predictors, only caregiver-

reported problems significantly predicted cognitive outcomes (b=-0.0480, p<0.0001). **Impact:**

These findings underscore the importance of incorporating caregivers feedback in understanding

cognitive consequences of stroke.

Key words: stroke outcomes, stroke severity, caregiver assessment, cognition

INTRODUCTION

Stroke is a leading cause of death and disability globally.[1, 2] The prevalence of cognitive

impairment in the chronic phase ranges from 22%[3] to 57% after an individual's first stroke.[4, 5]

Stroke recovery is a highly variable and individualized process that does not adhere to a fixed

timeline. The first six months after a stroke, termed the "critical recovery window," are

characterized by the most dramatic improvements in the physical and cognitive functioning of

the survivior[6]. However, the degree of these improvements greatly differs among stroke

survivors after six months; some experience vast recovery, leading to fewer residual deficits,

while others may show more modest progress. The survivors who experience larger functional

improvements during this critical period are more likely to demonstrate better cognitive scores in

subsequent years[7]. This variable rate of recovery can contribute to a broad range of cognitive and

functional outcomes in post-stroke recovery.

To better improve the outcomes of stroke survivors, it is necessary to understand factors

that may be prognostic of post-stroke cognitive recovery. Presence of functional impairment due

to stroke severity has been shown to predict worse long-term functional outcomes[8, 9] including

occupational[10], depressive[11] or other psychosocial symptoms[12], and overall quality of life[13, 14] in

survivors. In previous studies, it has been demonstrated that initial stroke severity, indicated by

both volume and location of stroke, is the most consequential predictor of cognitive impairment,

with severe strokes predicting higher chances of cognitive impairment.[15-17]

Stroke severity in the acute phase is often determined through an objective evaluation of

neurological functioning at the time hospitalization and at discharge. A common objective

evaluation of stroke severity is the Glasgow Coma Scale (GCS) which describes the extent of

impaired consciousness according to the patient's ability to respond commands[18]. Severity is also

commonly determined through an assessment of the patient's functional status that measures

capacity to carry out activities of daily living (ADLs), one such functional assessment of stroke

is the modified Rankin Scale (mRS)[19]. Both the GCS[20] and mRS[21] play crucial roles in guiding

clinical decision-making and treatment planning, this is partially due to their ability to predict

long-term stroke outcomes. However, these types of measures have some limitations in their

ability to predict functional outcomes, particularly regarding subtle cognitive or physical

changes, which may warrant the exploration of additional sources of information[22, 23].

Caregivers, who spend the most time with stroke survivors, are in a unique position to

provide valuable insight into survivors' functioning. Their observations can potentially

complement clinically administered assessments and contribute to a more comprehensive

understanding of stroke survivors' cognitive challenges and recovery process. Integrating

caregiver reports into the assessment process may help to identify cognitive deficits and areas of

difficulty that might otherwise be overlooked by formal assessments.

For this investigation, it is hypothesized that more severe stroke, as measured by

clinically administered scales at hospital discharge, is predictive of lower longitudinal cognitive

scores in stroke survivors. Additionally, a higher number of impairments and problems reported

by the caregivers is hypothesized to be associated with lower cognitive scores in stroke survivors

over time. The inclusion of caregiver reports, particularly after the six-month critical recovery

window, may provide a more comprehensive picture of the patient's cognitive and functional

status and improve the accuracy of predictions of long-term outcomes following a stroke.

102   **METHODS**

103   **Participant Recruitment**

104   All participants were recruited via the REasons for Geographic And Racial Differences in

Stroke (REGARDS) study. REGARDS is an ongoing national epidemiologic cohort study

investigating stroke incidence and mortality that initially enrolled 30,239 participants from 2003

to 2007[24]. The REGARDS study enrolled participants aged 45 or older and included 30% from

the Stroke Belt (encompassing Southeastern U.S. states), 20% from the Stroke Buckle (a subset

of the Stroke Belt, specifically the coastal plain region of North Carolina, South Carolina, and

Georgia), and the remainder from elsewhere in the continental U.S. Each region maintained a

balanced demographic, with roughly equal representation of White and Black individuals, and a

nearly equal gender distribution within each race-region group. The REGARDS study

intentionally oversampled Black individuals due to the study's specific interest in the comparison

of those two racial groups. Once enrolled in REGARDS, participants were administered bi-

annual cognitive testing over the phone to measure several cognitive domains such as memory

and verbal fluency/executive functioning. All relevant measures and assessments are detailed

below. The methodologies for participant sampling, recruitment, and data collection in the

REGARDS study have been described further in prior studies[24-26].

CARES Enrollment

120   Stroke survivors that were initially enrolled in the REGARDS project were then

recruited, along with their primary caregivers, to participate in the ancillary project CARES

(Caring for Adults Recovering from the Effects of Stroke). Study enrollment for CARES

occurred from 2005 to 2016. Eligibility for the CARES study required stroke survivors to (1) be

living in the community nine months post-stroke, excluding those in nursing homes or other

institutional settings; (2) have a family member or close friend acting as an informal caregiver, or

had one immediately following stroke; and (3) have the caregiver also agree to participate in the

CARES project[27]. Once participants were enrolled in the CARES study, the stroke survivor and

caregiver dyads were administered the initial baseline CARES interview[28]. The CARES

interviews were conducted via telephone approximately nine months following the stroke event.

Of the stroke survivor group, all self-reported a first-time stroke during a follow-up call. The

survivors' medical records were also examined independently to collect measures of stroke

severity at the time of hospitalization discharge, confirm a stroke event, and determine the type

of stroke when possible. Roth, et al. [28] provides a detailed overview of the data collection

procedures employed in the CARES study. Both projects were reviewed and approved by the

Institutional Review Boards of the University of Alabama at Birmingham and each participating

institution.

Demographics

Demographic information was collected from both the stroke survivors and their

caregivers during their enrollment into the REGARDS study as well as during the initial CARES

interview. The stroke survivor and caregiver demographic variables used in this analysis,

including race, educational level, age, and gender, were collected at CARES enrollment. Sex

(male or female), race (Black/AA or White) and marital status (married or unmarried) were

dichotomous variables based on self-report.

Measures

Stroke Survivor Severity

GCS[18] and mRS[29] scores were collected from hospital chart abstraction at time of

discharge. Both measures are clinically administered measures that are commonly utilized in

inpatient hospital settings as indicators of acute phase stroke severity and may be administered

by a physician or attending nurse. The GCS specifically evaluates a patient's level of

consciousness by assessing their visual, verbal, and motor responses, providing a quantitative

measure of neurological impairment. Scores can range from a minimum of three, indicating

severe neurologic symptoms, to a maximum of fifteen, indicating fully awake, responsive, and

oriented[18]. As a measure of functional ability, the mRS assesses the degree of disability or

dependence in daily activities. The mRS offers a broader perspective on stroke-related functional

limitations, with possible scores ranging from zero, indicating no symptoms, to six, representing

severe disability or death[29].

Each caregiver reported the total number of stroke-related impairments at CARES

enrollment, an average of nine months after stroke. This 28-item instrument collected caregiver-

reported problems that interfered with their activities of daily life (ADL) and instrumental

activities of daily living (IADL) over the previous week. The instrument items were drawn from

commonly used instruments including the Frenchay Activities Index [30], Barthel index [31], and the

Revised Memory and Behavior Problems Checklist [32]. This measure was used in a previous

CARES analysis to determine severity of stroke impairments and problems [33] and has shown

good concurrent validity (correlating over .62 with the Barthel Index and Rankin scores) with

observer-assessed indicators of cognitive impairment in stroke [34]. In previous analysis,

coefficient alpha for the scale was 0.92.[34]

Cognitive Assessment

The cognitive assessment is a brief telephone battery that is administered to each

REGARDS participant every two years. This battery was used to assess cognition at each time

point, including the most recent pre-stroke battery results and all post-stroke results. The

intervals for collection of this battery are less frequent to lower participant burden in collection

and was administered to REGARDS participants during follow-up. This battery includes the

following short cognitive assessments to assess the cognitive domains of learning, memory, and

executive functioning: a) Word List Learning and Delayed Recall, from the Consortium to

Establish a Registry for Alzheimer's Disease (CERAD) battery; b) Animal (Semantic) Fluency;

c) Letter-F (Phonemic) Fluency; and d) Registration, recall, and orientation from the Montreal

Cognitive Assessment (MoCA)[35] as a part of the National Institute of Neurological Disorders

and Stroke - Canadian Stroke Network Vascular Cognitive Impairment Harmonization Standards

(NINDS/CSN) 5-minute Battery[36]. The scores for each assessment have been normed to z-scores

according to the patient's age and sex. The cognitive composite score, the average of the

assessment scores at each assessment, served as the outcome variable.

Statistical Analysis

The data analyses for this paper were performed using SAS© software Studio version

3.81[37]. Descriptive statistics were reported for the participant dyads included means and standard

deviations for continuous measures and percentages for categorical variables. Age, gender, race,

and education for each stroke survivor were collected at the CARES baseline interview and

included as covariates in all models. P values < 0.05 were considered statistically significant.

The three predictor variables utilized in this analysis are GCS scores, mRS scores, and

total number of stroke-related problems as reported by their caregiver. This analysis included the

most recent pre-stroke evaluation for each participant as a measure of baseline cognitive

functioning. To assess the relationship between post-stroke functional ability and long-term

cognitive trajectories, separate mixed effects regression models were used for each predictor. A

combined model including all three predictors was also included in this analysis to assess the

independent predictive ability of each measure in the presence of the other predictors. These

models were structured to examine logarithmic trajectories of cognition over time (number of

years post-stroke). Logarithmic trajectories are recommended over linear trajectories when

assessing longitudinal cognitive scores in stroke survivors[38, 39]. Each mixed model included in

this analysis were covariate adjusted for pre-stroke cognitive scores, time after stroke (years),

age, race, sex, and education. Timepoints greater than five years were removed to minimize

survivorship bias. Random intercepts and slopes for time at the subject level were included with

an unstructured covariance matrix.

202   Figures 1-3 display the logarithmic relationship between the three measures of stroke

severity and cognitive composite scores over time. Participants were divided into two groups

based on the 50th percentile (median) of each predictor as a cut-off point. The division of

participants into two groups based on the median score allows for a visualization of contrasting

cognitive outcomes when grouped by severity of stroke symptoms as classified by each

predictor. Least squares means were used to calculate the predicted cognitive outcomes. In all

three figures, the blue group represents lower stroke severity, as determined by the assigned

predictor variable, while the red group represents worse stroke severity. The model for each

figure includes baseline cognition, age at CARES enrollment, gender, race, and education as

covariates. As these models use the natural logarithm of time, the first timepoint is estimated at

0.01 years (~4 days) post stroke.

213   **RESULTS**

214   The original CARES study recruited approximately 360 stroke survivors initially.

However, several exclusions were made to meet the specific criteria for this analysis. Stroke

survivors were excluded if the caregiver could not be enrolled (n=31). Dyads were excluded if

the stroke survivors did not have both GCS and mRS scores from medical record abstraction

(n=78). Additionally, dyads were excluded if the stroke survivors did not have a pre-stroke

baseline cognitive assessment (n=42) or if they did not have more than one cognitive assessment

over the course of five years following their stroke (n=52). The final sample for this analysis

consisted of 157 stroke survivor and caregiver dyads that met these criteria. All p-values and

statistical significances reported in this section are based on 95% confidence intervals.

Table 1 displays the demographics of the participants in this sample at the time of the

CARES baseline interview, including overall demographics and characteristics of each predictor

group. Of the 157 stroke survivors, there was a mean age of 75.74 years (SD=7.32) at the time of

CARES interview. The mean age at the time of stroke was 74.91 years (SD=7.31). The stroke

survivor group consisted of 86 women (55%) and included 97 White participants (62%) and 60

Black/African American participants (38%).

Regarding stroke type, 132 (84%) of the participants experienced an ischemic stroke, 13

(8.5%) had a hemorrhagic stroke, and for 12 (7.5%) participants the stroke type was not specified

in the reviewed medical records. Of the 157 caregivers, the mean age at enrollment was 61.82

years (SD=13.39), and the majority were female (77%; n=121). The racial composition of the

caregiver group paralleled that of the stroke survivor group, with 97 caregivers (62%) identifying

as White, 59 (37%) as Black/African American, and one (1%) as "Other." The mean GCS score

was 4.38 (SD=0.72, range = 3 to 5), mean mRS score was 1.82 (SD=1.34, range = 0 to 5), and

mean number of caregiver-reported problems and impairments was 7.09 (SD=5.69, range = 0 to

28).

GCS and mRS Main Effects

Covariates of age, race, sex, education, baseline scores, and time after stroke were

included in all models. In the first model, the GCS score at the time of stroke hospitalization

discharge was used as the primary predictor. This model demonstrated a significant main effect

of GCS score on cognitive outcomes (b=0.3380, p=0.0009). Lower GCS scores (more severe

stroke) are associated with lower cognitive composite scores. In the second model, the mRS

score at the time of stroke hospitalization discharge was utilized as the predictor variable. This

model revealed a significant main effect on cognitive outcomes (b=-0.2119, p=0.0002). Higher

mRS scores (more severe stroke) are associated with lower cognitive composite scores. This

confirms the study hypothesis that more severe stroke, as measured by clinically administered

scales in the acute phase of stroke, is predictive of lower cognitive scores in stroke survivors.

There was no main effect of time after stroke on cognitive outcomes in any model included in

this analysis.

Caregiver-Reported Problems Main Effects

The third model included the total number of caregiver-reported problems as the primary

predictor variable for cognitive outcomes. In this model, there was a significant main effect of

caregiver-reported post-stroke functional ability on cognitive outcomes (b=-0.0671, p<0.0001).

This confirms the study hypothesis that fewer caregiver-reported problems after stroke predict

higher cognitive composite scores. The main effect of each separate model and their interaction

by time after stroke is displayed in **Table 2**.

Comparing Predictors

A final model included all three predictors and covariates to assess the contribution of

each predictor, as well as all previously included covariates. When all predictors were considered

simultaneously, caregiver-reported problems continued to be a significant predictor of cognitive

outcomes (b=-0.0480, p<0.0001). Neither GCS (b=0.02511, p=0.8742) nor mRS (b=-0.1079,

p=0.2216) showed a statistically significant effect on cognitive outcomes in the presence of other

predictors.

Standardized beta coefficients for the three predictors were also calculated to provide

insight into their relationship to longitudinal cognitive scores. The standardized betas for the

predictors are as follows: caregiver report (B=-0.4341), GCS (B=0.2773), and mRS (B=-0.3250).

These were calculated using the mean and standard deviation of each predictor and the pre-stroke

baseline cognitive score (M=-0.2669, SD=0.8790).

270   **DISCUSSION**

The current investigation revealed significant main effects of all three predictors on

longitudinal cognitive outcomes. The positive relationship between GCS and cognition indicates

that a higher GCS score, representing less severe injury, is associated with a higher cognitive

score over time. Conversely, the negative relationship between mRS and cognitive composites

demonstrates that as mRS scores increase, representing worse stroke-related disability, cognitive

scores tend to decrease. This finding supports the current literature and demonstrates that

clinically administered measures of stroke severity are prognostic of longitudinal cognitive

scores in stroke survivors, emphasizing the importance of considering acute phase functioning

measures when evaluating cognitive recovery[40].

This analysis also supported the link between caregiver-reported functioning and stroke

survivor cognition over time. Specifically, higher numbers of caregiver-reported problems were

associated with lower cognitive composite scores over time. These findings support both study

hypotheses and indicate that both caregiver reports and clinical evaluations have value in

predicting cognitive outcomes in stroke survivors. The findings also align with previous research

on the critical role of caregivers in understanding and monitoring care recipient outcomes[41, 42]. In

all models, there was a significant main effect of baseline composite scores on cognitive

outcomes. This finding is expected, as pre-stroke cognitive function is known to precipitate

cognitive outcomes after stroke.

289   A comparison of the absolute values of the standardized betas revealed the relative

strength of each predictor. Amongst the three predictors, the caregiver-reported measure of

functional impairment had the strongest association with cognitive outcomes, followed by the

mRS, a measure of functional impairment, and then GCS scores, an objective measure of

consciousness based on observable clinical features of injury. These results suggest that

assessing functional impairment, as measured by both caregiver-reported problems and mRS

scores, is crucial for understanding cognitive outcomes in stroke survivors. Notably, caregiver-

reported problems demonstrated the strongest relationship with cognitive outcomes, emphasizing

the importance of incorporating caregivers' perspectives in stroke assessment and recovery

monitoring.

299   The strong association between caregiver-reported functional impairments and cognitive

outcomes may be explained by several factors. Caregiver-reported measures provide a more

comprehensive assessment of the impact of stroke on daily functioning, including problems with

ADLs and IADLs, which are significantly affected by impairments cognitive abilities. Also, it is

important to note that the caregiver-reported problems were collected an average of nine months

after the stroke event, while the other two measures were collected from the medical records at

the time of stroke hospitalization discharge. Therefore, incorporating caregiver-reported

measures at later stages can provide a more comprehensive and nuanced understanding of

cognitive outcomes in stroke survivors. This information is especially valuable as it comes after

the six-month critical recovery window, a time when substantial functional changes can still

occur[43].

These results suggest that integrating caregiver-reported problems with clinically

administered measures, such as GCS and mRS scores in the acute phase, can provide a more

comprehensive understanding of stroke survivors' cognitive recovery. This will help tailor

rehabilitation interventions to better address the specific needs of individual stroke survivors.

These results also emphasize the importance of monitoring stroke survivors longitudinally using

telephone cognitive assessments. Regular assessments following hospitalization can monitor the

cognition of stroke survivors while also capturing the dynamic relationship between caregiver-

reported problems and cognitive outcomes. Collecting caregiver reports is also cost-effective and

time-efficient compared to other assessments, as caregivers can provide expedited updates

without the need for repeated clinical visits, especially in a population that commonly has limited

insight into their own functioning. Incorporating caregiver reports of stroke impairment into the

assessment of stroke survivors provides valuable information for clinicians and researchers

designing interventions.

This analysis has several strengths, such as the inclusion of multiple longitudinal

assessments for many of the stroke survivors. These longitudinal assessments provide a

comprehensive insight into the time-dependent relationship of stroke severity and recovery.

Another strength is the inclusion of pre-stroke cognitive scores as well as demographic variables

as covariates that ensure a rigorous control for potential confounding variables. Also, the use of a

community-based sample enhances the generalizability of the findings due to the diverse and

large sample size.

A limitation of this analysis is that it relies on secondary data, as the original research

design was not specifically tailored to address the research questions of this analysis. Also, the

original CARES study involved purely observational methods. Therefore, the conclusions that

can be drawn are correlative in nature only and causation of effects cannot be determined

through analysis of this study data. A potential source of bias stems from the subjectivity of the

caregiver-reported measure of post-stroke functioning. However, this is counterbalanced by the

inclusion of both clinician-administered severity measures from hospitalization discharge along

with the caregiver-reported problems. These collectively provide a more comprehensive

evaluation of stroke impairment than solely relying on hospital-based assessments. The timing of

the caregiver-report allowed for a more detailed assessment of stroke-related impairments

compared to the post-hospitalization measures collected in the acute phase, as it is outside of the

critical recovery window.

342   Future reports from this analysis will seek to understand factors that moderate the

relationship between stroke severity and cognition. Future investigations should also consider the

multidirectional relationship between the caregiver's experience and the patient's cognitive

outcomes. Additionally, these studies should examine the longitudinal relationship between

caregiver-reported impairments, acute phase stroke severity measures, and longitudinal cognitive

outcomes in a larger, more diverse sample of stroke survivors.

Conclusion

349   This study highlights the critical role of caregiver-reported impairments, alongside other

stroke severity measures, in predicting cognitive outcomes in stroke survivors. These findings

highlight the value of ongoing assessments that extend beyond the traditionally emphasized six-

month window of stroke recovery. Integrating both clinical assessments and caregiver

perspectives will contribute to more accurate prediction of cognitive outcomes and facilitate

tailoring interventions to the evolving needs of stroke survivors. Ultimately, this analysis

enhances the understanding of the complex interplay of factors influencing cognitive

impairment, reaffirming the need for ongoing assessments and caregiver involvement in

understanding the cognitive consequences of stroke.

Acknowledgements

This research project is supported by cooperative agreement U01 NS041588 co-funded by the

National Institute of Neurological Disorders and Stroke (NINDS) and the National Institute on

Aging (NIA), National Institutes of Health, Department of Health and Human Service. The

content is solely the responsibility of the authors and does not necessarily represent the official

views of the NINDS or the NIA. Representatives of the NINDS were involved in the review of

the manuscript but were not directly involved in the collection, management, analysis, or

interpretation of the data. The authors thank the other investigators, the staff, and the

participants of the REGARDS study for their valuable contributions. A full list of participating

REGARDS investigators and institutions can be found at:

https://www.uab.edu/soph/regardsstudy/

Author Contributions

D.L.R. & O.J.C. designed and implemented the original experiment. J.A.B. conceptualized this

manuscript, performed the data analysis, and interpreted the data. Analysis was assisted

primarily by D.L.L., S.E.J. & O.J.C.

J.A.B. wrote the manuscript, all additional authors provided supporting conceptualization and

manuscript revisions. D.L.R. secured partial funding for the initial CARES ancillary study.

Sources of Funding

Funding for the initial CARES ancillary study was provided by an investigator-initiated grant

(R01 NS045789, David Roth, PhD, PI) and by a cooperative agreement (U01 NS041588) from

the NINDS and the NIA, National Institutes of Health, Department of Health and Human

Services. There was no external funding for the analysis of this manuscript.

Conflict of Interest

For work unrelated to this manuscript, D.L.L. received investigator-initiated research support

from Amgen, Inc. Otherwise, the authors declare no conflicts of interest.

REFERENCES

1. Benjamin EJ, Muntner P, Alonso A, Bittencourt MS, Callaway CW, Carson AP, et al. Heart disease and stroke statistics-2019 update: A report from the american heart association. *Circulation*. 2019;139:e56-e528

2. Virani SS, Alonso A, Benjamin EJ, Bittencourt MS, Callaway CW, Carson AP, et al. Heart disease and stroke statistics-2020 update: A report from the american heart association. *Circulation*. 2020;141:e139-e596

3. Douiri A, Rudd AG, Wolfe CD. Prevalence of poststroke cognitive impairment: South london stroke register 1995-2010. *Stroke*. 2013;44:138-145

4. Ihle-Hansen H, Thommessen B, Wyller TB, Engedal K, Oksengard AR, Stenset V, et al. Incidence and subtypes of mci and dementia 1 year after first-ever stroke in patients without pre-existing cognitive impairment. *Dement Geriatr Cogn Disord*. 2011;32:401-407

5. Leys D, Hénon H, Mackowiak-Cordoliani M-A, Pasquier F. Poststroke dementia. *The Lancet Neurology*. 2005;4:752-759

6. Duncan PW, Min Lai S. Stroke recovery. *Topics in stroke Rehabilitation*. 1997;4:51-58

7. Bonita R, Ford M, Stewart A. Predicting survival after stroke: A three-year follow-up. *Stroke*. 1988;19:669-673

8. Huang Y, Yang S, Jia J. Factors related to long-term post-stroke cognitive impairment in young adult ischemic stroke. *Medical science monitor: international medical journal of experimental and clinical research*. 2015;21:654

9. Jokinen H, Melkas S, Ylikoski R, Pohjasvaara T, Kaste M, Erkinjuntti T, et al. Post-stroke cognitive impairment is common even after successful clinical recovery. *European journal of neurology*. 2015;22:1288-1294

10. Kauranen T, Turunen K, Laari S, Mustanoja S, Baumann P, Poutiainen E. The severity of cognitive deficits predicts return to work after a first-ever ischaemic stroke. *Journal of Neurology, Neurosurgery & Psychiatry*. 2013;84:316-321

11. Shi Y, Yang D, Zeng Y, Wu W. Risk factors for post-stroke depression: A meta-analysis. *Frontiers in aging neuroscience*. 2017;9:218

12. Greenop KR, Almeida OP, Hankey GJ, Van Bockxmeer F, Lautenschlager NT. Premorbid personality traits are associated with post-stroke behavioral and psychological symptoms: A three-month follow-up study in perth, western australia. *International psychogeriatrics*. 2009;21:1063-1071

13. De Haan R, Limburg M, Van der Meulen J, Jacobs H, Aaronson N. Quality of life after stroke: Impact of stroke type and lesion location. *Stroke*. 1995;26:402-408

14. Kapoor A, Lanctot KL, Bayley M, Herrmann N, Murray BJ, Swartz RH. Screening for post-stroke depression and cognitive impairment at baseline predicts long-term patient-centered outcomes after stroke. *Journal of geriatric psychiatry and neurology*. 2019;32:40-48

15. Tatemichi T, Desmond D, Stern Y, Paik M, Sano M, Bagiella E. Cognitive impairment after stroke: Frequency, patterns, and relationship to functional abilities. *Journal of Neurology, Neurosurgery & Psychiatry*. 1994;57:202-207

16. Patel MD, Coshall C, Rudd AG, Wolfe CD. Cognitive impairment after stroke: Clinical determinants and its associations with long-term stroke outcomes. *Journal of the American Geriatrics Society*. 2002;50:700-706

17. Coco DL, Lopez G, Corrao S. Cognitive impairment and stroke in elderly patients. *Vascular health and risk management*. 2016;12:105

18. Teasdale G, Jennett B. Assessment of coma and impaired consciousness: A practical scale. *The Lancet*. 1974;304:81-84

19. Van Swieten J, Koudstaal P, Visser M, Schouten H, Van Gijn J. Interobserver agreement

for the assessment of handicap in stroke patients. *stroke*. 1988;19:604-607
20.   van Dijck JT, Reith FC, van Erp IA, van Essen TA, Maas AI, Peul WC, et al. Decision
making in very severe traumatic brain injury (glasgow coma scale 3-5): A literature
review of acute neurosurgical management. *J Neurosurg Sci*. 2018;62:153-177
21.   Kasner SE. Clinical interpretation and use of stroke scales. *The Lancet Neurology*.
2006;5:603-612
22.   Hand B, Page SJ, White S. Stroke survivors scoring zero on the nih stroke scale score
still exhibit significant motor impairment and functional limitation. *Stroke research and*
treatment. 2014;2014
23.   Kauranen T, Laari S, Turunen K, Mustanoja S, Baumann P, Poutiainen E. The cognitive
burden of stroke emerges even with an intact nih stroke scale score: A cohort study.
Journal of Neurology, Neurosurgery & Psychiatry. 2014;85:295-299
24.   Howard VJ, Cushman M, Pulley L, Gomez CR, Go RC, Prineas RJ, et al. The reasons
for geographic and racial differences in stroke study: Objectives and design.
Neuroepidemiology. 2005;25:135-143
25.   Roth DL, Haley WE, Wadley VG, Clay OJ, Howard G. Race and gender differences in
perceived caregiver availability for community-dwelling middle-aged and older adults.
The Gerontologist. 2007;47:721-729
26.   Howard G, Safford MM, Meschia JF, Moy CS, Howard VJ, Pulley L, et al. Stroke
symptoms in individuals reporting no prior stroke or transient ischemic attack are
associated with a decrease in indices of mental and physical functioning. *Stroke*.
2007;38:2446-2452
27.   Roth DL, Haley WE, Clay OJ, Perkins M, Grant JS, Rhodes JD, et al. Race and gender
differences in 1-year outcomes for community-dwelling stroke survivors with family
caregivers. *Stroke*. 2011;42:626-631
28.   Roth DL, Haley WE, Owen JE, Clay OJ, Goode KT. Latent growth models of the
longitudinal effects of dementia caregiving: A comparison of african american and white
family caregivers. *Psychology and aging*. 2001;16:427
29.   Rankin J. Cerebral vascular accidents in patients over the age of 60: Ii. Prognosis.
Scottish medical journal. 1957;2:200-215
30.   Margaret H, Skilobeck CE. An activities index for use with stroke patients. *Age and*
ageing. 1983;12:166-170
31.   Mahoney FI. Functional evaluation: The barthel index. *Maryland state medical journal*.
1965;14:61-65
32.   Teri L, Truax P, Logsdon R, Uomoto J, Zarit S, Vitaliano PP. Assessment of behavioral
problems in dementia: The revised memory and behavior problems checklist.
Psychology and aging. 1992;7:622
33.   Grant JS, Clay OJ, Keltner NL, Haley WE, Wadley VG, Perkins MM, et al. Does
caregiver well-being predict stroke survivor depressive symptoms? A mediation analysis.
Topics in stroke rehabilitation. 2013;20:44-51
34.   Haley WE, Allen JY, Grant JS, Clay OJ, Perkins M, Roth DL. Problems and benefits
reported by stroke family caregivers: Results from a prospective epidemiological study.
Stroke. 2009;40:2129-2133
35.   Nasreddine ZS, Phillips NA, Bédirian V, Charbonneau S, Whitehead V, Collin I, et al.
The montreal cognitive assessment, moca: A brief screening tool for mild cognitive
impairment. *Journal of the American Geriatrics Society*. 2005;53:695-699
36.   Hachinski V, Iadecola C, Petersen R, Breteler M, Nyenhuis D, Black S, et al. Ninds-csn
vascular cognitive impairment harmonization standards. *Stroke*. 2006;37:2220-2241
37.   Inc SI. Sas studio 3.81. 2016
38.   Koyama T, Matsumoto K, Okuno T, Domen K. A new method for predicting functional
recovery of stroke patients with hemiplegia: Logarithmic modelling. *Clinical rehabilitation*.

2005;19:779-789
39.   Suzuki M, Sugimura Y, Yamada S, Omori Y, Miyamoto M, Yamamoto J-i. Predicting
recovery of cognitive function soon after stroke: Differential modeling of logarithmic and
linear regression. *PloS one*. 2013;8:e53488
40.   Quinn T, Dawson J, Walters M, Lees K. Functional outcome measures in contemporary
stroke trials. *International Journal of Stroke*. 2009;4:200-205
41.   Allen RS, Haley WE, Small BJ, McMillan SC. Pain reports by older hospice cancer
patients and family caregivers: The role of cognitive functioning. *The Gerontologist*.
2002;42:507-514
42.   Bergström AL, Eriksson G, von Koch L, Tham K. Combined life satisfaction of persons
with stroke and their caregivers: Associations with caregiver burden and the impact of
stroke. *Health and quality of life outcomes*. 2011;9:1-10
43.   Ballester BR, Maier M, Duff A, Cameirão M, Bermúdez S, Duarte E, et al. A critical time
window for recovery extends beyond one-year post-stroke. *Journal of neurophysiology*.
2019
502

504   **Primary Figures and Tables**

Table 1. Baseline characteristics of REGARDS-CARES participants

Demographics	All Participants	High GCS	Low GCS	Low mRS	High mRS	Fewer CGRP	Higher CGRP
	N=157	N=83	N=74	N=79	N=78	N=73	N=84
Age at Stroke	74.91 (*SD*=7.3)	74.50 (*SD*=7.8)	75.39 (*SD*=6.7)	74.22 (*SD*=7.8)	75.63 (*SD*=6.8)	74.74 (*SD*=7.5)	75.07 (*SD*=7.2)
Age at CARES interview	75.77 (*SD*=7.3)	75.38 (*SD*=7.9)	76.21 (*SD*=6.7)	75.11 (*SD*=7.8)	76.45 (*SD*=6.8)	75.59 (*SD*=7.5)	75.93 (*SD*=7.2)
Age of Caregiver	61.82 (*SD*=13.4)	61.30 (*SD*=14.2)	62.41 (*SD*=12.5)	61.53 (*SD*=14.4)	62.12 (*SD*=12.4)	66.00 (*SD*=12.2)	58.23 (*SD*=13.4)
Gender (*n*, %)							
Females	86 (55%)	42 (51%)	43 (59%)	39 (49%)	47 (60%)	33 (45%)	53 (63%)
Race (*n*, %)							
White	97 (62%)	53 (64%)	44 (59%)	51 (65%)	46 (59%)	49 (67%)	48 (57%)
Black/AA	60 (38%)	30 (36%)	30 (41%)	28 (35%)	32 (41%)	24 (33%)	36 (43%)
Stroke Type (*n*, %)							
Ischemic	132 (84%)	66 (80%)	66 (89%)	62 (78%)	70 (90%)	63 (86%)	69 (82%)
Hemorrhagic	13 (8.5%)	6 (7%)	7 (10%)	6 (8%)	7 (9%)	5 (7%)	8 (10%)
Not Specified	12 (7.5%)	11 (13%)	1 (1%)	11 (14%)	1 (1%)	5 (7%)	7 (8%)
Stroke Severity							
CGRP at 9 months	7.09 (*SD*=5.7)	6.16 (*SD*=5.0)	8.13 (*SD*=6.3)	6.18 (*SD*=5.1)	8.01 (*SD*=6.2)	2.44 (*SD*=1.7)	11.13 (*SD*=4.8)
GCS at discharge	4.38 (*SD*=0.7)	5.0 (*SD*=0.0)	3.68 (*SD*=0.5)	5.0 (*SD*=0.0)	3.74 (*SD*=0.5)	4.47 (*SD*=0.7)	4.30 (*SD*=0.8)
mRS at discharge	1.82 (*SD*=1.3)	0.72 (*SD*=0.6)	3.05 (*SD*=0.8)	0.66 (*SD*=0.5)	3.00 (*SD*=0.8)	1.56 (*SD*=1.3)	2.04 (*SD*=1.3)

GCS=Glasgow Coma Scale; mRS=modified Rankin Scale; CGRP=caregiver reported problems
Participants were divided into two groups for each predictor based on the 50th percentile (median) score as a cut-off point. The groups of Low GCS, High mRS, and Higher CGRP indicate worse stroke severity. Means and standard deviations are shown for continuous variables. Frequency of categorical variables are shown as a percentage for each group.

Table 2. Separate Mixed Models for Each Predictor by Cognition

Effects in Longitudinal Model	Estimate (b)	Standard Error	t	p
GCS	0.338	0.100	3.38	0.0009
GCS x Time	-0.098	0.066	-1.48	0.1408
mRS	-0.212	0.055	-3.89	0.0002
mRS x Time	0.066	0.037	1.80	0.0737
Caregiver-Reported Problems	-0.067	0.013	-5.22	<.0001
Caregiver-Report x Time	0.017	0.009	1.80	0.0742

GCS = Glasgow Coma Scale; mRS = modified Rankin Scale; Time = time of each assessment post-stroke in years

This table includes the results from separate mixed models for each predictor. These models are adjusted for the covariates of baseline cognition, time after stroke, age, sex, race, and education. Low GCS scores, high mRS scores, and higher numbers of Caregiver-Reported Problems indicate worse stroke severity

509
510

Figure 1. Covariate Adjusted, Logarithmic Trajectories of Cognitive Scores by GCS

511
512   *Figure 1.* The cutoff for 'less severe stroke' included any participants with GCS scores less than the
513   median value (5.0). Cutoff for 'more severe stroke' were scores greater than or equal to the median value.
514   This model includes baseline cognition, age at enrollment, gender, race, and education as covariates. As
515   this model uses logarithm of time, the first timepoint is estimated at 0.01 years (~4 days) post stroke.
516

517
518
519
520

Figure 2. Covariate Adjusted, Logarithmic Trajectories of Cognitive Scores by mRS

521
522   *Figure 2.* The cutoff for 'less severe stroke' included any participants with mRS scores less than the
523   median value (1.0). Cutoff for 'more severe stroke' was greater than the median value. This model
524   includes baseline cognition, age at enrollment, gender, race, and education as covariates. As this model
525   uses logarithm of time, the first timepoint is estimated at 0.01 years (~4 days) post stroke.
526
527

528
529
530
531

Figure 3. Covariate Adjusted, Logarithmic Trajectories of Cognitive Scores by Caregiver Report

532
533   *Figure 3.* The cutoff for 'fewer problems' included all participants whose total number of caregiver-
534   reported problems were less than the median value (6.0). Cutoff for 'more problems' include individuals
535   with reported problems greater than or equal to the median value. This model includes baseline cognition,
536   age at enrollment, gender, race, and education as covariates. As this model uses logarithm of time, the
537   first timepoint is estimated at 0.01 years (~4 days) post stroke.

Figure 1. Covariate Adjusted, Logarithmic Trajectories of Cognitive Scores by GCS

Figure 2. Covariate Adjusted, Logarithmic Trajectories of Cognitive Scores by mRS

Figure 3. Covariate Adjusted, Logarithmic Trajectories of Cognitive Scores by Caregiver Report