Reliably quantifying the severity of social symptoms in children with autism using ASDSpeech

Marina Eni[1,3]✉, Michal Ilan[2,3], Analya Michaelovski[3,4], Hava M. Golan[5,3], Gal Meiri[2,3], Idan Menashe[6,3], Ilan Dinstein[7,8,3], and Yaniv Zigel[1,3]

1. Department of Biomedical Engineering, Ben-Gurion University of the Negev, Beer-Sheva, Israel.
2. Pre-School Psychiatry Unit, Soroka University Medical Center, Beer-Sheva, Israel.
3. Azrieli National Centre for Autism and Neurodevelopment Research, Ben-Gurion University of the Negev, Beer-Sheva, Israel.
4. Zusman Child Development Center, Soroka University Medical Center, Israel.
5. Department of Physiology and Cell Biology, Faculty of Health Sciences, Ben-Gurion University of the Negev, Beer-Sheva, Israel.
6. Department of Public Health, Ben-Gurion University of the Negev, Beer-Sheva, Israel.
7. Department of Psychology, Ben-Gurion University of the Negev, Beer-Sheva, Israel.
8. Department of Brain & Cognitive Sciences, Ben-Gurion University of the Negev, Beer-Sheva, Israel.

✉email: marinamu@post.bgu.ac.il.

Abstract

Several studies have demonstrated that the severity of social communication problems, a core symptom of Autism Spectrum Disorder (ASD), is correlated with specific speech characteristics of ASD individuals. This suggests that it may be possible to develop speech analysis algorithms that can quantify ASD symptom severity from speech recordings in a direct and objective manner. Here we demonstrate the utility of a new open-source AI algorithm, ASDSpeech, which can analyze speech recordings of ASD children and reliably quantify their social communication difficulties across multiple developmental timepoints. The algorithm was trained and tested on the largest ASD speech dataset available to date, which contained 99,193 vocalizations from 197 ASD children recorded in 258 Autism Diagnostic Observation Schedule, 2nd edition (ADOS-2) assessments. ASDSpeech was trained with acoustic and conversational features extracted from the speech recordings of 136 children, who participated in a single ADOS-2 assessment, and tested with independent recordings of 61 additional children who completed two ADOS-2 assessments,

separated by 1–2 years. Estimated total ADOS-2 scores in the test set were significantly correlated
with actual scores when examining either the first ($r(59) = 0.544$, $P < 0.0001$) or second ($r(59) =$
0.605, $P < 0.0001$) assessment. Separate estimation of social communication and restricted and
repetitive behavior symptoms revealed that ASDSpeech was particularly accurate at estimating
social communication symptoms (i.e., ADOS-2 social affect scores). These results demonstrate
the potential utility of ASDSpeech for enhancing basic and clinical ASD research as well as
clinical management. We openly share both algorithm and speech feature dataset for use and
further development by the community.

Introduction

Autism Spectrum Disorder (ASD) is diagnosed by the presence of social communication
difficulties and the existence of Restricted and Repetitive Behaviors (RRBs)[1]. Most ASD children
exhibit language delays during early childhood[2], with 25–30% remaining minimally verbal (i.e.,
use < 50 words) throughout childhood[3]. However, core ASD symptoms are not necessarily evident
in the amount of speech produced by an individual and may instead be evident in the way they
speak. Some ASD children exhibit poorer fluency[4], echolalia (i.e., speech repetition)[5], mix
pronouns[6], and use atypical articulation and prosody[7,8] that are apparent in the acoustic features of
their vocalizations[9,10]. Studies have reported, for example, that verbal ASD children tend to speak
with higher pitch and larger pitch variability than typically developing (TD) children[8,9]. ASD
children also exhibit significantly fewer phoneme vocalizations[11], fewer conversational turns (i.e.,
reciprocating in a conversation)[11–13], more non-speech vocalizations[12,14], more distressed
vocalizations (crying, screaming)[15], and a lower ratio of syllables to vocalizations[16] than TD
children.

Several studies have used automated speech analysis techniques to classify ASD and TD
children based on extracted speech features[17–24]. In some studies, diagnostic classification was
based on linguistic features such as vocabulary and fluency[24] while in others it was based on
acoustic features such as pitch[18–20,22,23], jitter[20,23], shimmer[20,23], energy[18,19], Zero-Crossing Rate
(ZCR)[18,19], and Mel-Frequency Cepstral Coefficients (MFCCs)[19].

Three recent studies have extended this research by training machine and deep learning
algorithms to estimate ASD severity according to extracted speech features. In all these studies
ground truth was established by clinicians using the Autism Diagnostic Observation Schedule 2nd
edition (ADOS-2), a semi-structured assessment where clinicians score the behavior of children
during specific tasks/games[25]. The ADOS-2 yields a total severity score as well as separate Social
Affect (SA) and Restricted and Repetitive Behaviors (RRB) scores that quantify social difficulties
and RRB symptoms, respectively. In the first study, the authors extracted vocalization rates and
durations from speech recordings of 33 ASD children during an ADOS-2 assessment and reported
that a trained synthetic random forest model was able to accurately estimate their ADOS-2 Social

Affect (SA) scores[26]. Another study extracted hundreds of conversational, acoustic, and lexical
speech features from speech recordings of 88 adolescents and adults with ASD during an ADOS
assessment (1st edition) and reported that a trained Deep Neural Network (DNN) was able to
accurately estimate scores of four specific ADOS items that quantify the ability to maintain a
mature social conversation[27]. Finally, in the third study, from our group, we extracted acoustic
features such as pitch and energy, and conversational features such as turn-taking and speech rate
from speech recordings of 72 children (56 with ASD) during an ADOS-2 assessment[28]. We
demonstrated that a trained Convolutional Neural Network (CNN) model was able to accurately
estimate total ADOS-2 scores across multiple train-test subsamples.

While these results are encouraging, algorithms developed so far were trained and tested with
relatively small ASD samples that are not likely to represent the large heterogeneity of speech
styles and characteristics in the broad ASD population[29]. Moreover, previous studies examined
only a single timepoint of data from each participant, thereby limiting the ability to assess the
reliability of algorithms to assess ASD symptom severity at different developmental timepoints.
Previous studies also did not compare the ability of deep learning models to successfully estimate
the severity of social ASD symptoms versus RRB symptoms. Most importantly, previous studies
did not share their algorithms and data in a transparent manner that would enable re-production of
results and further development of algorithms by the research community.

To address these limitations, we created the largest speech recording dataset available to date,
which contained 99,193 vocalizations from 197 ASD children recorded in 258 ADOS-2
assessments, with 61 of the children participating in two ADOS-2 assessments that were separated
by 1-2 years. This comprehensive dataset enabled us to train and test the ASDSpeech algorithm
on different subsets of children and compare its accuracy across two developmental timepoints as
well as sex and age sub-groups. In addition, we also examined the ability to estimate ADOS-2 SA
versus RRB scores (i.e., social difficulties versus RRB symptoms). We intentionally used raw
ADOS-2 scores, which have a considerably wider range than ADOS-2 calibrated severity scores
[30,31], thereby increasing the potential sensitivity of the algorithm. Finally, we openly share the
algorithm and speech feature dataset to promote transparency and enable further use and
development by the research community.

Methods

Participants and setting. We analyzed data collected at the Azrieli National Centre for Autism
and Neurodevelopment Research (ANCAN), a collaboration between Ben-Gurion University of
the Negev (BGU) and eight partner clinical centers where ASD is diagnosed throughout Israel.
ANCAN manages the national autism database of Israel with data from > 3000 children in 2023
and growing[32,33]. All recordings used in the current study were performed in a single ANCAN
assessment room located at Soroka University Medical Center (SUMC), the largest partner clinical

site. A total of 197 children (1–7-years-old) who completed at least one ADOS-2 assessment between 2015 and 2021 and received an ASD diagnosis were included in this study (Table 1). Of the participating children, 136 completed a single ADOS-2 assessment and 61 completed two ADOS-2 assessments at two timepoints separated by 10–29 months, yielding 258 ADOS-2 assessments in total. All ADOS-2 assessments were performed by a clinician with research reliability. In addition, all participating children had ASD diagnoses that were confirmed by both a developmental psychologist and either a child psychiatrist or a pediatric neurologist, according to Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5) criteria[34]. Informed consent was obtained from all parents, and the study received approval from the SUMC Helsinki committee.

Table 1. Participating children's characteristics

	Single assessment	Two assessments	
		T1	T2
	(*N* = 136)	(*N* = 61)	(*N* = 61)
Mean (SD)			
Age (years)	4.26 (1.34)	3.67 (0.98)	4.95 (0.94)
ADOS-2 Total	14.99 (5.88)	14.92 (5.85)	14.92 (5.61)
ADOS-2 SA	10.75 (4.96)	11.26 (5.14)	10.57 (4.55)
ADOS-2 RRB	4.24 (1.88)	3.66 (1.66)	4.34 (1.87)
N (%)			
Sex			
Male	108 (79)	42 (69)	42 (69)
Female	28 (21)	19 (31)	19 (31)
Module			
Module T	17 (13)	9 (15)	0 (0)
Module 1	45 (33)	329 (48)	28 (46)
Module 2	40 (29)	21 (34)	19 (31)
Module 3	3 (25)	2 (3)	14 (23)

ADOS-2 assessments. ADOS-2 is a semi-structured behavioral assessment where a clinician administers specific tasks, observes the behavior of the child, and scores their behavior[35]. The total ADOS-2 score (range: 0–30) is the sum of the Social Affect (SA, range: 0–22) and Restricted and Repetitive Behavior (RRB, range: 0–8) scores, with higher scores indicating more severe symptoms.

Recording setup. All ADOS-2 recordings were performed using a single microphone (CHM99, AKG, Vienna) located on a wall, ~1–2m from the child, and connected to a sound card (US-16x08, TASCAM, California). Each ADOS-2 session lasted ~40-minutes (40.75 ± 11.95 min) and was recorded at a sampling rate of 44.1 kHz, 16 bits/sample (down-sampled to 16 kHz).

Detection of child vocalizations. We manually labeled segments with child vocalizations in each of the audio recordings. These segments included speech, laughing, moaning, crying, and screaming. The child segments often contained multiple vocalizations (e.g., multiple utterances) separated by silence. We separated each segment into multiple vocalizations using energy thresholds of 2.79dB and 0.4dB above the background noise to define the beginning and end of each vocalization, respectively[28] (Supplementary Figure S1). Vocalizations that were shorter than 110ms were excluded from further analysis (too short to contain an utterance).

Features. We extracted 49 speech features from the child vocalizations that were categorized into nine groups: pitch, formants, jitter, voicing, energy, Zero-Crossing Rate (ZCR), spectral slope, duration, and quantity/number of vocalizations. All features, except duration and quantity, were first extracted in 40ms windows (window overlap of 75%), resulting in a vector of feature values per vocalization. The minimum, the maximum, and the mean pitch of the voiced vocalizations (across windows) were computed, deriving one value for each vocalization. We then selected a group of 10 consecutive vocalizations and computed the mean and variance across vocalizations for relevant features (Supplementary Table S1). We also computed the mean duration of vocalizations and the overall number of vocalizations in the recording. Taken together, these steps yielded a vector with 49 values corresponding to the 49 features per 10 vocalizations. We performed this procedure 100 times, selecting random groups of ten consecutive vocalizations from the recording. Combining these 100 samples yielded a features matrix of 100×49 per child (Supplementary Figure S2), with the last column (quantity of vocalizations) containing the same value across all rows. Features included:

Frequency related features:

- *Pitch (F0):* Vocal cords vibration frequency (the fundamental frequency) that exists only in voiced speech (e.g., vowels). Voiced Vocalization (VV) was defined as a vocalization where most of its frames ($\geq 60\%$)[10] were voiced (voicing threshold 0.45).

- *Formants:* The resonant frequencies of the vocal tract that shape vowel sounds[36]. The first two formants (F1 and F2) relate to tongue position (vertical and horizontal) and influence vowel quality. Their bandwidths affect the clarity of speech.

- *Jitter:* Variation across adjacent pitch values representing frequency instability[37].

- *Voicing*: Pitch peak amplitude as determined by the autocorrelation function.

Pitch and formants were calculated using the PRAAT software[38], with a pitch range set to 60–1600 Hz (a wide range to increase sensitivity to atypical vocal characteristics).

Energy/amplitude related features:

- *Energy:* The energy ratio between each child's vocalization and the background noise. The background noise energy was calculated from the energy values extracted from the lowest 5% of the recording's frames.

Spectral features:

- *Zero-Crossing Rate (ZCR):* The number of zero-crossings apparent in audio segments with child vocalizations[39].

- *Spectral slope:* The slope of the linear regression on the logarithmic power spectrum within the frequency bands of 20–500 Hz (lower band) and 500–1500 Hz (higher band)[40,41].

Conversational features:

- *Duration:* Child's mean vocalization length.

- *Quantity*: The total number of vocalizations.

All features, except for Pitch and Formants, were extracted with custom-written code in Matlab (Mathworks, Inc.).

Training and testing ASDSpeech. Training was performed with data from the 136 children who completed a single ADOS-2 session only. Feature matrices were used to train two deep learning models with an identical CNN architecture (Supplementary Figure S3). The first model estimated ADOS-2 SA scores and the second estimated ADOS-2 RRB scores. Training was based on minimizing the Mean Squares Error (MSE) of a regression analysis between estimated and

actual scores, using the RMSprop (Root Mean Square Propagation) as the optimization
algorithm[42]. The training process was preformed 25 times, creating 25 different SA and 25 RRB
models that were trained with different combinations of training data sub-samples and learning
parameters. We considered this analogous to having 25 clinicians, each with a different learning
style and different clinical experience. First, we performed the feature extraction procedure
described above 5 times for each child. Since feature extraction included a random selection of
consecutive vocalizations, this resulted in 5 different sub-samples of the data. When training each
model (separately for SA and RRB) we split the training data into a training-set (80%) and
validation set (20%) and applied a random search algorithm to optimize the following learning
parameters: batch size, number of epochs, and learning rate, while applying early stopping of
patience after 20 epochs to reduce overfitting[43]. Optimal learning hyper-parameters were selected
based on the highest concordance correlation coefficient[44], between estimated and actual ADOS-
2 scores in the training and validation sets respectively. This procedure was performed 5 times
using different selections of validation data (i.e., 5-fold cross validation), yielding 5 models with
different learning parameters per data sub-sample and 25 models in total for SA and RRB scores
separately.

Testing was performed with an entirely independent dataset of 61 ASD children who
completed two ADOS-2 assessments. For each of these children we estimated a separate SA and
RRB score from each of the 25 models described above and then computed their mean, yielding a
single SA and RRB score per child. This is analogous to a clinical consensus across the 25 models.
Accuracy of ASDSpeech estimation was measured using Pearson correlation and NRMSE (RMSE
/ (y_{max} - y_{min}), where y is the actual ADOS-2 score), which were calculated between the estimated
and actual ADOS-2 scores in the testing dataset, separately for the first and second ADOS-2
assessments (i.e., T1 and T2).

Hardware. All model training, optimization, and training were performed using custom-written
code in Python 3.9.13 using a Keras API 2.6.0 with TensorFlow (version 2.6.0) backend. The
training was conducted on an IntelI XI(R) Gold 6140 CPU @ 2.30GHz and NVIDIA GPU Tesla
T4.

Statistical Analysis. All statistical analyses were conducted using custom-written code in
Python. Associations between speech features and ADOS-2 scores were assessed using Pearson
correlations. To evaluate their statistical significance, we performed a random permutation test. In
this test, we randomly shuffled the actual ADOS-2 scores across children and calculated the
correlation between each feature and the shuffled scores. This randomization procedure was
performed 1,000 times, generating a null distribution of random correlation values as computed
from the original data that is not necessarily normally distributed as assumed by parametric
statistical tests. For a correlation between a speech feature and ADOS-2 score to be considered
significant, the actual correlation value had to be higher than the 97.5 percentile of the null

distribution. We used an equivalent analysis to assess the statistical significance of correlations
between actual and ASDSpeech estimated ADOS-2 scores. We also performed a similar analysis
with NRMSE values, where we assessed whether the actual NRMSE value was smaller than the
2.5 percentile of the null distribution. This statistical test, therefore, assessed whether correlation
values were higher than expected by chance and NRMSE values were lower than expected by
chance.

Data sharing. The ASDSpeech algorithm source-code and associated dataset are available at
https://github.com/Dinstein-Lab/ASDSpeech.

222 **Results**

Using the data from the 136 ASD children in the training dataset, we examined the relationships
between each of the 49 features and ASD symptom severity as defined clinically by the children's
ADOS-2 scores. Thirty-one features exhibited significant Pearson correlation coefficients with
total ADOS-2 scores (i.e., sum of SA and RRB scores), 31 with ADOS-2 SA scores, and 28 with
ADOS-2 RRB scores (Figure 1). While some features, such as the number of vocalizations,
exhibited a stronger correlation with SA than RRB score, others, such as mean jitter, exhibited the
opposite (Supplementary Figure S4). Hence, different features seem to carry distinct information
regarding each of the two core ASD symptoms, demonstrating the potential opportunity for a deep
learning algorithm to learn relevant associations.

Figure 1. Pearson correlation coefficients between each of the extracted features and ADOS-2 scores from the 136 children in the training dataset. Correlation coefficients are presented for total ADOS-2 scores (**a**), ADOS-2 SA scores (**b**), and ADOS-2 RRB scores (**c**). Each color represents a different group of features. **Asterisks**: significant Pearson correlation (∗ < 0.05, ∗∗ ≤ 0.01, ∗3 ≤ 0.001, ∗4 ≤ 0.0001).

Longitudinal stability of ADOS-2 scores

The 61 ASD children in the test dataset exhibited similar ADOS-2 scores across their two assessments, which were separated by 1-2 years, indicating overall stability in severity over time. Significant correlations were apparent across first and second assessments for ADOS-2 total ($r(59)$ = 0.743, $P < 0.001$), ADOS-2 SA ($r(59)$ = 0.666, $P < 0.001$), and ADOS-2 RRB ($r(59)$ = 0.5, $P < 0.001$) scores (Figure 2).

Figure 2. **Scatter plots demonstrating overall stability in ADOS-2 scores across first and second assessments (T1 and T2).** (**a**) ADOS-2 SA scores. (**b**) ADOS-2 RRB scores. (**c**) Total ADOS-2 scores (sum of SA and RRB scores). **Asterisk**: statistical significance of the Pearson correlation coefficient ($P < 0.0001$). **Shaded areas**: 95% confidence intervals. Children located below the diagonal (dashed line) exhibited lower ASD severity at T2 (improvement), while children above the diagonal exhibited the opposite.

Training and testing the ASDSpeech algorithm

We trained the ASDSpeech algorithm with data from 136 ASD children in the training dataset. The algorithm included two separate CNN models that were trained to estimate ADOS-2 SA and RRB scores independently, given that different speech features were associated with each symptom domain. The accuracy of the algorithm was tested with data from two independent ADOS-2 recordings of the 61 children in the testing dataset where ASDSpeech estimated the SA, RRB, and total ADOS-2 (sum of SA and RRB) scores of each child per recording (Figure 3).

Estimated total ADOS-2 scores were significantly correlated with actual scores at T1 ($r(59) = 0.544$, $P < 0.0001$) and T2 ($r(59) = 0.605$, $P < 0.0001$). Similarly, estimated ADOS-2 SA scores were significantly correlated with actual scores at T1 ($r(59) = 0.502$, $P < 0.0001$) and T2 ($r(59) = 0.592$, $P < 0.0001$). In contrast, estimated ADOS-2 RRB scores were not significantly correlated with actual RRB scores at T1 ($r(59) = 0.093$, $P = 0.474$), exhibiting significant correlations only at T2 ($r(59) = 0.332$ $P = 0.009$) with a relatively weaker effect size.

Normalized Root Mean Squared Error (NRMSE) between estimated and actual total ADOS-2 scores was significantly smaller than expected by chance when computed at T1 (NRMSE = 0.189, $P < 0.0001$) and T2 (NRMSE = 0.172, $P = 0.0001$). Similarly, NRMSE between estimated and actual ADOS-2 SA scores was significantly smaller than expected by chance when computed at T1 (NRMSE = 0.200, $P < 0.0001$) and T2 (NRMSE = 0.170, $P < 0.0001$). In contrast, NRMSE between estimated and actual ADOS-2 RRB scores was not significantly smaller than expected by

chance at T1 (NRMSE = 0.219, P = 0.460), exhibiting significant results only at T2 (NRMSE =
0.225, P = 0.006).

The statistical significance of the NRMSE results was determined with a randomization
analysis where we randomly shuffled ADOS-2 scores across children before computing NRMSE
values. We computed 1,000 random permutations to generate a null NRMSE distribution and
assessed statistical significance by determining whether the actual NRMSE value was smaller than
the 2.5 percentile of the null distribution (see Methods).

Figure 3. Accuracy of ASDSpeech. Scatter plots demonstrating the fit between estimated and
actual scores for the children at T1 (left column) and T2 (right column). **(a-b)** Total ADOS-2
scores (sum of SA and RRB scores). **(c-d)** ADOS-2 SA scores. **(e-f)** ADOS-2 RRB scores. Pearson
correlation coefficients and NRMSE values are noted in each panel. **Solid line:** Linear fit. **Dashed**

line: diagonal (unity line). **Asterisks:** statistical significance as determined by randomization test
($P < 0.05$).

Differences across age and sex subgroups
Next, we examined whether ASDSpeech accuracy differed across age and sex subgroups
(Figure 4). Estimated total ADOS-2 scores were significantly correlated with actual scores when
examining children above the median age at T1 ($r(28) = 0.604$, $P = 0.0004$) or T2 ($r(25) = 0.612$,
$P = 0.0007$) and children below the median age at T1 ($r(29) = 0.485$, $P = 0.006$) or T2 ($r(32) =$
0.657, $P < 0.0001$). There were no significant differences in the algorithm's accuracy between
younger and older children at T1 ($P = 0.540$) or T2 ($P = 0.780$) as tested with a randomization
analysis. Similarly, estimated total ADOS-2 scores were significantly correlated with actual scores
when examining males at T1: ($r(40) = 0.631$, $P < 0.0001$) or T2 ($r(40) = 0.601$, $P < 0.0001$).
Estimated ADOS-2 scores were also significantly correlated with actual scores when examining
females at T2 ($r(17) = 0.627$, $P = 0.004$), but the correlation did not reach statistical significance
at T1 ($r(17) = 0.363$, $P = 0.127$). Nevertheless, there were no significant differences in the
algorithm's accuracy between males and females at T1 ($P = 0.198$) or T2 ($P = 0.930$) as tested
with a randomization analysis.

Comparison of NRMSE across subgroups showed similar results. NRMSE between the
estimated and actual ADOS-2 scores was significantly smaller than expected by chance when
examining younger children at T1 (NRMSE = 0.199, $P = 0.008$) or T2 (NRMSE = 0.173, $P <$
0.0001) as well as older children at T1 (NRMSE = 0.178, $P < 0.0001$) or T2 (NRMSE = 0.172, P
< 0.0001). There were no significant differences in the algorithm's accuracy between younger and
older children at T1 ($P = 0.434$) or T2 ($P = 0.992$). NRMSE were also significantly smaller than
expected by chance when examining males at T1 (NRMSE = 0.172, $P < 0.0001$) or T2 (NRMSE
$= 0.166$, $P < 0.0001$). For females this was the case only at T2 (NRMSE = 0.186, $P = 0.006$) and
not at T1 (NRMSE = 0.222, $P = 0.140$). Nevertheless, there were no significant differences in the
algorithm's accuracy between males and females at T1 ($P = 0.094$) or T2 ($P = 0.588$) as tested
with a randomization test.

Figure 4. ASDSpeech accuracy as a function of sex and age at T1 and T2. (**a**,**b**) Pearson correlation values (**c**,**d**) Normalized Root Mean Squared Error (NRMSE) values. (**a**, **c**) comparison between younger and older children (median split according to age at each timepoint). (**b**, **d**) comparison between males and females. **Asterisks:** statistical significance as determined by randomization test ($P < 0.05$).

Discussion

Our results demonstrate the ability of ASDSpeech to quantify the severity of social symptoms in ASD children from recordings of their speech during ADOS-2 assessments. The algorithm, trained with recordings from 136 ASD children, was able to accurately estimate total ADOS-2 and ADOS-2 SA scores in an entirely independent sample of 61 ASD children, who were recorded at two different developmental timepoints separated by 1-2 years (Figure 3). It is remarkable that ASDSpeech was able to achieve this despite the large heterogeneity in language fluency and speech articulation abilities apparent across ASD children[45] as well as the large developmental changes that take place in speech abilities during the examined period of early childhood[46]. Moreover, the robust accuracy of ASDSpeech in estimating ADOS-2 SA scores is remarkable

given that the social difficulties assessed during the ADOS-2 assessment manifest themselves in
behaviors that have little to do with speech including difficulties with eye contact, imitation, joint
attention, and other social behaviors[3,47]. This suggests that combining ASDSpeech with analysis
of eye tracking[48–50], facial expressions[51], and body movement[52] data from the same children will
enable even higher accuracy and reliability in estimating ASD symptoms.

Separate estimation of social and RRB symptoms demonstrated that ASDSpeech was
considerably more accurate at estimating social ASD symptoms captured by the ADOS-2 SA
scores in contrast to the RRB symptoms captured by the ADOS-2 RRB scores (Figure 3). Note
that accurate estimation of total ADOS-2 scores (Figure 3) was likely based on the accurate
estimation of SA scores that account for two-thirds of the total scores. We believe there may be
several reasons for the more accurate estimation of SA scores. First, the limited range of the
ADOS-2 RRB scale (0–8) relative to the SA scale (0–22) may make it difficult for the algorithm
to identify differences across children. Indeed, a recent study reported that the limited number of
items on the RRB scale resulted in poor scale reliability across participants[53]. Second, the selected
speech features in the current study exhibited weaker correlations with RRB than SA scores
(Figure 1). Extraction of additional speech features, such as phrase or intonation repetitions
(indicative of echolalia) may improve the accuracy of ADOS-2 RRB score estimates. Regardless,
our results motivate separate modeling of social and RRB symptom domains as each of them is
likely associated with distinct features of speech.

In the current study we estimated raw ADOS-2 scores rather than calibrated severity scores
(CSS), which were developed to standardize ASD symptom severity measurements across
different ages and language abilities[30,31]. While ADOS-2 CSS are important for longitudinal
assessments of coarse changes in severity[54,55], their restricted scoring range (children with ASD
receive scores of 4-10) limits the sensitivity of deep learning algorithms in identifying differences
across children. By demonstrating that ASDSpeech achieves robust accuracy in estimating raw
ADOS-2 SA scores across different age groups and developmental timepoints we show that
severity estimations are independent of these factors, thereby justifying the use of raw scores.

Diagnostic classification with speech analysis algorithms

A variety of previous studies have reported that individuals with ASD, on average, speak
differently than TD individuals[4,8–16]. According to these studies, ASD individuals exhibit atypical
speech characteristics, including significantly fewer phonemes per utterance[11], fewer
conversational turns[13], higher pitch[9,19], and larger pitch range and variability[8,9] than TD children.
Differences in these and other speech characteristics have enabled the development of machine
and deep learning classification algorithms that can identify ASD and TD individuals with reported
accuracy rates of 75-98%[17–23].

However, these relatively high classification accuracies are likely to be inflated due to the
small sample size of most studies (<40 ASD participants) that are not likely to capture the true
heterogeneity of ASD symptoms or speech styles/characteristics of the broad ASD population.
Indeed, even "gold standard" clinical tests such as the ADOS-2, exhibit ~80% accuracy in
identifying children who will eventually receive an ASD diagnosis[56]. This is because establishing
an ASD diagnosis requires clinicians to incorporate additional information from parent interviews
and other clinical assessments[57]. Clinicians also report high diagnosis certainty in only ~70% of
ASD children because the presentation of ASD symptoms is equivocal in ~30% of cases[58]. These
studies suggest an expected upper limit of 70–80% accuracy when attempting to identify ASD
using digital phenotyping techniques such as speech analysis. Nevertheless, it is highly
encouraging that speech features contain information enabling the separation of ASD and TD
children.

Quantifying ASD severity with speech analysis algorithms

A more complex task is to develop machine and deep learning algorithms that can quantify the
severity core ASD symptoms. Results presented in the current and previous study from our lab[28]
demonstrated that multiple speech features were significantly correlated with SA and/or RRB
ADOS-2 scores (Figure 1), suggesting that distinct combinations of speech features are associated
with each of the two core ASD symptoms.

Three recent studies have attempted to use these relationships to predict ADOS-2 scores by
analyzing speech recordings of ASD individuals[26–28]. The first trained a synthetic random forest
model to estimate ADOS-2 SA scores according to vocalization rate and turn-taking features
extracted from ADOS-2 recordings of 33 ASD children. The algorithm was able to predict ADOS-
2 SA scores that were significantly correlated with actual scores ($r = 0.634$). The second study
utilized a DNN model to estimate four ADOS (first edition) item scores using hundreds of
conversational and acoustic features extracted from speech recordings of 88 high-functioning ASD
adolescents/adults during an ADOS assessment[27]. This algorithm was able to estimate scores that
exhibited significant Spearman correlations with the actual scores ($\rho = 0.519–0.645$). Finally, in a
previous study from our lab[28], we demonstrated that a CNN model was able to estimate ADOS-2
total scores that were significantly correlated with actual scores ($r = 0.718$) when using 60
conversational and acoustic features extracted from speech recordings of 72 children (56 of them
with ASD) during ADOS-2 assessment.

The current study extends previous work in several critical ways. First, we utilized a
considerably larger dataset (258 ADOS-2 recordings) that was at least three times larger than the
ones used to date. This was important for training ASDSpeech with speech recordings form a large
cohort with heterogeneous language abilities. Second, the 61 ASD children in our testing dataset
were recorded twice during two ADOS-2 assessments separated by 1–2 years. This enabled us to
test the robustness of ASDSpeech across two developmental timepoints. Third, we trained

ASDSpeech to estimate ADOS-2 SA and ADOS-2 RRB scores using separate CNN models. The
results demonstrated that this separation was critical with accurate performance apparent primarily
for the ADOS-2 SA scores. Fourth, the large sample size enabled us to demonstrate that
ASDSpeech accuracy was similar across age and sex subgroups. Fifth, the recordings utilized in
the current study were performed over a 6-year period in a busy public healthcare medical center
that services a population of ~1 million people. Recordings were performed with a wall mounted
microphone (see Methods) in "real world" noisy conditions (e.g., announcement system in the
hallway). This demonstrates the robustness of ASDSpeech to variable recording conditions.

ASDSpeech achieved similar accuracy to that reported in previous studies. The important
advance in the current study is in demonstrating that this accuracy is robust to age and
developmental stage of the examined children when examining a large heterogeneous population
within an active clinical setting. Most importantly, we openly share ASDSpeech and its associated
dataset with the research community.

Limitations

The current study had several limitations. First, we did not examine the language content of the
recordings, which is likely to improve the estimation of ASD symptom severity[4,24]. Second, we
did not identify echolalia, crying, or shouting events that are likely to be informative of RRB
symptoms. Indeed, our weaker results estimating RRB scores suggest that different speech features
are necessary for estimating severity in this domain. Third, we did not apply any noise reduction
or multi-speaker analysis techniques to improve the quality of the analyzed vocal segments.
Finally, our sample had a 4:1 male to female ratio, which is equivalent to the sex ratio in the
national ASD population of Israel[59]. Hence, higher ASDSpeech accuracy for males at T1 may be
due to the larger number of males in the training and testing datasets. This could be rectified by
future studies.

Conclusions

This study adds to accumulating evidence demonstrating that speech recordings contain reliable
information about the social symptom severity of ASD children. We demonstrate the ability of the
ASDSpeech algorithm to quantify these symptoms in a robust manner across two developmental
timepoints with recordings that were performed within a busy community healthcare center. We
openly share the algorithm and its associated dataset for further use, testing, and development by
the research community and are confident that future versions of the algorithm will achieve even
higher and more robust accuracy rates, yielding a transformative new tool for clinical and basic
ASD research.

References

Lord C, Elsabbagh M, Baird G, Veenstra-Vanderweele J. Autism spectrum disorder. *The Lancet* 2018; **392**: 508–520.

Gabbay-Dizdar N, Ilan M, Meiri G, Faroy M, Michaelovski A, Flusser H *et al.* Early diagnosis of autism in the community is associated with marked improvement in social symptoms within 1–2 years. *Autism* 2022; **26**: 1353–1363.

Tager-Flusberg H, Kasari C. Minimally Verbal School-Aged Children with Autism Spectrum Disorder: The Neglected End of the Spectrum. *Autism Research* 2013; **6**: 468–478.

Salem AC, MacFarlane H, Adams JR, Lawley GO, Dolata JK, Bedrick S *et al.* Evaluating atypical language in autism using automated language measures. *Sci Rep* 2021; **11**. doi:10.1038/S41598-021-90304-5.

Chi NA, Washington P, Kline A, Husic A, Hou C, He C *et al.* Classifying Autism From Crowdsourced Semistructured Speech Recordings: Machine Learning Model Comparison Study. *JMIR Pediatr Parent* 2022; **5**: e35406.

Mostek J. Cognitive Development and Language Acquisition in Autistic Children. *Science Insights* 2022; **41**: 719–724.

Loukusa S. Autism Spectrum Disorder. In: Cummings L (ed). *Handbook of Pragmatic Language Disorders*. Springer International Publishing: Cham, 2021, pp 45–78.

Bonneh YS, Levanon Y, Dean-Pardo O, Lossos L, Adini Y. Abnormal Speech Spectrum and Increased Pitch Variability in Young Autistic Children. *Front Hum Neurosci* 2011; **4**: 237.

Asghari SZ, Farashi S, Bashirian S, Jenabi E. Distinctive prosodic features of people with autism spectrum disorder: a systematic review and meta-analysis study. *Sci Rep* 2021; **11**: 23093.

Oller DK, Niyogi P, Gray S, Richards JA, Gilkerson J, Xu D *et al.* Automated vocal analysis of naturalistic recordings from children with autism, language delay, and typical development. *Proceedings of the National Academy of Sciences* 2010; **107**: 13354–13359.

Moffitt JM, Ahn YA, Custode S, Tao Y, Mathew E, Parlade M *et al.* Objective measurement of vocalizations in the assessment of autism spectrum disorder symptoms in preschool age children. *Autism Research* 2022; : 1–10.

Ferguson EF, Nahmias AS, Crabbe S, Liu T, Mandell DS, Parish-Morris J. Social language opportunities for preschoolers with autism: Insights from audio recordings in urban classrooms. *Autism* 2020; **24**: 1232–1245.

13   Warren SF, Gilkerson J, Richards JA, Oller DK, Xu D, Yapanel U *et al.* What Automated
Vocal Analysis Reveals About the Vocal Production and Language Learning Environment
of Young Children with Autism. *J Autism Dev Disord* 2010; **40**: 555–569.

14   Warlaumont AS, Richards JA, Gilkerson J, Oller DK. A Social Feedback Loop for Speech
Development and Its Reduction in Autism. *Psychol Sci* 2014; **25**: 1314–1324.

15   Plumb AM, Wetherby AM. Vocalization Development in Toddlers With Autism Spectrum
Disorder. *Journal of Speech, Language, and Hearing Research* 2013; **56**: 721–734.

16   Tenenbaum EJ, Carpenter KL, Sabatos-DeVito M, Hashemi J, Vermeer S, Sapiro G *et al.*
A Six-Minute Measure of Vocalizations in Toddlers with Autism Spectrum Disorder.
Autism Research 2020; **13**: 1373–1382.

17   Pokorny FB, Schuller BW, Marschik PB, Brueckner R, Nyström P, Cummins N *et al.*
Earlier identification of children with autism spectrum disorder: An automatic vocalisation-
based approach. *Proceedings of the Annual Conference of the International Speech
Communication Association, INTERSPEECH* 2017; **2017-Augus**: 309–313.

18   Mohanta A, Mittal VK. Classifying Speech of ASD Affected and Normal Children Using
Acoustic Features. In: *2020 National Conference on Communications (NCC)*. IEEE, 2020,
pp 1–6.

19   Mohanta A, Mittal VK. Analysis and classification of speech sounds of children with autism
spectrum disorder using acoustic features. *Comput Speech Lang* 2022; **72**: 101287.

20   Asgari M, Shafran I. Robust and Accurate Features for Detecting and Diagnosing Autism
Spectrum Disorders. *Proceedings of the Annual Conference of the International Speech
Communication Association, INTERSPEECH* 2013; : 191–194.

21   Yankowitz LD, Schultz RT, Parish-Morris J. Pre- and Paralinguistic Vocal Production in
ASD: Birth Through School Age. *Curr Psychiatry Rep* 2019; **21**: 126.

22   Lee S, Yeo EJ, Kim S, Chung M. Knowledge-driven speech features for detection of
Korean-speaking children with autism spectrum disorder*. *Phonetics and Speech Sciences*
2023; **15**: 53–59.

23   Briend F, David C, Silleresi S, Malvy J, Ferré S, Latinus M. Voice acoustics allow
classifying autism spectrum disorder with high accuracy. *Transl Psychiatry* 2023; **13**: 250.

24   MacFarlane H, Salem AC, Chen L, Asgari M, Fombonne E. Combining voice and language
features improves automated autism detection. *Autism Res* 2022; **15**: 1288–1300.

25   Lord C, Rutter M, Di Lavore P, Risi S, Gotham K, Bishop S. Autism and Diagnostic
Observation Schedule, Second Edition (ADOS-2) Manual (Part I): Modules 1-4. 2012.

26 Sadiq S, Castellanos M, Moffitt J, Shyu M, Perry L, Messinger D. Deep Learning Based
Multimedia Data Mining for Autism Spectrum Disorder (ASD) Diagnosis. In: *2019*
International Conference on Data Mining Workshops (ICDMW). 2019, pp 847–854.

27 Chen C-P, Gau SS-F, Lee C-C. Learning Converse-Level Multimodal Embedding to Assess
Social Deficit Severity for Autism Spectrum Disorder. In: *2020 IEEE International*
Conference on Multimedia and Expo (ICME). IEEE, 2020, pp 1–6.

28 Eni M, Dinstein I, Ilan M, Menashe I, Meiri G, Zigel Y. Estimating Autism Severity in
Young Children From Speech Signals Using a Deep Neural Network. *IEEE Access* 2020;
8: 139489–139500.

29 Asghari SZ, Farashi S, Bashirian S, Jenabi E. Distinctive prosodic features of people with
autism spectrum disorder: a systematic review and meta-analysis study. *Sci Rep* 2021; **11**:
23093.

30 Esler AN, Bal VH, Guthrie W, Wetherby A, Weismer SE, Lord C. The Autism Diagnostic
Observation Schedule, Toddler Module: Standardized Severity Scores. *J Autism Dev Disord*
2015; **45**: 2704–2720.

31 Gotham K, Pickles A, Lord C. Standardizing ADOS scores for a measure of severity in
autism spectrum disorders. *J Autism Dev Disord* 2009; **39**: 693–705.

32 Dinstein I, Arazi A, Golan HM, Koller J, Elliott E, Gozes I *et al.* The National Autism
Database of Israel: a Resource for Studying Autism Risk Factors, Biomarkers, Outcome
Measures, and Treatment Efficacy. *Journal of Molecular Neuroscience* 2020; **70**: 1303–
1312.

33 Meiri G, Dinstein I, Michaelowski A, Flusser H, Ilan M, Faroy M *et al.* The Negev Hospital-
University-Based (HUB) Autism Database. *J Autism Dev Disord* 2017; **47**: 2918–2926.

34 American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*.
5th ed. American Psychiatric Publishing: Arlington, VA, 2013
doi:10.1176/appi.books.9780890425596.744053.

35 Lord C, Rutter M, Di Lavore P, Risi S, Gotham K, Bishop S. Autism and Diagnostic
Observation Schedule, Second Edition (ADOS-2) Manual (Part I): Modules 1-4. 2012.

36 Padmini P, Gupta D, Zakariah M, Alotaibi YA, Bhowmick K. A Simple Speech Production
System Based on Formant Estimation of a Tongue Articulatory System Using Human
Tongue Orientation. *IEEE Access* 2021; **9**: 4688–4710.

37 Rusz J, Benova B, Ruzickova H, Novotny M, Tykalova T, Hlavnicka J *et al.* Characteristics
of motor speech phenotypes in multiple sclerosis. *Mult Scler Relat Disord* 2018; **19**: 62–69.

Boersma P, van Heuven V. Speak and unSpeak with Praat. *Glot International* 2001; **5**: 341–347.

G Pillai L, Sherly E. A Deep Learning Based Evaluation of Articulation Disorder and Learning Assistive System for Autistic Children. *International Journal on Natural Language Computing* 2017; **6**: 19–36.

Tamarit L, Goudbeek M, Scherer K. Spectral Slope Measurements in Emotionally Expressive Speech. In: *Proceedings of ISCA Tutorial and Research Workshop (ITRW) on Speech Analysis and Processing for Knowledge Discovery*. 2008, pp 1–4, paper 007.

Eyben F, Scherer KR, Schuller BW, Sundberg J, Andre E, Busso C *et al.* The Geneva Minimalistic Acoustic Parameter Set (GeMAPS) for Voice Research and Affective Computing. *IEEE Trans Affect Comput* 2016; **7**: 190–202.

Taqi AM, Awad A, Al-Azzo F, Milanova M. The Impact of Multi-Optimizers and Data Augmentation on TensorFlow Convolutional Neural Network Performance. *Proceedings - IEEE 1st Conference on Multimedia Information Processing and Retrieval, MIPR 2018* 2018; **April**: 140–145.

Prechelt L. Early Stopping — But When? In: *Neural Networks: Tricks of the Trade*. 2012, pp 53–67.

MacFarlane H, Salem AC, Bedrick S, Dolata JK, Wiedrick J, Lawley GO *et al.* Consistency and reliability of automated language measures across expressive language samples in autism. *Autism Research* 2023; **16**: 802–816.

Tager-Flusberg H. Defining language phenotypes in autism. *Clin Neurosci Res* 2006; **6**: 219–224.

Tager-Flusberg H, Rogers S, Cooper J, Landa R, Lord C, Paul R *et al.* Defining Spoken Language Benchmarks and Selecting Measures of Expressive Language Development for Young Children With Autism Spectrum Disorders. *Journal of Speech, Language, and Hearing Research* 2009; **52**: 643–652.

Vivanti G, Bottema-Beutel K, Turner-Brown L. Understanding and Addressing Restricted and Repetitive Behaviors in Children with Autism. In: *Clinical Guide to Early Interventions for Children with Autism*. 2020, pp 61–77.

Avni I, Meiri G, Bar-Sinai A, Reboh D, Manelis L, Flusser H *et al.* Children with autism observe social interactions in an idiosyncratic manner. *Autism Research* 2020; **13**: 935–946.

Chong E, Clark-Whitney E, Southerland A, Stubbs E, Miller C, Ajodan EL *et al.* Detection of eye contact with deep neural networks is as accurate as human experts. *Nat Commun* 2020; **11**: 6386.

50 Jones W, Klaiman C, Richardson S, Aoki C, Smith C, Minjarez M *et al.* Eye-Tracking-
Based Measurement of Social Visual Engagement Compared With Expert Clinical
Diagnosis of Autism. *JAMA* 2023; **330**: 854–865.

51 Perochon S, Di Martino JM, Carpenter KLH, Compton S, Davis N, Eichner B *et al.* Early
detection of autism using digital behavioral phenotyping. *Nat Med* 2023; **29**: 2489–2497.

52 Budman I, Meiri G, Ilan M, Faroy M, Langer A, Reboh D *et al.* Quantifying the social
symptoms of autism using motion capture. *Sci Rep* 2019; **9**: 7712.

53 Frazier TW, Whitehouse AJO, Leekam SR, Carrington SJ, Alvares GA, Evans DW *et al.*
Reliability of the Commonly Used and Newly-Developed Autism Measures. *J Autism Dev*
Disord 2023. doi:10.1007/s10803-023-05967-y.

54 Gabbay-Dizdar N, Ilan M, Meiri G, Faroy M, Michaelovski A, Flusser H *et al.* Early
diagnosis of autism in the community is associated with marked improvement in social
symptoms within 1–2 years. *Autism* 2022; **26**: 1353–1363.

55 Waizbard-Bartov E, Ferrer E, Young GS, Heath B, Rogers S, Wu Nordahl C *et al.*
Trajectories of Autism Symptom Severity Change During Early Childhood. *J Autism Dev*
Disord 2021; **51**: 227–242.

56 Maddox BB, Brodkin ES, Calkins ME, Shea K, Mullan K, Hostager J *et al.* The Accuracy
of the ADOS-2 in Identifying Autism among Adults with Complex Psychiatric Conditions.
J Autism Dev Disord 2017; **47**: 2703.

57 Bishop SL, Lord C. Commentary: Best practices and processes for assessment of autism
spectrum disorder - the intended role of standardized diagnostic instruments. *J Child*
Psychol Psychiatry 2023; **64**. doi:10.1111/JCPP.13802.

58 Klaiman C, White S, Richardson S, McQueen E, Walum H, Aoki C *et al.* Expert Clinician
Certainty in Diagnosing Autism Spectrum Disorder in 16-30-Month-Olds: A Multi-site
Trial Secondary Analysis. *J Autism Dev Disord* 2022. doi:10.1007/S10803-022-05812-8.

59 Raz R, Weisskopf MG, Davidovitch M, Pinto O, Levine H. Differences in autism spectrum
disorders incidence by sub-populations in Israel 1992-2009: a total population study. *J*
Autism Dev Disord 2015; **45**: 1062–1069.

Acknowledgments

598 This study was supported by the Israeli Science Foundation (Grant no. 1150/20) and the Israel
599 Ministry of Science & Technology (Grant no. 3-17422).

Author contributions

M.E. collected the data, performed the experiments, built the models, analyzed the data, and wrote the paper. I.D., and Y.Z. designed the study, guided data collection and analysis, and wrote the paper. M.I., A.M., H.M.G., G.M., and I.M. contributed to participant recruitment, behavioral assessments, data collection, and interpretation of the findings. All authors approved the final manuscript.

Additional information

Competing interests. The authors declare no competing interests.