Costs, catastrophic out-of-pocket payments and impoverishment related to accessing surgical

care among households in rural Ethiopia

Short title: Economic impact of accessing surgical care in Ethiopia

Yohannes Hailemichael[1], Tigist Eshetu[2], Hanna Negussie[2], Andualem Deneke[3], Amezene

Tadesse[3], Ahmed Abdella[4], Abebe Bekele[3], Andrew Leather[5], Girmay Medhin[6], Martin Prince[7],

Charlotte Hanlon[8,2]*

Affiliations

[1]Armauer Hansen Research Institute, Addis Ababa, Ethiopia;

[2]Department of Psychiatry, School of Medicine, College of Health Sciences, Addis Ababa University,
Addis Ababa, Ethiopia;

[3]Department of Surgery, School of Medicine, College of Health Sciences, Addis Ababa University, Addis
Ababa, Ethiopia;

[4]Department of Obstetrics and Gynaecology, School of Medicine, College of Health Sciences, Addis
Ababa University, Addis Ababa, Ethiopia;

[5]King's Centre for Global Health and Health Partnerships, School of Life Course and Population Sciences,
King's College London, London, UK;

[6;]Aklilu-Lemma Institute of Pathobiology, Addis Ababa University, Addis Ababa, Ethiopia;

[7]King's Global Health Institute, King's College London, London, UK;

[8]Centre for Global Mental Health, Health Service and Population Research Department, Institute of
Psychiatry, Psychology & Neuroscience, King's College London, London, UK

Authors' email addresses

Yohannes Hailemichael: yohannes.h.michael@gmail.com;

Tigist Eshetu: tigisteshetu.g@gmail.com;

Hanna Negussie: hanna.abnet@yahoo.com;

Andualem Deneke: andualemdb94@yahoo.com;

Amezene Tadesse: amezenet@yahoo.com;

Ahmed Abdella: ahmedamu@yahoo.com;

Abebe Bekele: abekele@ughe.org;

Andrew Leather: andy.leather@kcl.ac.uk;

Girmay Medhin: gtmedhin@yahoo.com;

Martin Prince: martin.prince@kcl.ac.uk;

Charlotte Hanlon: charlotte.hanlon@kcl.ac.uk.

* Correspondence: Charlotte Hanlon charlotte.hanlon@kcl.ac.uk, Department of Psychiatry, 7th Floor,
College of Health Sciences Administration Building, Black Lion Hospital Compound, Addis Ababa, PO
9086, Ethiopia
37

Abstract

Background

The objective of this study was to assess the costs, catastrophic out-of-pocket (OOP) health expenditure,

impoverishment and coping mechanisms used to pay for surgical care in a predominantly rural area of

Ethiopia.

Methods

We conducted a community-based, cross-sectional household survey of 182 people who had undergone

a surgical procedure. Participants were interviewed in their homes at six weeks post-operation. Using a

contextually adapted version of the Study of global AGEing and adult health (SAGE) questionnaire, we

estimated direct and indirect costs of surgical care from a household perspective. Catastrophic out-of-

pocket (OOP) health expenditure was estimated using thresholds of 10% and 25% of annual household

consumption expenditure. Impacts of surgical care payments on poverty levels was estimated by

comparing pre- and post-operative OOP payments. Analysis of variance, t-test and a logit model were

used to assess factors associated with catastrophic OOP health expenditure.

Results

Most surgical patients were women (87.9%), with 65% receiving obstetric surgical care. Direct costs

dominated expenditure: direct medical costs Birr 1649.5 (44.6%), direct non-medical costs Birr 1226.5

(33.2%), indirect costs Birr 821.9 (22.2%). Catastrophic OOP surgical care expenditure was experienced

by 69.2% households at the 10% threshold and 45.6% at the 25% threshold. The increase in average

normalized poverty gap due to OOP surgical care expenditure was higher in non-obstetric (14.1%)

compared to obstetric (5.8%) procedures, and for non-emergency (13.3%) compared to emergency care

(6.3%). To pay for surgical care, 38% of households had sold assets and 5% had borrowed money.

Conclusions

Due to surgical care, households faced severe financial burdens leading to impoverishment. Households

implemented hardship coping strategies to mitigate the financial constraints. Provision of free maternal

care reduced but did not eliminate these burdens. There is a pressing need to tailor financial risk

protection mechanisms to achieve universal coverage for surgical care.

Key words: catastrophic healthcare expenditure; impoverishment, universal health coverage, global

surgery

Background

In low-income and lower-middle-income countries, over nine out of ten people cannot access safe,

timely and affordable surgical and anaesthesia care [1]. Key barriers include the direct and indirect costs

related to surgical services [2, 3]. Surgical conditions additionally can have a severe impact on the

person's day-to-day activities and result in a loss of income to the household, either in the form of wage

losses from the patient or caregivers, or due to losses in agricultural output or other earnings. Shrime et

al. (2016) reported that half of the world population, or 3.7 billion people (posterior credible interval:

3.2-4.2 billion), are at risk of catastrophic expenditure if they were to need surgery because they do not

have financial risk protection [4]. Each year, an estimated 81 million people worldwide are driven to

financial catastrophe due to the cost of surgery and its associated indirect costs, with the biggest burden

borne by people in low- and middle-income countries [5]. In patients who underwent surgery at a

government hospital in Uganda, 31% faced catastrophic expenditure [6]. In Malawi, 94% of district and

87% of central hospital patients experienced catastrophic expenditure for surgical care [7] .

In Ethiopia pre-payment and other financial risk protection mechanisms are limited. In recognition of

this, in recent years the country has introduced health care financing reforms to increase affordable

access to health services and achieve universal health coverage (UHC) [8]. In Ethiopia, efforts to enhance

financial risk protection for accessing essential health services include provision of high-impact

interventions free of charge through an exemption program; subsidization of more than 80% of the cost

of care in public health facilities; implementation of community-based health insurance (CBHI) schemes;

and full subsidization of the very poor through fee waivers [9]. However, the coverage of CBHI is low

(28%)[10], while the coverage of social health insurance for the formal sector is less than 3% [11] and

fee waivers are based on a quota system, covering only 10% of the poorest population [12]. Therefore,

for the majority, care provision is based on "fees for service". Thus, out-of-pocket health expenditure

(OOPHE) still dominates all sources of expenditures for health care and constituted 35% of total health

expenditure in 2018 [13]. As a result, OOPHE has impoverishing effects on households of those who

experience ill-health, especially among the poorest [14, 15]. In addition, due to the impact of OOP

payments, a large treatment gap is likely to exist. In a recent study from Ethiopia, Kiros et al. (2020)

estimated that, on average, 0.9 % (range 0.1-5.0%) of the population becomes impoverished every year

because of the cost of accessing health care services [16]. Thus, where healthcare payments are made

mostly through OOP payments, it becomes a trade-off for patients and families to get care and suffer

catastrophic expenditure or accept the consequences of not accessing care. As is the case in Ethiopia,

many households may risk not seeking care [17].

Several studies in Ethiopia have investigated and compared the burden of OOPHE and impoverishment

for specific health conditions [18-20] and between different regional states [15, 16]. However, we were

not able to find many studies on the magnitude of catastrophic health care expenditure due to surgical

care. It is important for policy makers to understand the financial impact of surgical care on the

economic welfare of households to guide the national health policy to achieve equitable, accessible and

affordable surgical services. In light of this, the objective of this study was to quantify direct and indirect

costs, investigate the magnitude of catastrophic out-of-pocket payments and impoverishment related to

accessing surgical care and to identify the coping strategies implemented by affected households.

Materials and Methods

We conducted a community-based, cross-sectional household survey of people who had undergone a

surgical procedure and were interviewed in their homes at six weeks after the operation. This study was

conducted as part of the pre-implementation phase of the ASSET programme (Health System

Strengthening in sub-Saharan Africa). The goal of ASSET was to identify system bottlenecks to accessing

quality surgical, maternal and primary healthcare in Ethiopia, Sierra Leone, South Africa and Zimbabwe

[21]. In this paper, we present findings from Ethiopia.

Study setting

The study was conducted in Meskan, Misrak Meskan, Sodo and South Sodo districts and Butajira town of

the Gurage zone located in the Southern Nations, Nationalities and People's Region of Ethiopia. Health

services are provided through governmental and non-governmental health facilities. In the five districts

there are 94 health posts and 16 primary health care units (health centers) for preventive and curative

health services. There is one general and one primary hospital where surgical care is available. In the

general hospital, there is one surgeon, one obstetrician/gynaecologist and two Integrated Emergency

Surgical Officers (IESOs) who provide both emergency and elective surgical care. In the primary hospital,

there are two IESOs providing only emergency surgical care. The common surgical procedures

performed in the hospitals are Caesarean section, abdominal hysterectomy, management of pelvic

organ prolapse, tubal ligation, repair of hernias, management of acute appendicitis,

haemorrhoidectomy, fracture and dislocation management and tooth extractions. Most surgical care

services are based on a "fee for service" model, but obstetric care is an exempted service.

Participants

Participants were all surgical patients who had undergone a surgical procedure between 3rd June and

27th September 2019 at the general or primary hospital and were available for a community-based

survey 6-8 weeks following the operation. The aim was to investigate the costs incurred when accessing

surgical care and to estimate the proportion of people facing catastrophic OOP payments and

impoverishment (pushing the household below the poverty threshold) [22, 23].

Recruitment procedure

Participants were approached by research staff while in hospital once their condition had stabilized.

They were invited to participate in the study and provide informed consent for research staff to visit

their home for an interview on expenditures related to surgical care.

Sample size

We estimated that we would recruit 150-200 participants over the four month study period. This was a

pragmatic decision based on what was feasible and comparable with previous studies from Malawi [7]

and Ghana [24].

Outcome variables

The primary outcome was cost of surgical care, with secondary outcomes of catastrophic out-of-pocket

healthcare expenditure, the poverty impact of accessing surgical care and coping strategies

implemented to manage financial constraints.

Measurements

A contextually adapted and abbreviated version of the World Health Organization SAGE (Study of global

AGEing and adult health) survey instrument was used. SAGE was used previously in a study on health

and ageing in six LMICs [25] and in Ethiopia [14]. The SAGE instrument is used to collect information on a

variety of individual and household socio-economic attributes such as consumption expenditure,

income, assets and household demographics (detailed below). Cost of illness was derived from

consumption expenditure data and income loss of patients and caregivers associated with seeking

surgical care and providing care for the person who had undergone a surgical procedure.

Type of surgery was categorised as emergency versus elective with the assumption that this would be

relevant to the cost of care. Moreover, we classified the type of surgical care as obstetric vs. non-

obstetric procedures for the reason that obstetric care alone was exempted from full costs. These

classifications were based on information obtained from hospital records. Sociodemographic data were

measured in terms of age, sex, residential area (urban/rural), marital status, religion and education.

Information on the availability of financial risk protection mechanisms for the patient was obtained from

the hospital record and classified as having community-based health insurance (CBHI), a 'free

certificate' (fee waiver) or not covered by any financial risk protection mechanism.

Consumption expenditures

Consumption expenditure variables were collected at the household level. Food expenditure was

estimated by summing consumption of the household's own food products and expenditure on food

items over the seven days preceding the survey**.** Non-food expenditure was calculated by summing

expenditure on rent, electricity, transport, clothes and health care for the 30 days preceding the survey,

and expenditure on education, health aids, hospitalization and long-term health care for the 12 months

preceding the survey. To obtain comparable time periods, all consumption expenditures data were

annualised and then converted to a 30-day time period. Monthly household consumption expenditure

was calculated as the sum of food, non-food and health expenditure.

Out-of-pocket health expenditure

Out-of-pocket health expenditures were those made by households at the point of receiving health

services and payments directly related to seeking care (transportation, food, accommodation and

caregiving) as reported by respondents. We further classified OOP payments into 'catastrophic' and

'non-catastrophic' categories. Catastrophic OOPHE was defined in relation to a household's

consumption expenditures. OOP health care payment was taken to be financially catastrophic when it

exceeded 10% or 25% of total household consumption [26, 27].

For cost of surgical care, data were collected on expenditures incurred as:

(i) direct medical costs, including registration, consultation, medicines, diagnostics (investigations and

procedures), medical supplies and hospital charges; (ii) direct non-medical costs, like transportation to

seek care for the affected person and accompanying household members, food, accommodation, hiring

of someone as a care provider/attendant; and (iii) indirect costs, which was income that was foregone

due to the patient's inability to work or other income foregone by other household members because

they accompanied the patient to the facility or spent time caring for the patient at home [28]. In

estimating productivity losses, we followed recommended practice to use the actual income losses

rather than the potential losses. The rationale is that, in agricultural societies or for people engaged in

informal labour there are seasons in which work intensity is high and others in which work intensity is

low. In addition, household use coping strategies that may mitigate these potential losses. Thus, days of

ill-health do not necessarily translate neatly into days of lost work [29, 30]. All costs were expressed in

Ethiopian Birr and exchange rate to USD and PPP (the purchasing power parity conversion rate to

International dollar was provided at the foot note of the tables). [31, 32].

Incidence and intensity of catastrophic out-of-pocket health expenditure

The incidence of catastrophic OOPHE is measured by catastrophic head count which is the percentage of

households incurring OOPHE in excess of a specified threshold in one year [33]. In this study, two

thresholds were used: greater than or equal to 10% and 25% of total consumption expenditure. Both

thresholds are indicators linked to Sustainable Development Goals for monitoring Universal Health Care

[34].

The intensity of catastrophic OOPHE captures how much a household's OOPHE exceeds the catastrophic

threshold and is estimated through the catastrophic overshoot and the mean positive overshoot

(MPO)[33]. For calculation of catastrophic OOPHE**,** direct medical and non-medical costs were used.

Poverty impact of out-of-pocket health expenditure

The poverty impact of OOPHE was measured in terms of:

(i) poverty head count, which is the percentage of households living below the poverty line;

(ii) poverty gap, or the average amount by which resources fall short of the poverty line, measures

poverty depth or intensity of poverty (the amount by which poor households fall short of the poverty

line).

(iii) normalized poverty gap, obtained by dividing the poverty gap by the poverty line for pre- and post-

OOPHE.

The poverty line was set based on median consumption expenditure, a relative poverty line of half and

two-thirds of median consumption per capita [35, 36] .

Statistical analysis

OOP health care expenditure and poverty impact of surgical care expenditures were analyzed using one-

way analysis of variance, Student's t-test and Kruskal Wallis test depending on the required statistical

assumptions and number of categories to be compared. Categorical data were cross-tabulated and the

statistical significance of observed differences in proportions was investigated using Pearson's Chi-

square (χ^2) or alternatively Fisher's exact test, as appropriate. Further analysis was conducted to

investigate the effect of OOPHE on household food and non-food consumption expenditure using a

linear probability model.

Results

Out of 224 people who underwent surgical procedures during the study period, 182 (81.3%) participated

in the survey in the community after 6-8 weeks. See Figure 1 for a flow chart of participants.

Figure 1: Flow chart of participants

Background characteristics

See Table 1 for demographic characteristics of respondents. The mean (standard deviation; SD) age of

respondents was 33.1 (SD 13.5) years, with 84.1% within the 19–40 year age group. Most (about 88%) of

household heads were men, 56.0% (n=102) were Muslim, 81 (44.5%) had attended primary level

education and 55 (30.2%) had no formal education. The majority (87.9%) of respondents were married.

In terms of financial protection, 14 (8.4%) of the study participants were members of CBHI and 25

(13.8%) had a poverty certificate (fee waiver) to access health services without payment.

Table1: Household characteristics

Characteristics	Frequency	Percentage
Age group (years)		
19-40	153	84.1
41-50	8	4.4
51-60	9	4.9
61 or above	12	6.6
Gender		
Male	22	12.1
Female	160	87.9
Residence		
Urban	96	54.2
Rural	81	45.8
Education attainment		
No formal education	55	30.2
Primary education	81	44.5
Secondary education	20	11.0
Above secondary	26	16.3
Marital		
Never married	11	6.0
Married	160	87.9
Separated/ Divorced/ Widowed	11	6.0
Religion		
Muslim	102	56.0

Orthodox	68	37.4
Protestant	12	6.6
Member of Community-Based Health Insurance		
Yes	14	8.4
No	153	91.6
Free health care certificate(fee waiver)		
Yes	25	13.8
No	156	86.2
Either CBHI and fee waiver		
Yes	39	21.4
No	153	78.6
Surgery by condition		
Emergency	122	67.1
Non-emergency	60	32.9
Surgery by type		
Obstetric	120	65.9
Non-obstetric	62	34.1
Length of Hospital stay(days)		
1-3	84	46.9
4-5	72	40.2
6 or more	23	12.9

Consumption expenditures

Supplementary table 1 shows the monthly household consumption expenditure by consumption

categories. On average, household food consumption was Birr 5970.3 (SD 4400.3), representing 43.7%

of total consumption expenditure, with Birr 2423.3 (SD 7981.7) spent on household utilities, Birr 1694.9

(SD 6830.9) on big purchases and Birr 2876.1 (SD 3848.9) on healthcare.

Table 2 provides the detailed household consumption expenditure by household characteristics and

surgery type (emergency vs. elective; obstetric vs. non-obstetric). The share of OOP payments for health

care as a percentage of total consumption expenditure was 16.4% for obstetric surgical procedures

compared to 35.6% for non-obstetric procedures (p<0.001). Health care costs for rural residents were

about 13% (p< 0.001) higher than health care costs for urban residents. The cost of treatment for

women was about 19.7% (p<0.001) less in terms of OOP health care expenditure compared to cost of

treatment for men.

Table 2. Mean consumption expenditures (in Birr) by household characteristics, catastrophic health expenditure and type of surgical care

Household characteristics	Food Mean (%)	Housing utilities Mean (%)	Big expenditures Mean (%)	Healthcare Mean (%)	All expenditures
Urban	6804.5(46.7)	3590.3(24.5)	1700.5(11.6)	2540.1 (17.3)	14635.4
Rural	4905.8(46.2)	1016.1(9.6)	1439.4(13.6)	3241.4 (30.6)***	10602.7
Male	6699.4(44.5)	1210.5(8.0)	1229.2(8.1)	5904.9(39.2)	15044.0
Female	5795.4(46.8) **	2514.4(20.3)**	1589.4(12.8)	2459.7(19.9) ***	12358.9
Obstetric	5923.9(45.7)	2979.6(23.0)	1927.7(14.9)	2124.3(16.4)	12955.5
Non-obstetric	5867.5(48.2)	1151.4(9.5)	806.9(6.7)	4331.4(35.6) ***	12157.2
Emergency	6234.6(49.8)	2073.5(16.6)	1575.6(12.6)*	2615.3(17.3)	12499.0
Non-emergency	5234.0(40.0)**	2932.6(22.4)	1485.5(11.3)	3406.5(26.0)**	13058.6
Catastrophic	5263.8(45.7)***	1481.6(12.8)***	805.5(6.9)***	3958.3(34.3)***	11509.2
Non catastrophic	7275.0(47.1)	4458.3(28.9)	3128.8(20.2)	562.6(3.6)	15424.7

1 USD = 29.07 Birr (2019, purchasing power parity, 1 international dollars=10.74 Birr)

ANOVA and t-test for means and Pearson's χ2 for categorical variables. P<0.05,P<0.01**;P<0.001***.*

Households that faced catastrophic OOPHE had significantly lower expenditure on food (p=0.002), housing utilities (p=0.008) and big item expenditures (p=0.012) compared to non-catastrophic households.

Direct and indirect costs of surgical care

Table 3 shows the mean expenditure on surgical care. Direct medical costs contributed the largest proportion (44.6%) of the economic cost of surgical care, comprising an average of Birr 1649.5 per respondent. Of the direct medical costs, 36.2% was spent on medication and 29.3% on other medical items.

265   **Table 3. Cost of surgical care by cost categories**

Cost categories	Mean(SD)	Percentage share
A. Direct medical	1649.5(2600.6)	44.6
Registration	7.3(8.8)	0.4
Medications	596.8(1137.3)	36.2
Medical supplies	264.0(1012.2)	16.0
Hospital bed	225.4(743.8)	13.7
Investigations	73.3(201.3)	4.4
Other medical items	482.6(1212.1)	29.2
B. Direct non-medical	1226.5(1956.1)	33.2
Transportation	227.3(611.7)	18.5
Food items	596.8(937.6)	48.7
Care giver/attendants	402.5(1097.4)	32.8
C. Indirect cost (Lost income)	821.9(1698.6)	22.2

1 USD = 29.07 Birr (2019, purchasing power parity, 1 international dollars=10.74 Birr)

Following direct medical costs, the cost of direct non-medical and indirect costs accounted for an

average of Birr 1226.5 and Birr 821.9 per respondent, respectively. The major contributor to direct non-

medical expenses was expenditure on food items (mean Birr 596.8). Income loss to the patient and

family member accounted for 22.2% of all total costs of surgical care.

Table 4 shows the distribution of health care cost categories across households.

Table 4. Mean cost of surgical care by cost category, surgical type and household
characteristics

Characteristics	Frequency	Direct medical cost Mean (%)	Direct non-medical cost Mean (%)	Indirect cost Mean (%)	Total cost
Male	22	3682.2(54.4)	2222.7(32.9)	856.8 (12.7)	6761.7
Female	160	1370.1(41.8)	1089.6 (33.2)	817.2(25.0)	3276.9
P value of t-test		<0.001	0.017	0.918	
Urban	96	1516.6(46.2)	1023.5 (31.2)	740.5(22.6)	3280.6
Rural	81	1754.1(41.8)	1487.3 (35.4)	956.9 (22.8)	4198.3
P value of t-test		0.269	0.060	0.202	
Obstetric	120	1129.7(38.8)	994.6(34.2)	786.6 (27.0)	2910.9

Non- obstetric	62	2655.8(50.9)	1675.5 (32.0)	890.3(17.1)	5221.6
P-value t-test		<0.001	0.025	0.697	
Emergency	122	1387.2(39.5)	1228.2(35.0)	891.9(25.4)	3507.2
Non-Emergency	60	2183.1(53.4)	1223.4(30.0)	679.8(16.6)	4086.3
P value of t-test		0.026	0.506	0.785	
Catastrophic	124	2351.1 (47.3)	1607.2 (32.4)	1008.9 (20.3)	4967.2
Non-catastrophic	58	149.8(15.2)	412.8(32.0)	422.2 (32.8)	984.8
P value of t-test		<0.001	<0.001	0.029	
1-3 days	84	905.4(32.8)	1093.5(39.6)	761.0(27.6)	2759.9
4-5 days	72	1819.2(48.1)	1144.1(30.3)	818.8(21.6)	3782.1
6 or more days	23	3908.3(55.8)	2022.8(28.9)	1075.4(15.3)	7006.5
P value of ANOVA		<0.001	0.557	0.389	

1 USD = 29.07 Birr (2019, purchasing power parity, 1 international dollars=10.74 Birr)

There was a similar pattern of distribution with the direct medical, direct non-medical and indirect costs,

with males, rural residents, and those undergoing non-obstetric surgery spending more compared to

their counterparts. Length of hospitalization had an effect on treatment cost. For every additional day of

hospitalization on average patient incur Birr 376.0 for direct medical cost.

Catastrophic out-of-pocket health care expenditure

Table 5 shows the distribution of catastrophic OOPHE across consumption expenditure quintiles. The

proportion of households whose healthcare expenditure was above the two set thresholds for

catastrophic spending (i.e., 10% and 25%) was high. For the 10% threshold, catastrophic OOP healthcare

expenditure was significantly higher for male patients compared with female patients (95.4% vs. 64.3%;

p<0.001). Similarly, catastrophic OOP health expenditure was higher in those with non-emergency

compared to emergency conditions (p=0.016) and in those undergoing non-obstetric surgical care

compared to obstetric care (p<0.001). Significantly more rural households experienced catastrophic OOP

health expenditure compared to urban residents (83.9% vs. 54.1%; p<0.001). Catastrophic OOP health

expenditure was higher in participants who were hospitalized for 6 or more days (93.1%) compared to

those who hospitalized for 1-3 days (57.1%; p=0.005).

Table 5. Household experiencing catastrophic OOPHE by socio-economic status and surgical type

Characteristics	Catastrophic OOPHE measures					
	>=10% threshold(95% CI)			>=25% (95% CI)		
	Headcount	Overshoot	MPO	Headcount	Overshoot	MPO
Male	95.4***(72.6-99.4)	31.3***(21.0-41.6)	32.8(22.5-43.1)	68.1**(45.9-84.3)	19.4**(10.7-28.1)	28.5(18.9-38.0)
Female	64.3(56.5-71.4)	14.8(11.7-17.9)	23.0(18.9-27.0)	37.5(30.2-45.3)	7.5(5.1-9.8)	20.0(15.0-24.9)
Urban	54.1(44.0-63.9)	10.9(7.6-14.2)	20.1(15.4-24.9)	31.2(22.7-41.2)	5.0(2.8-7.2)	16.0(10.6-21.4)
Rural	83.9***(74.1-90.5)	22.6***(17.5-27.6)	26.9(21.4-32.3)	50.6**(39.7-61.4)	12.7**(8.5-16.8)	25.1(18.8-31.3)
Obstetric	56.6(47.5-65.3)	11.4(8.2-14.5)	20.1(15.5-24.8)	27.5 (20.1-36.2)	15.7(10.7-20.7)	19.8(13.5-26.2)
Non-obstetric	90.3***(79.8-95.6)	27.2***(21.3-33.0)	30.1(24.1-36.1)	67.7*** (55.0-70.2)	15.7(10.7-20.7)	23.2(17.0-29.4)
Emergency	62.2(53.2-70.5)	13.5(10.3-16.7)	21.6(17.5-25.8)	34.4(26.4-43.3)	6.6(4.3-8.8)	19.1(14.4-23.9)
Non-emergency	80.0*(67.7-88.3)	23.5**(17.0-29.9)	29.3(22.2-36.5)	55.0**(42.1-67.1)	13.7(8.3-19.1)	24.9(16.9-33.0)
CBHI(Yes)	67.9(60.0-74.9)	14.8(3.7-25.9)	20.8(6.5-35.0)	35.7(15.0-63.5)	7.3(0.9-15.6)	20.5(0.2-43.1)
CBHI(No)	71.4(42.6-89.3)	17.7(14.2-21.6)	26.0(21.8-30.2)	43.7(36.0-51.8)	9.6(6.9-12.3)	22.0(17.3-26.7)
Granted free certificate	68.0(47.2-83.4)	13.9(4.7-23.2)	20.5(7.9-33.2)	32.0(16.5-52.7)	7.1(0.3-14.7)	22.4(0.5-45.4)
No free certificate	67.9(60.1-74.8)	17.1(13.8-20.4)	25.2(21.2-29.2)	42.3(34.7-50.2)	9.2(6.6-11.7)	21.7(17.3-26.2)
Financial protection,(Yes)	66.2(58.0-73.5)	15.4(12.1-18.6)	23.2(19.1-27.4)	40.0(32.2- 48.2)	7.8(5.3-10.3)	19.5(14.5-24.4)
(No)	75.6(59.1-87.0)	22.2(14.2-30.2)	29.4(20.3-38.5)	45.9(30.5-62.1)	13.4(7.0-19.7)	29.2*(19.8-38.5)
Length of stays (1-3 days)	57.1(46.2-67.3)	11.6(7.7-15.5)	20.4(14.6-26.1)	30.9(21.9-41.7)	5.5(2.6-8.4)	17.8(10.0-25.7)
4-5 days	72.2(60.6-81.4)	17.6(12.7-22.5)	24.4(18.6-30.1)	44.4(33.3-56.1)	9.2(5.4-13.0)	20.8(14.1-27.5)
6 or more days	91.3*(70.2-97.9)	33.0***22.6-43.4)	36.2*(25.9-46.5)	69.5(47.8-85.0)	20.8***(11.8-29.8)	29.9*(20.0 -39.8)

293   *P<0.05*,P<0.01**;P<0.001***.P is from Pearson's χ2 and Kruskal walls test*

294   Financial risk protection included, CBHI(community based health insurance) and those granted free certificate to access health care without
295   payment;

Impoverishment effect of out-of-pocket payments

As shown in Supplementary table 2, using half and two-thirds of household per capita median

consumption expenditures as the poverty line, out-of-pocket surgical care expenditures led to a 10.4%

and 19.2% rise in poverty head count ratios, respectively. The normalized poverty gap increased by 6.0%

(a 90.9% increment) for a poverty line of half median consumption expenditure, and by 8.6% (an 82.7%

increment) for the poverty line of two-thirds median consumption. Similarly, using half of median

consumption, the poverty gap increased by Birr 289.9, representing a 92.0% relative increment. A

similar pattern was observed when the two-thirds of median total consumption threshold was used as

the poverty line.

A stratified analysis of poverty impact of OOP health expenditure for surgical care is presented in Table

6. The analysis shows that using a poverty line of half of median consumption expenditure, 3.9% of

obstetric, 17.6% of non-obstetric, 6.6% of emergency and 18.3% of non-emergency patients were

pushed below the poverty line after paying for surgical care. The poverty gap also increased by Birr

2042.1 for obstetric surgical intervention, Birr 2119.9 for non-obstetric surgical intervention, Birr 2134

for emergency cases and Birr 2022.8 for non-emergency cases.

**Table 6. Poverty impact of OOPHE using 1/2 and2/3 of annual median consumption as a poverty line
by surgical type**

Poverty measures	Surgical type			
	Obstetric	Non- Obstetric	Emergency	Non-emergency
I. using 1/2 of median consumption				
Headcount				
Prepayment head count	17.5%	20.9%	14.7%	26.7%
Post payment head count	21.4%	38.7%	21.3%	45.0%
Absolute point change	3.9%	17.6%***	6.6%	18.3%*
Relative change	22.2%	84.2%	44.8%	68.5%
Poverty gaps				
Prepayment poverty gap (Birr)	315.6	313.9	268.2	410.2
Post payment poverty gap (Birr)	493.5	820.6	454.7	910.2

Absolute gap change(Birr)	177.9	506.6*	186.5	500.0*
Relative change	56.3%	161.3%	69.5%	121.8%
Normalized gaps				
Prepayment normalized gap^X	6.6%	6.5%	5.6%	8.6%
Mean positive pre-payment poverty gap(Birr)	1803.4	1497.2	1817.9	1538.4
Post-payment normalized gap ^{Y}	10.3%	17.2%	9.5%	19.0%
Mean positive post-payment poverty gap (Birr)	2042.1	2119.9	2134.0	2022.8
Absolute percentage point change (impact) $^Z_{(=Y-X)}$	3.7%	10.6%*	3.9%	10.4%*
Relative percentage change (=Z/X*100)	56.0%	163.0	69.6%	120.9%
II. Using 2/3 of median consumption				
Headcount				
Prepayment head count	30.0%	27.4%	24.6%	38.3%
Post payment head count	44.1%	56.4%	42.6%	60.0%
Absolute point change	14.1%	29.0%*	18.0%	21.7%
Relative change	47%	105.8%	73.1%	56.6%
Poverty gaps				
Prepayment poverty gap	655.3	1025.8	539.3	925.2
Post payment poverty gap(Birr)	688.2	1586.8	943.5	1772.9
Absolute gap change(Birr)	132.9	561.0	404.2	847.7
Relative change	20.2%	54.6%	74.9%	91.6%
Normalized gaps				
Prepayment normalized gap^X	10.3%	10.8%	8.4%	14.5%
Mean positive pre-payment poverty gap(Birr)	2184.4	2510.2	2193.3	2413.5
Post-payment normalized gap ^{Y}	16.1%	29.4%	14.8%	27.9%
Mean positive post-payment poverty gap(Birr)	2322.7	2810.9	2213.6	2954.8
Absolute percentage point change (impact) $^Z_{(=Y-X)}$	5.8%	14.1%*	6.3%	13.3%*
Relative percentage change (=Z/X*100)	56.3%	130.5%	75.0%	91.7%

1 USD = 29.07 Birr (2019, purchasing power parity, 1 international dollars=10.74 Birr)

Kruskal-Wallis comparing pre and post payment by surgical care sub groups

Out-of-pocket payments for surgical care raised the average poverty gap (normalized gap) for

households of obstetric, non-obstetric, emergency and non-emergency cases by 5.8%, 14.1%, 6.3% and

13.3%, respectively.

At the poverty line of two-thirds of consumption expenditure, 14.1% of obstetric surgical procedures,

29.0% of non- obstetric surgical procedures, 18.0% of emergency and 21.7% of non-emergency patients

fell into poverty due to out-of-pocket payments for healthcare. We also found significant differences in

the poverty gap after paying for health care. The poverty gap increased by Birr 2184.4 for obstetric

surgical procedures, Birr 2510.2 for non-obstetric surgical procedures, Birr 2193.3 for emergencies and

Birr 2413.5 for non-emergencies.

Coping strategies implemented for financial constraints

Supplementary table 3 shows the source of payments for surgical care. More than half (53.3%) of those

who received surgical care paid for the service from their income, with over one-third (37.9%) of

households needing to sell items, 15.3% having care paid for by relatives and 4.9% needing to borrow

money.

Factors associated with catastrophic OOP health expenditure

Table 7 presents factors associated with catastrophic OOP health expenditure in excess of either 10% or

25% of the household's consumption expenditure. Patients who received obstetric surgical care had

much lower odds of catastrophic OOP health expenditure (adjusted Odds Ratio (AOR) 0.1, 95% CI: 0.03,

0.7; p=0.047) compared to non-obstetric surgical patients. Rural residents had significantly higher odds

of catastrophic OOP health expenditure than urban residents at the 10% threshold (AOR 4.2, 95%CI: 1.8,

9.7; p=0.002) and at the 25% threshold (AOR 1.8, 95%CI: 1.2, 3.9; p=0.049. At the 10% and 25%

thresholds, the odds of facing catastrophic expenditure for those who had no financial risk protection

mechanism were (AOR 3.3, 95%CI: 0.7-15.4) and (AOR 2.6, 95%CI: 0.7-8.1) times higher, respectively,

compared to those who had financial risk protection. However, these differences are not statistically

significant.

Table 7. Factors associated with catastrophic OOPHE

Characteristics	Catastrophic thresholds			
	>=10%		>=25%	
	COR(95%,CI)	AOR(95%,CI)	COR(95%,CI)	AOR(95%,CI)
Gender				
Male	Reference	Reference	Reference	Reference
Female	0.08* (0.01- 0.6)	0.5(0.04-6.0)	0.2(0.1-0.7)	0.6(0.3-5.0)
Residence				
Urban	Reference	Reference	Reference	Reference
Rural	4.4*** (2.1-9.0)	4.2** (1.8-9.7)	2.2**(1.2-4.1)	1.8* (1.2-3.9)
Surgical condition				
Emergency	Reference	Reference	Reference	Reference
Non- emergency	2.42* (1.16- 5.0)	1.0(0.3-3.2)	2.3**(1.2-4.3)	1.6(0.2-2.0)
Surgical type				
Obstetric	0.1* (0.05-0.3)	0.1* (0.03-0.7)	0.1***(0.09-0.3)	0.1** (0.04-0.5)
Non-obstetric	Reference	Reference	Reference	Reference
Users by payment status				
CBHI				
Yes	Reference	Reference	Reference	Reference
No	0.8(0.2-2.8)	3.1(0.5-18.1)	1.4(0.4-4.3)	4.1(0.8-19.9)
Free certificate				
Yes	Reference	Reference	Reference	Reference
No	0.9(0.4-2.4)	2.8(0.4-24.1)	1.5(0.6-3.8)	2.1(0.3-12.4)
Length of stays				
1-3 days	0.1(0.02-0.5)	0.3(0.05-2.0)	0.1**(0.03-0.4)	0.1* (0.07-0.5)
4-5 days	0.2(0.05-1.1)	0.5(0.1-3.8)	0.5*(0.01-0.8)	0.3* (0.1-0.9)
6 or more days	Reference	Reference	Reference	Reference
Financial risk protection				
Yes	Reference	Reference	Reference	Reference
No	0.6(0.2-1.4)	3.3(0.7-15.4)	0.7(0.3-1.6)	2.6(0.7-8.1)

Financial risk protection included, CBHI(community based health insurance) or granted free certificate to access

health care without payment; p<0.05*,p<0.01**,p<0.001***

Discussion

This study examined the extent of cost of surgical care among service users in primary and general

hospitals in rural Ethiopia. Even though our service-using sample was likely to include households who

were already of better socioeconomic status than those not accessing care at all, we found that the cost

of surgical care was catastrophic for households.

More than two-thirds of surgical patients faced catastrophic OOP health expenditure at the 10%

threshold of total consumption expenditure. Our finding is lower than reports from Malawi [7] and

Madagascar [37], while being greater than the 59% for Vietnam [38], 12% for Sierra Leone [39] and 31%

reported for Uganda [6]. All surgical care is considered free in Malawi and Uganda, and in Madagascar

there is a form of insurance for the poor [37]. In Ethiopia, surgical services are based on payments at the

point of care, with the exception of obstetric services which are meant to be provided free of charge

through the exemption programme [9, 40, 41]. Beyond these differences in healthcare financial

strategies across countries, it is likely that the proportion of those in need of surgical care who even

manage to access hospital care may vary, expected to be lower in this rural, low-income Ethiopian

setting. Thus, those accessing surgical care represent households of relatively higher socio-economic

status or women accessing obstetric care. A further reason for varying estimates of catastrophic OOP

health expenditure is due to methodological differences in estimating catastrophic expenditure. In some

of these studies, wage loss was included in the estimation of catastrophic health expenditure whereas

we considered that this cost is not an OOP payment.

Despite obstetric care being considered an essential and fee-exempted health service at government

health facilities in Ethiopia, 56.6% and 27.5% of women that received obstetric surgical care

encountered catastrophic OOP payments at the 10% and 25% thresholds, respectively. Earlier studies

from Ethiopia reported that maternal services are supposed to be provided free of charge in Ethiopia

but costs incurred by the households to assess care are substantial [40, 41]. Similarly, in our analysis,

despite having financial protection mechanisms either in the form of CBHI or a free certificate to access

health care, about 68% of surgical patients faced catastrophic OOP payments findings. The free health

initiative in Sierra Leone provided more protection than in Ethiopia [39]. Nonetheless, our finding is

similar to the findings from Ghana that reported 58–87% of insured patients face financial catastrophe

when accessing surgical care [24]. There have been similar reports from a previous study from Ethiopia

on general health care among the members of CBHI [42]. However, it remains unclear why households

that are members of health insurance programs are paying OOP expenses for services and facing

catastrophic health expenditure. Probably the frequent stock out of medicines and medical supplies in

government facilities may force patients to buy these items from private retail pharmacies. On the other

hand the fee waiver targeting poor individuals proved inadequate, benefiting only 10% of the poor

households [43].

Our findings demonstrated that OOP payment was a major cause of poverty in patients receiving

surgical care. Thus, using half and two-thirds of median consumption expenditure as poverty lines,

paying for surgical care pushed 10.4 and 19.2 percent households into poverty, respectively. These

findings are higher than the nine percent reported for Sierra Leone [44] and the three percent for

Uganda [6]. One reason could be in Sierra Leone patients were from high wealth quintiles and less likely

to be affected by OOP payments. The other reason for the difference could be due to the differences in

the poverty measures used. We used a relative poverty line while the Sierra Leone and the Uganda

studies used international poverty lines. Using a relative poverty threshold might better identify

individuals with limited resources that are at risk of having adverse socio-emotional outcomes [45]. Our

study might have overestimated the poverty impact of OOP payment for surgical care; the reference

household expenditure for calculating catastrophic healthcare expenditure was based on expenditure

after surgery which may already have reduced as a financial coping strategy.

In a stratified analysis of poverty impact, both poverty headcount and poverty gap were much higher for

non-obstetric and non-emergency patients compared to obstetric and emergency patients, resulting in a

major impoverishment burden. The findings for non-emergency vs. emergency surgical conditions

reflect that most emergency surgical conditions are obstetric and thus exempted from point-of-care

costs. The poverty head count impact of OOP payment using half of median-consumption expenditure

for those who had obstetric or non-obstetric surgical care were 3.9 and 17.6%, respectively, while for

emergency vs. non-emergency surgical care it was 6.6 and 18.3 percent, respectively. When a poverty

line of two-thirds of median- consumption expenditure was used, poverty impact of OOP payment

showed similar trends with higher numbers of households being pushed below the poverty line.

The poverty gap following OOP payments increased for all surgical types. However, it was three times

higher for non-obstetric than obstetric and twice as high for non-emergency compared to emergency

surgical care. This finding indicates that all types of surgical care should be considered in priority setting

and financial risk protection mechanisms.

Health care expenditure was the second largest consumption expenditure category for the household

both in absolute terms and as a proportion of total consumption expenditure. Thus, spending a large

share of household budget on health care might impact expenditure on other subsistence needs of the

household. This result is similar to findings from Vietnam [46] and China [47].

In our analysis all respondents incurred OOP spending, irrespective of their payment status (i.e member

of CBHI, free certificate and payer). Overall, patients spent a higher share of surgical care cost on direct

medical costs (44.6%), while the indirect cost (lost income) contributed the smallest share (22.2%). The

interviews took place quite early in the recovery phase which might under-estimate the share of indirect

cost. A study of surgical in-patient care costs in Sierra Leone reported that direct medical costs

accounted for 63% of surgical care costs whereas indirect costs were only 21% [44]. Medications costs

were the largest expenditure category of direct medical costs which accords with findings from Rwanda

[48], Ghana [24] and Malawi [49].

Income losses of patients and their caregivers due to surgical illness were substantial. However, the

observed loss of income was smaller than that reported in previous studies [7, 44]. This could be due to

the high proportion of our sample that was women (88%) and housewives, thus reporting less income in

our study or differences in wage rates. Transportation costs for the patient and accompanying family

members required to stay with the patient during their hospital stay also contributed to catastrophic

OOP payments. In a review of surgical care costs, Okoroh (2021) identified that transportation cost and

wage loss were major drivers of catastrophic expenditures for surgery [50].

Inequitable impacts of surgical care costs were evident in our study. The ratio of health expenditure to

total consumption expenditure was highest (30.6%) in rural compared to urban (17.3%) households. This

indicates that rural households are disproportionately burdened by cost of surgical care. This is

explained by the high transport cost to access care from rural areas and lower total household

consumption of rural residents. In a household survey of catastrophic health expenditure in India [51]

and Zambia [52], the highest proportion of catastrophic expenditure was reported among the poorest

and rural residents, although all economic status groups experienced high catastrophic costs. Therefore

financial risk protection mechanisms must do more to target rural households.

In our study, non-emergency or non-obstetric surgical patients incurred significantly more costs

compared to emergency or obstetric cases. However, the majority of our sample had undergone

obstetric procedures and were exempted from costs at the point of care and had in most cases had

presented as an emergency. For such women, transport costs were subsidized or free if they were able

to access an ambulance. Our finding of lower surgical care costs for obstetric care is similar with the

study from Uganda that reported patients who received Cesarean sections spent less [6], which is

explained by the shorter duration of obstetrics admission in the hospital. In our analysis, women paid

lower surgical care costs compared to their male counterparts. Longer hospital stays were associated

with increased direct medical costs. The longer the stay the higher the costs of medicines, hospital bed

and the medical supplies used by the patient. In similar studies prolonged hospitalization was associated

with increased costs [44, 53].

Direct medical, direct non-medical and indirect costs were significantly higher for households that faced

catastrophic OOP payment for surgical care compared to those households that did not face

catastrophic payments. Therefore, these costs are likely to compromise the consumption of food and

other subsistence needs and impact negatively on the quality of life of these households.

In practice, in Ethiopia, surgical care service charges are similar regardless of economic status. Hence,

the poor pay the same amount for surgical care despite having less income. Therefore, even relatively

low spending on surgical care by the poor is catastrophic and thus likely to compromise access to

services. Moreover, in Ethiopia where one-fifth of the population is living below the poverty line[9], OOP

payment is a major obstacle to the use of essential surgical care and exacerbates impoverishment.

In order to pay for surgical care households used different coping strategies. The majority used their

income, which may not be an option for poorer households who could thus not access surgical care at

all. In addition, use of income for meeting health expenses reduces current consumption of other goods

and services. Moreover, the other coping strategies used were borrowing and selling assets, which are

deeply regressive and could induce long-term household economic impact and impoverishment.

Previous studies on coping strategies for paying of surgical care reported similar findings [7, 44, 50].

Although we believe this is the first study in Ethiopia to investigate surgical care costs, catastrophic

expenditure, and poverty impact of OOP payments, our study has the following limitations. Only two

hospitals were chosen and these may not be representative of all hospitals in Ethiopia. The sample size

was relatively small and thus may have limited the statistical power to detect differences. The possibility

of underestimation of non-medical costs due to recall bias cannot be ruled out. In subsistence farming

estimating lost income and consumption expenditure is difficult and might lead to under-estimation of

these costs for the respondents. We did not adjust consumption expenditure for household composition

(i.e., age and size) as data were not collected on these variables. Thus, estimates were calculated per

household which may affect interpretation of the difference between urban and rural households.

Lastly, the survey was conducted a few weeks after the patient was discharged from hospital and

households may have already reduced expenditure on non-essential (and some essential) items to

manage the catastrophic costs. So this might have over-estimated the degree to which the surgical costs

were catastrophic.

Conclusions

Due to surgical care, households faced severe financial burdens of treatment costs and lost income

leading to impoverishment. This was particularly marked for non-obstetric procedures which were not

exempt from payment at the point of care. Households were forced to use hardship coping mechanisms

to pay treatment cost. Therefore, surgical care should be accessible to all without financial hardship to

achieve universal health coverage. In addition, strategies need to be explored by the Ethiopian health

insurance service to include direct non-medical expenses like transportation costs that are not covered

currently.

Supplementary material

Supplementary table 1: Household consumption expenditure cost categories

Supplementary table 2: Poverty impact of OOP payments for surgical care

Supplementary table 3: Sources of payment for surgical care

Abbreviations

ASSET: Health System Strengthening in sub-Saharan Africa Health System Strengthening in sub-Saharan

Africa; OOP: Out-of-pocket; SAGE: Study on global Ageing and adult health; SDGs: Sustainable

development goals; UHC: Universal health coverage; WHO: World health Organization

Acknowledgments

We would like to thank all participants in the study for their participation. For the purposes of open

access, the author has applied a Creative Commons Attribution (CC BY) licence to any Accepted Author

Manuscript version arising from this submission.

Ethics approval and consent to participation

This study was conducted according to the principles expressed in the Declaration of Helsinki. Ethical

approval was obtained from the Institutional Review Board of the College of Health Sciences, Addis

Ababa University (026/18/psy; 16th May 2018), and the Research Ethics Committee of King's College

London (RESCM-17/18-6144; 18th August 2018). All participants gave written, informed consent. For

people who were non-literate, a witness confirmed that the information sheet had been read out

correctly and the participant provided a fingerprint.

Authors' contributions

Conceptualization (CH, MP and AL). YH analyzed the data, drafted the manuscript and wrote

subsequent versions. CH and MP provided guidance and inputs to all versions of the

manuscript. All co-authors gave critical feedback on drafts of the manuscript, read and

approved the final version of the manuscript.

Availability of data and materials

Available from the corresponding author on request.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no conflict of interests.

References

1.   Meara JG, Leather AJ, Hagander L, Alkire BC, Alonso N, Ameh EA, et al. Global Surgery 2030: evidence and

solutions for achieving health, welfare, and economic development. Lancet (London, England).

2015;386(9993):569-624. Epub 2015/05/01. doi: 10.1016/s0140-6736(15)60160-x. PubMed PMID: 25924834.

2.   Grimes CE, Bowman KG, Dodgion CM, Lavy CBD. Systematic Review of Barriers to Surgical Care in Low-

Income and Middle-Income Countries. World journal of surgery. 2011;35(5):941-50. doi: 10.1007/s00268-011-

1010-1.

3.   Willott C, Boyd N, Wurie H, Smalle I, Kamara TB, Davies JI, et al. Staff recognition and its importance for

surgical service delivery: a qualitative study in Freetown, Sierra Leone. Health policy and planning. 2021;36(1):93-

100. Epub 2020/11/28. doi: 10.1093/heapol/czaa131. PubMed PMID: 33246332; PubMed Central PMCID:

PMCPMC7938499.

4.   Shrime MG, Dare A, Alkire BC, Meara JG. A global country-level comparison of the financial burden of

surgery. Br J Surg. 2016;103(11):1453-61. Epub 2016/07/19. doi: 10.1002/bjs.10249. PubMed PMID: 27428044.

5.   Shrime MG, Bickler SW, Alkire BC, Mock C. Global burden of surgical disease: an estimation from the

provider perspective. The Lancet Global health. 2015;3 Suppl 2:S8-9. Epub 2015/05/01. doi: 10.1016/s2214-

109x(14)70384-5. PubMed PMID: 25926322.

6.   Anderson GA, Ilcisin L, Kayima P, Abesiga L, Portal Benitez N, Ngonzi J, et al. Out-of-pocket payment for

surgery in Uganda: The rate of impoverishing and catastrophic expenditure at a government hospital. PloS one.

2017;12(10):e0187293. Epub 2017/11/01. doi: 10.1371/journal.pone.0187293. PubMed PMID: 29088302; PubMed

Central PMCID: PMCPMC5663485 manuscript have the following competing interests: MGS has received speaking

fees from Ethicon Inc. however Ethicon was not involved in any portion of this study. This does not alter our

adherence to PLOS ONE policies on sharing data and materials.

7.   Bijlmakers L, Wientjes M, Mwapasa G, Cornelissen D, Borgstein E, Broekhuizen H, et al. Out-of-pocket

payments and catastrophic household expenditure to access essential surgery in Malawi - A cross-sectional patient

survey. Ann Med Surg (Lond). 2019;43:85-90. Epub 2019/07/16. doi: 10.1016/j.amsu.2019.06.003. PubMed PMID:

31304010; PubMed Central PMCID: PMCPMC6580231.

8.   Alebachew A, Hatt L, Kukla M. Monitoring and evaluating progress towards Universal Health Coverage in

Ethiopia. PLoS Med. 2014;11(9):e1001696. Epub 2014/09/23. doi: 10.1371/journal.pmed.1001696. PubMed PMID:

25244146; PubMed Central PMCID: PMCPMC4171462.

9.   FMOH. Health Sector Transformational Plan II 2020/2021-2024/2025. Addis Ababa: Federal Ministry of

Health, 2021.

10.   Ethiopian Public Health Institute - EPHI, Federal Ministry of Health - FMoH, ICF. Ethiopia Mini

Demographic and Health Survey 2019. Addis Ababa, Ethiopia: EPHI/FMoH/ICF, 2021.

11.   FMOH. 7th National Health Accounts 2016/17; Statistical Report. Addis Ababa,Ethiopia: Federal

Democratic Republic of Ethiopia Ministry of Health, 2019.

12.   Agency EHI. CBHI Members' Registration and Contribution (2011-2020): . Addis Ababa, Ethiopia: Ethiopian

Health Insurance Agency, 2020.

13.   WHO. Global Health expenditure data base. Geneva: World Health Organization, 2019.

14.   Hailemichael Y, Hanlon C, Tirfessa K, Docrat S, Alem A, Medhin G, et al. Catastrophic health expenditure

and impoverishment in households of persons with depression: a cross-sectional, comparative study in rural

Ethiopia. BMC public health. 2019;19(1):930. Epub 2019/07/13. doi: 10.1186/s12889-019-7239-6. PubMed PMID:

31296207; PubMed Central PMCID: PMCPMC6625021.

15.   Obse AG, Ataguba JE. Assessing medical impoverishment and associated factors in health care in Ethiopia.

BMC Int Health Hum Rights. 2020;20(1):7. Epub 2020/04/02. doi: 10.1186/s12914-020-00227-x. PubMed PMID:

32228634; PubMed Central PMCID: PMCPMC7106681.

16.   Kiros M, Dessie E, Jbaily A, Tolla MT, Johansson KA, Norheim OF, et al. The burden of household out-of-

pocket health expenditures in Ethiopia: estimates from a nationally representative survey (2015-16). Health policy

and planning. 2020;35(8):1003-10. Epub 2020/08/11. doi: 10.1093/heapol/czaa044. PubMed PMID: 32772112;

PubMed Central PMCID: PMCPMC7553759.

17.   Hailemariam M, Fekadu A, Prince M, Hanlon C. Engaging and staying engaged: a phenomenological study

of barriers to equitable access to mental healthcare for people with severe mental disorders in a rural African

setting. International journal for equity in health. 2017;16(1):156. Epub 2017/08/31. doi: 10.1186/s12939-017-

0657-0. PubMed PMID: 28851421; PubMed Central PMCID: PMCPMC5576237.

18.   Tolla MT, Norheim OF, Verguet S, Bekele A, Amenu K, Abdisa SG, et al. Out-of-pocket expenditures for

prevention and treatment of cardiovascular disease in general and specialised cardiac hospitals in Addis Ababa,

Ethiopia: a cross-sectional cohort study. BMJ Glob Health. 2017;2(2):e000280. Epub 2017/12/16. doi:

10.1136/bmjgh-2016-000280. PubMed PMID: 29242752; PubMed Central PMCID: PMCPMC5584490.

19.   Memirie ST, Metaferia ZS, Norheim OF, Levin CE, Verguet S, Johansson KA. Household expenditures on

pneumonia and diarrhoea treatment in Ethiopia: a facility-based study. BMJ Glob Health. 2017;2(1):e000166. Epub

2017/06/08. doi: 10.1136/bmjgh-2016-000166. PubMed PMID: 28589003; PubMed Central PMCID:

PMCPMC5321393.

20.   Kasahun GG, Gebretekle GB, Hailemichael Y, Woldemariam AA, Fenta TG. Catastrophic healthcare

expenditure and coping strategies among patients attending cancer treatment services in Addis Ababa, Ethiopia.

BMC public health. 2020;20(1):984. Epub 2020/06/24. doi: 10.1186/s12889-020-09137-y. PubMed PMID:

32571275; PubMed Central PMCID: PMCPMC7310089.

21.   Seward N, Hanlon C, Abdulahi A, Abrams Z, Alem A , Araya R. HeAlth System StrEngThening in four sub-

Saharan African countries (ASSET) to achieve high-quality, evidence-informed surgical, maternal and new born, and

primary care: protocol for pre-implementation phase studies. Research Square Preprint 2021.

22.   Seward N, Hanlon C, Abdella A, Abrahams Z, Alem A, Araya R, et al. HeAlth System StrEngThening in four

sub-Saharan African countries (ASSET) to achieve high-quality, evidence-informed surgical, maternal and newborn,

and primary care: protocol for pre-implementation phase studies. Global health action. 2022;15(1):1987044. Epub

2022/01/18. doi: 10.1080/16549716.2021.1987044. PubMed PMID: 35037844; PubMed Central PMCID:

PMCPMC8765245.

23. Seward N, Murdoch J, Hanlon C, Araya R, Gao W, Harding R, et al. Implementation science protocol for a

participatory, theory-informed implementation research programme in the context of health system strengthening

in sub-Saharan Africa (ASSET-ImplementER). BMJ open. 2021;11(7):e048742. Epub 2021/07/11. doi:

10.1136/bmjopen-2021-048742. PubMed PMID: 34244274; PubMed Central PMCID: PMCPMC8268893.

24. Okoroh J, Sarpong DO, Essoun S, Riviello R, Harris H, Weissman JS. Does insurance protect individuals from

catastrophic payments for surgical care? An analysis of Ghana's National Health Insurance Scheme at Korle-Bu

teaching Hospital. BMC Health Serv Res. 2020;20(1):45. Epub 2020/01/19. doi: 10.1186/s12913-020-4887-2.

PubMed PMID: 31952520; PubMed Central PMCID: PMCPMC6969414.

25. WHO. World Health Organization study on global Ageing and adult health (SAGE). Geneva: World Health

Organization, 2013.

26. Wagstaff A, Flores G, Hsu J, Smitz MF, Chepynoga K, Buisman LR, et al. Progress on catastrophic health

spending in 133 countries: a retrospective observational study. The Lancet Global health. 2018;6(2):e169-e79.

Epub 2017/12/19. doi: 10.1016/s2214-109x(17)30429-1. PubMed PMID: 29248367.

27. Bank WatW. Tracking Universal Health Coverage: The 2017 Global Monitoring Report. Geneva: World

Health Organization and the International Bank for Reconstruction and Development / The World Bank, 2017.

28. Russell S. The economic burden of illness for households in developing countries: a review of studies

focusing on malaria, tuberculosis, and human immunodeficiency virus/acquired immunodeficiency syndrome. Am J

Trop Med Hyg. 2004;71(2 Suppl):147-55. Epub 2004/08/28. PubMed PMID: 15331831.

29. WHO. WHO guide to identifying the economic consequences of disease and injury. Geneva: World Health

Organization, 2009.

30. Chisholm D, Stanciole AE, Tan Torres Edejer T, Evans DB. Economic impact of disease and injury: counting

what matters. BMJ (Clinical research ed). 2010;340:c924. Epub 2010/03/04. doi: 10.1136/bmj.c924. PubMed

PMID: 20197323.

31. Ethiopia NBo. Inter-bank daily foreign exchange rate in (USD). Addis Ababa: national bank of Ethiopia,

2020.

32.   Bank W. World development indicators 2022 [1st June,2022]. Available from:

https://data.worldbank.org/indicator/PA.NUS.PPP?end=2019&start=1990.

33.   Wagstaff A, van Doorslaer E. Catastrophe and impoverishment in paying for health care: with applications

to Vietnam 1993-1998. Health Econ. 2003;12(11):921-34. Epub 2003/11/06. doi: 10.1002/hec.776. PubMed PMID:

14601155.

34.   UN. Transforming Our World: The 2030 Agenda for Sustainable Development'. New York: United Nations,

2015.

35.   Wagstaff A, Flores G, Smitz MF, Hsu J, Chepynoga K, Eozenou P. Progress on impoverishing health

spending in 122 countries: a retrospective observational study. The Lancet Global health. 2018;6(2):e180-e92.

Epub 2017/12/19. doi: 10.1016/s2214-109x(17)30486-2. PubMed PMID: 29248366.

36.   and JLA, Bennett NG, Conley DC, Li J. The Effects of Poverty on Child Health and Development. Annual

Review of Public Health. 1997;18(1):463-83. doi: 10.1146/annurev.publhealth.18.1.463. PubMed PMID: 9143727.

37.   Bruno E, White MC, Baxter LS, Ravelojaona VA, Rakotoarison HN, Andriamanjato HH, et al. An Evaluation

of Preparedness, Delivery and Impact of Surgical and Anesthesia Care in Madagascar: A Framework for a National

Surgical Plan. World journal of surgery. 2017;41(5):1218-24. doi: 10.1007/s00268-016-3847-9.

38.   Nguyen H, Ivers R, Jan S, Pham C. Cost of surgery and catastrophic expenditure in people admitted to

hospital for injuries: estimates from a cohort study in Vietnam. Lancet (London, England). 2015;385 Suppl 2:S50.

Epub 2015/08/28. doi: 10.1016/s0140-6736(15)60845-5. PubMed PMID: 26313100.

39.   van Duinen AJ, Westendorp J, Ashley T, Hagander L, Holmer H, Koroma AP, et al. Catastrophic expenditure

and impoverishment after caesarean section in Sierra Leone: An evaluation of the free health care initiative. PloS

one. 2021;16(10):e0258532. Epub 2021/10/16. doi: 10.1371/journal.pone.0258532. PubMed PMID: 34653191;

PubMed Central PMCID: PMCPMC8519447.

40.   Abadi T, Mebratie AD. Cost of Treating Maternal Complications and Associated Factors in Mekelle General

Hospital, Northern Ethiopia. Risk management and healthcare policy. 2021;14:87-95. Epub 2021/01/16. doi:

10.2147/rmhp.s285793. PubMed PMID: 33447112; PubMed Central PMCID: PMCPMC7802893.

41.   Akalu T, Guda A, Tamiru M, Haile Mariam D. Examining out of pocket payments for maternal health in

rural Ethiopia: Paradox of free health care un-affordability. Ethiopian Journal of Health Development. 2012;26:251-

7.

42.   Agency EHI. Evaluation of Community-Based Health Insurance Pilot Schemes in Ethiopia: Final Report. .

Addis Ababa,Ethiopia: Ethiopian health Insurance Agency(EHIA), 2015.

43.   Demissie B, Gutema Negeri K. Effect of Community-Based Health Insurance on Utilization of Outpatient

Health Care Services in Southern Ethiopia: A Comparative Cross-Sectional Study. Risk management and healthcare

policy. 2020;13:141-53. Epub 2020/03/12. doi: 10.2147/rmhp.s215836. PubMed PMID: 32158291; PubMed Central

PMCID: PMCPMC7049267.

44.   Phull M, Grimes CE, Kamara TB, Wurie H, Leather AJM, Davies J. What is the financial burden to patients

of accessing surgical care in Sierra Leone? A cross-sectional survey of catastrophic and impoverishing expenditure.

BMJ open. 2021;11(3):e039049. Epub 2021/05/20. doi: 10.1136/bmjopen-2020-039049. PubMed PMID: 34006018;

PubMed Central PMCID: PMCPMC7942261.

45.   Lee K, Zhang L. Cumulative Effects of Poverty on Children's Social-Emotional Development: Absolute

Poverty and Relative Poverty. Community mental health journal. 2022;58(5):930-43. Epub 2021/11/10. doi:

10.1007/s10597-021-00901-x. PubMed PMID: 34750684.

46.   Nguyen KT, Khuat OT, Ma S, Pham DC, Khuat GT, Ruger JP. Effect of health expenses on household

capabilities and resource allocation in a rural commune in Vietnam. PloS one. 2012;7(10):e47423. Epub

2012/10/19. doi: 10.1371/journal.pone.0047423. PubMed PMID: 23077612; PubMed Central PMCID:

PMCPMC3471826.

47.   Fang K, Jiang Y, Shia B, Ma S. Impact of illness and medical expenditure on household consumptions: a

survey in western China. PloS one. 2012;7(12):e52928. Epub 2013/01/04. doi: 10.1371/journal.pone.0052928.

PubMed PMID: 23285229; PubMed Central PMCID: PMCPMC3532419.

48.   Rickard JL, Ngarambe C, Ndayizeye L, Smart B, Majyambere JP, Riviello R. Risk of Catastrophic Health

Expenditure in Rwandan Surgical Patients with Peritonitis. World journal of surgery. 2018;42(6):1603-9. Epub

2017/11/17. doi: 10.1007/s00268-017-4368-x. PubMed PMID: 29143091.

49.   Cornelissen D, Mwapasa G, Gajewski J, McCauley T, Borgstein E, Brugha R, et al. The Cost of Providing

District-Level Surgery in Malawi. World journal of surgery. 2018;42(1):46-53. Epub 2017/08/10. doi:

10.1007/s00268-017-4166-5. PubMed PMID: 28791448; PubMed Central PMCID: PMCPMC5740194.

50.   Okoroh JS, Riviello R. Challenges in healthcare financing for surgery in sub-Saharan Africa. Pan Afr Med J.

2021;38:198. Epub 2021/05/18. doi: 10.11604/pamj.2021.38.198.27115. PubMed PMID: 33995804; PubMed

Central PMCID: PMCPMC8106793.

51.   Chowdhury S, Gupta I, Trivedi M, Prinja S. Inequity & burden of out-of-pocket health spending: District

level evidences from India. The Indian journal of medical research. 2018;148(2):180-9. Epub 2018/11/02. doi:

10.4103/ijmr.IJMR_90_17. PubMed PMID: 30381541; PubMed Central PMCID: PMCPMC6206772.

52.   Masiye F, Kaonga O, Kirigia JM. Does User Fee Removal Policy Provide Financial Protection from

Catastrophic Health Care Payments? Evidence from Zambia. PloS one. 2016;11(1):e0146508. Epub 2016/01/23.

doi: 10.1371/journal.pone.0146508. PubMed PMID: 26795620; PubMed Central PMCID: PMCPMC4721670

interests.

53.   Nwanna-Nzewunwa O, Oke R, Agwang E, Ajiko MM, Yoon C, Carvalho M, et al. The societal cost and

economic impact of surgical care on patients' households in rural Uganda; a mixed method study. BMC Health Serv

Res. 2021;21(1):568. Epub 2021/06/11. doi: 10.1186/s12913-021-06579-x. PubMed PMID: 34107950; PubMed

Central PMCID: PMCPMC8190862.

Figure 1