Artificial Intelligence and Patient Education: Examining the Accuracy and Reproducibility of

Responses to Nutrition Questions Related to Inflammatory Bowel Disease by GPT-4

Short: ChatGPT and nutrition

Jamil S. Samaan[a], MD, Kelly Issokson[a], RD, MS, Erin Feldman[a], RD, CSP, Christina Fasulo[a], RD,

MS, CNSC, Wee Han Ng[b], Nithya Rajeev[c], BS, Barbara Hollander[a], MD, RD, Yee Hui Yeo[a], MD,

MSc, Eric Vasiliauskas[d], MD.

[a]Karsh Division of Digestive and Liver diseases, Department of Medicine, Cedars-Sinai Medical

Center, 8700 Beverly Blvd, Los Angeles, CA 90048.

[b]Bristol Medical School, University of Bristol, 5 Tyndall Ave, Bristol BS8 1UD, United Kingdom.

[c]Keck School of Medicine of USC, 1975 Zonal Ave, Los Angeles, CA 90033

[d]Inflammatory Bowel and Immunobiology Research Institute, Karsh Division of Digestive and

Liver Diseases, Department of Medicine, Cedars-Sinai Medical Center, Los Angeles, CA, USA.

Correspondence to:

Jamil S. Samaan, MD

Karsh Division of Digestive and Liver diseases,

8700 Beverly Blvd

Los Angeles, California, USA

Email: jamil.samaan@gmail.com

Phone: (310) 423-3277

Manuscript word count: 3118

Abstract word count: 288/250

ABSTRACT

Background and Aims: Generative Pre-trained Transformer-4 (GPT-4) is a large language model

(LLM) trained on a variety of topics, including the medical literature. Nutrition plays a critical

role in managing inflammatory bowel disease (IBD), with an unmet need for nutrition-related

patient education resources. The aim of this study is to examine the accuracy and

reproducibility of responses by GPT-4 to patient nutrition questions related to IBD.

Methods: Questions were curated from adult IBD clinic visits, Facebook, and Reddit. Two IBD-

focused registered dieticians independently graded the accuracy and reproducibility of GPT-4's

responses while a third senior IBD-focused registered dietitian arbitrated. To ascertain

reproducibility, each question was inputted twice into the model. Descriptive analysis is

presented as counts and proportions.

Results: In total, 88 questions were included. The model provided correct responses to 73/88

questions (83.0%), with 61 (69.0%) graded as comprehensive. A total of 15/88 (17%) responses

were graded as mixed with correct and incorrect/outdated data. When examined by category,

the model provided comprehensive responses to 10 (62.5%) questions related to "Nutrition and

diet needs for surgery", 12 (92.3%) "Tube feeding and parenteral nutrition", 11 (64.7%)

"General diet questions", 10 (50%) "Diet for reducing symptoms/inflammation" and 18 (81.8%)

"Micronutrients/supplementation needs". The model provided reproducible responses to

81/88 (92.0%) questions.

Conclusion: GPT-4 provided comprehensive responses to the majority of questions,

demonstrating the promising potential of LLMs as supplementary tools for IBD patients seeking

nutrition-related information. However, 17% of responses contained incorrect information,

highlighting the need for continuous refinement and validation of LLMs prior to incorporation

into clinical practice. Future studies should focus on leveraging LLMs to enhance patient

outcomes. Furthermore, efforts promoting patient and healthcare professional proficiency in

using LLMs are essential to maximize their efficacy and facilitate personalized care.

Keywords: ChatGPT, GPT-4, nutrition, dietitian, IBD, gastroenterology.

INTRODUCTION

Inflammatory bowel disease (IBD), which includes Crohn's disease (CD) and ulcerative

colitis (UC), are chronic, often debilitating conditions impacting an estimated 3.1 million adults

and 70,000 children in the United States.(1,2) The global burden of IBD is also substantial, with

a rising incidence in newly industrialized nations that have adopted more Western lifestyles.(3)

Nutrition plays a pivotal role in the management of patients with IBD, not only influencing

disease activity but also impacting overall patient outcomes given the increased risk for

malnutrition and micronutrient deficiencies in this patient population. Proper nutritional

interventions, such as exclusive enteral nutrition (EEN), can alleviate symptoms, reduce the

need for medications, and promote mucosal healing, thus improving their quality of life.(4–7)

Furthermore, diet and lifestyle modifications can influence the course of these diseases by

modulating gut microbiota, inflammatory responses, and disease-related complications.(8)

The critical role of nutrition in IBD underscores a pressing need for effective patient

education in this domain. Patient surveys have consistently indicated a strong demand for

comprehensive nutrition guidance and education, both in general and within the IBD

community.(9) One study of IBD patients showed 15.4% of patients did not talk to any provider

about nutrition while 15.5% rarely entered discussion with their healthcare providers regarding

this subject. When asked the reason for the lack of discussion, 31.3% of patients felt their

provider did not have enough time during clinical visits while 20.9% reported feeling their

provider lacked the requisite expertise on the subject.(10) Moreover, studies show patients

frequently turn to external sources of information in lieu of their healthcare providers.(11,12)

More and more patients are seeking information about their disease from online sources, and

this trend is anticipated to grow.(13) Despite this demand, there is variable access to high

quality IBD-specific dietary information available to patients.(14) Furthermore, when navigating

online sources patients are confronted with both high-quality information as well as

misinformation that may be harmful, a dichotomy that often makes it challenging to distinguish

between valid medical guidance from erroneous or misleading advice.

The emergence and growing popularity of large language models (LLMs), such as

ChatGPT, may revolutionize the way information is accessed and disseminated. These models

have the potential in assisting patients with IBD by helping them navigate the complex

landscape of nutrition information. Personalized dietary recommendations can play a pivotal

role in managing IBD, and ChatGPT may offer an accessible platform for patients to receive

timely and pertinent information. While not a replacement for professional medical advice, it

can serve as a complementary resource by providing general dietary guidelines and addressing

common nutritional queries. There is a growing body of literature showing the impressive

ability of ChatGPT in answering clinical questions on a wide range of topics including cirrhosis,

bariatric surgery, laryngology and orthopaedic surgery.(15–18)

There is little to no literature examining the performance of ChatGPT or other LLMs

regarding nutrition-related questions. Examining the performance of LLMs, like ChatGPT, in the

realm of nutrition will be critical to understanding the future role and limitations of this

technology. Therefore, we examined the accuracy and reproducibility of GPT-4's responses to

nutrition questions related to IBD.

MATERIALS AND METHODS

Question Curation/Data Source

Questions were obtained from adult patient clinic visits at a tertiary academic hospital,

Facebook groups ("Crohn's and Colitis Food, recipes, Diet, and Nutrition", "Crohn's Disease &

Ulcerative Colitis Support Group", "Crohn's Disease Support Group") and Reddit pages

("r/CrohnsDisease", "r/IBD"). Questions were curated, screened, and approved by three

authors (CF, EF, KI) to evaluate their inclusion in the study. Only nutrition questions related to

IBD were included. Duplicate and similar questions from multiple sources were removed.

Questions requiring subjective or personalized responses (ex.) and questions that were vague

(ex.) were rephrased to a generic language format to allow for inclusion in the study. Other

questions were grammatically modified to eliminate ambiguity. A total of 88 questions were

included and used to generate responses from GPT-4. To better characterize GPT-4's

performance in various topics within IBD nutrition, questions were categorized into multiple

groups for statistical analysis purposes: 1. Nutrition and diet needs for surgery, 2. Tube feeding

and parenteral nutrition 3. General diet questions, 4. Diet for reducing symptoms/inflammation

and 5. Micronutrients/Supplementation needs. Approval from the institutional review board

was not sought given all questions are publicly available.

ChatGPT

ChatGPT is a LLM trained using an expansive dataset from various sources such as online

websites, books, and articles up to the year 2021, including the medical literature.(19) GPT-4, or

Generative Pre-trained Transformer 4, was released in March of 2023 and is the fourth iteration

of the GPT series. Based on a given input, the model generates responses that are east-to-

understand and conversational in nature.(20) By leveraging billions of sentences from diverse

internet sources, GPT-4 can engage in complex conversations, answer questions, write essays,

and even generate creative content like stories or poems. Its developers employed

Reinforcement Learning from Human Feedback (RLHF) to optimize the model, ensuring it

effectively responds to a wide range of commands and written inquiries. This fine-tuning was

driven by human preferences, which served as a reward signal, ensuring the model's alignment

with user intent and desired outcomes.(21) The model has been fine-tuned to resonate with

user goals and to reduce biased, or potentially harmful outputs. The exact origins of the data

used to train ChatGPT remain undisclosed. As a product of extensive research and refinement,

this model has found applications across various industries with growing literature highlighting

its potential use in the field of healthcare.

Response Generation

To produce responses, each question was prompted to the March 14[th] version of GPT-4.

Questions were entered as is and without additional prompting in order to emulate the manner

in which patients may potentially engage with the model. Every question was entered twice at

different times using the "new chat" feature, yielding two answers for each question. This

approach aimed to assess the reproducibility of responses to the same question.

Question Grading

Responses to the questions underwent an initial evaluation for both accuracy and

reproducibility by two actively practicing IBD-focused registered dietitians in a tertiary care

center (E.F., C.F.). The reviewers used information available up to 2021 as the benchmark for

assessing accuracy. Reproducibility was assessed by examining the similarity in accuracy

between the two responses provided for each question. The grading scale for response

accuracy was as follows:

- *Comprehensive:* Defined as accurate and comprehensive, nothing more a registered dietitian

would add if asked this question by a patient in clinic.

- *Correct but inadequate:* All information is correct but incomplete, a registered dietitian would

have more important information to add if asked this question by a patient in clinic.

- *Mixed with correct and incorrect/outdated data*

- *Completely incorrect*

In instances of disagreement regarding reproducibility or accuracy grading, a third

reviewer who has 10 years of experience as a clinical dietitian working with IBD patients at a

tertiary care center arbitrated (K.I.). These final grades were subsequently aggregated to gauge

GPT-4's proficiency in addressing questions related to nutrition in IBD.

Statistical Analysis

The proportions of responses earning each grade were calculated and shown as counts

and percentages. To assess reproducibility, both responses for each question were catagorized

into two categories based on the presence of incorrect information: grades comprehensive and

correct but inadequate formed the first category, while grades of mixed with correct and

incorrect/outdated data and completely incorrect made up the second. A pair of responses to a

question were deemed non-reproducible if their respective grades belonged to separate

categories. All analyses were conducted using Microsoft Excel. (version 16.69.1).

RESULTS

In total, 88 nutrition questions related to IBD were inputted into GPT-4. The model

performed well overall in terms of accuracy and provided correct responses to 73/88 questions

(83.0%), with 61 (69.0%) of responses graded as comprehensive. For example, the model

provided a comprehensive summary of EEN that is clear, concise, comprehensive, and accurate.

This included a summary of the route of administration, avoidance of consumption of solids,

nutritional content of EEN, its impact on mucosal healing, length of administration,

reintroduction of solids, and its efficacy in inducing remission particularly in children and

adolescents. Another example of a comprehensive response is an important and common

questions asked by patients regarding taking vitamins and minerals despite normal laboratory

values. The model first stressed the importance of monitoring nutrient deficiencies in patients

with IBD and the temptation by patients to discontinue these supplements when laboratory

tests reveal normal values. The model then advised the user to consult with their healthcare

professional who will consider the patient's medical history, medications, and IBD disease

status prior to making such a decision.

A total of 15/88 (17.0%) responses were graded as mixed with correct and

incorrect/outdated data while no responses were graded as completely incorrect. For example,

the model was asked about diet recommendations for IBD patients, which is one of the most

common questions encountered in the IBD clinic. The model offered various suggestions that

are not necessarily applicable to all patients with inflammatory bowel disease. For example, a

low-fat and lactose-free diet may only be suitable for a limited subset of these patients.

Furthermore, the model recommended a low residue diet, often conflated with "low fiber" diet,

which is not recognized as a diet by the Academy of Nutrition and Dietetics due to the lack of

definition and therefore inability to accurately measure from available data(22). This also

contradicts the current International Organization for the Study of IBD (IOIBD) Diet Guidelines,

which recommend patients with IBD consume more fiber. These recommendations may lead to

confusion or unnecessary nutritional restrictions for patients and ultimately to worse

outcomes. Another example is the model stating "a plant-based diet is not a guaranteed cure"

when asked if a plant-based diet is a cure for IBD. This is inaccurate as it suggests that a plant-

based diet may sometimes cure IBD.

When examined by question category, the model provided comprehensive responses to

10 (62.5%) questions related to "Nutrition and diet needs for surgery", 12 (92.3%) "Tube

feeding and parenteral nutrition", 11 (64.7%) "General diet questions", (50%) "Diet for reducing

symptoms/inflammation" and 18 (81.8%) to "Micronutrients/Supplementation needs" (Table 1,

Figure 1). The percentage of comprehensive responses was lowest among questions regarding

"Diet for reducing symptoms/inflammation" at 50% and highest among questions regarding

"Tube feeding and parenteral nutrition" at 92.3%.

Table 1: Grading of responses generated by GPT-4 to nutrition questions related to

inflammatory bowel disease stratified by question category.

Question Category and Grading Scale	Number of responses (%)
Nutrition and Diet Needs for Surgery (N=16)	
1. Comprehensive	10 (62.5)
2. Correct but inadequate	4 (25.0)
3. Mixed with correct and incorrect/outdated data	2 (12.5)
4. Completely incorrect	0
Tube Feeding and Parenteral Nutrition (N=13)	
1. Comprehensive	12 (92.3)
2. Correct but inadequate	0
3. Mixed with correct and incorrect/outdated data	1 (7.7)
4. Completely incorrect	0
General Diet Questions (N=17)	
1. Comprehensive	11 (64.7)

2. Correct but inadequate	5 (29.4)
3. Mixed with correct and incorrect/outdated data	1 (5.8)
4. Completely incorrect	0
Diet for Reducing Symptoms/Inflammation (N=20)	
1. Comprehensive	10 (50.0)
2. Correct but inadequate	2 (10.0)
3. Mixed with correct and incorrect/outdated data	8 (40.0)
4. Completely incorrect	0
Micronutrients/Supplementation Needs (N=22)	
1. Comprehensive	18 (81.8)
2. Correct but inadequate	1 (4.5)
3. Mixed with correct and incorrect/outdated data	3 (13.6)
4. Completely incorrect	0

206
207

Figure 1: Graphical illustration of grading of responses generated by GPT-4 to nutrition

questions related to inflammatory bowel disease stratified by question category.

The model provided overall high reproducibility in accuracy with 81/88 (92.0%) of

questions generating reproducible responses. When examined by question category, the model

provided reproducible responses to 100% of questions related to "Tube feeding and parenteral

nutrition" (100%), 94.1% "General diet questions", 90.0% "Diet for reducing

symptoms/inflammation", 95.4% "Micronutrients/Supplementation needs", and 81.3%

"Nutrition and diet needs for surgery" (Table 2).

Table 2. Proportion of nutrition questions related to inflammatory bowel disease with

reproducible grading of responses generated by GPT-4 stratified by question category.

Question Categories	Number of responses (%)
Nutrition and Diet Needs for Surgery (N=16)	13 (81.3%)
Tube Feeding and Parenteral Nutrition (N=13)	13 (100.0%)
General Diet Questions (N=17)	16 (94.1%)
Diet for Reducing Symptoms/Inflammation (N=20)	18 (90.0%)
Micronutrients/Supplementation Needs (N=22)	21 (95.4%)

222   Reproducibility was defined as no difference in grading categories (Comprehensive and Correct

223   but inadequate vs mixed with correct and incorrect/outdated data and completely incorrect)

224   between the two responses for each question.

DISCUSSION

The increasing global prevalence of IBD, coupled with the critical role of nutrition in its management, underscores the importance of reliable and accessible nutrition-related patient educational resources. Our study evaluated the accuracy and reproducibility of GPT-4's responses to patient nutrition questions related to IBD. The model provided accurate and reproducible responses to the majority of questions prompted, highlighting the future potential of LLMs as a complementary source of information related to nutrition for IBD patients. However, the model provided inaccurate and incomplete responses to 17% of questions, underscoring the need for further improvement in performance prior to its introduction into clinical practice. With future iterations and improvements in performance, we see this technology as a supplementary tool rather than a substitute for advice from licensed healthcare professionals.

Individuals with IBD place significant emphasis on their diet, frequently implementing self-imposed dietary limitations(23–25). Subsequently, patients frequently seek nutritional guidance and turn to external sources in lieu of their healthcare providers with one study revealing that IBD patients were most confident in advice from online sources when experiencing active disease(11,12,23). The quality and reliability of information available online are either limited or frequently inconsistent. Using conventional online search engines provides patients with an abundance of information from a variety of sources, making distinguishing reliable information both difficult and time consuming. As the disease predominantly affects young adults with high computer literacy, the potential impact of online resources, like LLMs, becomes increasingly significant. This provides urgency in examining the role of LLMs in

nutrition related patient education in order to understand the strengths and limitations of this

technology. GPT-4 provided accurate and comprehensive responses to the majority of

questions in our study, highlighting a promising potential for LLMs in the realm of patient

education. The reproducibility of GPT-4's responses in our study, which exceeded 80% across all

categories, is also noteworthy and crucial for building trust among users and ensuring

consistently accurate information. If validated, LLMs like ChatGPT may serve as an easy-to-

understand and efficient tool for patients to obtain reliable supplemental information based on

the medical literature.

Its also important to highlight the limitations of GPT-4, as seen in our study. The model

provided incorrect or outdated information in 17% of responses which demonstrates its lack of

readiness for clinical use in its current form. There are multiple possible reasons for this.

ChatGPT's training includes data from a wide range of sources, some of which may not be

accurate, leading to potential inaccuracies in its outputs. Moreover, the data sources used to

train the model have not been publicly disclosed making critical appraisal of its knowledge base

regarding a particular topic not possible. Even when accessing information from the medical

literature in its training data, the model may emphasize or utilize conflicting or outdated

information resulting in inaccurate or inconsistent outputs. Prompting strategies may also play

a role in the quality of responses. For our investigation, we pursued a pragmatic study design

where real world patient questions were curated from our IBD clinic and online sources and

subsequently inputted into GPT-4 without additional prompting. This is likely not the optimal

prompting method to produce the highest quality outputs from LLMs. We hypothesize that

more advanced prompt engineering would likely produce higher quality responses, both in

accuracy and comprehensiveness. In light of this, the design of intuitive, precise prompts for

both patients and healthcare providers may be an effective strategy in harnessing the full

potential of LLMs, as it ensures these tools deliver relevant, accurate information to specific

medical contexts. By educating both groups on effective prompt crafting, we can greatly

enhance the quality of interactions with LLMs, potentially leading to improved patient

outcomes, more efficient healthcare delivery, and a deeper integration of AI in healthcare.

If future iterations of this technology are validated for clinical use, they may have a

significant impact on the patient experience and healthcare outcomes. The timely accessibility

of healthcare providers, particularly dietitians and IBD specialists, remains a consistent hurdle

for many IBD patients.(26–28) This can not only exacerbate delays in care and result in

potentially detrimental outcomes, but also augment the challenges of patients navigating

through the sea of inaccurate or insufficient online information. These barriers may also disrupt

the proactive efforts of informed patients, leading to misguided actions based on flawed

knowledge. In the future, patients equipped with accurate and holistic knowledge from LLMs

may be able to streamline in-person appointments with more focused questions, allowing

healthcare providers to focus on patient-specific concerns.(29) Moreover, empowering patients

with more information can potentially foster a greater sense of autonomy among patients,

encouraging them to initiate discussions regarding nutrition with their healthcare providers.

This, in turn, may help promote the focus on nutrition during clinic visits, and potentially

increase the rate of referrals to dietitians.

LLMs may also help bridge healthcare disparities by democratizing access to health

information. Barriers to care, such as lengthy wait times and inaccessibility to a dietician or IBD

specialist, are even more pronounced in patients of minority backgrounds, which can further

exaggerate disparities and lead to delays in diagnosis and treatment.(30) LLMs may empower

these patients with knowledge regarding their disease, giving them a tool which helps them

proactively engage with their healthcare providers. LLMs may also prove beneficial to patients

who have language discordance with their healthcare providers, given disparities in healthcare

outcomes based on language preference have been previously shown.(31,32) ChatGPT's ability

to provide patient education in languages other than English is an active area of research with

some studies demonstrating its ability to respond to inquiries related to cirrhosis in Arabic,

Korean, Mandarin, and Spanish (33–35). LLMs can provide a valuable resource for these

patients, and potentially serve to mitigate disparities in outcomes.

302   Looking ahead, further avenues of research should aim towards ameliorating the

inaccuracies and limitations of this technology as revealed by our study. While ChatGPT can

provide answers to general IBD nutrition-related questions, it lacks the ability to

comprehensively assess individual health status, address behavioral needs, or monitor progress

over time. IBD can vary significantly from person to person in terms of symptoms, triggers, and

nutritional needs. Registered dietitians can assess individual cases, considering medical history,

current medications, symptom severity, and tailor nutrition advice to the individuals' specific

needs and goals. While the model's linguistic prowess is commendable, it can sometimes

produce responses that sound convincing but may be incorrect or nonsensical, a phenomenon

termed "hallucinations". Continuous feedback and iterative improvements are essential to

minimize these limitations. Considering the growing popularity of LLMs and their potential in

delivering patient education, it's imperative to further study and validate their utility. The role

of dieticians and physicians in managing patients with IBD remains critical to positive outcomes,

and further research should investigate ways in which collaborations between AI-models and

the healthcare team can benefit our patients.

CONCLUSION

GPT-4 provided comprehensive responses to the majority of questions, demonstrating the

promising potential of LLMs as supplementary tools for IBD patients seeking nutrition-related

information. Its important to note that 17% of responses contained incorrect information,

highlighting the need for continuous refinement and validation of LLMs prior to their

incorporation into clinical practice. Going forward, it's essential to approach the use of LLMs as

an adjunct to professional medical advice. Future studies should focus on leveraging LLMs to

enhance patient outcomes in the realm of IBD nutrition. Furthermore, efforts towards

promoting patient and healthcare professional proficiency in using LLMs are essential to

maximizing their impact and personalization.

Conflict of Interest

The authors declare that they have no conflict of interest.

Funding/Support

None.

Author contributions

J.S.S conceived and designed the study, manuscript preparation.

K.I. Question curation, response grading, edited the paper for important intellectual content.

E.F. Question curation, response grading, edited the paper for important intellectual content.

C.F. Question curation, response grading, edited the paper for important intellectual content.

W.N. Data management, statistical analysis, edited the paper for important intellectual content.

N.R. Assisted with manuscript preparation, edited the paper for important intellectual content.

B.H. Edited the paper for important intellectual content.

Y.Y. Edited the paper for important intellectual content.

E.V. Edited the paper for important intellectual content.

Ethical Approval

This article does not contain any studies with human participants or animals performed by any

of the authors.

Informed Consent Statement

Informed Consent does not apply.

Declaration of AI and AI-assisted technologies in the writing process

During the preparation of this work the authors used GPT-4 in order to improve readability and

language. After using this tool/service, the authors reviewed and edited the content as needed

and take full responsibility for the content of the publication.

REFERENCES

1. Xu F, Dahlhamer JM, Zammitti EP, Wheaton AG, Croft JB. Health-Risk Behaviors and Chronic

Conditions Among Adults with Inflammatory Bowel Disease — United States, 2015 and

2016. MMWR Morb Mortal Wkly Rep. 2018 Feb 16;67(6):190–5.

2. Rosen MJ, Dhawan A, Saeed SA. Inflammatory Bowel Disease in Children and Adolescents.

JAMA Pediatr. 2015 Nov 1;169(11):1053.

3. Ng SC, Shi HY, Hamidi N, Underwood FE, Tang W, Benchimol EI, et al. Worldwide incidence

and prevalence of inflammatory bowel disease in the 21st century: a systematic review of

population-based studies. The Lancet. 2017 Dec;390(10114):2769–78.

4. Bischoff SC, Escher J, Hébuterne X, Kłęk S, Krznaric Z, Schneider S, et al. ESPEN practical

guideline: Clinical Nutrition in inflammatory bowel disease. Clinical Nutrition. 2020

Mar;39(3):632–53.

5. Richman E, Rhodes JM. Review article: evidence-based dietary advice for patients with

inflammatory bowel disease. Aliment Pharmacol Ther. 2013 Nov;38(10):1156–71.

6. Lee D, Albenberg L, Compher C, Baldassano R, Piccoli D, Lewis JD, et al. Diet in the

Pathogenesis and Treatment of Inflammatory Bowel Diseases. Gastroenterology. 2015

May;148(6):1087–106.

7. Whelan K, Murrells T, Morgan M, Cummings F, Stansfield C, Todd A, et al. Food-related

quality of life is impaired in inflammatory bowel disease and associated with reduced intake

of key nutrients. The American Journal of Clinical Nutrition. 2021 Apr;113(4):832–44.

8. Lewis JD, Abreu MT. Diet as a Trigger or Therapy for Inflammatory Bowel Diseases.

Gastroenterology. 2017 Jan;152(2):398-414.e6.

9. Limdi JK, Aggarwal D, McLaughlin JT. Dietary Practices and Beliefs in Patients with

Inflammatory Bowel Disease: Inflammatory Bowel Diseases. 2016 Jan;22(1):164–70.

10. Tinsley A, Ehrlich OG, Hwang C, Issokson K, Zapala S, Weaver A, et al. Knowledge, Attitudes,

and Beliefs Regarding the Role of Nutrition in IBD Among Patients and Providers:

Inflammatory Bowel Diseases. 2016 Oct;22(10):2474–81.

11. Bernstein KI, Promislow S, Carr R, Rawsthorne P, Walker JR, Bernstein CN. Information

needs and preferences of recently diagnosed patients with inflammatory bowel disease:

Inflammatory Bowel Diseases. 2011 Feb;17(2):590–8.

12. Limdi JK, Butcher RO. Information resources and inflammatory bowel disease: Inflammatory

Bowel Diseases. 2011 Aug;17(8):E89–90.

13. Cima RR, Anderson KJ, Larson DW, Dozois EJ, Hassan I, Sandborn WJ, et al. Internet use by

patients in an inflammatory bowel disease specialty clinic: Inflammatory Bowel Diseases.

2007 Oct;13(10):1266–70.

14. Prince AC, Moosa A, Lomer MCE, Reidlinger DP, Whelan K. Variable access to quality

nutrition information regarding inflammatory bowel disease: a survey of patients and

health professionals and objective examination of written information. Health Expect. 2015

Dec;18(6):2501–12.

15. Yeo YH, Samaan JS, Ng WH, Ting PS, Trivedi H, Vipani A, et al. Assessing the performance of

ChatGPT in answering questions regarding cirrhosis and hepatocellular carcinoma. Clin Mol

Hepatol [Internet]. 2023 Mar 22 [cited 2023 Apr 5]; Available from: http://www.e-

cmh.org/journal/view.php?doi=10.3350/cmh.2023.0089

16. Samaan JS, Yeo YH, Rajeev N, Hawley L, Abel S, Ng WH, et al. Assessing the Accuracy of

Responses by the Language Model ChatGPT to Questions Regarding Bariatric Surgery. OBES

SURG [Internet]. 2023 Apr 27 [cited 2023 May 29]; Available from:

https://link.springer.com/10.1007/s11695-023-06603-5

17. Lechien JR, Georgescu BM, Hans S, Chiesa-Estomba CM. ChatGPT performance in

laryngology and head and neck surgery: a clinical case-series. Eur Arch Otorhinolaryngol

[Internet]. 2023 Oct 24 [cited 2023 Oct 28]; Available from:

https://link.springer.com/10.1007/s00405-023-08282-5

18. Kaarre J, Feldt R, Keeling LE, Dadoo S, Zsidai B, Hughes JD, et al. Exploring the potential of

ChatGPT as a supplementary tool for providing orthopaedic information. Knee Surg Sports

Traumatol Arthrosc. 2023 Nov;31(11):5190–8.

19. openai. ChatGPT: Optimizing Language Models for Dialogue. 2023;

https://openai.com/blog/chatgpt/. Accessed 1/1/2023, 2023.

20. OpenAI. GPT-4 Technical Report. 2023 [cited 2023 Aug 11]; Available from:

https://arxiv.org/abs/2303.08774

21. Ouyang L, Wu J, Jiang X, Almeida D, Wainwright CL, Mishkin P, et al. Training language

models to follow instructions with human feedback. 2022 [cited 2023 Feb 10]; Available

from: https://arxiv.org/abs/2203.02155

22. Cunningham E. Are Low-Residue Diets Still Applicable? Journal of the Academy of Nutrition

and Dietetics. 2012 Jun;112(6):960.

23. Larussa T, Suraci E, Marasco R, Imeneo M, Abenavoli L, Luzza F. Self-Prescribed Dietary

Restrictions are Common in Inflammatory Bowel Disease Patients and Are Associated with

Low Bone Mineralization. Medicina. 2019 Aug 20;55(8):507.

24. Marsh A, Kinneally J, Robertson T, Lord A, Young A, Radford –Smith G. Food avoidance in

outpatients with Inflammatory Bowel Disease – Who, what and why. Clinical Nutrition

ESPEN. 2019 Jun;31:10–6.

25. Godala M, Gaszyńska E, Durko Ł, Małecka-Wojciesko E. Dietary Behaviors and Beliefs in

Patients with Inflammatory Bowel Disease. JCM. 2023 May 14;12(10):3455.

26. Habashi P, Bouchard S, Nguyen GC. Transforming Access to Specialist Care for Inflammatory

Bowel Disease: The PACE Telemedicine Program. Journal of the Canadian Association of

Gastroenterology. 2019 Oct 9;2(4):186–94.

27. Borren NZ, Conway G, Tan W, Andrews E, Garber JJ, Yajnik V, et al. Distance to Specialist

Care and Disease Outcomes in Inflammatory Bowel Disease: Inflammatory Bowel Diseases.

2017 Jul;23(7):1234–9.

28. Malter L, Jain A, Cohen BL, Gaidos JKJ, Axisa L, Butterfield L, et al. Identifying IBD Providers'

Knowledge Gaps Using a Prospective Web-based Survey. Inflammatory Bowel Diseases.

2020 Aug 20;26(9):1445–50.

29. Bickmore TW, Pfeifer LM, Paasche-Orlow MK. Using computer agents to explain medical

documents to patients with low health literacy. Patient Education and Counseling. 2009

Jun;75(3):315–20.

30. Liu JJ, Abraham BP, Adamson P, Barnes EL, Brister KA, Damas OM, et al. The Current State of

Care for Black and Hispanic Inflammatory Bowel Disease Patients. Inflammatory Bowel

Diseases. 2023 Feb 1;29(2):297–307.

31. Al Shamsi H, Almutairi AG, Al Mashrafi S, Al Kalbani T. Implications of Language Barriers for

Healthcare: A Systematic Review. Oman Med J. 2020 Mar;35(2):e122.

32. Divi C, Koss RG, Schmaltz SP, Loeb JM. Language proficiency and adverse events in US

hospitals: a pilot study. Int J Qual Health Care. 2007 Apr;19(2):60–7.

33. Yeo YH, Samaan JS, Ng WH, Ma X, Ting PS, Kwak MS, et al. GPT-4 outperforms ChatGPT in

answering non-English questions related to cirrhosis [Internet]. Gastroenterology; 2023

May [cited 2023 Jul 29]. Available from:

http://medrxiv.org/lookup/doi/10.1101/2023.05.04.23289482

34. Zhu Z, Ying Y, Zhu J, Wu H. ChatGPT's potential role in non-English-speaking outpatient clinic

settings. DIGITAL HEALTH. 2023 Jan;9:20552076231184091.

35. Samaan JS, Yeo YH, Ng WH, Ting PS, Trivedi H, Vipani A, et al. ChatGPT's ability to

comprehend and answer cirrhosis related questions in Arabic. Arab Journal of

Gastroenterology. 2023 Sep;S1687197923000588.