Social skills group training (KONTAKT[TM]) for Chinese autistic children: a feasibility wait-list RCT protocol

Authors and Affiliations

Uchong Lao[1□] · Huilin Zhu[1□*] · Fengjing Liang[1] · Wuxia Bai[12] · Peipei Yin[1] · Xiaoqian Huang[1] · Sonya Girdler[2345*] · Sven Bölte[234*] · Xiaobing Zou[1*]

[1] Child Development and Behavior Center, The Third Affiliated Hospital, Sun Yat-Sen University, China, 2693 Kai Chuang Avenue, Huangpu District, Guangzhou, 510530, China

[2] School of Allied Health, Curtin University, Kent Street, Bentley, Perth, WA, 6102, Australia

[3] Curtin Autism Research Group (CARG), Curtin University, Perth, WA, Australia

[4] Center of Neurodevelopmental Disorders (KIND), Centre for Psychiatry Research, Division of Neuropsychiatry, Department of Women's and Children's Health, Karolinska Institutet & Child and Adolescent Psychiatry, Stockholm Health Care Services, Stockholm County Council, Stockholm, Sweden

[5] School of Allied Health, University of Western Australia, Perth, WA, Australia

[□] Equal contributions

[*] Corresponding authors

ORCID ID:

Uchong Lao: 0000-0001-8658-880X

Huilin Zhu: 0000-0002-1052-708X

Sonya Girdler: 0000-0001-7992-0800

Sven Bölte: 0000-0002-4579-4970

Corresponding authors information:

Huilin Zhu

Email: zhuhlin6@mail.sysu.edu.cn

Sonya Girdler

Email: sonya.girdler@curtin.edu.au

Sven Bölte

Email: sven.blote@ki.se

Xiaobing Zou

Email: zouxb@mail.sysu.edu.cn

Abstract

Introduction

Challenges in social communication and interaction are common among school-aged autistic children in mainstream society settings. Although social skills group training (SSGT) is effective in promoting positive psycho-social functioning outcomes in autistic children, there is no standardised evidence-based SSGT in Mainland China to date. Therefore, the main aim of this study is to evaluate the feasibility and acceptability of the culturally-adapted 16-session version of the KONTAKT[TM] programme in Chinese autistic children. In addition, this study will provide preliminary effectiveness and cost-effectiveness data on the 16-session Chinese version of KONTAKT[TM].

Methods and analysis

This is a randomised, single-blinded, wait-list controlled feasibility study with 36 autistic children aged 8−12 years with Intelligence Quotient over 70. They will be assigned to either the immediate training group (ITG) or the delayed training group (DTG) based on gender. Feasibility will be measured by the mixed-methods data of acceptability and adherence of participants, parents, and providers. Preliminary effectiveness and cost-effectiveness will be assessed by quantitative data collected at five-time points. The main outcome is the improvement in social skills as measured by the Contextual Assessment of Social Skills (CASS). Other outcomes include autism traits, adaptive functioning, self-assertiveness, psycho-social functioning, parental reflective functioning, and school support. Common process factors and their effects on outcomes will also be explored.

Cost-effectiveness will consider both societal and healthcare perspectives.

Ethics and dissemination

The current study protocol has been reviewed and ethics approval obtained from the Ethical

Board Committee at the Third Affiliated Hospital of Sun Yat-sen University (II2023-119-01). The

trial has pre-registered in Chinese Clinical Trials (ChiCTR2300072136) on 05 June 2023. The

results of this trial will be actively disseminated through manuscript publications and conference

presentations.

Keywords

Social skills group training; Children; KONTAKT[TM]; Cross-cultural adaptation; Feasibility; Autism;

Chinese

Strengths and limitations of this study

● Comprehensive assessment via diverse outcome acquisition methods and multiple informants.

● Systematic cultural adaptation using the Integrative Cultural Adaptation Framework (ICAF)

ensures the appropriateness of the Chinese version of KONTAKT[TM].

● Collect data on readiness for change, providing insights into "help-seeker" effects.

● Limited generalisability due to a single study site, location-specific inclusion, and absence of

active comparator.

● Uncertain psychometric properties of certain measures due to a lack of Chinese population

data.

Introduction

Background and rationale

Autism[1] is a neuro-developmental condition that is characterised by persistent challenges in reciprocal social interaction and communication [1]. As a result, autistic children often struggle to fulfill the expected social norms, limiting their maximum functional capacity in everyday contexts [2]. The estimated prevalence of autism among children aged 6 to 12 in Mainland China is 0.7% (~ 1 in 143 children), with nearly half of the identified cases being enrolled in mainstream schools [3]. This data underscores the need for evidence-based interventions to enhance the social competence and well-being of cognitively capable Chinese autistic children.

Among interventions for verbal and cognitively capable school-aged autistic children, Social Skills Group Training (SSGT) stands out as an evidence-based group approach [4, 5, 6]. Multiple meta-analyses of SSGT programmes in randomised controlled trials (RCTs) consistently report moderate to large d-type effects in achieving personally meaningful social goals, enhancing social skills, social skills knowledge, social participation, and social functioning [4, 7, 8, 9].

However, the generalisation of SSGT effects remains undefined, primarily due to over-reliance on parent ratings and a lack of blinded observation [4, 7]. Results from previous meta-analyses [4, 7, 10, 11] inferred that improvements observed in parent and clinical ratings were primarily related to increased social skills knowledge or specific task-related skills, rather than real-life social improvements [7].

---

[1] Herein, identity-first language is used since this is the endorsed terminology among autistic advocates (Kenny, 2016).

Another conspicuous limitation is the limited input from autistic children themselves.

Understandably, up to 50% autistic population struggles with understanding and interpreting

emotion states, scilicet co-occurring alexithymia, making it challenging to gather their perspectives

[12]. Additionally, autistic children may encounter comprehension challenges in emotional

vocabulary and Likert scale metrics, raising concerns about the credibility of self-report measures

[13]. However, recent meta-analysis findings suggest that emotional self-awareness in school-aged

autistic children is comparable to their peers despite the high heterogeneity between the included

studies [14]. Furthermore, applying validated non-verbal psychometric measurements, such as

projection tests (Draw-A-Person Test [DAPT], [15, 16]), may be conducive to exploring autistic

children's mental states [17, 18, 19].

Another discernible limitation in the past SSGT studies was the sparseness of the

investigations in common process factors [20]. While a significant proportion of SSGT studies have

reported programme adherence and satisfaction [7], the data of these studies were descriptive. In

other words, they did not systematically examine these process factors about outcomes [20].

KONTAKT^{TM} programme is a well-established SSGT designed for children and adolescents

with neurodivergent conditions [21]. Originating in Germany [22], it has now been further

developed, standardised, culturally adapted, and evaluated in Sweden [23, 24] and Australia [25].

Three session-length variants (24-session, 16-session, and 12-session) have been standardized and

evaluated through RCTs, demonstrating efficacy in improving participants' social functioning,

adaptive functioning, emotional well-being, emotion recognition and regulation,

personal-meaningful goals, and gaze fixation [23, 24, 25]. Both quantitative and qualitative data

support KONTAKT^{TM} as an effective, safe, and well-received program with trivial adverse events

[23, 24, 25, 26, 27].

To our knowledge, there is no culturally-standardised SSGT program for school-aged autistic

children in Mainland China. KONTAKT^{TM}, with its diverse, individualised, and naturalistic session

design [21], is envisaged to effectively cultivate the social functioning and well-being of Chinese

autistic children, provided that due consideration is taken on Chinese cultural and social norms [28].

Cultural adaptation involves systematically modifying an evidence-based intervention to align

with the target population's needs, values, languages, conditions, and customs [29, 30, 31].

However, only a few cultural adaptation studies on SSGT have well-documented and

theoretically-based methodologies. Moreover, past SSGT studies often lack input from the target

population, despite language translation and the inclusion of culturally relevant content [6].

Ignoring cultural differences can significantly hamper the effectiveness of evidence-based

SSGTs for autism [28]. Since the core characteristics of autism involve social interaction and

communication, social-ecological variations between cultures will predominantly influence the

definition between "functioning" and "disorder" regarding autism [32]. These differences also affect

collaboration among various social service providers, including healthcare, schools, and

communities [33, 34, 35]. For example, Chinese parents are less likely to disclose their child's

autism diagnosis to the child and school, reflecting limited family-school collaboration [36].

Various cultural adaptation frameworks are available [29, 30, 37, 38], sharing similar steps

despite some methodological distinctions. This study adopts the integrative cultural adaptation

framework (ICAF) since it aligns with consensus in the field [29]. The ICAF involves five stages:

information gathering, preliminary design, preliminary testing, refinement, and final trial. This

study focuses on the first four stages to prepare for the final trial (the last stage).

Objectives and hypotheses

The overarching purpose of the present study is to assess the feasibility and cross-cultural

validity of the adapted 16-session version of KONTAKT^{TM} programme for Chinese autistic children

in preparation for a larger RCT trial. The specific objectives are:

1. Evaluating the integration of KONTAKT^{TM} into a tertiary hospital's routine activity;

2. Assessing the acceptability and feasibility of data collection;

3. Examining preliminary quantitative and qualitative effects of KONTAKT^{TM}; and,

4. Exploring the impact of common process factors on outcomes.

Our primary hypothesis is that the Chinese 16-session KONTAKT^{TM} version will be feasible

and culturally appropriate for Chinese autistic children with minor revisions, implemented

successfully in a tertiary clinic. We also hypothesize that participants in the immediate training

group (ITG), in comparison to the delayed training group (DTG) [wait-list], will experience greater

improvements in outcomes from Time 1 to Time 3 (waiting period of DTG). We expect that

participants and their parents with higher readiness for change and a stronger alliance in the ITG

will benefit more from KONTAKT^{TM}. Lastly, we anticipate that the training effects in the ITG will

persist at follow-up, and significant improvements in children's outcomes will occur in the DTG

from Time 3 to Time 5 (training endpoint of the DTG).

Trial Design

This study was approved by the Ethical Board Committee at the Third Affiliated Hospital of Sun Yat-sen University (II2023-119-01) and preregistered in Chinese Clinical Trials (ChiCTR2300072136).

Any modification to this protocol will be reported to the ethical board and updated in Chinese Clinical Trials. Following Thabane and Lancaster's guidelines in 2019 [39], the Chinese KONTAKT^{TM} CHILD protocol is planned and reported following the Standard and Protocol Items: Recommendations for Interventional Trials (SPIRT) [40]. The findings of this study will be reported adherent to the Consolidated Standards of Reporting Trials for social and psychological interventions (CONSORT-SPI) [41].

The Chinese KONTAKT^{TM} CHILD feasibility study has a prospective, wait-list controlled, single-site, stratified, randomised design. Blinding applies to data collectors, outcome adjudicators, and data analysts. Three major assessment time-points (ITG: Pre-Post-Follow Up, DTG: Pre-Pre-Post), with an additional mid-point cost and processing factors data collection for each group, are included. Measurements will be taken in person except for data from teachers and prescreening processes. Moreover, this study will employ a mixed-methods design, with between-group and within-participant comparisons, to explore the feasibility and acceptability of implementing a large-scale RCT. Randomisation is performed at a 1:1 allocation ratio, with variable block length and gender stratification.

Methods: participants, interventions, and outcomes

Study setting

All in-person assessments and interventions will take place at the outpatient department of the Child Development and Behaviour Centre (CDBC) of the tertiary academic hospital named the Third-affiliated Hospital of Sun Yat-sen University in Guangzhou, China. The CDBC, one of China's first established and leading developmental behaviour paediatric units, has over 20 years of experience diagnosing and promoting well-being on autistic populations. On the other hand, online assessments will be designed and distributed using the SurveyStar online survey platform (https://wjx.cn/).

The KONTAKT[TM] certified trainers are experienced CBDC clinicians (clinical psychologists, developmental-behavioural paediatricians, and therapists) working with Chinese autistic children (with 3-10 years working experiences). They have completed standardised online KONTAKT[TM] methodological training, including self-directed training and video-conference seminars, followed by a 13-session enhanced in-person training..

KONTAKT[TM] sessions, except for the excursion session, will occur in a spacious, well-lit room where participants and trainers form a circle. The room will also include a flipchart for session notes and a blackboard for group rules and session agenda. Excursions will be held at nearby cafes, bubble tea shop,s or convenience stores. Parent sessions will take place in a separate room with a projector for lectures, following a similar circle setup.

Recruitment

Figure 1 displays the flow of participants throughout the study. Thirty-six autistic children will
198

be recruited through the outpatient department of the CDBC. Recruitment avenues will be multiple,
199

which include word-of-mouth, referral by clinicians, posters with study information displayed at the
200

CDBC clinic, and advertising articles published on the WeChat[©] Official account
201

(https://www.wechat.com/).
202

Parents will complete an online registration questionnaire if they and their children desire to
203

participate. They will provide their children's demographic information, diagnostic statuses,
204

previous evaluations on the Wechsler Intelligence Scale for Children-IV Chinese version (WISC-IV)
205

[42], the Autism Diagnostic Observation Schedule-Generic (ADOS-G) [43], and screen for
206

behaviour problems using the Childhood Behavior Checklist-parent version (CBCL) [44], and their
207

contact information. Parents should expect to spend approximately 10-15 minutes completing the
208

questionnaire. Applying the step of online registration can facilitate the analysis of demand, and
209

research coordinators can monitor recruitment.
210

After parents submit the registration questionnaire, our research coordinators will assess
211

candidate eligibility. Prequalified candidates' parents will then undergo a 10-15 minute prescreening
212

phone interview, discussing their children's school lives, academic performances, and special needs,
213

and addressing any study-related questions. The interviews will also verify the candidate's cognitive
214

and verbal abilities for the course materials.
215

Figure 1 Chinese KONTAKT[TM] Child feasibility study flowchart.

If fulfilling the prescreening criteria, candidate participants and their parents will come to the

clinic for a 50-minute face-to-face intake interview with the research coordinator. During the intake

interview, both children and parents will each have about a 20-minute individual interview with the

research coordinator. If the family meets the eligibility criteria and agrees to participate, an written

informed consent form will be obtained.

Eligibility criteria

Inclusion criteria

(1) Aged 8-12 years;

(2) A clinical diagnosis of autism (confirmed by ADOS-G, module 3);

(3) A Full-scale IQ > 70 on the WISC-IV;

(4) Resided in Guangzhou city for the past 12 months.

Exclusion criteria

(1) Significant physical impairment or severe metabolic disease(s);

(2) Significant rule-breaking or aggressive behaviours within the last six months, as confirmed

by CBCL;

(3) Low intrinsic motivation to participate;

(4) Significant difficulties with Chinese characters reading comprehension or Mandarin

expression and comprehension;

(5) Prior or current behaviour or psychiatric conditions that could disrupt participation or

necessitate other treatment(s).

Participants with common co-occurring neurodevelopmental conditions in autistic individuals,

such as attention-deficiency hyperactivity disorder, depression, anxiety, and tic disorder, are eligible

if these conditions do not significantly interfere with participation. They can continue their regular

pharmacological treatments, psychological therapies, or interventions while participating in

KONTAKT^{TM}. Written informed consent will be obtained from participants and parents after a

thorough explanation of the study during the intake interview..

Intervention

KONTAKT^{TM} Intervention

Before the completion of this protocol, a preliminary adaptation of KONTAKT^{TM} (The first

step of the ICAF - Information gathering) has been conducted. The KONTAKT^{TM} programme

includes workbooks for trainers, children, and parents, ensuring intervention fidelity. To adapt the

16-session variant of KONTAKT^{TM} from Australian English to Mandarin Chinese, we followed a

rigorous process. First, we translated the materials and verified accuracy among authors. Then, a

workgroup of KONTAKT^{TM} trainers, who are also CDBC clinicians, provided feedback. This step

had led to a significant modification on the discussing topic regarding autism diagnosis in a more

general and implicit way (session 3). In addition, to facilitate parents discussing the autism

diagnosis with their child, information relating to "how to talk about the diagnosis with your child"

was added in the first parent session. This adaptation was made because study shows that over 60%

of autistic children do not notice their autism diagnoses, with 40% of parents of autistic children

prefer to remain the diagnosis undisclosed to their autistic children in China. Additionally, parents

received support in helping their children with homework, including role-play exercises and

demonstration videos. We included session-by-session feedback and progress reports for parents, as

Chinese parents often require more social coaching support. We modified and added questions,

games, activities, and scenarios in line with Chinese cultural norms. We also created a Chinese

name for KONTAKT[TM], emphasizing meaningful socialization (交得益). Finally, a family with an

autistic adolescent proofread the manual to enhance language fluency and localise terminology.

Additionally, we customised the logo and stationary to foster group cohesion among participants.

The KONTAKT[TM] programme integrates Cognitive Behaviour Therapy principles and settings

tailored for autistic children (see **Figure 2**) [21], including psychoeducation, observational learning,

modeling, behavioural activation, homework, computerised cognitive training, rehearsal, structured

agenda, and parental involvement. It employs mandatory, recurring, and variable activities to foster

participants generalising their personally meaningful social goals. Sessions aim to enhance

understanding of social norms, develop adaptive socio-emotional skills, and promote practical

applications. The first 11 sessions have consistent themes and assignments, while later sessions are

more individualised.

Figure 2 Elements, structures, and goals of KONTAKT[TM].

A group of 4-6 participants will attend 16 consecutive sessions of KONTAKT^{TM} led by 2-3

trainers over weekends, with each session lasting 60 minutes. Three parent sessions will take place

at the beginning, middle, and end of the training, offering individual feedback on their children's

progresses, autism knowledge and discussing autism diagnosis with their children, and facilitating

parents as social coaches for their children. Parents can ask questions, share feedback, and provide

mutual support during parent sessions. Parents are encouraged to assist with homework, and trainers

will offer children's session-by-session feedback after class. Refer to **Supplementary Table 1**

(**online supplemental file 1**) for the structure and content of the KONTAKT^{TM} Chinese version.

To safeguard the fidelity of the KONTAKT^{TM} treatment protocol, trainers will complete

session-by-session fidelity checklists and receive supervision from certified KONTAKT^{TM} trainers.

Description on the rationales of adapting the Chinese 16-session KONTAKT^{TM}

The reasons for choosing the 16-session KONTAKT^{TM} variant are four-fold. Firstly, while the

RCT efficacy study from Australia indicates that four additional sessions to the 12-session variant

of KONTAKT^{TM} showed no substantial improvement on autistic traits [25], the feasibility and

efficacy of the 16-session variant remain unestablished in autistic children. Additionally, previous

research found that the 12-session variant did not significantly outperform standard care [23], and

the 24-session variant had a higher attrition rate (26% vs 11%/14%) [23, 24, 25]. Thirdly, aligning

with the Chinese academic year structure, which consists of two terms with approximately five

months each, the 16-session version is more practical. Lastly, to ensure ethical and practicality, we

aim for a 4-5 month waiting period in the wait-list group, as extended waiting times can impact

adherence negatively [45].

Explanation for the choice of comparators

296   The waiting period of the DTG is limited to 4-5 months, roughly equivalent to one round of

297   KONTAKT^{TM} training. While an active comparator in an RCT is typically the most robust, due to

298   limited clinical resources in Mainland China, a wait-list control group is considered more practical.

299   Moreover, given that "non specific" care is common in Mainland China, a wait-list control group

300   allows for assessing the additional impact of KONTAKT^{TM} on usual care.

Intervention adherence

We will use a participant attendance sheet to track compliance, aiming for at least 80%

attendance (13 out of 16 sessions) as acceptable. We also record homework completion, with a

target of over 60% considered tolerable.

We adopt several approaches recommended by Robiner to enhance the wait-list group's

adherence [46]:

(1) Provide detailed study information during intake interviews;

(2) Emphasize the importance of adherence;

(3) Garner trust with families during intake interviews;

(4) Offer face-to-face assessments for regular clinical visits;

(5) Minimise waiting time to 4-5 months with a single-round intervention;

(6) Schedule appointments at convenient times.

Randomisation

After the initial visit to the clinic, the participant will be randomised to either ITG or DTG.

Randomisation will be conducted by a third party not involved in the Chinese KONTAKT^TM

CHILD trial using computer-generated random numbers. The permuted block numbers of 4-6 will

be used to restrict randomisation within the strata (gender). Each participant will have a unique code

to denote.

Assessments/measures

Outcomes from multiple informants (children, parents, teachers, clinical ratings) and multiple

means (questionnaires, interviews, observations, projective tests) will be acquired. The assessment

timeline is presented in **Table 1**.

[Table 1]

All online questionnaire fields are mandatory, eliminating the expectation of missing data. QR

codes will be used for distributing the registration and teacher questionnaires, completed via the

WeChat^© app. Each teacher questionnaire will assign unique research serial numbers for participant

identification.

The primary outcome of the Mandarin KONTAKT^TM CHILD study is social communication

and interaction skills improvement measured by the Contextual Assessment of Social Skills (CASS),

an observational measure. *Secondary outcomes* include autistic characteristics (Social

Responsiveness Scale-2 [SRS-2]), generalised social skills (Snack Time Assessment of

Relationships and Social Skills [STARSS]), adaptive functioning (Adaptive Behaviour Assessment

System-II), self-esteem and self-concept (Self-concept Scale Draw-A-Person [SCS-DAP], Child

Rosenberg Self-Esteem Scale [CRSES]), cognitive development and psychosocial well-being

(Koppitz Draw-A-Person Test, Child Outcome Rating Scale), perceived school support school level,

and parental reflective functioning. *Feasibility outcomes* include acceptability, appropriateness, and

negative effects from quantitative data (fidelity checklists, recruitment rates, attendance rates,

homework completion rates, Group Session Rating Scale, Theoretical Framework of Acceptability

Questionnaire, and Negative Effects Questionnaire) and qualitative data from focus groups

post-intervention. *Common process factors measurements* include treatment alliance (Therapeutic

Alliance Scales for Children) and readiness for change (Preparedness for Participating in

KONTAKT^{TM} Scale). *Cost-effectiveness outcomes* include healthcare consumption and productivity

loss (the tailored version of the Trimbos/iMTA questionnaire for patients with a psychiatric disorder

child version) and health-related quality of life (Child Health Utility 9D, EQ-5D-Y).

For details of these measures, see **online supplemental file 2**.

Retention

Assessment appointments will offer flexibility within the time frame, with reminders from

research assistants. Paper questionnaires will be provided for those unable to have in-person

assessments, with families returning them by post. All non-retention events will be documented and

reported.

Data management

In the Chinese KONTAKT^{TM} CHILD feasibility study, all paper data will be converted to

electronic format. Trained research assistants will input the data within two weeks after each

assessment. Both paper and electronic data will be stored in EpiData [47] with double-entry for data

integrity. Checks for referenced range values, valid values, and regular data consistency will be

applied. Non-numeric data, including open-ended questions and notes, will be entered as originally

written.

Video and audio recordings will be labeled with unique serial names, and their details recorded

in the EpiData repository. Paper documents will be securely stored in locked filing cabinets at the

clinic. Electronic textual data will be password-protected. Video and audio files will have dual

backups on encrypted hard disks. Only the principal investigator will have full data access.

Incremental data and recording files will be back-up daily.

Sample size

The primary outcome measure is CASS. Previous SSGT studies applying the CASS suggested

a medium effect size (ES) of 0.42 [48]. Additionally, a rough average ES of 0.40, as measured by

the SRS-2, is derived from the three prior RCT studies evaluating KONTAKT^{TM} [23, 24, 25],

informs the expected ES for the future Chinese KONTAKT^{TM} CHILD main trial. According to

Whitehead et al., a feasibility study with a sample size of 15 per treatment arm allows for detecting

a medium EF (0.3~0.7) with 90% power and two-sided 5% significance in the main trial [49].

Accounting for a 15% attrition rate from previous KONTAKT^{TM} evaluation studies, we require a

sample size of $n = 18$ per arm ($N = 36$ in total).

Statistical analysis

Statistical methods for primary and secondary outcomes

The primary outcome for this study is social-communication skills improvement as measured

by the CASS, with the follow-up being the primary endpoint. The analysis will follow the

intent-to-treat (ITT) principles [50], and all data from every participant will be analysed. Missing

data will be handled based on established guidelines for each measure or CONSORT-SPI statements.

No interim analysis will occur.

Data will be checked for normality and variance heterogeneity. Baseline demographics will be

compared between ITG and DTG using t-tests or Wilcoxon rank test (continuous variables) and

chi-square tests (categorical variables) to verify group comparability. Random mixed effects

regression models, which have the advantage of handling missing data, will compare ITG and DTG

for primary and secondary analyses, considering time (baseline, endpoint, follow-up), group (ITG,

DTG), age, gender, IQ, co-occurring conditions, process factors (PPKS, GSRS at baseline), and th

interaction variable (time × group). Participants' ID will be a random effect to account for

intra-individual correlations. Intervention effects will be reported as least-square mean differences.

ES will be reported by dividing the groups' mean difference (baseline to endpoint and baseline to

follow-up) by the pooled standard deviations, with positive values favoring intervention.

Feasibility measures will be reported descriptively. Additionally, qualitative data will be

transcribed verbatim, de-identified, then imported into Nvivo 12 for theme-based analysis.

Health economic analyses analysis

Cost-effectiveness analysis (CEA) will follow an ITT basis, considering societal and

healthcare perspectives. Cost differences will be statistically tested, while resource usage

differences will not be compared to prevent false positives. Quality-adjusted life year (QALY) will

be calculated from EQ-5D-Y and CHU9D scores using validated algorithms for Chinese children

and adolescents [51, 52]. Given potential non-normality in cost data, a non-parametric random

forest permutation model will handle missing data. The incremental cost-effectiveness ratio (ICER)

will be calculated as Δcost/ΔQALYs. Cost differences between ITG and DTG at different time

points will be analysed via linear regression with bootstrapping for 5,000 repetitions to generate

non-parametric confidence intervals (See **online supplemental file 3** for CEA details).

Discussion

The study commenced recruitment on June 5, 2023. This article presents the design of the

Chinese KONTAKT[TM] CHILD trial, a feasibility wait-list RCT featuring the Chinese 16-session

KONTAKT[TM] variant for autistic children. This design incorporates multiple data sources and

diverse outcome assessment methods, facilitating a comprehensive exploration of outcome data

feasibility from different perspectives. Furthermore, a systematic cultural adaptation process

ensures the suitability, acceptability, and comprehensiveness of the Chinese KONTAKT[TM] version.

Adhering to the ICAF framework, this study will guide the refinement of the Chinese

KONTAKT[TM] version and future larger-scale evaluations, addressing the question of whether SSGT

can be effectively implemented for Chinese school-aged autistic children.

While there are several strengths in the study design mentioned earlier, it's important to

acknowledge some limitations. Firstly, the study's single-site nature and its focus on autistic

children in Guangzhou city may limit the generalisability of the findings. Additionally, certain

measures planned for use in the study, such as the CRSES and the SCS-DAP, may have uncertain

psychometric properties due to the lack of Chinese population data. However, this feasibility study

aims to provide empirical data on the psychometrics of these measures. Another limitation is the

absence of an active comparator, as the wait-list control could potentially overestimate treatment

effects [53]. Nevertheless, this aligns with the prevailing situation in China. Furthermore, the study

will collect data on parents' and children's readiness for change, allowing for an exploration of the

"help-seeker" effects, following the recommendation by Cunningham and colleagues in 2013 [53].

As mentioned above, the final stage of the ICAF involves running a full-scale RCT trial to

evaluate the efficacy of the Chinese version of KONTAKT[TM]. Therefore, upon completing this

feasibility study, the investigators will integrate all quantitative and qualitative results to decide

whether a full-scale RCT trial will be progressed or redesigning another KONTAKT[TM] feasibility

study will be needed. The following aspects will be considered: (1) participants and parents'

acceptability and adherence rates, (2) participants and parents' feedback on demand for

KONTAKT[TM], (3) trainers' feedback on implementing KONTAKT[TM] in Chinese clinical setting, (4)

trainers' fidelity on implementing KONTAKT[TM], (5) the feasibility results of the processes for data

collecting, and (6) considering the potential capacity of the intervention setting (human resources,

materials, and equipment) on implementing a full-scale RCT regarding the sample size calculation

for the primary outcome.

Plans for dissemination of finding

434   The findings of the Chinese KONTAKT^{TM} CHILD trial will be submitted to peer-review

435   journals for consideration. Abstracts will also be submit to both local and international conferences,

436   and community-based lay abstracts will be written and disseminated to provide feedback for autism

437   stakeholders in China.

References

1. American Psychiatric Association. Diagnostic and statistical manual of mental

disorders. 5th ed., text rev.;2022.

2. Tillmann J, San Jose Caceres A, Chatham CH, et al. Investigating the factors

underlying adaptive functioning in autism in the EU-AIMS Longitudinal European Autism

Project. Autism Res 2019;12(4):645-57. https://doi.org/10.1002/aur.2081. Epub 2019 Feb

12.

3. Zhou H, Xu X, Yan W, et al. Prevalence of Autism Spectrum Disorder in China: A

Nationwide Multi-center Population-based Study Among Children Aged 6 to 12 Years.

Neurosci Bull 2020;36(9):961-71. https://doi.org/10.1007/s12264-020-00530-6. Epub 2020

Jul 02.

4. Gilmore R, Ziviani J, Chatfield MD, et al. Social skills group training in adolescents

with disabilities: A systematic review. Res Dev Disabil 2022;125:104218.

https://doi.org/10.1016/j.ridd.2022.104218. Epub 2022 Mar 21.

5. Kaat AJ, Lecavalier L. Group-based social skills treatment: A methodological review.

Res Autism Spectr Disord 2014;8(1):15-24. https://doi.org/10.1016/j.rasd.2013.10.007.

6. Afsharnejad B, Black MH, Falkmer M, et al. The Methodological Quality and

Intervention Fidelity of Randomised Controlled Trials Evaluating Social Skills Group

Programs in Autistic Adolescents: A Systematic Review and Meta-analysis. J Autism Dev

Disord 2023 https://doi.org/10.1007/s10803-023-05893-z. Epub 2023 Jan 22.

7. Gates JA, Kang E, Lerner MD. Efficacy of group social skills interventions for youth

with autism spectrum disorder: A systematic review and meta-analysis. Clin Psychol Rev

2017;52:164-81. https://doi.org/10.1016/j.cpr.2017.01.006. Epub 2017 Jan 29.

8. Wolstencroft J, Robinson L, Srinivasan R, et al. A Systematic Review of Group

Social Skills Interventions, and Meta-analysis of Outcomes, for Children with High

Functioning   ASD.   J   Autism   Dev   Disord   2018;48(7):2293-307.

https://doi.org/10.1007/s10803-018-3485-1. Epub 2018 Feb 10.

9. Zheng S, Kim H, Salzman E, et al. Improving Social Knowledge and Skills among

Adolescents with Autism: Systematic Review and Meta-Analysis of UCLA PEERS(R) for

Adolescents.   J   Autism   Dev   Disord   2021;51(12):4488-503.

https://doi.org/10.1007/s10803-021-04885-1. Epub 2021 Jan 30.

10. Miller A, Vernon T, Wu V, et al. Social Skill Group Interventions for Adolescents

with Autism Spectrum Disorders: a Systematic Review. Rev J Autism Dev Disord

2014;1(4):254-65. https://doi.org/10.1007/s40489-014-0017-6.

11. Soares EE, Bausback K, Beard CL, et al. Social Skills Training for Autism

Spectrum Disorder: a Meta-analysis of In-person and Technological Interventions. J Technol

Behav Sci 2021;6(1):166-80. https://doi.org/10.1007/s41347-020-00177-0. Epub 2020 Nov

24.

12. Kinnaird E, Stewart C, Tchanturia K. Investigating alexithymia in autism: A

systematic   review   and   meta-analysis.   Eur   Psychiatry   2019;55:80-89.

https://doi.org/10.1016/j.eurpsy.2018.09.004. Epub 2018 Nov 07.

13. Tavernor L, Barron E, Rodgers J, et al. Finding out what matters: validity of quality

of life measurement in young people with ASD. Child Care Health Dev 2013;39(4):592-601.

https://doi.org/10.1111/j.1365-2214.2012.01377.x. Epub 2012 Apr 21.

14. Huggins CF, Donnan G, Cameron IM, et al. Emotional self-awareness in autism: A

meta-analysis of group differences and developmental effects. Autism 2021;25(2):307-21.

https://doi.org/10.1177/1362361320964306. Epub 2020 Nov 07.

15. Lee A, Peter Hobson R. Drawing self and others: How do children with autism

differ from those with learning difficulties? Br J Dev Psychol 2006;24(3):547-65.

https://doi.org/10.1348/026151005X49881.

16. Papangelo P, Pinzino M, Pelagatti S, et al. Human Figure Drawings in Children

with Autism Spectrum Disorders: A Possible Window on the Inner or the Outer World. Brain

Sci 2020;10(6) https://doi.org/10.3390/brainsci10060398. Epub 2020 Jun 27.

17. Kotroni P, Bonoti F, Mavropoulou S. Children with autism can express social

emotions in their drawings. Int J Dev Disabil 2018;65(4):248-56.

https://doi.org/10.1080/20473869.2018.1434855. Epub 2018 Feb 13.

18. Schweizer C, Knorth EJ, Spreen M. Art therapy with children with Autism

Spectrum Disorders: A review of clinical case descriptions on 'what works'. The Arts in

Psychotherapy 2014;41(5):577-93. https://doi.org/10.1016/j.aip.2014.10.009.

19. Waller D. Art therapy for children: how it leads to change. Clin Child Psychol

Psychiatry 2006;11(2):271-82. https://doi.org/10.1177/1359104506061419. Epub 2006 Nov

08.

20. Albaum CS, Vashi N, Bohr Y, et al. A Systematic Review of Therapeutic Process

Factors in Mental Health Treatment for Autistic Youth. Clin Child Fam Psychol Rev

2023;26(1):212-41. https://doi.org/10.1007/s10567-022-00409-0. Epub 2022 Aug 24.

21. Bölte S. Social färDTGhetsträning i grupp med fokus på kommunikation och social

interaktion vid autismspektrumtillstånd enligt Frankfurtmodellen (KONTAKT). Göttingen,

Stockholm: Hogfrefe; 2018.

22. Herbrecht E, Poustka F, Birnkammer S, et al. Pilot evaluation of the Frankfurt

Social Skills Training for children and adolescents with autism spectrum disorder. Eur Child

Adolesc Psychiatry 2009;18(6):327-35. https://doi.org/10.1007/s00787-008-0734-4. Epub

2009 Jan 24.

23. Choque Olsson N, Flygare O, Coco C, et al. Social Skills Training for Children and

Adolescents With Autism Spectrum Disorder: A Randomized Controlled Trial. J Am Acad

Child Adolesc Psychiatry 2017;56(7):585-92. https://doi.org/10.1016/j.jaac.2017.05.001.

Epub 2017 Jun 26.

24. Jonsson U, Olsson NC, Coco C, et al. Long-term social skills group training for

children and adolescents with autism spectrum disorder: a randomized controlled trial. Eur

Child Adolesc Psychiatry 2019;28(2):189-201. https://doi.org/10.1007/s00787-018-1161-9.

Epub 2018 May 12.

25. Afsharnejad B, Falkmer M, Black MH, et al. KONTAKT(R) social skills group

training for Australian adolescents with autism spectrum disorder: a randomized controlled

trial. Eur Child Adolesc Psychiatry 2022;31(11):1695-713.

https://doi.org/10.1007/s00787-021-01814-6. Epub 2021 May 31.

26. Afsharnejad B, Falkmer M, Picen T, et al. "I Met Someone Like Me!": Autistic

Adolescents and Their Parents' Experience of the KONTAKT(R) Social Skills Group

Training.   J   Autism   Dev   Disord   2022;52(4):1458-77.

https://doi.org/10.1007/s10803-021-05045-1. Epub 2021 May 05.

27. Choque Olsson N, Rautio D, Asztalos J, et al. Social skills group training in

high-functioning autism: A qualitative responder study. Autism 2016;20(8):995-1010.

https://doi.org/10.1177/1362361315621885. Epub 2016 Feb 26.

28. Hodge DR, Jackson KF, Vaughn MG. Culturally sensitive interventions and health

and behavioral health youth outcomes: a meta-analytic review. Soc Work Health Care

2010;49(5):401-23. https://doi.org/10.1080/00981381003648398. Epub 2010 Jun 04.

29. Barrera M, Jr., Castro FG, Strycker LA, et al. Cultural adaptations of behavioral

health interventions: a progress report. J Consult Clin Psychol 2013;81(2):196-205.

https://doi.org/10.1037/a0027085. Epub 2012 Feb 01.

30. Bernal G, Bonilla J, Bellido C. Ecological validity and cultural sensitivity for

outcome research: issues for the cultural adaptation and development of psychosocial

treatments   with   Hispanics.   J   Abnorm   Child   Psychol   1995;23(1):67-82.

https://doi.org/10.1007/BF01447045. Epub 1995 Feb 01.

31. Faregh N, Lencucha R, Ventevogel P, et al. Considering culture, context and

community in mhGAP implementation and training: challenges and recommendations from

the field. Int J Ment Health Syst 2019;13:58. https://doi.org/10.1186/s13033-019-0312-9.

Epub 2019 Aug 30.

32. Bolte S, Lawson WB, Marschik PB, et al. Reconciling the seemingly irreconcilable:

The WHO's ICF system integrates biological and psychosocial environmental determinants

of autism and ADHD: The International Classification of Functioning (ICF) allows to model

opposed biomedical and neurodiverse views of autism and ADHD within one framework.

Bioessays 2021;43(9):e2000254. https://doi.org/10.1002/bies.202000254. Epub 2021 Apr

03.

33. Bishop-Fitzpatrick L, Kind AJH. A Scoping Review of Health Disparities in Autism

Spectrum   Disorder.   J   Autism   Dev   Disord   2017;47(11):3380-91.

https://doi.org/10.1007/s10803-017-3251-9. Epub 2017 Aug 02.

34. Kang-Yi CD, Grinker RR, Beidas R, et al. Influence of Community-Level Cultural

Beliefs about Autism on Families' and Professionals' Care for Children. Transcult Psychiatry

2018;55(5):623-47. https://doi.org/10.1177/1363461518779831. Epub 2018 Jul 05.

35. Ravindran N, Myers BJ. Cultural Influences on Perceptions of Health, Illness, and

Disability: A Review and Focus on Autism. J Child Fam Stud 2012;21(2):311-19.

https://doi.org/10.1007/s10826-011-9477-9.

36. Clark E, Zhou Z, Du L. Autism in China: Progress and challenges in addressing the

needs   of   children   and   families.   Int   J   Sch   Educ   Psychol   2019;7(2):135-46.

https://doi.org/10.1080/21683603.2019.1570885.

37. Aarons GA, Green AE, Palinkas LA, et al. Dynamic adaptation process to

implement an evidence-based child maltreatment intervention. Implement Sci 2012;7:32.

https://doi.org/10.1186/1748-5908-7-32. Epub 2012 Apr 20.

38. Highfield L, Hartman MA, Mullen PD, et al. Intervention Mapping to Adapt

Evidence-Based Interventions for Use in Practice: Increasing Mammography among African

American Women. Biomed Res Int 2015;2015:160103. https://doi.org/10.1155/2015/160103.

Epub 2015 Nov 21.

39. Thabane L, Lancaster G. A guide to the reporting of protocols of pilot and

feasibility trials. Pilot Feasibility Stud 2019;5:37.

https://doi.org/10.1186/s40814-019-0423-8. Epub 2019 Mar 13.

40. Chan AW, Tetzlaff JM, Altman DG, et al. SPIRIT 2013 statement: defining standard

protocol items for clinical trials. Ann Intern Med 2013;158(3):200-7.

https://doi.org/10.7326/0003-4819-158-3-201302050-00583. Epub 2013 Jan 09.

41. Grant S, Mayo-Wilson E, Montgomery P, et al. CONSORT-SPI 2018 Explanation

and Elaboration: guidance for reporting social and psychological intervention trials. Trials

2018;19(1):406. https://doi.org/10.1186/s13063-018-2735-z. Epub 2018 Aug 01.

42. Zhang H. The revision of WISC-IV Chinese version. Psychological Science

2009;32(5):1177-79. https://doi.org/

43. Lord C, Rutter M, Le Couteur A. Autism Diagnostic Interview-Revised: a revised

version of a diagnostic interview for caregivers of individuals with possible pervasive

developmental disorders. J Autism Dev Disord 1994;24(5):659-85.

https://doi.org/10.1007/BF02172145. Epub 1994 Oct 01.

44. Su L, Li X, Wan G, et al. The norms of achenbach child behavior checklist in hunan

province. Chin J Clin Psychol 1996;4(1):24-28. https://doi.org/

45. Thornton CS, Tsai WH, Santana MJ, et al. Effects of Wait Times on Treatment

Adherence and Clinical Outcomes in Patients With Severe Sleep-Disordered Breathing: A

Secondary Analysis of a Noninferiority Randomized Clinical Trial. JAMA Netw Open

2020;3(4):e203088. https://doi.org/10.1001/jamanetworkopen.2020.3088. Epub 2020 Apr

21.

46. Robiner WN. Enhancing adherence in clinical research. Contemp Clin Trials

2005;26(1):59-77. https://doi.org/10.1016/j.cct.2004.11.015.

47. Lauritsen J, Bruus M. EPIDATA: Data entry system for use with Stata, SPSS, SAS,

Excel, text formats. 2000 https://doi.org/

48. Rabin SJ, Laugeson EA, Mor-Snir I, et al. An Israeli RCT of PEERS(R):

Intervention Effectiveness and the Predictive Value of Parental Sensitivity. J Clin Child

Adolesc Psychol 2021;50(6):933-49. https://doi.org/10.1080/15374416.2020.1796681. Epub

2020 Aug 12.

49. Whitehead AL, Julious SA, Cooper CL, et al. Estimating the sample size for a pilot

randomised trial to minimise the overall trial sample size for the external pilot and main trial

for a continuous outcome variable. Stat Methods Med Res 2016;25(3):1057-73.

https://doi.org/10.1177/0962280215588241. Epub 2015 Jun 21.

50. McCoy CE. Understanding the Intention-to-treat Principle in Randomized

Controlled Trials. West J Emerg Med 2017;18(6):1075-78.

https://doi.org/10.5811/westjem.2017.8.35985. Epub 2017 Nov 01.

51. Yang P, Chen G, Wang P, et al. Psychometric evaluation of the Chinese version of

the Child Health Utility 9D (CHU9D-CHN): a school-based study in China. Qual Life Res

2018;27(7):1921-31. https://doi.org/10.1007/s11136-018-1864-x. Epub 2018 May 08.

52. Yang Z, Jiang J, Wang P, et al. Estimating an EQ-5D-Y-3L Value Set for China.

Pharmacoeconomics 2022;40(Suppl 2):147-55. https://doi.org/10.1007/s40273-022-01216-9.

Epub 2022 Nov 18.

53. Cunningham JA, Kypri K, McCambridge J. Exploratory randomized controlled

trial evaluating the impact of a waiting list control design. BMC Med Res Methodol

2013;13:150. https://doi.org/10.1186/1471-2288-13-150. Epub 2013 Dec 10.

Footnotes

Authors' contributions

LU, ZH, ZX, SB, and SG undertook the conceptualization and design of the study. LU, ZH, BW translated the manuals and workbooks. LU, ZH, SB, SG, and ZX selected the measures used. BW, FJ, YP, HX, ZH, LU, and ZX improved the content of the manual. FJ, YP, ZH, and ZX assist with administration of groups and offer clinical support. LU drafted the manuscript, with inputs from ZH, SB, SG, and ZX. All authors contributed to, read and approved the final manuscript.

Funding statement

This research received no specific grant from any funding agency in the public, commercial or non-profit sectors.

Competing interests

All authors declare that they have no conflicts of interest.

Ethics approval

This study has been approved by the Ethical Board Committee at the Third Affiliated Hospital of Sun Yat-sen University (II2023-119-01), and pre-registered in Chinese Clinical Trials (ChiCTR2300072136).

Availability of data and materials

The ethical consent and funding documents have been reviewed by the review board of the

Chinese Clinical Trial Registry and are available for review by the Editorial Office. Anonymised

data related to this study will be available from the corresponding author upon reasonable request

after study completion and publication of the study. The final dataset generated during this study

will upload to the Research Data Deposit of Sun Yat-sen University. Trial results will be reported

and published in scientific journals.

Acknowledgments

We acknowledge the generous support from the Swedish team at Karolinska

Institutet/Stockholm Health Services and the Australian team at Curtin University. They facilitated

the training, provided precious experiences and materials for the CDBC KONTAKT[TM] trainers . We

thanks the CDBC KONTAKT[TM] team for facilitating the process of this trial. We sincerely thank

the autistic individual, Yihan Liu, and her families for validating the translation and contents of the

Chinese KONTKAT[TM] manual. Additionally, we appreciate Yan Li for designing the KONTAKT[TM]

China tools.

 Tables

649 **Table 1**. Schedule of enrolment, interventions, and assessments for the study

						STUDY PERIOD						
	Involvement[a]					Enrolment	Allocation	Post-allocation				
	C	P	T	E	F							
TIME POINTS[b]						*-T1*	**0**	*T1*	*T2*	*T3*	*T4*	*T5*
ENROLMENT:												
Eligibility Screen				X		X						
Informed consent				X		X						
Allocation							X					
INTERVENTIONS (KONTAKT[TM]**):**												
Immediate Intervention	X	X	X	X	X			●━━━━━●				X
Delay Intervention	X	X	X	X	X			X		●━━━━●		
ASSESSMENTS:												
CBCL		X				X						
Screening WISC-IV	X			X		X						
ADOS-G	X			X		X						
STI		X		X		X						

	SFI	X	X		X		X						
	SES		X				X						
Primary **Outcome**	CASS (+CRS)	X			X					X		X		X
Secondary **Outcome**	SRS-2		X	X						X		X		X
	CRSES	X								X		X		X
	Koppitz DAPT	X			X					X		X		X
	SCS-DAP	X			X					X		X		X
	CORS	X								———	———	———	———	
	PRFQ		X							X		X		X
	ABAS-2		X	X						X		X		X
	PSS		X							X		X		X
	STARSS	X			X					———	———	———	———	
Processing **Factors**	TPOCS	X			X					X	X	X	X	X
	PPKS	X	X				X							
	TASC	X	X			X				X	X	X	X	X
Fidelity **Outcome**	Attendance	X	X			X				———	———	———	———	
	HCR	X				X				———	———	———	———	
	Focus Group	X	X			X						X		X

	KFC				X			←———→	←———→		
	GSRS	X						←———→	←———→		
	TFAQ		X					X	X	X	X	X
	NEQ		X							X		X
Cost-effectiv	TIC-PK		X					X	X	X	X	X
eness	CHU9D	X						X	X	X	X	X
Outcome	EQ-5D-Y		X					X	X	X	X	X

a: Involvement: C: Child; P: Parent; T: Teacher; E: Evaluator blinded to the hypothesis; F: Trainer.

b: Time points: T1: Time 1; T2: Time 2; T3: Time 3; T4: Time 4; T5: Time 5.

Measures: ABAS-2:Adaptive Behaviour Assessment System-II; ADOS-G:Autism Diagnostic Observation Scale-Generic; CASS+CRS: Contextual Assessment of Social Skills + Conversation Rating Scale; CBCL: Child Behaviour Checklist; CHU9D: Child Health Utility 9D; CORS: Child Outcome Rating Scale; GSRS: Group Session Rating Scale; KFC: KONTAKT[TM] Fedility Checklist; Kopptiz DAPT: Koppitz Draw-A-Person Test; NEQ: Negative Effects Questionnaire; PPKS: Preparedness for Participating KONTAKT[TM] Scale; PRFQ: Parental Reflective Functioning Questionnaire; PSS: Perceived school support; TASC: Therapeutic Alliance Scales for Children; TFAQ: Theoretical Framework of Acceptability Questionnaire; TCI-PK: Trimbos/iMTA questionnaire for patients with a psychiatric disorder child version; SCS-DAP: Self-Concept Scale-Draw-A-Person; SES: Socioeconomic Status Scale; SFI: Screening Face Interview; STARSS: Snack Time Assessment of Relationships and Social Skills; STI: Screening Telephone Interview;

SRS-2: Social Responsiveness Scale-2; WISC-IV: Wechsler Intelligence Scale for Children-IV

Chinese version.

Goals

Social communication and interaction skills

Social motivation

Social cognition

Social competence

Self-confidence

Awareness of self and others

Problem-solving strategies

Emotion recognition and expression

Session Structure

Opening Round

Homework Review

Thematic Discussion

Group Activities

Snack Time

New Homework Assignment

Closing Round

Overall Structure

16 weekly 60-minute sessions facilitated by 2-3 therapists with 4-6 children per group

3 sessions include parent meeting facilitated by one therapist

Session1-11 fixed themes and similar homework

Session 12-15 individualised themes and homework

Session 16 summary and graduation

Essential Elements

Principles of CBT

Behavioural activation

Psychoeducation

Observational learning

Computerised social cognition training

Parental involvement

KONTAKT™