Original Article

AMEBIASIS: AN ANALYSIS OF ITS DISTINCTIVE PRESENTATION IN

THE CONTEXT OF JORDANIAN CASES

Enas Al-khlifeh and Ahmad B. Hasanat

1- Department of Medical Laboratory Science, Al-Balqa Applied University, Jordan.

2- Faculty of Information Technology, Mutah University, Karak, Jordan.

Correspondence to: Enas Al-khlifeh, *https://orcid.org/0009-0000-4601-3796*

Parasitology laboratory, Department of Medical Laboratory Science, Faculty of Science, Al-Balqa

Applied University, Al-Salt (19117) Jordan.

Tel: +962795856110

E-mail: *Al-khlifeh.en@bau.edu.jo*

ABSTRACT

Amebiasis is a serious health problem, particularly in economically disadvantaged countries.

However, unresolved questions remain regarding the clinical symptoms and laboratory results of

amebiasis in Jordan, a developing country. Therefore, we assessed the prevalence and severity of

amebiasis in a sizable population in the center of Jordan in relation to potential risk variables such

as environment, age and sex of the patients, collecting data from the medical records of the patients

treated at the Al-Hussein Hospital in Al-Salt city between 2020 and 2022. Clinical manifestations

and laboratory findings, including fecal red blood cell and leukocyte counts, total and differential

leukocyte counts by age, and other information were included in this report. This study showed that

amebiasis is a common problem in Al-Salt city, accounting for 16.8% of all gastroenteritis cases.

Amebiasis affects individuals of all sexes and ages. When comparing males of different ages to

females of similar ages, both age and male sex showed an increased risk of amebiasis (P=0.014).

Infants and toddlers accounted for more than one-quarter of the cases, with more than half of them

diagnosed with watery diarrhea. Numerous patients of all age groups displayed age-related

neutrophilic leukocytosis and atypical WBC counts. The findings of this study suggest that

amebiasis is highly prevalent, affects a significant proportion of infants and toddlers, and causes

severe clinical symptoms. This is the first study on Al-Salt city population with respect to multiple

variables and demographic factors. These data are crucial for the decision-making of medical

professionals. The epidemiology of amebiasis in Jordan must be determined, and preventive

measures must be developed, especially for childcare at educational facilities.

Keywords: *E. histolytica*; Amebiasis; Jordan; diarrhea; laboratory finding; male sex; toddlers

1. INTRODUCTION

Parasitic gastroenteritis (GE) is a serious global health concern. Amebiasis is transmitted by the

fecal-oral route ("CDC - DPDx - Amebiasis," 2019). The transmission occurs when a parasitic cyst

in the infectious fecal matter is consumed via food or water. The acid-resistant cyst upon entering

the small intestine, excysts to produce the organism's motile and invasive form, trophozoites, which

travel to the large intestine, and colonize the colon lumen as commensal flora or infiltrate the

colonic epithelium, causing inflammation and amoebic colonic ulcers (Yue et al., 2021).

Proliferation of the parasite in the colon results in symptoms of GE, including abdominal pain,

dysentery, or mucous soft stools. However, the majority of infected people show no overt

symptoms. The infection becomes life threatening if *E. histolytica* becomes invasive and spreads

extra-intestinally to form liver abscesses because excess hepatic iron creates an ideal milieu for *E.*

histolytica proliferation in the liver (Salles et al., 2003).

Poor hand hygiene and contamination of water bodies and underground water with fecal matter are

the main risk factors for amebiasis ("CDC - DPDx - Amebiasis," 2019). Amebiasis can affect

people of any age, including children (Haque et al., 2006; Hegazi et al., 2013). Male sex is a known

risk factor for the development of an amoebic liver abscess, especially in young men, although

amebiasis can afflict women as well (Ghosh et al., 2014). According to previous studies, fecal

contamination during oral and anal sex with homosexual or bisexual men increases the risk of

amebiasis (Hung et al., 2011). Patients with compromised immune systems are extremely

vulnerable (Zanetti et al., 2021).

Past research conducted in Jordan has revealed that amebiasis is a prevalent infection. However,

several questions remain unanswered. Along with uncertainties regarding the clinical and laboratory

signs, the severity of amebiasis in Jordanian patients of all ages and sexes remains unknown. This

study aimed to determine the prevalence of amebiasis in a significant population in Jordan's central

region in relation to potential risk factors to understand the epidemiological profile of amebiasis in

Jordan. In addition, we used random forest machine learning algorithm to classify the important

laboratory features and risk factors associated with the severity of infection

2. MATERIALS AND METHODS

2.1. DATA COLLECTION

Data were collected from Jordan's Hakeem electronic health record system between January 20,

2021, and August 28, 2022. Since 2009, Hakeem has managed and collected medical data from

Jordan's public healthcare system ("Hakeem Program | Electronic Health Solutions," n.d.). The data

used for the analysis included wet-mount reports of the detection of intestinal parasites. Laboratory

tests were performed in accordance with the normal operating procedures of the Jordanian health

centers. Briefly, stool samples were collected in labeled, clean, dry, leak-proof, and sterile plastic

containers for a laboratory diagnosis from all patients with gastroenteritis (GE) with suspected

involvement of an intestinal parasite. Within 30 min of sample collection, direct stool examinations

were performed to examine for the presence of common intestinal parasites, including *E.*

histolytica/dispar, *Giardia lamblia*, and worms such as *Ascaris lumbricoides*, *Hymenolepis*

nana, *Enterobius vermicularis*, *Strongyloides stercoralis*, and *Trichuris trichiura*, hookworms

besides *Taenia* and other cestode species. After stool examinations were completed, the infected

patients were treated according to the national guidelines.

Positive parasitology results for *E. histolytica* were exclusively included in this study because it is

the predominant protozoan reported. A few sporadic cases of *Giardia* (less than 10) were reported

during the study period but were not included in our analysis. Hakeem's sociodemographic data and

laboratory results were examined and compiled using a worksheet specifically designed for this

purpose. The dataset contained the date of sample collection, sex, age (years), diagnosis (ordered

categorical), and stool examination results with severity indicators such as stool-red (RBCs) and

leukocytosis or white blood cell (WBCs) counts. The WBC count was categorized into three based

on the number of cells/high field: low = 0 to 3, medium = 3 to 6, and high = more than 6. In

addition, results of the complete blood count (CBC) were used to obtain WBCs and neutrophil

counts as it is considered a sign of the reaction to the infection and severity. The patients were

categorized into three age groups: toddlers (less than five years), school-going age (represents 5–18

years), active workforce (19–59 years old), and adults (older than 60 years).

2.2.DATA PROCESSING, ANALYSIS, AND VISUALIZATION

The data were curated after extraction from Hakeem using the Pandas Python package (2023). The

python packages numpy (Harris et al., 2020), scipy.stats (Virtanen et al., 2020) and statsmodels

(Seabold and Perktold, 2010) were used to categorize the data and establish the association between

variables and display their distribution patterns. In order to compare qualitative data, a Chi-square

test was performed, which clarified the distributions of categorical variables, specifically the

relationship between risk factors for amebiasis, including sex, age, and other laboratory findings

that indicated the severity of infection, including the severity of diarrhea or the presence of RBC or

WBC in the stool. Correlations among age, WBC count, neutrophil count, and neutrophil

percentage were tested using one-way analysis of variance. Student's t-test was used to compare

numerical variables (WBC count, neutrophil count, and neutrophil percentage) and to determine

whether there was a significant difference in the mean between groups. The analysis adhered to a

significance threshold, wherein a P-value of less than 0.05 was deemed statistically significant,

ensuring the reliability and validity of the results. The code used for the analysis is included in

Supplementary File (S1). Graphical data representations were generated using Matplotlib. pyplot

package (Hunter, 2007) and Seaborn in Python (Waskom et al., 2017). We have also included a

brief excerpt from the dataset in the Supplementary File (S2).

RESULTS

We performed a retrospective study on patients' parasitology analysis data from Al-Hussein/Salt

Hospital in Jordan, collected between September 2020 and December 2022. To identify intestinal

parasites in the study area, 4429 stool tests were performed over the study period on patients with

GE symptoms. Among these, 763 were confirmed to have amebiasis (16.98%), with 381 male

(49.9%) and 382 (50.1%) female. The study participants were between the ages of less than one

month and 99 years. The median age of female patients was 24 years, whereas that of male patients

was 19 yrs. There was no significant difference between the median ages of females and males (P =

0.43, Figure 2). The prevalence of amebiasis among males and females across various age groups

was not significantly different (P = 0.39). However, among the toddlers and preschoolers, there

were more males than females (Figure 2A). In contrast, there were considerably more females than

males in the adult group (aged 19 to 59).

Among all age groups studied, adults aged 20–59 years had the highest infection rate, whereas those

older than 60 years had the lowest infection rate (Figure 2A). The percentages of amoebic infections

across age groups did not differ significantly (P= 0.06). According to the clinical signs, the intensity

of the symptoms was as follows: abdominal pain without diarrhea, 145 (19%); semisoft stools, 189

(24.7%); watery diarrhea, 426 (55.8%); rectal bleeding, 3 (0.3%). Individuals aged 1 month to 4

years accounted for 27% of the total, with 107/206 cases (51.9%) diagnosed with watery diarrhea.

As expected, WBC and neutrophil counts showed a correlation with the diagnosis. When patients

were diagnosed with watery diarrhea, these measurements increased (Figure 3). The correlation

between other laboratory features and the diagnosis is shown in Figure (2B).

The following results were found when the risk

variables and diarrhea severity were compared: each diagnosis had a different median age for males

and females (Figure 1A). Both males and females under 48 years of age accounted for the majority

(75%) of the recorded cases, but many older individuals had watery diarrhea.

There were no significant variations in the diagnosis of loose stools according to age group.

However, a significant difference was found in males according to age when sex, age group, and

diagnosis were analyzed (P=0.014; Figure 1). All severity features, including the stool-RBC, stool-

WBC, WBC, and neutrophil count for the age and sex groups, were statistically insignificant (P ≥

0.05) (Figures 2).

Figure 1 Clinical signs and laboratory findings with respect to age among patients with confirmed case of of *E. histolytica* gastroenteritis. P value < 0.05 indicates a significant difference. *n*= 763.

Figure 2 Clinical signs and laboratory findings with respect to sex among patients with

confirmed case of of *E. histolytica* gastroenteritis. P value < 0.05 indicates a significant

difference. *n*= 763.

3. DISCUSSION

During the study, 4429 cases of GE diagnoses and requests for stool tests were recorded in the main

hospitals of Al-Salt city. A total of 763 cases had confirmed amebiasis diagnoses, accounting for

16.8% of all GE cases. This demonstrates that GE-associated amoebiasis is a common public health

concern in Al-Salt city.

The primary technique for verifying the presence of amoebiasis was conventional microscopic

analysis of stool samples. Compared with other detection methods, such as the *E. histolytica* antigen

detection test, it has been shown to have higher sensitivity and specificity for identification (Al-

Dalabeeh et al., 2020). However, according to our assessment of computerized data, confirmed

instances of amoebiasis displayed GE symptoms, including vomiting, abdominal pain, diarrhea, and

bloody stools. There are seven species of Entamoeba that can infect people, *E. histolytica* is

commonly acknowledged as a disease-causing organism. Whereas *E. dispar,* which is

morphologically identical to *E. histolytica,* is mostly non-pathogenic and associated with

asymptomatic infection. At present, it is advised to use molecular techniques to precisely

differentiate between pathogenic species of Entamoeba. Therefore, the confirmation of the exact

species responsible of amoebiasis is outside the scope of this study, since we relied on retrospective

analysis of lab test results.

An overall prevalence of 0.1–80.7% for amebiasis has been reported in the majority of earlier

studies conducted in different regions of Jordan (Nawafleh et al., 2014; Abdel-Dayem et al., 2014).

Although this claim was supported by the results of molecular typing and serology, the prevalence

of the pathogenic type *E. histolytica* might be overestimated when using microscopy (Hijjawi et al.,

2021; Al-Dalabeeh et al., 2020). There were disparities in the prevalence of infection between our

findings and earlier reports from other Jordanian locations. A diagnosis is traditionally made based

on the evaluation of a spherical cyst (~10 mm) or trophozoit and is considered to be subjective.

Additionally, geographical elements such as the source and nature of drinking water may have an

impact on the presence of variance between various locales.

Demographic factors such as age and male sex pre-disposed to amebiasis infection. Males appear to

acquire amebiasis at an earlier age than females. Our results agree with those of the previous

Jordanian (Nawafleh et al., 2014) and global studies (Nasrallah et al., 2022). Collectively, these

studies demonstrate a correlation between age and male sex with amebiasis. This bias is mostly due

to high-risk sexual behaviors (Hung et al., 2011) or differences in immunological responses to

infection between the sexes. For example, men have immunological characteristics, such as sex

chromosomal complement, immune cell receptors, and sex hormones that make them more

vulnerable to serious infections (Jacobsen and Klein, 2021; Lotter et al., 2013; Bernin et al., 2014).

Young children and toddlers under the age of five were discovered to make up more than a quarter

of the patients in this study, with more than half of them having been diagnosed with watery

diarrhea, which indicates a high severity of infection, as well as the presence of a significant

proportion of aberrant WBC and neutrophil counts for the age. This could have severe effects on

children of this age as diarrheal infections linked to amebiasis are more likely to cause dehydration,

stunted growth, and induce malnutrition (Mondal et al., 2006). Severe complications caused by

dehydration necessitate immediate hospitalization. The significant prevalence of amebiasis that we

discovered in this study is higher than what was 20 years ago for Jordanian children of the same age

group (Youssef et al., 2000). Furthermore, amebiasis in infants younger than one year was more

prevalent than predicted. This may be considered an unusual manifestation of amebiasis at this age,

given that the transmission of intestinal parasites is frequently associated with the contamination of

water and food and that infants are not normally expected to develop amebiasis. However, the risk

of infection is increased by inadequate breastfeeding, mixed feeding, and daycare (Hegazi et al.,

2013). This finding emphasizes the importance of carefully observing the GE infection dynamics in

Jordanian children to ascertain whether the increasing frequency of amebiasis in children is also

prevalent in other Jordanian regions. There have been a few occasional reports with inconsistent

results. Specifically, a prior surveillance research conducted in the capital city of Jordan discovered

that the frequency of intestinal parasites (including *E. histolytica*) was the highest in children under

the age of five (Chazal and Adi, 2007). Additionally, children under the age of 15 make up more

than 60% of affected patients in northern Jordanian cities (Jaran, 2017). In contrast, adult patients in

southern sites have *E. histolytica* at higher rates than children (Nawafleh et al., 2014). It is essential

to obtain more information on the socioeconomic situation, water quality, dietary background,

nursery, and daycare, to understand the causes of amebiasis in Jordan's toddlers and young children.

amebiasis is a disease with varying degrees of severity because it frequently causes silent and

chronic infections in contagious patients. In this study, some patients were diagnosed with

abdominal pain without diarrhea. It has been proposed that Entamoeba sp. including *E. histolytica*

can survive, reproduce, and disseminate without virulence (Marie and Petri, 2014). The parasite

may, however, have genetically distinct isolates. One of these genotypes may have caused the

severe manifestations observed in the studied cases. *E. histolytica* genotyping falls beyond the

scope of this study. The host's immune response, history of infection, and previous exposure may

have an impact on the severity of infection in addition to the genotype of the parasite. Thus, further

research is required to pinpoint the underlying genetic and immunological mechanisms responsible

for the aggressive presentation of amebiasis in our study.

LIMITATIONS OF THE STUDY

This study had some limitations. This study was performed in the Jordanian city of Al-Salt and was

primarily focused on a single community. A more comprehensive, multicentric study involving

different cohorts of patients from diverse ethnic, socio-economic, and regional backgrounds would

provide more concrete data and reproducible patterns on the distribution of potentially diverse

genotypes of the parasite and respective differences in the pathogenesis and immune responses

upon infection. However, a sufficient sample of the local population was reflected in the number of

patients admitted to city-only hospitals. Investigation of the immunological characteristics such as

cytokine measurements and genotyping of the isolated Entamoeba species would have been

beneficial in elucidating the potential mechanisms underlying the severity of amebiasis in Jordanian

patients.

CONCLUSIONS AND RECOMMENDATIONS

Amebiasis was found to be highly prevalent among patients with GE across all age and sex groups

in the population under investigation. However, younger men were susceptible to amebiasis and

exhibited severe clinical symptoms such as diarrhea. Typical clinical manifestations and laboratory

findings of invasive amebiasis, such as significant WBC counts and neutrophilic leukocytosis, were

observed in all age and sex groups. As per our findings, amebiasis can have a very high chance of

recurrence depending upon the environmental and host variables primarily linked to sex, inadequate

childcare, and irregularities in the water supply. As discussed before, the epidemiology of amebiasis

in Jordan must be determined on a large scale, and preventive measures must be devised to curb the

incidence in high-risk areas such as childcare and educational facilities. Additionally, to lower the

prevalence of infection with amebiasis and other infectious GE agents in Jordan, periodic

inspections should be performed during the transport and distribution of commercial water because

the majority of Jordanians use it.

ACKNOWLEDGMENTS:

We acknowledge Jordan's AL-Hussein/Salt Hospital for allowing access to medical records from

the microbiology laboratory section. In addition, we gratefully acknowledge Mrs. Qamar Al-Mimi

Geographic Information System Specialist from Nature Conservation Monitoring Center for

providing a map of the study area.

FUNDING: This research received no specific grant from any funding agency in the public,

commercial, or not-for-profit sectors.

ETHICAL APPROVAL: This study was approved by the scientific and administration committee

of scientific research at Al-Balqa applied university and Jordan's AL-Hussein/Salt Hospital.

Approval number/ID: 35/4/1205. The material is the authors' own original work, which has not

been previously published elsewhere.

DATA AVAILABILITY: The dataset is constantly being updated, and a subset can be made

available on request.

SUPPLEMENTARY MATERIALS: Supplementary material associated with this article can be

found, in the online version, at doi:

REFERENCES

Abdel-Dayem, M., Al Zou'bi, R., Hani, R. B., & Amr, Z. S. (2014). Microbiological and parasitological

investigation among food handlers in hotels in the Dead Sea area, Jordan. *Journal of Microbiology,*

Immunology and Infection, 47(5), 377–380. https://doi.org/10.1016/j.jmii.2013.06.001

Al-Dalabeeh, E. A., Irshaid, F. I., Roy, S., Ali, I. K. M., & Al-Shudifat, A. M. (2020). Identification of Entamoeba

histolytica in Patients with Suspected Amebiasis in Jordan Using PCR-based Assays. *Pakistan Journal*

of Biological Sciences: PJBS, 23(2), 166–172. https://doi.org/10.3923/pjbs.2020.166.172

Alvarado-Esquivel, C., Hernandez-Tinoco, J., & Sanchez-Anguiano, L. F. (2015). Seroepidemiology of

Entamoeba histolytica Infection in General Population in Rural Durango, Mexico. *Journal of Clinical*

Medicine Research, *7*(6), 435–439. https://doi.org/10.14740/jocmr.v7i6.2131

Bernin, H., Marggraff, C., Jacobs, T., Brattig, N., An, L. V., Blessmann, J., & Lotter, H. (2014). Immune

markers characteristic for asymptomatically infected and diseased Entamoeba histolytica

individuals and their relation to sex. *BMC Infectious Diseases*, *14*, 621.

https://doi.org/10.1186/s12879-014-0621-1

CDC - DPDx—Amebiasis. (2019, October 15). https://www.cdc.gov/dpdx/amebiasis/index.html

Chazal, A. M., & Adi, H. K. (2007). THE PREVALENCE OF INTESTINAL PARASITES IN AMMAN, JORDAN.

Bulletin of Pharmaceutical Sciences. Assiut, *30*(2), 235–239.

https://doi.org/10.21608/bfsa.2007.64203

dos Santos Zanetti, A., Malheiros, A. F., de Matos, T. A., dos Santos, C., Battaglini, P. F., Moreira, L. M.,

Lemos, L. M. S., Castrillon, S. K. I., da Costa Boamorte Cortela, D., Ignotti, E., & Espinosa, O. A. (n.d.).

Diversity, geographical distribution, and prevalence of Entamoeba spp. in Brazil: A systematic

review and meta-analysis. *Parasite*, *28*, 17. https://doi.org/10.1051/parasite/2021028

Estimates of global, regional, and national morbidity, mortality, and aetiologies of diarrhoeal diseases: A

systematic analysis for the Global Burden of Disease Study 2015. (2017). *The Lancet. Infectious*

Diseases, *17*(9), 909–948. https://doi.org/10.1016/S1473-3099(17)30276-1

Flaih, M. H., Khazaal, R. M., Kadhim, M. K., Hussein, K. R., & Alhamadani, F. A. B. (2021). The epidemiology

of amoebiasis in Thi-Qar Province, Iraq (2015-2020): Differentiation of Entamoeba histolytica and

Entamoeba dispar using nested and real-time polymerase chain reaction. *Epidemiology and Health*,

43, e2021034. https://doi.org/10.4178/epih.e2021034

Ghosh, S., Sharma, S., Gadpayle, A. K., Gupta, H. K., Mahajan, R. K., Sahoo, R., & Kumar, N. (2014). Clinical,

Laboratory, and Management Profile in Patients of Liver Abscess from Northern India. *Journal of*

Tropical Medicine, *2014*, e142382. https://doi.org/10.1155/2014/142382

Haghighi, A., Riahi, S. M., Taghipour, A., Spotin, A., Javanian, M., Mohammadi, M., Esmaeili Dooki, M., &

Rostami, A. (2018). Amoebiasis in Iran: A systematic review and meta-analysis. *Epidemiology and*

Infection, *146*(15), 1880–1890. https://doi.org/10.1017/S0950268818001863

Hakeem Program | Electronic Health Solutions. (n.d.). Retrieved September 2, 2023, from

https://ehs.com.jo/hakeem-program

Haque, R., Mondal, D., Duggal, P., Kabir, M., Roy, S., Farr, B. M., Sack, R. B., & Petri, W. A. (2006).

Entamoeba histolytica Infection in Children and Protection from Subsequent Amebiasis. *Infection*

and Immunity, *74*(2), 904–909. https://doi.org/10.1128/IAI.74.2.904-909.2006

Harris, C. R., Millman, K. J., van der Walt, S. J., Gommers, R., Virtanen, P., Cournapeau, D., Wieser, E., Taylor,

J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., van Kerkwijk, M. H., Brett, M., Haldane, A., del

Río, J. F., Wiebe, M., Peterson, P., … Oliphant, T. E. (2020).

Hegazi, M. A., Patel, T. A., & El-Deek, B. S. (2013). Prevalence and characters of Entamoeba histolytica

infection in Saudi infants and children admitted with diarrhea at 2 main hospitals at south Jeddah:

A re-emerging serious infection with unusual presentation. *The Brazilian Journal of Infectious*

Diseases, *17*(1), 32–40. https://doi.org/10.1016/j.bjid.2012.08.021

Hijjawi, N., Zahedi, A., & Ryan, U. (2021). Molecular characterization of Entamoeba, Blastocystis and

Cryptosporidium species in stool samples collected from Jordanian patients suffering from

gastroenteritis. *Tropical Parasitology*, *11*(2), 122–125. https://doi.org/10.4103/tp.TP_106_20

Hung, C.-C., Wu, P.-Y., Chang, S.-Y., Ji, D.-D., Sun, H.-Y., Liu, W.-C., Wu, C.-H., & Chang, S.-C. (2011).

Amebiasis among Persons Who Sought Voluntary Counseling and Testing for Human

Immunodeficiency Virus Infection: A Case-Control Study. *The American Journal of Tropical Medicine*

and Hygiene, *84*(1), 65–69. https://doi.org/10.4269/ajtmh.2011.10-0238

Hunter, J. D. (2007). Matplotlib: A 2D Graphics Environment. *Computing in Science & Engineering*, *9*(3), 90–

95. https://doi.org/10.1109/MCSE.2007.55

Jacobsen, H., & Klein, S. L. (2021). Sex Differences in Immunity to Viral Infections. *Frontiers in Immunology*,

12, 720952. https://doi.org/10.3389/fimmu.2021.720952

Jaran, A. S. (2017). Prevalence and seasonal variation of human intestinal parasites in patients attending

hospital with abdominal symptoms in northern Jordan. *Eastern Mediterranean Health Journal = La*

Revue De Sante De La Mediterranee Orientale = Al-Majallah Al-Sihhiyah Li-Sharq Al-Mutawassit,

22(10), 756–760. https://doi.org/10.26719/2016.22.10.756

Kumanan, T., Sujanitha, V., & Sreeharan, N. (2020). Amoebic liver abscess: A neglected tropical disease. *The*

Lancet Infectious Diseases, *20*(2), 160–162. https://doi.org/10.1016/S1473-3099(19)30696-6

Lin, F.-H., Chen, B.-C., Chou, Y.-C., Chien, W.-C., Chung, C.-H., Hsieh, C.-J., & Yu, C.-P. (2022). The

Epidemiology of Entamoeba histolytica Infection and Its Associated Risk Factors among Domestic

and Imported Patients in Taiwan during the 2011–2020 Period. *Medicina*, *58*(6), 820.

https://doi.org/10.3390/medicina58060820

Lotter, H., Helk, E., Bernin, H., Jacobs, T., Prehn, C., Adamski, J., González-Roldán, N., Holst, O., & Tannich, E.

(2013). Testosterone Increases Susceptibility to Amebic Liver Abscess in Mice and Mediates

Inhibition of IFNγ Secretion in Natural Killer T Cells. *PLoS ONE*, *8*(2), e55694.

https://doi.org/10.1371/journal.pone.0055694

Marie, C., & Petri, W. A. (2014). Regulation of Virulence of Entamoeba histolytica. *Annual Review of*

Microbiology, *68*, 493–520. https://doi.org/10.1146/annurev-micro-091313-103550

Mercado, E. H., Ochoa, T. J., Ecker, L., Cabello, M., Durand, D., Barletta, F., Molina, M., Gil, A. I., Huicho, L.,

Lanata, C. F., & Cleary, T. G. (2011). Fecal Leukocytes in Children Infected with Diarrheagenic

Escherichia coli⬚. *Journal of Clinical Microbiology*, *49*(4), 1376–1381.

https://doi.org/10.1128/JCM.02199-10

Mondal, D., Petri, W. A., Sack, R. B., Kirkpatrick, B. D., & Haque, R. (2006). Entamoeba histolytica-associated

diarrheal illness is negatively associated with the growth of preschool children: Evidence from a

prospective study. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, *100*(11),

1032–1038. https://doi.org/10.1016/j.trstmh.2005.12.012

Nakada-Tsukui, K., & Nozaki, T. (2016). Immune Response of Amebiasis and Immune Evasion by Entamoeba

histolytica. *Frontiers in Immunology*, 7.

https://www.frontiersin.org/articles/10.3389/fimmu.2016.00175

Nasrallah, J., Akhoundi, M., Haouchine, D., Marteau, A., Mantelet, S., Wind, P., Benamouzig, R., Bouchaud,

O., Dhote, R., & Izri, A. (2022). Updates on the worldwide burden of amoebiasis: A case series and

literature review. *Journal of Infection and Public Health*, *15*(10), 1134–1141.

https://doi.org/10.1016/j.jiph.2022.08.013

Nawafleh, H., Alhroob, A., Kawafha, M., & Altaif, K. (2014). EPIDEMIOLOGICAL STUDY: LABORATORY DATA

MINING IN SOUTH OF JORDAN. *American Journal of Infectious Diseases*, *10*, 137–141.

https://doi.org/10.3844/ajidsp.2014.137.141

Nimri, L. F., & Meqdam, M. (2004). Enteropathogens associated with cases of gastroenteritis in a rural

population in Jordan. *Clinical Microbiology and Infection*, *10*(7), 634–639.

https://doi.org/10.1111/j.1469-0691.2004.00891.x

Salles, J. M., Moraes, L. A., & Salles, M. C. (2003). Hepatic amebiasis. *The Brazilian Journal of Infectious*

Diseases: An Official Publication of the Brazilian Society of Infectious Diseases, *7*(2), 96–110.

https://doi.org/10.1590/s1413-86702003000200002

Seabold, S., & Perktold, J. (2010). Statsmodels: Econometric and Statistical Modeling with Python.

Proceedings of the 9th Python in Science Conference, 2010.

Servián, A., Helman, E., Iglesias, M. del R., Panti-May, J. A., Zonta, M. L., & Navone, G. T. (2022). Prevalence

of Human Intestinal Entamoeba spp. in the Americas: A Systematic Review and Meta-Analysis,

1990–2022. *Pathogens*, *11*(11), 1365. https://doi.org/10.3390/pathogens11111365

Shirley, D.-A. T., Farr, L., Watanabe, K., & Moonah, S. (2018a). A Review of the Global Burden, New

Diagnostics, and Current Therapeutics for Amebiasis. *Open Forum Infectious Diseases*, *5*(7), ofy161.

https://doi.org/10.1093/ofid/ofy161

Singh, A., Banerjee, T., Khan, U., & Shukla, S. K. (2021). Epidemiology of clinically relevant Entamoeba spp.

(E. histolytica/dispar/moshkovskii/bangladeshi): A cross sectional study from North India. *PLOS*

Neglected Tropical Diseases, *15*(9), e0009762. https://doi.org/10.1371/journal.pntd.0009762

Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson,

P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson, J., Millman, K. J., Mayorov, N.,

Nelson, A. R. J., Jones, E., Kern, R., Larson, E., … van Mulbregt, P. (2020). SciPy 1.0: Fundamental

algorithms for scientific computing in Python. *Nature Methods*, *17*(3), Article 3.

https://doi.org/10.1038/s41592-019-0686-2

Waskom, M., Botvinnik, O., O'Kane, D., Hobson, P., Lukauskas, S., Gemperline, D. C., Augspurger, T.,

Halchenko, Y., Cole, J. B., Warmenhoven, J., Ruiter, J. de, Pye, C., Hoyer, S., Vanderplas, J., Villalba,

S., Kunter, G., Quintero, E., Bachant, P., Martin, M., … Qalieh, A. (2017). *mwaskom/seaborn: V0.8.1*

(September 2017) [Computer software]. Zenodo. https://doi.org/10.5281/zenodo.883859

Water, sanitation and hygiene | UNICEF Jordan. (n.d.). Retrieved August 28, 2023, from

https://www.unicef.org/jordan/water-sanitation-and-hygiene

Youssef, M., Shurman, A., Bougnoux, M., Rawashdeh, M., Bretagne, S., & Strockbine, N. (2000). Bacterial,

viral and parasitic enteric pathogens associated with acute diarrhea in hospitalized children from

northern Jordan. *FEMS Immunology and Medical Microbiology*, *28*(3), 257–263.

https://doi.org/10.1111/j.1574-695X.2000.tb01485.x

Yue, B., Meng, Y., Zhou, Y., Zhao, H., Wu, Y., & Zong, Y. (2021). Characteristics of endoscopic and

pathological findings of amebic colitis. *BMC Gastroenterology*, *21*(1), 367.

https://doi.org/10.1186/s12876-021-01941-z

Zanetti, A. dos S., Malheiros, A. F., Matos, T. A. de, Santos, C. dos, Battaglini, P. F., Moreira, L. M., Lemos, L.

M. S., Castrillon, S. K. I., Cortela, D. da C. B., Ignotti, E., & Espinosa, O. A. (2021). Diversity,

geographical distribution, and prevalence of Entamoeba spp. in Brazil: A systematic review and

meta-analysis. *Parasite*, *28*, 17. https://doi.org/10.1051/parasite/2021028