Cross-sectional associations between prenatal maternal per- and poly-fluoroalkyl substances and bioactive lipids in three Environmental influences on Child Health Outcomes (ECHO) cohorts

Himal Suthar[a], Tomás Manea[a], Dominic Pak[a], Megan Woodbury[b], Stephanie M. Eick[c], Amber Cathey[d], Deborah J. Watkins[d], Rita S. Strakovsky[e,f], Brad A. Ryva[e,g,h], Subramaniam Pennathur[i,j], Lixia Zeng[i], David Weller[k], June-Soo Park[l], Sabrina Smith[l], Erin DeMicco[m], Amy Padula[m], Rebecca C. Fry[n], Bhramar Mukherjee[o], Andrea Aguiar[p,q], Sarah Dee Geiger[q,r], Shukhan Ng[q], Gredia Huerta-Montanez[b], Carmen Vélez-Vega[s], Zaira Rosario[t], Jose F. Cordero[s], Emily Zimmerman[u], Tracey J. Woodruff[m], Rachel Morello-Frosch[m,v], Susan L. Schantz[p,q], John D. Meeker[d], Akram Alshawabkeh[b], Max T. Aung[a] and on behalf of Program Collaborators for Environmental Influences on Child Health Outcomes*

Affiliations:

[a]Department of Population and Public Health Sciences, University of Southern California, Los Angeles, CA, USA

[b]Department of Civil and Environmental Engineering, Northeastern University, Boston, MA, USA

[c]Gangarosa Department of Environmental Health, Emory University Rollins School of Public Health, Atlanta, GA, USA

[d]Department of Environmental Health Sciences, University of Michigan School of Public Health, Ann Arbor, MI, USA

[e]Institute for Integrative Toxicology, Michigan State University, East Lansing, MI, USA

[f]Department of Food Sciences and Human Nutrition, Michigan State University, East Lansing, MI, USA

[g]Department of Pharmacology and Toxicology, Michigan State University, East Lansing, MI, USA

[h]College of Osteopathic Medicine, Michigan State University, East Lansing, MI, USA

[i]Department of Internal Medicine-Nephrology, University of Michigan, Ann Arbor, MI, USA

[j]Department of Molecular and Integrative Physiology, University of Michigan, Ann Arbor, MI, USA

[k]NSF International, Ann Arbor, MI, USA

[l]Environmental Chemistry Laboratory, Department of Toxic Substances Control, California Environmental Protection Agency, Berkeley, CA, USA

[m]Program on Reproductive Health and the Environment, University of California, San Francisco, San Francisco, CA, USA

[n]Department of Environmental Sciences and Engineering, University of North Carolina, Chapel Hill, Gillings School of Global Public Health, Chapel Hill, NC, USA

[o]Department of Biostatistics, University of Michigan School of Public Health, Ann Arbor, MI, USA

[p]Beckman Institute for Advanced Science and Technology, University of Illinois Urbana-Champaign, Illinois, USA

[q]Department of Comparative Biosciences, University of Illinois Urbana-Champaign, IL, USA

[r]Department of Kinesiology and Community Health, University of Illinois at Urbana-Champaign, Champaign, Illinois, USA

[s]Department of Epidemiology and Biostatistics, University of Georgia, Athens, Georgia, USA

^tUniversity of Puerto Rico Graduate School of Public Health, San Juan, PR, USA

^uDepartment of Communication Sciences and Disorders, Northeastern University, Boston, MA, USA

^vDepartment of Environmental Science, Policy and Management and School of Public Health, University of California, Berkeley, Berkeley, CA, USA

Corresponding author:

Max T. Aung

University of Southern California, Keck School of Medicine

Department of Population and Public Health Sciences

SSB 225R

1845 N Soto St, Los Angeles, CA, 90032

Email: maxaung@usc.edu

** See Acknowledgements for full listing of collaborators

ABSTRACT

Background: Per- and poly-fluoroalkyl substances (PFAS) exposure can occur through

ingestion of contaminated food and water, and inhalation of indoor air contaminated with these

chemicals from consumer and industrial products. Prenatal PFAS exposures may confer risk for

pregnancy-related outcomes such as hypertensive and metabolic disorders, preterm birth, and

impaired fetal development through intermediate metabolic and inflammation pathways.

Objective: Estimate associations between maternal pregnancy PFAS exposure (individually and

as a mixture) and bioactive lipids.

Methods: Our study included pregnant women in the Environmental influences on Child Health

Outcomes Program: Chemicals in our Bodies cohort (CiOB, n=73), Illinois Kids Developmental

Study (IKIDS, n=287), and the ECHO-PROTECT cohort (n=54). We measured twelve PFAS in

serum and 50 plasma bioactive lipids (parent fatty acids and eicosanoids derived from

cytochrome p450, lipoxygenase, and cyclooxygenase) during pregnancy (median 17 gestational

weeks). Pairwise associations across cohorts were estimated using linear mixed models and

meta-analysis. Associations between the PFAS mixture and individual bioactive lipids were

estimated using quantile g-computation.

Results: PFDeA, PFOA, and PFUdA were associated ($p<0.05$) with changes in bioactive lipid

levels in all three enzymatic pathways (cyclooxygenase [n=6 signatures]; cytochrome p450 [n=5

signatures]; lipoxygenase [n=7 signatures]) in at least one combined cohort analysis. The

strongest signature indicated that a doubling in PFOA corresponded with a 24.3% increase (95%

CI [7.3%, 43.9%]) in PGD2 (cyclooxygenase pathway) in the combined cohort. In the mixtures

analysis, we observed nine positive signals across all pathways associated with the PFAS

mixture. The strongest signature indicated that a quartile increase in the PFAS mixture was

associated with a 34% increase in PGD2 (95% CI [8%, 66%]), with PFOS contributing most to

the increase.

Conclusions: Bioactive lipids were revealed as biomarkers of PFAS exposure and could provide

mechanistic insights into PFAS' influence on pregnancy outcomes, informing more precise risk

estimation and prevention strategies.

Keywords: PFAS; Mixtures; Bioactive lipids; Eicosanoids

Abbreviations:

CiOB, Chemicals in Our Bodies;

CV, coefficient of variation;

Et-PFOSA-AcOH, 2-(N-Ethyl-perfluorooctane sulfonamido) acetic acid;

IKIDS, Illinois Kids Development Study;

Me-PFOSA-AcOH, 2-(N-Methyl-perfluorooctane sulfonamido) acetic acid;

NHANES, National Health and Nutrition Examination Study;

PFAS, Per- and poly-fluoroalkyl substances;

PFBS, Perfluorobutanesulfonic acid;

PFDeA, Perfluorodecanoic acid;

PFDoA, Perfluorododecanoic acid;

PFHpA, Perfluoroheptanoic acid;

PFHxS, Perfluorohexane sulfonate;

PFNA, Perfluorononanoic acid;

PFOA, Perfluorooctanoic acid;

PFOS, Perfluorooctanesulfonic acid;

PFOSA, Perfluorooctanesulfonamide;

PFUdA, Perfluoroundecanoic acid;

R^2, regression coefficients;

RSD, relative standard deviation;

UIUC, University of Illinois Urbana-Campaign;

WQS, weighted quantile sum.

1. Introduction

Widespread environmental contamination of per- and poly-fluoroalkyl substances (PFAS) poses a major public health concern. Human exposure to PFAS can occur through ingestion of contaminated food and water, as well inhalation of indoor air from contaminated by consumer products (e.g., clothing, furniture, cookware) (Sunderland et al., 2019). Biomonitoring in the National Health and Nutrition Examination Study (NHANES) has reported over 98% detection of multiple PFAS compounds in serum among individuals in the United States. (Calafat et al., 2007). Importantly, PFAS metabolism and excretion is slow due to their strong carbon-fluorine bonds, with half-lives in the human body spanning from several months to years (Chiu et al., 2022). Increasing experimental and epidemiological evidence indicates adverse health effects attributable to PFAS exposure, including liver toxicity, kidney dysfunction, hormone disruption, and reproductive and developmental toxicity (Fenton et al., 2021). Additionally, biomonitoring of PFAS in NHANES and the Environmental influences on Child Health Outcomes Program indicates moderate correlation between individual compounds, highlighting the need to consider PFAS as a mixture to evaluate cumulative health effects and ameliorate residual confounding (Calafat et al., 2007, Padula et al. 2023).

Pregnancy is a sensitive period of the life course, during which prenatal PFAS exposures have been identified as potential risk factors for adverse birth outcomes and pregnancy complications alike. For example, a systematic review of prenatal PFAS exposures identified select compounds associated with increased odds of preterm birth and miscarriage (Gao et al., 2021). There is also evidence indicating disparities in associations between PFAS exposure in women and chronic health outcomes. For example, a previous study identified that Black women (median age 49) had higher risk of developing hypertension compared to White women (Ding et

al. 2023). Our team also identified increased depressive symptoms in association with higher

PFAS exposure among immigrant pregnant women compared to U.S. born pregnant women

(median age 34) in the Chemicals in Our Bodies cohort (Aung et al. 2023). Widespread evidence

of environmental and human exposure, coupled with associated health effects, warrants detailed

investigation into intermediate mechanisms of PFAS toxicity to inform risk assessment and

identify potential intervention targets.

Bioactive lipids, including poly-unsaturated fatty acids such as arachidonic acid are

metabolized by conserved families of enzymes (e.g. cytochrome p450s, lipoxygenases, and

cyclooxygenases) to yield secondary eicosanoid metabolites with important downstream

physiological functions (Yuan et al., 2018). For example, eicosanoids partly regulate

inflammation and influence cardiovascular and renal function, and perturbations in circulating

eicosanoids may be important biomarkers of adverse pregnancy outcomes (Eek et al., 2012; Ma

et al., 2017; Patel et al., 2004). In a previous metabolomic study in Atlanta, Georgia, lipid

metabolism pathways were associated with gestational age at birth, including bioactive lipid

compounds such as linoleic acid (Taibl et al., 2023). We have also shown in a previous study in

the LIFECODES cohort that several eicosanoid metabolites were associated with increased risk

of spontaneous preterm birth (Aung et al., 2019). Another study in the LIFECODES cohort

identified eicosanoid metabolites from the cytochrome p450 and lipoxygenase pathway

associated with greater risk of being born small for gestational age (Welch et al., 2020).

A review of PFAS exposures and non-targeted metabolomics in epidemiologic studies

identified substantial evidence that PFAS are associated with changes to several biological

pathways, including metabolism of bioactive lipids, amino acids, and xenobiotic detoxification

(Guo et al., 2022). These findings are supported by experimental studies indicating that PFAS

can interfere with cytochrome p450 signaling, which is critical for cellular metabolism (Hvizdak

et al., 2023; Zanger and Schwab, 2013). PFAS can also interfere with homeostasis of

intracellular calcium gradients, which in turn can impact calcium dependent enzymatic activity

and alter systemic bioactive lipid profiles (Cao and Ng, 2021). Given the increasing evidence

that PFAS are linked to whole pathways of lipid metabolism, there is a need for deeper

investigation of targeted lipid metabolites to determine precise molecular signatures of PFAS

exposure and implications for pregnancy outcomes and fetal development.

The objective of this study was to quantify maternal PFAS exposures and bioactive lipids

during pregnancy and estimate the individual and cumulative associations of PFAS on these

biomarkers. There is a need for modern epidemiology studies to integrate greater racial,

socioeconomic, and geographic diversity to better inform risk estimation across historically

marginalized communities and for varied contexts of PFAS exposure levels. Thus, in the current

study, we utilized a diverse study sample across three birth cohorts in the Environmental

influences on Child Health Outcomes Program. Our study hypothesis was that higher

concentrations of PFAS are associated with increased bioactive lipid signatures across enzymatic

groups that govern eicosanoid synthesis.

2. Methods

2.1 Study Populations

The Environmental influences on Child Health Outcomes (ECHO) Program comprises 69

ongoing and new pregnancy and pediatric cohorts. The ECHO Program's goal is to advance

research that increases understanding of how environmental factors spanning from preconception

through childhood influences child health and development (Knapp et al. 2023). This study

integrates data across three ECHO cohorts (**Table 1**): Chemicals in Our Bodies (CiOB), Illinois

Kids Development Study (IKIDS), and the ECHO-PROTECT cohort (Eick et al., 2021;

Ferguson et al., 2019). The CiOB cohort participants are based in San Francisco, CA, and were

recruited during the second trimester of pregnancy at three University of California San

Francisco hospitals. Women included in the CiOB cohort had to be at least 18 years of age,

speak English or Spanish as a primary language, and be having a singleton pregnancy. The

IKIDS cohort participants were recruited between 10 and 14 weeks' gestation from two obstetric

clinics in Champaign-Urbana, Illinois. Women included in the IKIDS cohort had to be between

18 and 40 years of age, have English fluency, have a singleton pregnancy, be $\leq$ 15 weeks

gestation at enrollment, not have a child already in the IKIDS cohort, reside within a 30 minute

drive from the University of Illinois Urbana-Campaign (UIUC) campus, and plan to remain in

the area until at least the child's first birthday. The ECHO-PROTECT cohort participants were

recruited before 20 weeks' gestation from two hospitals and five clinics in the Northern Karst

aquifer region in Puerto Rico. Enrollment of the cohort began in 2011 and is ongoing. Inclusion

criteria for the women in the ECHO-PROTECT cohort included being between 18 and 40 years

of age, residing in the Northern Karst aquifer region, not have used oral contraceptives during

the three months pre-pregnancy, not have undergone in-vitro fertilization, and have no major

pre-existing conditions. Detailed information on study recruitment and data collection on

sociodemographic and health information for each cohort has been previously described (Eick et

al., 2021; Ferguson et al., 2019). Participants in each cohort provided written informed consent

to be included in this study and local institutional review boards for each cohort reviewed and

approved study protocols. Sample sizes for each cohort are described in greater detail below, and

the flow diagram for selection of the final analytical dataset used in statistical analyses is

illustrated in **Supplemental Figure 1**.

2.2. PFAS Exposure Assessment

152   In the CiOB cohort (n = 73, median gestational age of sample collection = 23 weeks) and

IKIDS (n = 287, median gestational age of sample collection = 17 weeks) cohorts, twelve PFAS

compounds were quantified in maternal serum during pregnancy: Perfluorononanoic acid

(PFNA), Perfluoroheptanoic acid (PFHpA), Perfluorodecanoic acid (PFDeA),

Perfluorododecanoic acid (PFDoA), Perfluorooctanoic acid (PFOA),

Perfluorooctanesulfonamide (PFOSA), 2-(N-Methyl-perfluorooctane sulfonamido) acetic acid

(Me-PFOSA-AcOH), 2-(N-Ethyl-perfluorooctane sulfonamido) acetic acid (Et-PFOSA-AcOH),

Perfluoroundecanoic acid (PFUdA), Perfluorohexane sulfonate (PFHxS),

Perfluorooctanesulfonic acid (PFOS), and Perfluorobutanesulfonic acid (PFBS). After sample

collection, serum was frozen at -80°C. Samples were processed at the Environmental Chemical

Laboratory at the California Department of Toxic Substances Control using a previously

described analytical protocol (Morello-Frosch et al., 2016). PFAS quantification was achieved

using internal standards in each serum sample and an automated on-line solid phase extraction

method coupled to liquid chromatography and tandem mass spectrometry. The limits of

detection for individual PFAS compounds were equal to three times the standard deviation of the

blank negative control sample (Morello-Frosch et al., 2016). In each batch of serum PFAS

analyses, analytes were quantified using a constructed calibration, and regression coefficients

(R^2) of 0.98 to 0.99 were generally obtained. We utilized in-house quality control materials that

were prepared by spiking a known amount of PFAS compounds in blank bovine serum at low

and high levels. We utilized standard reference materials (SRM 1958) from the National Institute

of Standards and Technology (Gaithersburg, MD), and quality control samples spiked with

known PFAS concentrations from the United States Centers for Disease Control and Prevention

as reference materials. Blank samples of bovine serum (Hyclone/GE Healthcare LifeSciences)

were also processed with each batch of samples, and no PFAS were detected above their

respective limited of detection in these blank samples.

Maternal serum samples from the ECHO-PROTECT cohort (n = 54, median gestational

age of sample collection =26 weeks) were used to quantify nine PFAS compounds (PFNA,

PFHpA, PFDeA, PFOA, PFOSA, Me-PFOSA-AcOH, PFUdA, PFHxS, PFOS). After sample

collection, serum was frozen at -80°C and processed at NSF International (Ann Arbor, MI,

USA). PFAS quantification was also performed by liquid chromatography and tandem mass

spectrometry. The method simulates the Centers for Disease Control and Prevention's

polyfluoroalkyl chemicals laboratory procedure method No: 6304.1. Standards of known purity

and identity were used during preparation of the calibration, quality control, and internal

standards. The validated analyte calibration curve correlation coefficient (R^2) ranges were 0.996

or greater. The method accuracy (% nominal concentration) and precision (% relative standard

deviation [RSD]) were determined through six replicate analyses of analytes spiked at three

different concentrations in pooled human serum across validation runs on three separate days (n

= 18), which reflects both the intra-day and inter-day variability of the assay. The accuracy (%

nominal concentration) range across all analytes was 95.1–103% with precision (%RSD) range

for the serum quality control samples across all analytes being 2.3–16%.

For samples with values below the limit of detection, we used machine read values (if

reported) or concentrations imputed using the limit of detection divided by the square root of 2

(Hornung and Reed, 1990). Of the PFAS compounds measured, we selected to analyze those

with a detection rate of ≥ 70%, which included PFNA, PFHpA, PFDeA, PFOA, Me-PFOSA-

AcOH, Et-PFOSA-AcOH, PFUdA, PFHxS, and PFOS in the CiOB cohort; PFNA, PFDeA,

PFOA, Me_PFOSA-AcOH, PFUdA, PFHxS, and PFOS in the IKIDS cohort; and PFNA and

PFOS in the ECHO-PROTECT cohort (**Table 2A**).

2.3. Bioactive Lipids Assay

Maternal plasma samples were used to quantify a targeted panel of 50 bioactive lipids (**Table

2B)** in the CiOB (n = 73, median gestational age of sample collection = 24 weeks), IKIDS (n =

287, median gestational age of sample collection = 17 weeks), and PROTECT (n = 54, median

gestational age of sample collection = 26 weeks) cohorts. For ECHO-PROTECT and CiOB

cohorts, plasma was collected using ethylenediaminetetraacetic acid plasma tubes and

temporarily stored at +4°C for less than four hours. Blood was subsequently centrifuged for 20

minutes and stored at -80°C. For participants in the IKIDS cohort, women provided a fasted (10

to 12 hours) blood sample collected in green-top sodium heparin tubes. IKIDS participant blood

samples were kept at room temperature for two hours prior to processing and centrifuged at room

temperature for 20 minutes. The resulting plasma was aliquoted immediately into cryovials for

storage at -80°C. The targeted bioactive lipids panel consisted of five parent fatty acid

compounds and 45 eicosanoid metabolites derived from three enzymatic groups (lipoxygenases,

cytochrome p450s, and cyclooxygenases), and full names and abbreviations are documented in

Table 2B. Quantification of bioactive lipid concentrations was achieved using a 6490 Triple

Quadrupole mass spectrometer (Agilent, New Castle, DE, USA). In this setting, the mass

spectrometer was set to a targeted multiple reaction monitoring mode and individual biomarkers

were identified based on metabolite-specific fragmentation and retention time. Limits of

detection were not calculated across individual instrument analysis cycles and all measured

bioactive lipids had machine read values. Further information on instrumental parameters and

quality control/assurance have been previously documented (Afshinnia et al., 2020). Briefly, we

conducted sequential dilution of each internal standard in duplicate to establish linearity and

estimate the coefficient of variation (CV) of the measurements at various concentrations. We

also ran a pool of study samples at the beginning of each batch and then after each 12 samples

during mass spectrometry to assess drift in measurements over time as well as the batch-to-batch

variability.

2.4. Statistical Analyses

We utilized a conceptual model (**Figure 1**) to inform our statistical analyses. We tabulated

distributions of key covariates from our conceptual model for each of the cohorts in our study.

Spearman correlations were estimated between all high-detect (i.e., $\geq 70\%$ of observations

exceeded minimum level of detection in cohort) PFAS compounds and bioactive lipids within

each of the cohorts. Combined-cohort correlations were not conducted due to differences in high-

detect PFAS across cohorts. Based on which PFAS were highly detected in each cohort (**Table**

2A), associations were estimated in combined cohort samples with two (CiOB and IKIDS) or

three cohorts (CiOB, IKIDS, and ECHO-PROTECT) using linear mixed effect regressions

(random intercept for cohort), and meta-analysis. Significance was evaluated at a level of $\alpha \leq$

0.05. Associations between PFAS mixtures and bioactive lipids were estimated using quantile g-

computation in the combined cohort consisting of IKIDS and CiOB participants (Keil et al.,

2020). All analyses are described in greater detail below and were performed with R (version

4.3.0).

2.4.1. Combined cohort analysis

Linear mixed effects models were utilized to test combined-cohort pairwise associations

between all sampled bioactive lipids and common high-detect PFAS, using natural log-

transformed lipid levels as outcomes and natural log-transformed PFAS as exposures. These

linear mixed effects models included a random intercept for the cohort identity to partly

ameliorate bias from different laboratory analysis of PFAS and for heterogeneity in demographic

characteristics between cohorts. All effect estimates were back-transformed to enhance

interpretability and represent a percent change in a given bioactive lipid corresponding to a

doubling (or 100% increase) of individual PFAS. Unadjusted and adjusted random intercept

models were generated for all three cohorts using PFAS common to all three cohorts: PFNA and

PFOS (**Table 2A**). Unadjusted and adjusted random intercept models were also generated

combining the CiOB and IKIDS cohorts, using the high-detect PFAS common to those two

cohorts as exposures: ME-PFOSA-AcOH, PFDEA, PFHxS, PFNA, PFOA, PFOS, and PFUDA

(**Table 2A**). Adjusted models included variables selected *a priori* based on our previous

investigation of PFAS and biomarkers of lipid peroxidation and oxidative stress (Taibl et al.,

2022). The covariates included in the adjusted models were maternal age in years, education

(categorized into less than high school, high school/GED/some college, bachelor's degree, and

graduate degree), pre-pregnancy BMI, parity (categorized into 0 and 1 or more), and gestational

age in weeks at sampling collection for bioactive lipids. Missing observations for covariates were

omitted from adjusted models. Q-values were calculated for all pairwise models to control for

false discovery rates (Storey et al., 2015)

2.4.2. Meta-analysis

Pairwise effect estimates for PFAS and bioactive lipids from each individual cohort were

estimated using linear regression, adjusted for the same covariates modeled in the combined

cohort analysis with the exception of ECHO-PROTECT, which did not include parity as a

covariate due to 0% variance in the variable within the ECHO-PROTECT cohort. We combined

these effect estimates by meta-analysis using the METAL method (Willer et al., 2010), yielding

overall effect estimates and *p* values for pairwise associations between individual bioactive lipids

and PFAS. This method, used previously in genome-wide association studies, utilizes an inverse

variance calculation to weigh all beta coefficients by their standard errors. In doing so, this

approach allows researchers to obtain effect estimates across models with different underlying

populations and covariates, without requiring access to the raw data. Meta-analysis was

implemented for all three cohort-specific models, and then for the IKIDS and CiOB cohort-

specific models. Interpretation of cohort-specific effect estimates were treated as a sensitivity

analysis to evaluate consistencies in combined cohort and meta-analysis and reported in

supplementary materials.

2.4.3. PFAS mixtures analysis

To address the study goal of estimating the cumulative effect of multiple PFAS on

bioactive lipids, we used quantile g-computation (Keil et al., 2020), a generalization of the

weighted quantile sum (WQS) regression method (Carrico et al., 2015; Czarnota et al., 2015),

which does not assume directional homogeneity of the exposures' effects on the outcome. This

method outputs an unbiased estimate of the effect on a particular bioactive lipid associated with

simultaneously increasing all PFAS exposures by one quartile. Quantile g-computation was

implemented in the combined CiOB and IKIDS sample to measure the mixtures effect of log-

transformed high-detect PFAS common to both the CiOB and IKIDS cohorts (Et-PFOSA-AcOH,

PFDEA, PFHxS, PFNA, PFOA, PFOS, and PFUDA**; Table 2A**) on all log-transformed bioactive

lipids, while controlling for the same covariates included in the adjusted combined CiOB and

IKIDS random intercept model. The cohort itself was included as an additional fixed effect

covariate in the models to account for any between-cohort differences (Taibl et al., 2022).

3. Results

3.1. Descriptive statistics

Table 1 shows the distribution of maternal race/ethnicity, age, education, pre-pregnancy

BMI, parity, household income category, and gestational age at plasma collection for bioactive

lipids across all three cohorts. White women represented the largest group in CiOB (47%) and

IKIDS (78%), and ECHO-PROTECT had only Hispanic women (100%). Women in CiOB were

the oldest with a median age of 33 years, followed by IKIDS (median age = 32 years), and

ECHO-PROTECT (median age = 28 years) ($p < 0.01$). Women in IKIDS had the highest

educational attainment, with 86% having completed a bachelor's or graduate degree, followed by

CiOB (72%) and ECHO-PROTECT (47%; $p < 0.01$). All mothers in ECHO-PROTECT reported

having at least one previous birth, which was much higher compared to CiOB (49%) and IKIDS

(46%, $p < 0.01$). The annual household income for women in CiOB was the highest with 65%

reporting annual household incomes of ≥ \$100,000, followed by women in IKIDS (34%), while

93% of women in ECHO-PROTECT reported a household income < \$50,000 ($p < 0.01$). Income

was not adjusted for local cost of living or median salary. Pre-pregnancy BMI was similar across

all three cohorts (median 24 kg/m^2 in CIOB and 25 kg/m^2 in IKIDS and ECHO-PROTECT, $p =$

0.2).

Distributions of PFAS and bioactive lipids by cohort are reported in **Table 2A** and

Supplemental Table 1, respectively. Among the high-detect PFAS common to all three cohorts,

PFNA (median 0.20 ng/mL) and PFOS (median 1.78 ng/mL) concentrations were the lowest in

the ECHO-PROTECT cohort, ($p < 0.01$) while the CiOB cohort had the highest median

concentration of PFNA (0.53 ng/mL), and the IKIDS cohort had the highest median

concentration of PFOS (3.27 ng/mL). Among the high-detect PFAS common to only the CiOB

and IKIDS cohorts, the greatest differences were observed for PFUdA (median 0.18 ng/mL in

CiOB and median 0.06 ng/mL in IKIDS; $p < 0.01$) and PFHxS (median 0.55 ng/mL in CiOB and

median 0.76 ng/mL in IKIDS; $p = 0.02$). Among the bioactive lipids, we observed the greatest

difference across cohorts for linoleic acid concentrations (median 668 µMol/L in CiOB, 1,185

µMol/L in IKIDS, and 102 µMol/L in ECHO-PROTECT; $p < 0.01$) and 20-carboxy arachidonic

acid (CAA) concentrations (median 185 nMol/L in CiOB, 302 nMol/L in IKIDS, and 74 nMol/L

in ECHO-PROTECT; $p < 0.01$).

3.2. Within-Cohort Correlations

Within-cohort Spearman correlations between bioactive lipids and PFAS are shown in

Figure 2A for the CiOB cohort, **Figure 2B** for the IKIDS cohort, and **Figure 2C** for the ECHO-

PROTECT cohort. In CiOB, correlation coefficients between bioactive lipids and PFAS ranged

between -0.31 and 0.38, with the strongest negative correlation between DHA and PFOS, and the

strongest positive correlation between 13S-HODE and PFUDA. For bioactive lipids in the CiOB

cohort, correlation coefficients ranged from -0.47 (16-HETE and 13-oxoODE) and 1 (12(13)-

EpoME and 13S-HODE), while correlation coefficients for PFAS were between -0.08 (PFDeA

and Et-PFOSA-AcOH) and 0.81 (PFNA and PFOS). In IKIDS, correlation coefficients between

bioactive lipids and PFAS ranged between -0.14 and 0.22, with the strongest negative correlation

between 15-HETE and PFHxS, and the strongest positive correlation between arachidonic acid

and PFNA. Correlation coefficients for bioactive lipids in the IKIDS cohort were between -0.75

(13,14-D-PGJ2 and 18-HETE) and 0.89 (9S-HODE and 13S-HODE), and between 0.05 (PFOA

and Me-PFOSA-AcOH) and 0.75 (PFNA and PFOA) for PFAS. In ECHO-PROTECT,

correlation coefficients between bioactive lipids and PFAS ranged between -0.37 and 0.32, with

the strongest negative correlation between 9,10-DiHOME and PFNA, and the strongest positive

correlation between 13,14-D-PGJ2 and PFNA. Bioactive lipid correlation coefficients within the

ECHO-PROTECT cohort were between 0.59 (PGA2 and 11-HETE) and 0.8 (9(10)-EpoME and

12(13)-EpoME), and the correlation coefficients between PFNA and PFOS in the ECHO-

PROTECT cohort was 0.57.

3.3. Pair-wise associations between individual PFAS and bioactive lipids

All coefficients representing the pairwise associations between bioactive lipids and PFAS from

the adjusted random intercept models and meta-analyses are presented in a heatmap in **Figure 3**

as the effect per doubling of PFAS concentration. Additional output of these models, including

exact *p* values and standard errors can be found in **Supplemental Tables 2-5**. Additionally,

effect estimates and 95% confidence intervals from significant associations found in at least one

of the four models are reported in **Table 3**. With the exception of the associations between 9-

oxoODE (cytochrome p450 pathway) and PFNA, 15-HETE (lipoxygenase pathway) and PFHxS,

and αLA and PFUdA (parent compounds), all significant associations observed in at least one of

the four combined models were positive.

In the cyclooxygenase pathway, 15 significant associations between bioactive lipids and

PFAS were observed in at least one of the four combined cohort models. Of these, the

association between BCPGE1 and PFNA was significant in all four models (doubling effect in

PFNA ranging from 10.3% to 12.8% increase in BCPGE1 across models). Additionally, the

associations between 9-oxoODE and PFNA (doubling effect in PFNA ranging from 2.3% to

2.4% decrease in 9-oxoODE), PGD2 and PFOS (doubling effect in PFOS ranging from 11.9% to

17.0% increase in PGD2), and PGD3 and PFOS (doubling effect in PFOS ranging from 10.0% to

11.5% increase in PGD3) were significant in three of the four models. In the cytochrome p450

pathway, no significant associations were found which were common to all four models;

however, there were significant positive associations between 14(15)-EET and PFDeA (doubling

effect ranging from 6.4% to 6.6%) and between 8(9)-EET and Me-PFOSA-AcOH (doubling

effect ranging from 5.4% to 5.8%) observed in the CiOB and IKIDS random intercept model and

in the meta-analysis integrating effect estimates from CiOB and IKIDS. Within the lipoxygenase

pathway, significant positive associations were observed between 12-oxoETE and PFOS

(doubling effect ranging from 7.1% to 8.2%) and 15-oxETE and PFOS (doubling effect ranging

from 10.3% to 10.4%) across all four combined cohort models. Among the parent compounds,

significant positive associations were observed for arachidonic acid and PFNA (doubling effect

ranging from 8.7% to 10.9%) and arachidonic acid and PFOS (doubling effect ranging from

8.0% to 9.7%) across all four combined cohort models.

We explored multiple testing comparison adjustments in the linear mixed effects models

(**Supplemental Tables 2** and **3**). While none were below a threshold of 0.1 in the two cohort or

three cohort models, there were eight PFAS and bioactive lipid pairs (PGD2 and PFOA; PGD3

and PFUdA; 13,14DHK-PGF2a* and PFUdA; 12-oxoETE and PFUdA; 15-oxoETE and PFUdA;

15-oxoETE and PFOS; arachidonic acid and PFNA; arachidonic acid and PFOS) below the 0.2

threshold in the two cohort models. Our sensitivity analysis of within-cohort pairwise

associations can be found in **Supplemental Tables 6-11**. While individual cohort analyses are

underpowered to detect associations, directions of pair-wise associations between individual

cohorts and combined cohorts were largely consistent.

3.4. PFAS Mixture Associations

Quantile g-computation utilizing the CiOB and IKIDS cohorts indicated that

simultaneously increasing all log-transformed PFAS (PFNA, PFDeA, PFoA, Me-PFOSA,

PFUdA, PFHxS, and PFOS) in the mixture by one quartile corresponded to increases in

BCPGE1, PGD2, PGD3, and PGE3 in the cyclooxygenase pathway; 12,13-DiHOME and 5(6)-

EET in the cytochrome p450 pathway; 12-oxoETE, 15-oxoETE, and 5-ETE in the lipoxygenase

pathway, and arachidonic acid and linoleic acid parent compounds. The largest increase in the

cyclooxygenase pathway was observed in PGD2 (34% increase, 95% CI [8%, 66%]), with

decomposition of the quantile g-computation effect estimate for PGD2 indicating that PFOS

exhibited the greatest positive weight to the overall mixture effect relative to the other PFAS

compounds in the overall sample (**Supplemental Table 12**). In the cytochrome p450 pathway,

the largest increase associated with a simultaneous one quartile increase in all PFAS in the

mixture was in 5(6)-EET (31% increase, 95% CI [5%, 64%]), with PFHxS contributing the most

to this increase. The largest increase in the lipoxygenase pathway was observed in 12-oxoETE

(16% increase, 95% CI [4%, 30%]), with PFUdA having the largest contribution to this increase.

In the parent compounds, linoleic acid had the largest increase (18% increase, 95% CI [1%,

38%]), driven predominantly by PFUdA. No significant decreases in bioactive lipids were

observed. These effects are visualized in **Figure 4** and reported in detail in **Supplemental Table**

12.

4. Discussion

4.1. Summary of findings across statistical approaches

In the present study, we advanced mechanistic insight into PFAS exposures during

pregnancy by estimating associations between PFAS and bioactive lipids in single pollutant and

mixture models. We utilized multiple statistical approaches to estimate associations, including

cohort stratified analyses, combined cohort analyses, and meta-analyses. Combined cohort

analyses and meta-analyses identified mostly positive associations of PFAS with parent fatty

acid compounds and their secondary eicosanoid metabolites derived from the lipoxygenase,

cytochrome p450, and cyclooxygenase pathways. Individual cohorts have limited power to

detect significant associations, which underscores the utility of exploring combined cohort

analyses and meta-analyses. Additionally, mixtures analysis using quantile g-computation

revealed that the entire PFAS mixture exhibited predominantly positive associations with

bioactive lipids across all three enzymatic pathways (BCPGE1, PGD2, PGD3, PGE3, 12(13)-

DiHOME, 5(6)-EET, 12-oxoETE, 15-oxoETE, 5-HETE, arachidonic acid, and linoleic acid),

and, with the exception of 12(13)-DiHOME, at least one individual PFAS was associated with

these bioactive lipids in the single pollutant models for either combined cohort analyses or meta-

analyses. Largely consistent results between mixtures analysis and pair-wise associations

strengthen confidence in targeted bioactive lipids as a potential mechanistic biomarker of PFAS

exposure and provide insight into addressing the health effects of PFAS as an entire class. Doing

so can inform more precise risk estimation and ameliorate the influence of residual confounding

resulting from co-occurring PFAS compounds. Although our estimated associations are

susceptible to false-positive associations, our findings still allow for prioritization of pairs of

PFAS and bioactive lipids for future hypothesis testing and replication in independent samples.

4.2. Biological context of associations in bioactive lipid enzymatic pathways

The physiological implications of our findings vary based on the metabolic pathway

under investigation. In our study, we observed the strongest positive effect between the PFAS

mixture and cyclooxygenase derived metabolite PGD2, which was consistent with individual

PFAS compound analyses for PFDeA, PFOA, and PFOS in the CiOB and IKIDS meta-analyses.

Systemic inflammation and oxidative stress are both antecedent physiological states that increase

the risk of adverse pregnancy outcomes such as spontaneous preterm birth and preeclampsia

(Ferguson and Chin, 2017; Gomez-Lopez et al., 2022). Cyclooxygenases have been implicated in

promoting inflammation through the production of prostaglandins (Ricciotti and FitzGerald,

2011). Prostaglandin production is also sensitive to imbalances in reactive oxygen species and

oxidative stress (Burdon et al., 2007). Previous mechanistic studies indicate that

cyclooxygenases are also important for regulating reproductive health and fetal development,

where animal models have found that deficiencies in the genes encoding these enzymes result in

altered implantation, increased mortality, and impaired organ development in offspring (Xu et

al., 2007). Animal studies indicate that disruptions in cyclooxygenase function and prostaglandin

synthesis can also lead to altered neurodevelopment and behavior (Davis-Bruno and Tassinari,

2011; Wong et al., 2019). For example, a previous study in an experimental rat model found that

the PGD2 signaling pathway is involved in neuroinflammation, and induction of this pathway

results in neurodegenerative pathologies (Corwin et al., 2018). Evidence from previous

experimental mechanistic studies in combination with our findings that PFAS are linked to

altered eicosanoid concentrations within the cyclooxygenase pathway underscore the need to

further investigate this pathway as a potential mediator between PFAS and adverse pregnancy

outcomes.

In our study, we observed that the PFAS mixture was associated with two cytochrome

p450 derived eicosanoids: 12,13-DiHOME and 5(6)-EET. Cytochrome p450 enzymes have

varied regulatory roles, including biosynthesis of endogenous hormones, detoxification of

xenobiotics, and cellular metabolism (Zanger and Schwab, 2013). Importantly, *in vitro*

experiments indicate that multiple PFAS compounds can directly interfere and inhibit

cytochrome p450 activity (Hvizdak et al., 2023). 12,13-DiHOME has been classified as an

oxylipin derived from linoleic acid, and is involved with inflammation, endocrine signaling, and

adipogenesis (Hildreth et al., 2020; Macêdo et al., 2022). In a previous case-control study that

our research team led in the LIFECODES cohort using the same bioactive lipids panel measured

(median 26 weeks' gestation) in this present study, we reported that 12,13-DiHOME was

associated with increased risk of spontaneous preterm birth (n_{cases} = 31, $n_{controls}$ = 115) (Aung et

al., 2019). Further, placental 5(6)-EET has been detected at higher levels in preeclamptic women

compared to controls, linking 5(6)-EET to regulation pathways associated with preeclampsia

(Dalle Vedove et al., 2016; Herse et al., 2012). Additionally, a study of 146 adult women

reported that single nucleotide polymorphisms of cytochrome p450 genes amplified cancer risk

attributable to PFAS exposures (Ghisari et al., 2014). While bioactive lipids were not measured

in that study, this is consistent biological inference based on their findings of increased breast

cancer risk in association with higher PFOS and PFOA and polymorphisms in cytochrome p450

genes. Therefore, cytochrome p450 derived eicosanoids may link PFAS to adverse maternal

health outcomes and serve as early signals to inform precise prevention efforts.

Lipoxygenases are calcium dependent enzymes that catalyze the formation of

hydroperoxides from poly-unsaturated fatty acids and have been associated with several adverse

health outcomes including asthma, skin disorders, and cancers (Mashima and Okuyama, 2015).

In our study, we observed that the PFAS mixture was associated with higher levels of the

lipoxygenase derived eicosanoids 12-oxoETE, 15-oxoETE, and 5-HETE. These oxylipins may

be important biomarkers for metabolic and cardiovascular disorders. For example, 12-oxoETE

has been explored as a biomarker of diabetic macular edema (Rhee et al., 2021) and

experimental models indicate that 15-oxoETE may influence atherosclerosis (Ma et al., 2017).

Additionally, our previous study in the LIFECODES cohort identified higher levels of 12-

oxoETE and 5-HETE in association with spontaneous preterm birth (Aung et al., 2019). Our

present study findings showing that increases in 12-oxoETE and 15-oxoETE were associated

with increased PFAS exposure are further aligned with previous experimental evidence showing

that PFAS can interfere with intracellular calcium gradients which can influence the catalytic

activity of lipoxygenases (Cao and Ng, 2021). Therefore, future studies should continue to

investigate this pathway as a mechanistic link between PFAS exposure and adverse maternal and

child health outcomes.

We observed differences in bioactive lipid compound distributions between individual

cohorts, which may be driven by the variation in genetic makeup of the populations caused by

heterogeneous ethnic composition of the cohorts (Sergeant et al., 2012), or differences in

environmental exposures of the cohorts due to diet (Saadatian-Elahi et al., 2009). Among the

parent poly-unsaturated fatty acids, the PFAS mixture was associated with increased

concentrations of arachidonic acid and linoleic acid. These findings are aligned with the positive

signatures we observed for the secondary eicosanoid metabolites described above that are

derived from these parent compounds. Our reported findings can partially be contextualized with

previous metabolomics studies to confer biological inference of PFAS-induced effects on lipid

metabolism. One metabolomics study investigated a mixture of six common PFAS (PFOS,

PFHxS, PFHpS, PFOA, PFNA, and PFDA) in children and adolescents (n = 137) based in Los

Angeles, CA, and observed positive associations of the mixture with arachidonic acid and

linoleic acid (Goodrich et al., 2023). Another study of 267 maternal-newborn dyads in Atlanta,

GA, reported that maternal PFAS exposures were associated with newborn metabolomic

signatures for bioactive lipid metabolism (including leukotrienes), cytochrome p450 pathway,

and linoleic acid (Taibl et al., 2023). Collectively, metabolomics studies underscore the

importance of lipid metabolism as a potential intermediate mechanism of PFAS exposure.

4.3. Strengths and limitations

Our study has notable strengths. First is the diversity of our study population, which

included pregnant women from three distinct geographic areas, with high heterogeneity across

demographics, socioeconomic status, and PFAS exposures. Second is our statistical approach.

We performed combined-cohort analysis utilizing two methods (linear mixed effects models and

meta-analysis), and a mixtures analysis using quantile g-computation to assess cumulative

associations with PFAS mixtures. The combination of methods applied and the consistency in

results across methods strengthens inference in identified associations. Third was our selection of

outcomes, which was a targeted assay of bioactive lipids that has not been previously tested for

associations with PFAS. This targeted assay directly complements existing studies that have

utilized non-targeted metabolomics by deepening knowledge of specific lipid metabolite features

to contextualize larger biological pathways and processes observed in past studies. Combined,

this resulted in a robust investigation, substantiated by the corroboration of our key findings both

within our study and with previous literature.

Our study also has limitations to consider. First is the cross-sectional nature of data

collection, as serum PFAS and plasma bioactive lipids were measured during the same visit.

Single time point assessment is more susceptible to measurement error and reverse causation

than longitudinal studies. Because of their long half-lives, PFAS measures may be relatively

stable during pregnancy, however, future studies should consider repeated measures of both

PFAS and bioactive lipid levels to reduce measurement error and evaluate more precise windows

of vulnerability during pregnancy. Additionally, some confounding variables (e.g. maternal

education and parity) were heterogenous across cohorts. There are also unmeasured confounders

that we did not model in the present study such as consumer product use and dietary intake

which may influence both PFAS and bioactive lipid concentrations. However, exploration of

these confounders warrants more comprehensive assessment of the contribution that diet and

consumer products have on both PFAS and bioactive lipids, which was beyond the scope of the

present study. In terms of statistical approaches, we recognize there are multiple important

approaches for analyzing chemical mixtures in health studies. Our study focused on cumulative

PFAS mixture associations. However, future studies may consider alternative approaches such as

investigation of non-linear effects, high-order interactions across multiple exposure variables,

latent clustering of correlated exposure variables, and dimension reduction through development

of risk scores or summation based on sources of exposures. Our study evaluated 12 serum PFAS

compounds, and there are thousands of PFAS compounds that humans may be exposed to.

Therefore, we are likely underestimating the effect of PFAS as a whole class on bioactive lipids.

Further, limited sample size and low detection rates of PFAS chemicals in the PROTECT cohort

limited our ability to examine differences in associations between Hispanic and White women

exposed to PFAS. Additionally, we have previously documented associations between other

classes of endocrine disrupting chemicals (e.g., phthalates, phenols, and parabens) and bioactive

lipids during pregnancy in the LIFECODES cohort (N = 173) (Aung et al. 2021). It is possible

that these chemicals are correlated with prenatal PFAS exposures, contributing to residual

confounding and potential interactions, which may influence the effects of PFAS on maternal

bioactive lipid profiles. Therefore, future investigations should consider broader chemical

mixtures analyses with bioactive lipids and pregnancy outcomes.

4.4. Conclusions

In conclusion, we observed associations between PFAS exposure and bioactive lipids during

pregnancy in all three enzymatic pathways (cytochrome p450, lipoxygenase, and

cyclooxygenase), most of which were positive. Our study complements recent efforts in

disentangling endogenous biomarker signatures of PFAS exposures by identifying specific lipid

metabolites as biomarkers of exposure and potential mechanistic targets. Future studies should

confirm these findings among cohorts of different demographic and socioeconomic makeup,

including consideration of meta-analysis and PFAS mixture analysis to strengthen confidence in

reported associations. These findings may be utilized to develop hypothesis driven mediation

analyses to link PFAS exposures to downstream maternal and child health outcomes. Further,

these findings can inform precise risk estimation aimed at reducing the harm of the PFAS

exposures on vulnerable populations.

CRediT authorship contribution statement:

Funding acquisition (MTA, AA, JDM, JFC, RMF, TJW, SLS, AA); Conceptualization (MTA,

HS, AA); Data curation (ED, SN, MW, SME, AC, SP, LZ, JSP, DW, SS, AA, SDG, TJW, RMF,

SLS, JDM, AA, MTA, HS); Formal analysis (HS); Investigation (HS, MTA); Methodology (HS,

MTA, SME); Literature Review (TM, DP, HS, BAR, MTA); Writing – original draft (HS,

MTA); Writing – Reviewing and editing (HS, TM, DP, MW, SME, AC, DJW, RSS, BAR, SP,

LZ, DW, JSP, SS, ED, AP, RCF, BM, AA, SDG, SN, GHM, CVV, ZR, JFC, EZ, TJW, RMF,

SLS, JDM, AA, MTA).

Declaration of competing interests: The authors declare they have no competing financial or

non-financial interests as defined by Environment International, or other interests that might be

perceived to influence the results and/or discussion reported in this paper.

Data and code availability: R-script used for the present study are found in supplemental

materials.

Select de-identified data from the ECHO Program are available through NICHD's Data and

Specimen Hub (DASH). Information on study data not available on DASH, such as some

Indigenous datasets, can be found on the ECHO study DASH webpage.

Acknowledgements:

The authors thank our Environmental Influences on Child Health Outcomes (ECHO) colleagues;

the medical, nursing, and program staff; and the children and families participating in the ECHO

cohorts. Furthermore, we acknowledge the contribution of the following ECHO program

collaborators:

ECHO Components—Coordinating Center: Duke Clinical Research Institute, Durham, North

Carolina: Smith PB, Newby LK; Data Analysis Center: Johns Hopkins University Bloomberg

School of Public Health, Baltimore, Maryland: Jacobson LP; Research Triangle Institute,

Durham, North Carolina: Catellier DJ; Person-Reported Outcomes Core: Northwestern

University, Evanston, Illinois: Gershon R, Cella D.

Funding: Research reported in this publication was supported by the Environmental Influences

on Child Health Outcomes (ECHO) Program, Office of the Director, National Institutes of

Health (NIH), under award numbers U2COD023375 (Coordinating Center), U24OD023382

(Data Analysis Center), U24OD023319 with co-funding from the Office of Behavioral and

Social Science Research (Pro-Core), U2COD023375 (Opportunities and Infrastructure Fund),

UH3OD023251 (Alshawabkeh), UH3OD023272 (Schantz and Woodruff). This work was

additionally supported by the Children's Environmental Health and Disease Prevention Research

Center (grant numbers ES022848 and RD83543401), the Molecular Phenotyping and

Metabolomics Core, Michigan Nutrition and Obesity Center (5P30DK089503), US

Environmental Protection Agency (grant number RD83543301), National Institute of

Environmental Health Sciences (NIEHS; grant numbers P42ES017198, P50ES026049,

P01ES022841, P30ES019776, P30ES030284, and P30ES007048). M.T.A and S.M.E. were also

supported in part by the JPB Foundation Environmental Health Fellows Program.

The content of this study is solely the responsibility of the authors and does not necessarily

represent the official views of the NIH.

Role of the Funder: The sponsor, NIH, participated in the overall design and implementation of

the ECHO Program, which was funded as a cooperative agreement between NIH and grant

awardees. The sponsor approved the Steering Committee-developed ECHO protocol and its

amendments including COVID-19 measures. The sponsor had no access to the central database,

which was housed at the ECHO Data Analysis Center. Data management and site monitoring

were performed by the ECHO Data Analysis Center and Coordinating Center. All analyses for

scientific publication were performed by the study statistician, independently of the sponsor. The

lead author wrote all drafts of the manuscript and made revisions based on co-authors and the

ECHO Publication Committee (a subcommittee of the ECHO Steering Committee) feedback

without input from the sponsor. The study sponsor did not review nor approve the manuscript for

submission to the journal.

Appendix A. Supplementary Figure

Appendix B. Supplementary Tables

References

Afshinnia, F., Zeng, L., Byun, J., Wernisch, S., Deo, R., Chen, J., Hamm, L., Miller, E.R., Rhee, E.P., Fischer, M.J., Sharma, K., Feldman, H.I., Michailidis, G., Pennathur, S., CRIC Study Investigators, 2020. Elevated lipoxygenase and cytochrome P450 products predict progression of chronic kidney disease. Nephrol. Dial. Transplant. 35, 303–312. https://doi.org/10.1093/ndt/gfy232

Aung, M.T., Yu, Y., Ferguson, K.K., Cantonwine, D.E., Zeng, L., McElrath, T.F., Pennathur, S., Mukherjee, B., Meeker, J.D., 2019. Prediction and associations of preterm birth and its subtypes with eicosanoid enzymatic pathways and inflammatory markers. Scientific Reports. 9, 1–17.

Aung M.T., Eick S.M., Padula A.M., Smith S., Park J.S., DeMicco E., Woodruff T.J., Morello-Frosch R., 2023. Maternal per- and poly-fluoroalkyl substances exposures associated with higher depressive symptom scores among immigrant women in the Chemicals in Our Bodies cohort in San Francisco. Environ. Int. 172, 107758. doi: 10.1016/j.envint.2023.107758.

Aung M.T., Yu Y., Ferguson K.K., Cantonwine D.E., Zeng L., McElrath T.F., Pennathur S., Mukherjee B., Meeker J.D., 2021. Cross-sectional estimation of endogenous biomarker associations with prenatal phenols, phthalates, metals, and polycyclic aromatic hydrocarbons in single-pollutant and mixtures analysis approaches. Environ. Health. Perspect. 129, 37007. doi: 10.1289/EHP7396.

Burdon, C., Mann, C., Cindrova-Davies, T., Ferguson-Smith, A.C., Burton, G.J., 2007. Oxidative stress and the induction of cyclooxygenase enzymes and apoptosis in the murine placenta. Placenta. 28, 724–733. https://doi.org/10.1016/j.placenta.2006.12.001

Calafat, A.M., Wong, L.-Y., Kuklenyik, Z., Reidy, J.A., Needham, L.L., 2007. Polyfluoroalkyl

chemicals in the US population: data from the National Health and Nutrition Examination

Survey (NHANES) 2003–2004 and comparisons with NHANES 1999–2000. Environ.

Health Perspect. 115, 1596–1602.

Cao, Y., Ng, C., 2021. Absorption, distribution, and toxicity of per-and polyfluoroalkyl

substances (PFAS) in the brain: a review. Environ. Sci.: Processes & Impacts. 23, 1623–

1640.

Carrico, C., Gennings, C., Wheeler, D.C., Factor-Litvak, P., 2015. Characterization of weighted

quantile sum regression for highly correlated data in a risk analysis setting. J Agric. Biol.

Environ. Stat. 20, 100–120. https://doi.org/10.1007/s13253-014-0180-3

Chiu, W.A., Lynch, M.T., Lay, C.R., Antezana, A., Malek, P., Sokolinski, S., Rogers, R.D.,

2022. Bayesian estimation of human population toxicokinetics of PFOA, PFOS, PFHxS,

and PFNA from studies of contaminated drinking water. Environ. Health Perspect.. 130,

127001. https://doi.org/10.1289/EHP10103

Corwin, C., Nikolopoulou, A., Pan, A.L., Nunez-Santos, M., Vallabhajosula, S., Serrano, P.,

Babich, J., Figueiredo-Pereira, M.E., 2018. Prostaglandin D2/J2 signaling pathway in a

rat model of neuroinflammation displaying progressive parkinsonian-like pathology:

potential novel therapeutic targets. J. Neuroinflammation. 15, 272.

https://doi.org/10.1186/s12974-018-1305-3

Czarnota, J., Gennings, C., Wheeler, D.C., 2015. Assessment of weighted quantile sum

regression for modeling chemical mixtures and cancer risk. Cancer Inform. 14, 159–171.

https://doi.org/10.4137/CIN.S17295

Davis-Bruno, K., Tassinari, M.S., 2011. Essential fatty acid supplementation of DHA and ARA

and effects on neurodevelopment across animal species: a review of the literature. Birth

Defects Res. B. Dev. Reprod. Toxicol. 92, 240–250. https://doi.org/10.1002/bdrb.20311

Dalle Vedove, F., Fava, C., Jiang, H., Zanconato, G., Quilley, J., Brunelli, M., Guglielmi, V.,

Vattemi, G., Minuz, P., 2016. Increased epoxyeicosatrienoic acids and reduced soluble

epoxide hydrolase expression in the preeclamptic placenta. J. Hypertens. 34, 1364-1370.

doi: 10.1097/HJH.0000000000000942.

Ding, N., Karvonen-Gutierrez, C.A., Zota, A.R., Mukherjee, B., Harlow, S.D., Park, S.K., The

role of exposure to per- and polyfluoroalkyl substances in racial/ethnic disparities in

hypertension: Results from the study of Women's health across the nation. Environ. Res.

227, 115813. doi: 10.1016/j.envres.2023.115813.

Eek, P., Järving, R., Järving, I., Gilbert, N.C., Newcomer, M.E., Samel, N., 2012. Structure of a

calcium-dependent 11R-lipoxygenase suggests a mechanism for Ca2+ regulation. J. Biol.

Chem. 287, 22377–22386. https://doi.org/10.1074/jbc.M112.343285

Eick, S.M., Enright, E.A., Geiger, S.D., Dzwilewski, K.L.C., DeMicco, E., Smith, S., Park, J.-S.,

Aguiar, A., Woodruff, T.J., Morello-Frosch, R., Schantz, S.L., 2021. Associations of

maternal stress, prenatal exposure to per- and polyfluoroalkyl substances (PFAS), and

demographic risk factors with birth outcomes and offspring neurodevelopment: an

overview of the ECHO.CA.IL prospective birth cohorts. Int. J. Environ. Res. Public

Health. 18, 742. https://doi.org/10.3390/ijerph18020742

Fenton, S.E., Ducatman, A., Boobis, A., DeWitt, J.C., Lau, C., Ng, C., Smith, J.S., Roberts,

S.M., 2021. Per☐and polyfluoroalkyl substance toxicity and human health review:

Current state of knowledge and strategies for informing future research. Environ.

Toxicol. and Chem.. 40, 606–630.

Ferguson, K.K., Chin, H.B., 2017. Environmental chemicals and preterm birth: Biological

mechanisms and the state of the science. Curr. Epidemiol. Rep. 4, 56–71.

https://doi.org/10.1007/s40471-017-0099-7

Ferguson, K.K., Rosario, Z., McElrath, T.F., Vélez Vega, C., Cordero, J.F., Alshawabkeh, A.,

Meeker, J.D., 2019. Demographic risk factors for adverse birth outcomes in Puerto Rico

in the PROTECT cohort. PLoS One. 14, e0217770.

https://doi.org/10.1371/journal.pone.0217770

Gao, X., Ni, W., Zhu, S., Wu, Y., Cui, Y., Ma, J., Liu, Y., Qiao, J., Ye, Y., Yang, P., Liu, C.,

Zeng, F., 2021. Per- and polyfluoroalkyl substances exposure during pregnancy and

adverse pregnancy and birth outcomes: A systematic review and meta-analysis. Environ.

Res. 201, 111632. https://doi.org/10.1016/j.envres.2021.111632

Ghisari, M., Eiberg, H., Long, M., Bonefeld-Jørgensen, E.C., 2014. Polymorphisms in Phase I

and Phase II genes and breast cancer risk and relations to persistent organic pollutant

exposure: a case–control study in Inuit women. Environ. Health. 13, 19.

https://doi.org/10.1186/1476-069X-13-19

Gomez-Lopez, N., Galaz, J., Miller, D., Farias-Jofre, M., Liu, Z., Arenas-Hernandez, M., Garcia-

Flores, V., Shaffer, Z., Greenberg, J.M., Theis, K.R., Romero, R., 2022. The

immunobiology of preterm labor and birth: intra-amniotic inflammation or breakdown of

maternal-fetal homeostasis. Reproduction. 164, R11–R45. https://doi.org/10.1530/REP-

22-0046

Goodrich, J.A., Walker, D.I., He, J., Lin, X., Baumert, B.O., Hu, X., Alderete, T.L., Chen, Z.,

Valvi, D., Fuentes, Z.C., Rock, S., Wang, H., Berhane, K., Gilliland, F.D., Goran, M.I.,

Jones, D.P., Conti, D.V., Chatzi, L.,2023. Metabolic signatures of youth exposure to

mixtures of per- and polyfluoroalkyl substances: A multi-cohort study. Environ. Health

Perspect.. 131, 027005. https://doi.org/10.1289/EHP11372

Guo, P., Furnary, T., Vasiliou, V., Yan, Q., Nyhan, K., Jones, D.P., Johnson, C.H., Liew, Z.,

2022. Non-targeted metabolomics and associations with per- and polyfluoroalkyl

substances (PFAS) exposure in humans: A scoping review. Environ. Int. 162, 107159.

https://doi.org/10.1016/j.envint.2022.107159

Herse, F., Lamarca, B, Hubel, C.A., Kaartokallio, T., Lokki, A.I., Ekholm, E., Laivuori, H.,

Gauster, M., Huppertz, B., Sugulle, M., Ryan, M.J., Novotny, S., Brewer, J., Park, J.K.,

Kacik, M., Hoyer, J., Verlohren, S., Wallukat, G., Rothe, M., Luft, F.C., Muller, D.N.,

Schunck, W.H., Staff, A.C., Dechend, R., 2012. Cytochrome P450 subfamily 2J

polypeptide 2 expression and circulating epoxyeicosatrienoic metabolites in

preeclampsia. Circulation. 126, 2990-2999. doi:

10.1161/CIRCULATIONAHA.112.127340.

Hildreth, K., Kodani, S.D., Hammock, B.D., Zhao, L., 2020. Cytochrome P450-derived linoleic

acid metabolites EpOMEs and DiHOMEs: A review of recent studies. J. Nutr. Biochem.

86, 108484. https://doi.org/10.1016/j.jnutbio.2020.108484

Hornung, R.W., Reed, L.D., 1990. Estimation of average concentration in the presence of

nondetectable values. Appl. Occup. and Environ. Hyg. 5, 46–51.

https://doi.org/10.1080/1047322X.1990.10389587

Hvizdak, M., Kandel, S.E., Work, H.M., Gracey, E.G., McCullough, R.L., Lampe, J.N., 2023.

Per- and polyfluoroalkyl substances (PFAS) inhibit cytochrome P450 CYP3A7 through

direct coordination to the heme iron and water displacement. J. Inorg. Biochem. 240,

112120. https://doi.org/10.1016/j.jinorgbio.2023.112120

Keil, A.P., Buckley, J.P., O'Brien, K.M., Ferguson, K.K., Zhao, S., White, A.J., 2020. A

quantile-based g-computation approach to addressing the effects of exposure mixtures.

Environ. Health Perspect. 128, 047004. https://doi.org/10.1289/EHP5838

Knapp, E.A., Kress, A.M., Parker, C.B., Page, G.P., McArthur, K., Gachigi, K.K., Alshawabkeh,

A.N., Aschner, J.L., Bastain, T.M., Breton, C.V., Bendixsen, C.G., Brennan, P.A., Bush,

N.R., Buss, C., Camargo, C.A. Jr., Catellier, D., Cordero, J.F., Croen, L., Dabelea, D.,

Deoni, S., . . ., Jacobson, L.P., Influences on Child Health Outcomes OBOPCFE., 2023.

The Environmental Influences on Child Health Outcomes (ECHO)-Wide Cohort. Am. J.

Epidemiol. 192, 1249-1263. doi: 10.1093/aje/kwad071.

Ma, G., Pan, B., Ren, S., Guo, C., Guo, Y., Wei, L., Zheng, L., Chen, B., 2017. 15-

oxoeicosatetraenoic acid mediates monocyte adhesion to endothelial cell. Lipids Health

Dis. 16, 137. https://doi.org/10.1186/s12944-017-0518-2

Macêdo, A.P.A., Muñoz, V.R., Cintra, D.E., Pauli, J.R., 2022. 12,13-diHOME as a new

therapeutic target for metabolic diseases. Life Sci. 290, 120229.

https://doi.org/10.1016/j.lfs.2021.120229

Mashima, R., Okuyama, T., 2015. The role of lipoxygenases in pathophysiology; new insights

and future perspectives. Redox Biol. 6, 297–310.

https://doi.org/10.1016/j.redox.2015.08.006

Morello-Frosch, R., Cushing, L.J., Jesdale, B.M., Schwartz, J.M., Guo, W., Guo, T., Wang, M.,

Harwani, S., Petropoulou, S.-S.E., Duong, W., Park, J.-S., Petreas, M., Gajek, R.,

Alvaran, J., She, J., Dobraca, D., Das, R., Woodruff, T.J., 2016. Environmental chemicals

in an urban population of pregnant women and their newborns from San Francisco.

Environ. Sci. Technol. 50, 12464–12472. https://doi.org/10.1021/acs.est.6b03492

Padula, A.M., Ning, X., Bakre, S., Barrett, E.S., Bastain, T., Bennett, D.H., Bloom, M.S., Breton,

C.V., Dunlop, A.L., Eick, S.M., Ferrara, A., Fleisch, A., Geiger, S., Goin, D.E., Kannan,

K., Karagas, M.R., Korrick, S., Meeker, J.D., Morello-Frosch, R., O'Connor, T.G., . . .

Woodruff, T.J.; program collaborators for Environmental influences on Child Health

Outcomes., 2023. Birth outcomes in relation to prenatal exposure to per- and

polyfluoroalkyl substances and stress in the Environmental influences on Child Health

Outcomes (ECHO) Program. Environ. Health Perspect. 131, 37006. doi:

10.1289/EHP10723.

Patel, N.S.A., Cuzzocrea, S., Chatterjee, P.K., Di Paola, R., Sautebin, L., Britti, D.,

Thiemermann, C., 2004. Reduction of renal ischemia-reperfusion injury in 5-

lipoxygenase knockout mice and by the 5-lipoxygenase inhibitor zileuton. Mol.

Pharmacol. 66, 220–227. https://doi.org/10.1124/mol.66.2.220

Rhee, S.Y., Jung, E.S., Suh, D.H., Jeong, S.J., Kim, K., Chon, S., Yu, S.-Y., Woo, J.-T., Lee,

C.H., 2021. Plasma amino acids and oxylipins as potential multi-biomarkers for

predicting diabetic macular edema. Sci. Rep. 11, 9727. https://doi.org/10.1038/s41598-

021-88104-y

Ricciotti, E., FitzGerald, G.A., 2011. Prostaglandins and inflammation. Arterioscler. Thromb.

Vasc. Biol. 31, 986–1000. https://doi.org/10.1161/ATVBAHA.110.207449

Saadatian-Elahi, M., Slimani, N., Chajès, V., Jenab, M., Goudable, J., Biessy, C., Ferrari, P.,

Byrnes, G., Autier, P., Peeters, P.H.M., Ocké, M., Bueno de Mesquita, B., Johansson, I.,

Hallmans, G., Manjer, J., Wirfält, E., González, C.A., Navarro, C., Martinez, C., Amiano,

. . . Riboli, E., 2009. Plasma phospholipid fatty acid profiles and their association with

food intakes: results from a cross-sectional study within the European Prospective

Investigation into Cancer and Nutrition. Am. J. Clin. Nutr. 89, 331–346.

https://doi.org/10.3945/ajcn.2008.26834

Sergeant, S., Hugenschmidt, C.E., Rudock, M.E., Ziegler, J.T., Ivester, P., Ainsworth, H.C.,

Vaidya, D., Case, L.D., Langefeld, C.D., Freedman, B.I., Bowden, D.W., Mathias, R.A.,

Chilton, F.H., 2012. Differences in arachidonic acid levels and fatty acid desaturase

(FADS) gene variants in African Americans and European Americans with

diabetes/metabolic syndrome. Br. J. Nutr. 107, 547–555.

https://doi.org/10.1017/S0007114511003230

Storey, J.D., Bass, A.J., Dabney, A. and Robinson, D., 2015. qvalue: Q-value estimation for false

discovery rate control. R package version, 2(0), pp.10-18129.

Sunderland, E.M., Hu, X.C., Dassuncao, C., Tokranov, A.K., Wagner, C.C., Allen, J.G., 2019. A

review of the pathways of human exposure to poly-and perfluoroalkyl substances

(PFASs) and present understanding of health effects. J. Expo. Sci. Environ. Epidemiol.

29, 131–147.

Taibl, K.R., Dunlop, A.L., Barr, D.B., Li, Y.-Y., Eick, S.M., Kannan, K., Ryan, P.B., Schroder,

M., Rushing, B., Fennell, T., Chang, C.-J., Tan, Y., Marsit, C.J., Jones, D.P., Liang, D.,

2023. Newborn metabolomic signatures of maternal per- and polyfluoroalkyl substance

exposure and reduced length of gestation. Nat. Commun. 14, 3120.

https://doi.org/10.1038/s41467-023-38710-3

Taibl, K.R., Schantz, S., Aung, M.T., Padula, A., Geiger, S., Smith, S., Park, J.-S., Milne, G.L.,

Robinson, J.F., Woodruff, T.J., Morello-Frosch, R., Eick, S.M., 2022. Associations of

per- and polyfluoroalkyl substances (PFAS) and their mixture with oxidative stress

biomarkers during pregnancy. Environ. Int. 169, 107541.

https://doi.org/10.1016/j.envint.2022.107541

Welch, B.M., Keil, A.P., Erve, T.J. van 't, Deterding, L.J., Williams, J.G., Lih, F.B.,

Cantonwine, D.E., McElrath, T.F., Ferguson, K.K., 2020. Longitudinal profiles of plasma

eicosanoids during pregnancy and size for gestational age at delivery: A nested case-

control study. PLOS Medicine 17, e1003271.

https://doi.org/10.1371/journal.pmed.1003271

Willer, C.J., Li, Y., Abecasis, G.R., 2010. METAL: fast and efficient meta-analysis of

genomewide association scans. Bioinformatics. 26, 2190–2191.

https://doi.org/10.1093/bioinformatics/btq340

Wong, C.T., Bestard-Lorigados, I., Crawford, D.A., 2019. Autism-related behaviors in the

cyclooxygenase-2-deficient mouse model. Genes Brain Behav. 18, e12506.

https://doi.org/10.1111/gbb.12506

Xu, Y., Wang, Q., Cook, T.J., Knipp, G.T., 2007. Effect of placental fatty acid metabolism and

regulation by peroxisome proliferator activated receptor on pregnancy and fetal

outcomes. J. Pharm. Sci. 96, 2582–2606. https://doi.org/10.1002/jps.20973

Yuan, Z.-X., Majchrzak-Hong, S., Keyes, G.S., Iadarola, M.J., Mannes, A.J., Ramsden, C.E.,

2018. Lipidomic profiling of targeted oxylipins with ultra-performance liquid

chromatography-tandem mass spectrometry. Anal. Bioanal. Chem. 410, 6009–6029.

Zanger, U.M., Schwab, M., 2013. Cytochrome P450 enzymes in drug metabolism: Regulation of

gene expression, enzyme activities, and impact of genetic variation. Pharmacol. Ther.

138, 103–141. https://doi.org/10.1016/j.pharmthera.2012.12.007

Table 1

Demographic profile of CiOB, IKIDS, and ECHO-PROTECT cohorts.

Characteristic		Overall	CiOB, N = 73[a]	IKIDS, N = 287[a]	ECHO-PROTECT, N = 54[a]	p value[b]
				Cohort		
Mother's Race						<0.001
	Non-Hispanic White	253 (62%)	34 (47%)	219 (78%)	0	
	Black	19 (54.6%)	-	14 (5.0%)	0	
	Asian	48 (12%)	14 (19%)	34 (12%)	0	
	Hispanic	78 (19%)	17 (23%)	7 (2.5%)	54 (100%)	
	Other	11 (2.7%)	-	8 (2.8%)	0	
	Missing	5	0	5	0	
Maternal age (years)		31.8 (28.8, 34.5)	33.1 (29.9, 36.0)	31.8 (29.1, 34.3)	28.0 (24.0, 33.0)	<0.001
	Missing	6	1	0	5	
Maternal education						<0.001
	< HS	11 (2.7%)	-	-	-	
	HS/GED/some college	77 (19%)	16 (22%)	39 (14%)	22 (45%)	
	Bachelors	142 (35%)	15 (21%)	111 (39%)	16 (33%)	
	Graduate	179 (44%)	37 (51%)	135 (47%)	7 (14%)	
	Missing	5	0	0	5	
Pre-pregnancy BMI		25 (22, 29)	24 (22, 27)	25 (22, 30)	25 (21, 29)	0.2
	Missing	20	11	1	8	
Parity						<0.001
	0	191 (47%)	35 (51%)	156 (54%)	0 (0%)	
	1 or more	214 (53%)	34 (49%)	131 (46%)	49 (100%)	
	Missing	9	4	0	5	
Maternal household						<0.001
	<$50,000	110 (28%)	18 (26%)	52 (18%)	40 (93%)	
	$50,000-$99,999	145 (36%)	6 (9%)	137 (48%)	-	
	>$100,000	143 (36%)	45 (65%)	97 (34%)	-	
	Missing	16	4	1	11	
Gestational age at visit		17.4 (16.6, 18.9)	23.1 (19.7, 25.7)	17.0 (16.4, 17.7)	25.7 (24.4, 27.9)	<0.001
	Missing	10	0	0	10	

[a]n (%), cells with counts ≤ 5 have been masked; Median (IQR)

[b]Kruskal-Wallis rank sum test; Pearson's Chi-squared test

Table 2A

PFAS detection rates and distributions across cohorts.

PFAS (ng/mL)	CiOB, N = 73			IKIDS, N = 287			ECHO-PROTECT, N = 54			
Full name (abbreviation)	MDL[a]	% Detected	Median (IQR)	MDL[a]	% Detected	Median (IQR)	MDL[a]	% Detected	Median (IQR)	p value[b]
Perfluorononanoic acid (PFNA)	**0.04; 0.03**	**99%**	**0.53 (0.34, 0.72)**	**0.03**	**100%**	**0.40 (0.26, 0.59)**	**0.1**	**81%**	**0.20 (0.16, 0.29)**	<0.001
Perfluoroheptanoic acid (PFHpA)	**0.04; 0.02**	**82%**	**0.03 (0.02, 0.05)**	0.02	58%	0.03 (0.015, 0.03)	0.1	0%	0.10 (0.10, 0.10)	-
Perfluorodecanoic acid (PFDeA)	**0.06; 0.04**	**96%**	**0.17 (0.11, 0.29)**	**0.04**	**98%**	**0.11 (0.07, 0.19)**	0.1	31%	0.10 (0.10, 0.16)	<0.001
Perfluorododecanoic acid (PFDoA)	0.14; 0.04	66%	0.06 (0.02, 0.16)	0.08	55%	0.10 (0.01, 0.11)	-	-	-	-
Perfluorooctanoic acid (PFOA)	**0.04**	**99%**	**1.13 (0.66, 1.92)**	**0.04**	**100%**	**1.03 (0.62, 1.73)**	0.5	41%	0.50 (0.50, 0.96)	0.4
Perfluorooctanesulfonamide (PFOSA)	0.02; 0.01	66%	0.01 (0.004, 0.02)	0.01	55%	0.01 (0.009, 0.01)	0.1	0%	-	-
2-(N-Methyl-perfluorooctane sulfonamido) acetic acid (Me-PFOSA-AcOH)	**0.01**	**93%**	**0.06 (0.02, 0.09)**	**0.008**	**95%**	**0.07 (0.03, 0.15)**	0.1	0%	-	0.07
2-(N-Ethyl-perfluorooctane sulfonamido) acetic acid (Et-PFOSA-AcOH)	**0.009; 0.008**	**71%**	**0.01 (0.006, 0.01)**	0.008	40%	0.01 (0.01, 0.01)	-	-	-	-
Perfluoroundecanoic acid (PFUdA)	**0.04; 0.02**	**96%**	**0.18 (0.11, 0.29)**	**0.02**	**91%**	**0.06 (0.03, 0.12)**	0.1	24%	0.10 (0.10, 0.10)	<0.001
Perfluorohexane sulfonate (PFHxS)	**0.01**	**99%**	**0.55 (0.33, 0.98)**	**0.06**	**99%**	**0.76 (0.40, 1.42)**	0.1	50%	0.12 (0.10, 0.17)	0.016
Perfluorooctanesulfonic acid (PFOS)	**0.04**	**99%**	**2.96 (1.88, 5.52)**	**0.04**	**100%**	**3.27 (2.01, 5.28)**	**0.1**	**100%**	**1.78 (1.40, 2.28)**	0.8
Perfluorobutanesulfonic acid (PFBS)	0.02	48%	0.03 (0.02, 0.03)	0.02	40%	0.03 (0.01, 0.03)	-	-	-	-

Bolded measurements denote PFAS with detection rates ≥ 70% and used in analyses.

[a]Method Detection Limits (ng/mL).

[b]Kruskal-Wallis Rank Sum Test; performed only across cohorts which had ≥ 70% detection rates for specific PFAS.

842 **Table 2B**

843 Bioactive lipids sampled from maternal plasma.

Pathway	Full name	Abbreviation
Cyclooxygenase	13,14-dihydro-15-keto Prostaglandin F2α	13,14DHK-PGF2α
Cyclooxygenase	13,14-Dihydro-15-keto Prostaglandin J2	13,14-D-PGJ2
Cyclooxygenase	15-deoxy-Δ12,14-Prostaglandin J2	15DO12,14-PGJ2
Cyclooxygenase	Bicyclo Prostaglandin E2	BCPGE2
Cyclooxygenase	Bicyclo Prostaglandin E1	BCPGE1
Cyclooxygenase	Prostaglandin A2	PGA2
Cyclooxygenase	Prostaglandin B2	PGB2
Cyclooxygenase	Prostaglandin D2	PGD2
Cyclooxygenase	Prostaglandin D3	PGD3
Cyclooxygenase	Prostaglandin E1	PGE1
Cyclooxygenase	Prostaglandin E2	PGE2
Cyclooxygenase	Prostaglandin E3	PGE3
Cyclooxygenase	Prostaglandin J2	PGJ2
Cyclooxygenase	Thromboxane B2	TXB2
Cyclooxygenase	9-Oxooctadeca-dienoic acid	9-oxoODE
Cytochrome p450	20-carboxy Arachidonic Acid	CAA
Cytochrome p450	9,10-Dihydroxy-octadecenoic acid	9,10-DiHOME
Cytochrome p450	12,13-Dihydroxy-octadecenoic acid	12,13-DiHOME
Cytochrome p450	13,14-dihydro-15-keto Prostaglandin D2	13,14DHK-PGD2
Cytochrome p450	5,6-Dihydroxy-eicosatrienoic acid	5,6-DHET
Cytochrome p450	8,9-Dihydroxy-eicosatrienoic acid	8,9-DHET
Cytochrome p450	11,12-Dihydroxy-eicosatrienoic acid	11,12-DHET
Cytochrome p450	9,10-Epoxy-octadecenoic acid	9(10)-EpoME
Cytochrome p450	12,13-Epoxy-octadecenoic acid	12(13)-EpoME
Cytochrome p450	5,6-Epoxy-eicosatrienoic acid	5(6)-EET
Cytochrome p450	8,9-Epoxy-eicosatrienoic acid	8(9)-EET
Cytochrome p450	11,12-Epoxy-eicosatrienoic acid	11(12)-EET
Cytochrome p450	14,15-Epoxy-eicosatrienoic acid	14(15)-EET
Cytochrome p450	11-Hydroxy-eicosatetraenoic acid	11-HETE
Cytochrome p450	16-Hydroxy-eicosatetraenoic acid	16-HETE
Cytochrome p450	17-Hydroxy-eicosatetraenoic acid	17-HETE
Cytochrome p450	18-Hydroxy-eicosatetraenoic acid	18-HETE
Cytochrome p450	20-Hydroxy-eicosatetraenoic acid	20-HETE
Cytochrome p450	9S-Hydroxy-octadecadienoic acid	9S-HODE
Lipoxygenase	13S-Hydroxy-octadecadienoic acid	13S-HODE
Lipoxygenase	5-Hydroxy-eicosatetraenoic acid	5-HETE
Lipoxygenase	8-Hydroxy-eicosatetraenoic acid	8-HETE
Lipoxygenase	12-Hydroxy-eicosatetraenoic acid	12-HETE
Lipoxygenase	15-Hydroxy-eicosatetraenoic acid	15-HETE
Lipoxygenase	5-Oxoeicosatetraenoic acid	5-oxoETE
Lipoxygenase	12-Oxoeicosatetraenoic acid	12-oxoETE
Lipoxygenase	15-Oxoeicosatetraenoic acid	15-oxoETE
Lipoxygenase	13-Oxooctadeca-dienoic acid	13-oxoODE
Lipoxygenase	Resolvin D1	RVD1
Lipoxygenase	Resolvin D2	RVD2
Parent Compound	α-Linolenic Acid	αLA
Parent Compound	Arachidonic Acid	AA
Parent Compound	Docosahexaenoic Acid	DHA
Parent Compound	Eicosapentaenoic Acid	EPA
Parent Compound	Linoleic Acid	LA

884
885

Table 3

β estimates and 95% confidence intervals corresponding to percentage change in bioactive lipids associated ($p \leq 0.05$) with a doubling

in PFAS; for joint cohort and meta-analyses.

Bioactive Lipid Pathway/Group	Lipid	PFAS	All 3 cohorts, random intercept (N = 387)			CiOB & IKIDS, random intercept (N = 347)			All 3 cohorts, meta-analysis (N = 387)			CiOB & IKIDS, meta-analysis (N = 347)		
			β	95% CI lower	95% CI upper	β	95% CI lower	95% CI upper	β	95% CI lower	95% CI upper	β	95% CI lower	95% CI upper
Cyclooxygenase	13,14-D-PGJ2	Me-PFOSA-AcOH	-	-	-	-	-	-	-	-	-	3.8	0.1	7.7
Cyclooxygenase	13,14DHK-PGF2a*	PFNA	10.6	1.6	20.4	10.4	1.7	20.0	-	-	-	-	-	-
Cyclooxygenase	13,14DHK-PGF2a*	PFUdA	-	-	-	6.8	2.4	11.5	-	-	-	6.4	1.9	11.1
Cyclooxygenase	9-oxoODE	PFNA	-	-	-	-2.4	-4.6	-0.1	-2.28	-4.24	-0.29	-2.3	-4.3	-0.3
Cyclooxygenase	BCPGE1	PFNA	12.8	2.9	23.6	10.3	0.3	21.3	11.57	1.79	22.29	10.6	0.6	21.7
Cyclooxygenase	BCPGE1	PFOA	-	-	-	11.8	1.6	23.1	-	-	-	11.6	1.4	22.9
Cyclooxygenase	PGD2	PFDeA	-	-	-	-	-	-	-	-	-	13.2	1.4	26.4
Cyclooxygenase	PGD2	PFNA	16.5	1.9	33.1	-	-	-	15.6	2.2	30.8	-	-	-
Cyclooxygenase	PGD2	PFOA	-	-	-	24.3	7.3	43.9	-	-	-	21.7	7.3	38.1
Cyclooxygenase	PGD2	PFOS	11.9	0.4	24.7	-	-	-	16.6	5.6	28.8	17.0	5.8	29.3
Cyclooxygenase	PGD3	PFOS	11.5	1.5	22.5	10.4	0.4	21.4	10.0	1.2	19.6	-	-	-
Cyclooxygenase	PGD3	PFUdA	-	-	-	10.3	3.5	17.6	-	-	-	-	-	-
Cyclooxygenase	PGE3	PFNA	13.8	1.2	28.1	-	-	-	-	-	-	-	-	-
Cyclooxygenase	PGE3	PFOA	-	-	-	16.3	2.6	31.9	-	-	-	15.6	2.2	30.7
Cyclooxygenase	TXB2	Me-PFOSA-AcOH	-	-	-	5.4	0.2	10.8	-	-	-	5.4	0.2	10.9
Cytochrome p450	13,14DHK-PGD2	Me-PFOSA-AcOH	-	-	-	5.2	0.5	10.2	-	-	-	-	-	-
Cytochrome p450	14(15)-EET	PFDeA	-	-	-	6.4	1.0	12.0	-	-	-	6.6	1.3	12.2
Cytochrome p450	20-HETE	PFNA	6.9	0.5	13.6	-	-	-	-	-	-	-	-	-
Cytochrome p450	5(6)-EET	PFOA	-	-	-	-	-	-	-	-	-	20.6	4.4	39.3

Bioactive Lipid Pathway/Group	Lipid	PFAS	All 3 cohorts, random intercept (N = 387)			CiOB & IKIDS, random intercept (N = 347)			All 3 cohorts, meta-analysis (N = 387)			CiOB & IKIDS, meta-analysis (N = 347)		
			β	95% CI lower	95% CI upper	β	95% CI lower	95% CI upper	β	95% CI lower	95% CI upper	β	95% CI lower	95% CI upper
Cytochrome p450	5(6)-EET	PFUdA	-	-	-	9.3	0.9	18.5	-	-	-	-	-	-
Cytochrome p450	8(9)-EET	Me-PFOSA-AcOH	-	-	-	5.4	0.9	10.0	-	-	-	5.8	1.5	10.2
Cytochrome p450	8,9-DHET	PFNA	-	-	-	6.9	0.3	13.8	-	-	-	-	-	-
Cytochrome p450	8,9-DHET	PFUdA	-	-	-	3.6	0.2	7.1	-	-	-	-	-	-
Lipoxygenase	12-oxoETE	PFDeA	-	-	-	8.0	1.6	14.8	-	-	-	7.9	1.4	14.7
Lipoxygenase	12-oxoETE	PFNA	10.1	1.9	19.0	-	-	-	8.7	0.8	17.3	-	-	-
Lipoxygenase	12-oxoETE	PFOS	8.2	1.8	15.0	7.2	1.2	13.5	7.4	1.4	13.9	7.1	1.0	13.5
Lipoxygenase	12-oxoETE	PFUdA	-	-	-	5.9	1.7	10.2	-	-	-	5.9	1.7	10.3
Lipoxygenase	15-HETE	PFHxS	-	-	-	-7.8	-13.6	-1.5	-	-	-	-7.5	-13.5	-1.2
Lipoxygenase	15-oxoETE	PFDeA	-	-	-	8.5	1.6	15.8	-	-	-	7.5	0.7	14.8
Lipoxygenase	15-oxoETE	PFOA	-	-	-	9.6	1.0	19.0	-	-	-	9.1	0.6	18.2
Lipoxygenase	15-oxoETE	PFOS	10.3	1.9	19.3	10.4	3.9	17.4	10.3	3.8	17.3	10.3	3.8	17.2
Lipoxygenase	15-oxoETE	PFUdA	-	-	-	6.5	2.1	11.2	-	-	-	6.0	1.5	10.7
Lipoxygenase	5-HETE	PFDeA	-	-	-	6.7	1.5	12.1	-	-	-	5.8	0.6	11.3
Lipoxygenase	RVD2	PFUdA	-	-	-	7.3	1.9	13.0	-	-	-	6.8	1.3	12.6
Parent Compound	AA	PFNA	10.9	3.0	19.4	10.6	2.9	19.0	9.5	2.3	17.3	8.7	1.4	16.6
Parent Compound	AA	PFOS	9.7	3.4	16.3	9.1	3.2	15.3	8.2	2.5	14.2	8.0	2.3	14.0
Parent Compound	EPA	PFUdA	-	-	-	3.7	0.3	7.2	-	-	-	3.8	0.2	7.4
Parent Compound	LA	PFOA	-	-	-	11.4	0.1	24.0	-	-	-	-	-	-
Parent Compound	LA	PFUdA	-	-	-	6.6	1.0	12.6	-	-	-	7.9	2.0	14.2
Parent Compound	αLA	PFUdA	-	-	-	-4.9	-9.1	-0.5	-	-	-	-	-	-

Figure 1. Directed acyclic graph of the relationships between maternal per- and poly-fluoroalkyl substances and bioactive lipids

Figure 2A. Correlation Matrix of Bioactive Lipids and High-Detect PFAS in the CiOB Cohort

Figure 2B. Correlation Matrix of Bioactive Lipids and High-Detect PFAS in the IKIDS Cohort

Figure 2C. Correlation Matrix of Bioactive Lipids and High-Detect PFAS in the ECHO-PROTECT Cohort

Figure 3. Heatmap of β Estimates Corresponding to Percentage Change in Bioactive Lipids as a Result of Doubling Log-Transformed PFAS for Adjusted Joint Analyses and Meta-Analyses run on Adjusted Within-Cohort Models. The magnitude of effect estimates in each cell in the heatmap corresponds to the intensity of the color band in the legend. Non-significant values (p > .05) are marked with black "X". Sample sizes range from 343 to 383 (see Supplemental Tables 3, 4, 5, and 12 for exact pair-wise sample sizes).

Figure 4. Forest Plot of Quantile g-computation Effect Estimates in the Combined Cohort Analysis with IKIDS and CIOB Cohorts (N=343), showing of β Estimates Corresponding to Percentage Change in Bioactive Lipids as a Result of Simultaneous 1-Quartile Increase in All Log-Transformed PFAS. Model adjusted for Maternal Age, Maternal Education, Pre-Pregnancy BMI, Parity, Gestational Age at Visit, and Cohort.