Role of Cardiorespiratory Fitness and Mitochondrial Energetics in Reduced Walk Speed of Older Adults with Diabetes in the Study of Muscle, Mobility and Aging (SOMMA)

Sofhia V. Ramos[1], Giovanna Distefano[1], Li-Yung Lui[2,3], Peggy M. Cawthon[2,3], Philip Kramer[4], Ian J. Sipula[6], Fiona M. Bello[6], Theresa Mau[2,3], Michael J. Jurczak[6], Anthony J. Molina[5], Erin E. Kershaw[6], David J. Marcinek[7], Frederico G.S Toledo[6], Anne B. Newman[6], Russell T. Hepple[8], Stephen B. Kritchevsky[4], Bret H. Goodpaster[1], Steven R. Cummings[2,3], Paul M. Coen[1]

Affiliations:

[1]Translational Research Institute, Advent Health, Orlando, Florida, USA.
[2]San Francisco Coordinating Center, California Pacific Medical Center Research Institute, San Francisco, California USA.
[3]Department of Epidemiology and Biostatistics, University of California, San Francisco, California, USA.
[4]Department of Gerontology and Geriatric Medicine, Wake Forest University, School of Medicine, Winston-Salem, North Carolina, USA.
[5]Department of Medicine, University of California, San Diego, California, USA.
[6]Division of Endocrinology and Metabolism, Department of Medicine, School of Medicine, University of Pittsburgh, Pittsburgh, Pennsylvania, USA.
[7]Department of Radiology, University of Washington School of Medicine, Seattle, Washington, USA.
[8]Department of Physical Therapy, University of Florida, Gainesville, Florida, USA.

Email addresses: Giovanna.distefano@adventhealth.com, lily.lui@ucsf.edu, peggy.cawthon@ucsf.edu, pkramer@wakehealth.edu, iansipula@pitt.edu, fiona.bello@epfl.ch Theresa.Mau@ucsf.edu, jurczakm@pitt.edu, ajmolina@health.ucsd.edu, kershawee@upmc.edu, dmarc@uw.edu, toledofgs7@pitt.edu, anewman@pitt.edu, rthepple@phhp.ufl.edu, skritche@wakehealth.edu, bret.goodpaster@adventhealth.com, steven.cummings@ucsf.edu

Correspondence:
Paul M. Coen, PhD
AdventHealth, Orlando
Translational Research Institute
765-426-9743
paul.coen@adventhealth.com

Key Words: Diabetes, Aging, Cardiorespiratory Fitness, Mitochondrial Energetics, Walking Speed

Word count = 3954, Tables = 3, Figures = 1

ABSTRACT

Rationale – Cardiorespiratory fitness and mitochondrial energetics are associated with reduced
walking speed in older adults. The impact of cardiorespiratory fitness and mitochondrial
energetics on walking speed in older adults with diabetes has not been clearly defined.

Objective – To examine differences in cardiorespiratory fitness and skeletal muscle
mitochondrial energetics between older adults with and without diabetes. We also assessed the
contribution of cardiorespiratory fitness and skeletal muscle mitochondrial energetics to slower
walking speed in older adults with diabetes.

Findings – Participants with diabetes had lower cardiorespiratory fitness and mitochondrial
energetics when compared to those without diabetes, following adjustments for covariates
including BMI, chronic comorbid health conditions, and physical activity. 4-m and 400-m
walking speeds were slower in those with diabetes. Mitochondrial oxidative capacity alone or
combined with cardiorespiratory fitness mediated ~20-70% of the difference in walk speed
between older adults with and without diabetes. Further adjustments of BMI and co-morbidities
further explained the group differences in walk speed.

Conclusions – Skeletal muscle mitochondrial energetics and cardiorespiratory fitness contribute
to slower walking speeds in older adults with diabetes. Cardiorespiratory fitness and
mitochondrial energetics may be therapeutic targets to maintain or improve mobility in older
adults with diabetes.

ARTICLE HIGHLIGHTS

Why did we undertake this study?

• To determine if mitochondrial energetics and cardiorespiratory fitness contribute to
slower walking speed in older adults with diabetes.

What is the specific question(s) we wanted to answer?

• Are mitochondrial energetics and cardiorespiratory fitness in older adults with diabetes
lower than those without diabetes? How does mitochondrial energetics and
cardiorespiratory fitness impact walking speed in older adults with diabetes?

What did we find?

• Mitochondrial energetics and cardiorespiratory fitness were lower in older adults with
diabetes compared to those without diabetes, and energetics, and cardiorespiratory
fitness, contributed to slower walking speed in those with diabetes.

What are the implications of our findings?

• Cardiorespiratory fitness and mitochondrial energetics may be key therapeutic targets to
maintain or improve mobility in older adults with diabetes.

Introduction

The US population is rapidly aging, and in 2022 the percentage of Americans aged 65 and older diagnosed with diabetes remains high at 30% [1]. This represents a healthcare challenge as older adults with diabetes have a greater risk of cardiovascular disease and mobility disability [2, 3]. The contributors to impaired mobility in older individuals with diabetes include the presence of co-morbidity, increased body mass index (BMI) [4], and lower muscle strength and quality [5-7]. However, the relationship between diabetes and loss of mobility remains only partially explained, and other contributors also likely play a role. Our group and others reported that cardiorespiratory fitness and skeletal muscle mitochondrial energetics are linked to reduced walking speed in aging, which is a predictor of mobility disability [8-10]. For example, muscle bioenergetics assessed using [31]P-MRS was highly correlated with walking speed and partially explained age-related poorer performance in short (6-m) and long (400-m) walking tasks [10]. Several studies have found a close relationship between cardiorespiratory fitness and walking speed in older adults [11], which suggests that the decline in aerobic capacity also contributes to slower walking speed and mobility disability with age.

Muscle biopsies from individuals with diabetes have been shown to have lower oxidative phosphorylation (OXPHOS) capacity compared with muscle from healthy controls [12-14]. However, some studies [15, 16], but not all [12, 13, 17], indicate no differences between patients with diabetes and BMI-matched controls, indicating that obesity *per se* may underlie the lower muscle oxidative capacity. Indeed, other studies in people without diabetes have shown that BMI is strongly related to muscle oxidative capacity [18]. In addition, reports assessing mitochondrial energetics in older adults with diabetes tend to have small sample size and lack adjustment for critical confounding variables including physical activity and adiposity [13, 14, 17], and there are

few reports focusing on older adults with diabetes specifically. Lower fitness is a predictor of

insulin resistance and diabetes [19], and prevalence of diabetes is greater in individuals with low

fitness [20]. It is unknown whether diabetes associated changes in cardiorespiratory fitness

and/or muscle energetics explain slower walking speed seen in people with diabetes, independent

of other covariates.

In this analysis, we leveraged data from the Study of Muscle, Mobility and Aging

(SOMMA) [21], to investigate the association of diabetes with cardiorespiratory fitness and

mitochondrial energetics in a cohort of 879 older adults who were well phenotyped in terms of

the free living physical activity behaviors and body composition. We utilized ^{31P}MRS, to

measure muscle mitochondrial energetics *in vivo* (ATPmax) and high-resolution respirometry to

measure mitochondrial energetics in muscle biopsies, a complementary approach that

interrogates mitochondrial OXPHOS capacity at the myocellular level. Here, we tested the

hypothesis that diabetes would be associated with lower cardiorespiratory fitness and muscle

energetics independent of adiposity, physical activity levels, chronic comorbid health conditions,

and oral hypoglycemic medication use. We also tested whether cardiorespiratory fitness and

mitochondrial energetics mediate slower walking speed of older adults with diabetes.

RESEARCH DESIGN AND METHODS

Participant Recruitment

The Study of Muscle Mobility and Aging (SOMMA; http://sommaonline.ucsf.edu) is a

prospective longitudinal cohort study of older adults and was designed to understand the

biological basis of muscle aging [21]. Men and women (70+ yrs. old) were recruited between

April 2019 to December 2021. Participants were enrolled at University of Pittsburgh and Wake

Forest University School of Medicine if they were willing and able to complete a skeletal muscle

biopsy and undergo Magnetic Resonance Imaging (MRI) and Spectroscopy (MRS). Exclusion

criteria included an inability to walk one-quarter of a mile or climb a flight of stairs; had a

BMI>40kg/m2; had an active malignancy or dementia; or any medical contraindication to biopsy

or MR. Finally, participants must have been able to complete the 400-m walk test; those who

appeared as they might not be able to complete the 400-m walk at the in-person screening visit

completed a short distance walk (4-m) to ensure their walking speed was ≥ 0.6m/s. Presence of

diabetes was identified by self-report, use of prescribed hypoglycemic medication, or having an

HbA1c $\geq 6.5\%$. WIRB-Copernicus Group (WCG) Institutional Review Board (WCGIRB, study

number 20180764) approved the study as the single IRB and all participants provided written

informed consent.

Baseline Assessments

Detail on study design and methodology have been published elsewhere [21]. The

SOMMA Baseline Visit generally consisted of three days of assessments over several weeks: on

"Day 1" the 400-m walk plus most other in-person assessments; "Day 2" cardiopulmonary

exercise testing (CPET) and Magnetic Resonance Imaging (MRI) and Spectroscopy (MRS) were

assessed; "Day 3" muscle biopsy specimen was collected. Age, sex, and race were collected by

self-report. The self-reported medical conditions (heart disease, stroke, kidney disease/renal

failure, and peripheral vascular disease) were assessed to determine if participants had those

conditions or not. Participants were asked to bring all prescription medications they had taken in

the 30 days prior to their "Day 1" clinic visit. If a participant forgot to bring one or more

medications, clinic staff obtained this information over the telephone or at the return visit. The

prescription medication use was reviewed and updated at the additional baseline visit days where

CPET and tissue sampling were done. Body weight, height, and physical function assessments

were also measured and accelerometry devices were provided. Most SOMMA participants

completed cardiopulmonary exercise testing (CPET), magnetic resonance spectroscopy (MRS)

and provided a percutaneous biopsy of the vastus lateralis muscle which was performed under

fasted conditions. On the day of the muscle biopsy, participants prescribed oral hypoglycemic

medication were asked to take their medications following tissue sampling.

Mobility Assessments

Mobility was assessed by 400m and 4m walking tests. For the 400-m walk test,

participants were instructed to walk 10 complete laps around a set course at their usual pace and

without overexerting themselves. 4m walking speed was measured in 2 trials and the

participant's best time was used to calculate walking speed in m/sec.

Physical Activity Assessments

Two devices were used to assess activity: the thigh-worn activPal that recorded total

daily step count and the wrist-worn ActiGraph GT9x. Devices were placed on the participants at

the baseline "Day 1" visit with a data collection period of 7 full days. Data was included only if

the device was worn over a 24-hour period and had ≥17hour per day wear time. Activity levels

collected from ActiGraph were determined by cut points described by Montoye and colleagues

[22], and included total daily physical activity time, and time spent in sedentary, light, and

moderate to vigorous physical activity (MVPA).

Cardiopulmonary Exercise Testing Assessment

Participants walked for 5 minutes at a preferred walking speed, and progressive

symptom-limited exercise protocol ensued with increases in speed (0.5 mph) and incline (2.5%)

in 2-minute increments using a modified Balke protocol or a manual protocol [23, 24].

Cardiorespiratory fitness was determined as VO_2peak (mL/min) which was identified as the

highest 30-second average of VO_2 (mL/min) achieved.

Magnetic Resonance (MR) Imaging and Spectroscopy

[31]Phosphorous-MRS was used to assess *in-vivo* mitochondrial adenosine triphosphate

(ATP) generation in the quadriceps muscle during an acute bout of knee isometric extension

exercise [21, 24] . A 3 Tesla magnetic resonance magnet (Siemens Medical Systems – Prisma

[Pittsburgh] or Skyra [Wake Forest]) with a 2.5" surface RF coil was used for the assessment.

All data were analyzed using jMURI v7.0 [25]. Magnetic resonance imaging (MRI) was

performed with a 2.5" surface RF coil and used to assess abdominal subcutaneous adipose tissue

(ASAT), visceral adipose tissue (VAT), and quadricep muscle fat infiltration (QFI). The entire

body was scanned and images were analyzed using AMRA Researcher® (AMARA medical AB,

Linköping Sweden) [26].

Muscle Biopsy and Mitochondrial Respiration

Muscle biopsy was completed with a Bergström trocar (5 or 6mm) as previously

described [24]. Approximately 10 mg of muscle tissue was allocated for respirometry assays and

were immediately submerged in ice cold BIOPS buffer. Fiber bundles of approximately 2-3 mg

wet weight were gently teased apart and permeabilized as previously described [24] and weight

was measured using an analytical balance (Mettler Toledo, Columbus, OH). Permeabilized fiber

bundle (PmFB) oxygen consumption was measured using a high-resolution respirometer

(Oxygraph-2k, Oroboros Instruments, Innsbruck, Austria). PmFB were placed into chambers

containing MiR05 buffer supplemented with 25µM blebbistatin in the absence of light. Two

protocols were completed in duplicate measuring carbohydrate supported (Protocol 1: 5mM

pyruvate and 2mM malate) and fatty acid oxidation (FAO) supported (Protocol 2: 25μM

palmitoyl-carnitine and 2mM malate) respiration. For protocol 1, state 3 respiration was

stimulated with the addition of 4.2mM ADP ($OXPHOS_{CHO}$) followed by subsequent additions of

10mM glutamate and 10mM succinate sequentially ($maxOXPHOS_{CHO}$). Maximal uncoupled

respiration was stimulated with consecutive 0.5uM titrations of 2-[2-[4-

(trifluoromethoxy)phenyl] hydrazinylidene-propanedinitrile (FCCP) until maximum electron

transport system (ETS) capacity was attained ($maxETS_{CHO}$). For protocol 2, state 3 respiration

was stimulated with the addition of 4mM ADP ($OXPHOS_{FAO}$) followed by subsequent additions

of 10mM glutamate and 10mM succinate to further stimulate mitochondrial complexes I and II

($maxOXPHOS_{FAO}$). PmFB eliciting an oxygen consumption response <15% were not included in

the final analysis. Respiratory control ratio (RCR) for protocol 1 (7.11 ± 2.86) and protocol 2

(3.47 ± 1.74) were assessed for quality control. Respiration was analyzed using DatLab 7.4.0.4.

In SOMMA, not all participants completed both protocols due to time constraints. Protocol 1

(carbohydrates) was prioritized over Protocol 2 (fatty acids) when limited technician time was

available, which left fewer participants with complete data for Protocol 2.

Statistical Analysis

Differences in participant characteristics between older adults with and without diabetes

were compared using the T-test for continuous variables (or Kruskal-Wallis non-parametric test

for skewed variables) and chi-square test for dichotomized variables. Linear regressions without

adjustment were performed to determine differences in cardiorespiratory fitness, and

mitochondrial energetics between those with and without diabetes, and additional adjustments

for the following confounders: age, race, gender, site/technician, BMI, and chronic conditions

(model 1). Model 1 was further adjusted for steps per day from ActiPal or volume of adipose

tissue depots: Abdominal Subcutaneous Adipose Tissue (ASAT), Visceral Adipose Tissue

(VAT) and muscle Quadricep Fat Infiltration (QFI). Lastly, to determine the impact of

hypoglycemic medication use within the group of older adults with diabetes, multivariate linear

regression was performed adjusting for age, race, and site/technician. Those analyses were

completed using JMP® Software, Version 16.

Linear regression was also used to determine the association of diabetes with walking

speed. We evaluated the potential mediating influence of VO_2peak and/or mitochondrial

energetics. We did this by comparing the beta coefficient for the association of diabetes status on

walking speed from the 400m and 4m walk. The base model included the confounding variables:

gender, age, and race. model 2: base model plus technician and $maxOXPHOS_{CHO}$, model 3: base

model plus technician and $maxOXPHOS_{FAO}$, model 4: base model plus clinical site and

VO_2peak, model 5: base model plus technician, VO_2peak and $maxOXPHOS_{CHO}$, model 6: base

model plus technician, VO_2peak and $maxOXPHOS_{FAO}$. Fully adjusted models also included the

potential mediators/confounders BMI, and chronic conditions. Finally, we compared the percent

difference in the beta coefficient between the base model and subsequent models to understand

how each adjustment impacted group differences in walking speed. Those analyses were

performed using SAS version 9.4 (SAS institute Inc., Cary, NC).

RESULTS

Participant Characteristics

A total of 879 participants (59.2% women) with an average age of 76.5 ± 5.0 years were

enrolled in SOMMA [21]. A summary of participants with available data are included in

Supplemental Figure 1. Participant characteristics are summarized in **Table 1**. 18% of the

SOMMA cohort were classified as having diabetes. The proportion of women to men was

greater in the group without diabetes (p=0.02). The proportion of White participants compared to

non-White was greater in the group without diabetes (p<0.001). Older adults with diabetes had a

higher body weight, waist circumference, BMI, and HbA1c (all, p<0.001). Individuals with

diabetes tended to have a higher prevalence of heart disease and hypertension, whereas those

without diabetes more often had history of cancer diagnosis (p<0.001). Approximately 75% of

participants with diabetes reported use of a hypoglycemic agent where 83% used metformin 19%

used insulin, and 2% used thiazolidinediones (TZD). Total daily steps, and time in MVPA was

significantly lower (p<0.001 for both) and total sedentary time was significantly higher in those

with diabetes (p<0.001). Quadriceps muscle fat infiltration (QFI) was higher in those with

diabetes (p=0.002). In addition, both abdominal subcutaneous adipose tissue (ASAT) and

visceral adipose tissue (VAT) volume was significantly higher in older adults with diabetes

(p=0.005 for ASAT and p<0.001 for VAT) (**Table 1**).

Cardiorespiratory Fitness and Muscle Energetics in Diabetes

260   In the adjusted model (**Table 2**, model 1 + PA), we report that VO_2peak was

significantly lower in older adults with diabetes compared to those without (p= 0.004). State 3

(coupled) respiration in the presence of pyruvate + malate ($OXPHOS_{CHO}$, p= 0.12,) and with the

addition of both glutamate and succinate ($maxOXPHOS_{CHO}$, p= 0.10) was similar between

groups. Maximal electron transfer system (ETS) capacity measured by maximal uncoupled

respiration with CHO substrates ($maxETS_{CHO}$, p=0.03) was significantly lower in those with

diabetes (**Table 2**). State 3 (coupled) respiration in the presence of palmitoyl carnitine + malate

was similar between those with and without diabetes ($OXPHOS_{FAO}$, p= 0.10). Maximal oxidative

phosphorylation supported by fatty acid substrates (maxOXPHOS$_{FAO}$, p=0.006) remained

significantly lower in older adults with diabetes in model 1 + PA. Sensitivity analyses completed

in those with data for both Protocol 1 and Protocol 2 had similar results (data not shown) to

ensure that differences between protocols were not due to differences in sample size.

Adjustment for levels of QFI, ASAT and VAT did not impact group differences in

mitochondrial energetics and VO$_2$peak measurements with an exception for OXPHOS$_{CHO}$ (**Table**

2). Adjustment for VAT tended to explain some variance in OXPHOS$_{CHO}$ between those with

and without diabetes (p=0.06, **Table 2**). Within the diabetes group only, ATPmax was lower in

older adults taking any hypoglycemic medication (p=0.05) or metformin alone (p=0.01,

Supplemental table 1). These findings make it difficult to distinguish the impact of medications

versus diabetes on ATPmax and therefore, further analysis of mitochondrial energetics focused

on respiration measurements. VO$_2$peak and mitochondrial energetics assessed by respiration

were similar with the use of all hypoglycemic agents and metformin (**Supplemental Table 1**).

VO$_2$peak and Mitochondrial Energetics Mediate the Relationship between Diabetes and

Walking Speed

Lastly, 400-m and 4-m walking speeds were significantly lower in those with diabetes

compared to those without diabetes following adjustments for gender, age, and race (base model,

Table 3). To assess whether VO$_2$peak and mitochondrial energetics mediated the relationship

between diabetes and walk speed, we compared the β coefficient for mean 400-m and 4-m

walking speed from the base linear regression model with the β coefficient of models that

included VO$_2$peak and/or mitochondrial energetics variables. Additional individual adjustments

for maxOXPHOS$_{CHO}$ (400-m, p=0.02, model 2), maxOXPHOS$_{FAO}$ (400-m, p=0.02; model 3)

and VO$_2$peak (400-m p=0.01; model 4) resulted in no significant change in group differences in

400-m walking speed between those with and those without diabetes (**Table 3**). Alternatively,

for 4-m walking speed, further adjustments for $maxOXPHOS_{CHO}$ (4-m, p=0.12; model 2) and

$maxOXPHOS_{FAO}$ (4-m, p=0.51, model 3) significantly mediated differences between groups,

whereas VO_2peak (4-m, p=0.04, model 4) did not. Examining the impact of combinations of

VO_2peak and mitochondrial energetics revealed that VO_2peak with $maxOXPHOS_{CHO}$ (400-m

p=0.07, 4-m p=0.27; model 5) or $maxOXPHOS_{FAO}$ (400-m, p=0.20, 4-m, p=0.99) mediated

differences between those with and without diabetes. Full adjustments including potential

confounders/mediators BMI and chronic conditions explained the remaining variance in walking

speed between groups. (**Table 3**). Comparing the β coefficient for mean 400-m and 4-m walking

speed from the base linear regression model with the β coefficient of models that included both

VO_2peak and mitochondrial energetics variables revealed that VO_2peak and mitochondrial

energetics explained an additional ~46-100% of the variance in 400-m and 4-m walking speed

between groups (**Figure 1**).

DISCUSSION

In this study of 879 well-phenotyped older adults we found that the 159 participants with

diabetes had slower 4- and 400-m walk speed. We report for the first time that the association of

diabetes status with slow walking speed was mediated by VO_2peak and skeletal muscle

mitochondrial energetics. We also examined differences in cardiovascular fitness and

mitochondrial energetics between those with and without diabetes while controlling for physical

activity and adiposity. Interestingly, accounting for these potential confounders did not explain

differences in VO_2peak and mitochondrial respiration between those with and without diabetes,

suggesting other factors may impact cardiorespiratory fitness and energetics in those with

diabetes. We also found that older adults taking hypoglycemic medications (insulin, TZDs,

and/or metformin) or metformin alone had significantly lower ATPmax (but not respirometry

measures of mitochondrial energetics) compared to older adults with diabetes but not taking

medications, providing evidence that aligns with reports in the literature that metformin can

impair mitochondrial energetics.

The well-phenotyped SOMMA cohort provided an opportunity to assess the impact of

diabetes status on VO_2peak and mitochondrial energetics in older adults while controlling for

objectively measured indices of physical activity, adiposity, and chronic conditions in addition to

demographic variables. Many of the previous studies reporting lower muscle oxidative capacity

measured in permeabilized fiber bundles from patients with diabetes have not rigorously

assessed and controlled for the participants' physical activity [12-14, 17]. This is important as

physical activity interventions (such as walking) can improve mitochondrial energetics in type 2

diabetes [27] and during weight loss in obesity [28], and in older adults [29]. Here, we report that

lower levels of objectively assessed physical activity partially explains lower respiration in older

adults with diabetes. Interestingly, comparing our respirometry protocols with distinct substrate

combinations, we found that CI+II FAO supported maxOXPHOS respiration remains lower in

the diabetes group, while CI+II CHO only supported maxOXPHOS did not differ following

adjustments for physical activity. This suggests that CHO supported coupled respiration may be

more sensitive to levels of physical activity compared to FAO supported respiration. This is also

in line with evidence indicating reduced rates of lipid oxidation in skeletal muscle from

individuals with diabetes, potentially due to mitochondrial overload and incomplete fatty acid

oxidation [30].

The influence of adiposity on mitochondrial energetics has been considered in prior

studies of diabetes, typically by matching control subjects for BMI [14]. Some studies [15, 16],

but not all [12, 13, 17], indicate no differences between patients with diabetes and BMI-matched

controls, suggesting that obesity *per se* may underlie the lower muscle oxidative capacity. Here,

we examined the influence that BMI and specific adipose depots (ASAT, VAT, and QFI) had on

muscle mitochondrial energetics in diabetes. However, regardless of whether adiposity was

adjusted for BMI alone or combined with individually objectively measured adipose depots,

mitochondrial respiration generally remained significantly lower in those with diabetes. Taken

together, these findings indicate that adiposity and physical activity largely (there were some

exceptions) do not entirely explain lower mitochondrial energetics in older adults with diabetes.

Metformin is one of the most commonly prescribed medications for diabetes, and

although its mechanism of action remains highly debated, there is evidence of an inhibitory

effect on complex I of the electron transport chain [31]. Both pre-clinical and clinical studies

report maladaptation of mitochondrial function following metformin treatment, increased muscle

atrophy and reduced exercise capacity, specifically in older adults [31, 32]. Conversely, others

have reported that metformin can improve mitochondrial function via effects on mitophagy,

autophagy or AMPK activation [33]. Our understanding of how metformin impacts skeletal

muscle mitochondrial function of older adults with diabetes is limited. Here, we reveal lower

ATPmax in older adults with diabetes taking hypoglycemic medication or metformin alone,

compared to older adults with diabetes not taking medications. Interestingly, we did not observe

an association of metformin with diabetes status in the respiration assays, suggesting different

sensitivities to metformin based on whether mitochondrial energetics are assessed in vivo

(ATPmax) compared to permeabilized fiber bundle preparations ex vivo. However, an important

caveat is that participants were asked to withhold medication prior to the muscle biopsy and

respirometry assays but not on the day of ^{31}PMRS assessments. In addition, others have reported

no effect on mitochondrial respiration after a two-week metformin treatment compared to control

[34], or in long term diabetes patients [35]. Given that metformin is water soluble, it is also

plausible that the inhibitory effect of metformin is lost when the muscle fiber bundles are washed

in preparation for the respirometry assay. Further work is needed to decipher the impact

metformin has on skeletal muscle mitochondrial energetics in older adults with diabetes.

Zhao and colleagues reported that adults with diabetes were 31-34% less likely to

participate in physical activity, which in turn may contribute to lower VO_2peak [36]. In line with

these reports, SOMMA participants with diabetes were significantly less active and significantly

more sedentary. Our results show that additional adjustments for physical activity did not resolve

differences in VO_2peak between groups, suggesting that lower VO_2peak in older adults with

diabetes is independent of physical activity in this cohort. In addition, we report that independent

of adiposity measured by BMI or combined BMI with ASAT, VAT and QFI, VO_2peak remained

significantly lower in those with diabetes compared to those without. Taken together, our work

suggests that physical activity and adiposity may not be implicated in lower cardiorespiratory

fitness in older adults with diabetes and that other factors including perhaps genetics and

heritability are more important determinants to low cardiorespiratory fitness in diabetes.

Previous work from our laboratory and others revealed a significant relationship between

walking speed with both VO_2peak and mitochondrial energetics in older adults [9, 37, 38]. Here,

we explored the contribution of both mitochondrial energetics and cardiorespiratory fitness to

mobility in older adults with diabetes. We report that mitochondria respiration, both

independently (~40-70%) and combined with VO_2peak (~55-100%), mediated the variance in 4-

m walking speed in those with diabetes. Differences between groups are further explained by

additional adjustments for potential confounders/mediators BMI, and comorbidities. Reports

indicate that muscle mass is lost at an accelerated rate in older adults with diabetes [7],

contributing to reductions in muscle strength and quality [6], and ultimately reducing physical

function and mobility [5, 39]. We extended these findings by highlighting how mitochondrial

energetics may independently and combined with cardiorespiratory fitness also contribute to

lower walk speed in older adults with diabetes. This finding is of particular interest because

slower walking speed is an important indicator of health and has been shown to associate with

survival in older adults [38]. This data raises the possibility that cardiorespiratory physiology and

mitochondrial energetics could be targeted to prevent mobility loss in older adults with diabetes.

There are some limitations to the study that should be considered. First, the duration of

diabetes status of SOMMA participants and the length of time that participants have been taking

oral hypoglycemic agents were not recorded. We acknowledge that our study participants were

predominantly (85.6%) non-Hispanic White which limits our ability to generalize to other

race/ethnic groups where mobility disability is more prevalent [40]. In addition, all participants

recruited were able to complete 4-m walk test at a gait speed of greater than or equal to 0.6m/s

and which may be faster than those with more advanced diabetes who are unable to complete

mobility related tasks [2]. Nevertheless, we observe differences in physiological and functional

aspects of skeletal muscle signifying the impact that diabetes has in the SOMMA cohort. The

strengths of our study are the large sample size of participants and rigorous assessment of

objectively measured physical activity, fitness and body composition, and collection of muscle

biopsies for fresh tissue assays.

In summary, our findings highlight that mitochondrial energetics independently and

combined with cardiorespiratory fitness contribute to the difference in 400-m and 4-m walk

speed in older adults with diabetes. Additionally, future work should aim to decipher the impact

of diabetes medication on mitochondrial function as this remains to be a gap in the literature.

ACKNOWLEDGEMENTS

Personal thanks - We gratefully acknowledge and thank all participants who participated in this

study at the University of Pittsburgh and Wake Forest University. We acknowledge the dedicated

staff and investigators at both clinical sites as well as at the San Francisco Coordinating Center.

Collection and analyses of ^{31}Phosphorous Magnetic Spectroscopy data would not have been

possible without the support from Eric Shankland.

Funding and Assistance – The National Institute on Aging (NIA) funded the Study of Muscle,

Mobility, and Aging (SOMMA; R01AG059416). In part, infrastructure support for SOMMA

was funded by the NIA Claude D. Pepper Older American Independence Centers at the

University of Pittsburgh (Pitt) and Wake Forest University School of Medicine (Wake),

P30AG024827 and P30AG021332 respectively. More SOMMA infrastructure support from the

Clinical and Translational Science Institutes is funded by the National Center for Advancing

Translational Science at both Wake (UL1TR001420). PMCo is supported by R01AG060153 and

R01AG060542. GD was supported by the American Diabetes Association (1-19-PDF-006)

during data collection and analysis.

Conflict of Interest – SRC and PMCa are consultants to Bioage Labs. PMCa is a consultant to

and owns stock in MyoCorps. All other authors report no conflict of interest.

Author Contributions and Guarantor Statement – SVR and LYL completed statistical

analysis with final review from the statistical team at the coordinating center. SVR and PMCo

conceived the idea and co-wrote the manuscript. LYL, GD, SM, PMCo and PK completed

experiments or completed quality control, validation, and interpretation of data. LYL, GD,

PMCa, PK, TM, SM, MJJ, AJM, EEK, DJM, FGST, SRC, ABN, RTH, SBK, BHG edited the

manuscript. SRC, PMCa, ABN, SBK, RTH, and BHG enabled the study with either funding

acquisition, project administration, and/or conceptualization to the study.

Prior Presentations – Data from this manuscript has been presented at the following meetings:

American Diabetes Association 83[rd] Scientific Sessions May 30[th] - June 6[th] 2022, and the

American College of Sports Medicine Annual Meeting and World Congress, May 30[th] – June

2[nd], 2023.

REFERENCES

1. *Centers for Disease Control and Prevention. National Diabetes Statistics Report website.*
2. Gregg, E.W., et al., *Diabetes and physical disability among older U.S. adults.* Diabetes Care, 2000. **23**(9): p. 1272-7.
3. Ryerson, B., et al., *Excess physical limitations among adults with diabetes in the U.S. population, 1997-1999.* Diabetes Care, 2003. **26**(1): p. 206-10.
4. Gregg, E.W., et al., *Diabetes and incidence of functional disability in older women.* Diabetes Care, 2002. **25**(1): p. 61-7.
5. Volpato, S., et al., *Role of muscle mass and muscle quality in the association between diabetes and gait speed.* Diabetes Care, 2012. **35**(8): p. 1672-9.
6. Park, S.W., et al., *Decreased muscle strength and quality in older adults with type 2 diabetes: the health, aging, and body composition study.* Diabetes, 2006. **55**(6): p. 1813-8.
7. Park, S.W., et al., *Excessive loss of skeletal muscle mass in older adults with type 2 diabetes.* Diabetes Care, 2009. **32**(11): p. 1993-7.
8. Coen, P.M., et al., *Skeletal muscle mitochondrial energetics are associated with maximal aerobic capacity and walking speed in older adults.* J Gerontol A Biol Sci Med Sci, 2013. **68**(4): p. 447-55.
9. Gonzalez-Freire, M., et al., *Skeletal muscle ex vivo mitochondrial respiration parallels decline in vivo oxidative capacity, cardiorespiratory fitness, and muscle strength: The Baltimore Longitudinal Study of Aging.* Aging Cell, 2018. **17**(2).
10. Choi, S., et al., *31P Magnetic Resonance Spectroscopy Assessment of Muscle Bioenergetics as a Predictor of Gait Speed in the Baltimore Longitudinal Study of Aging.* J Gerontol A Biol Sci Med Sci, 2016. **71**(12): p. 1638-1645.
11. Newman, A.B., et al., *Walking performance and cardiovascular response: associations with age and morbidity--the Health, Aging and Body Composition Study.* J Gerontol A Biol Sci Med Sci, 2003. **58**(8): p. 715-20.
12. Phielix, E., et al., *Lower intrinsic ADP-stimulated mitochondrial respiration underlies in vivo mitochondrial dysfunction in muscle of male type 2 diabetic patients.* Diabetes, 2008. **57**(11): p. 2943-9.
13. Kelley, D.E., et al., *Dysfunction of mitochondria in human skeletal muscle in type 2 diabetes.* Diabetes, 2002. **51**(10): p. 2944-50.
14. Mogensen, M., et al., *Mitochondrial respiration is decreased in skeletal muscle of patients with type 2 diabetes.* Diabetes, 2007. **56**(6): p. 1592-9.
15. Boushel, R., et al., *Patients with type 2 diabetes have normal mitochondrial function in skeletal muscle.* Diabetologia, 2007. **50**(4): p. 790-6.
16. Larsen, S., et al., *Are substrate use during exercise and mitochondrial respiratory capacity decreased in arm and leg muscle in type 2 diabetes?* Diabetologia, 2009. **52**(7): p. 1400-8.
17. Schrauwen-Hinderling, V.B., et al., *Impaired in vivo mitochondrial function but similar intramyocellular lipid content in patients with type 2 diabetes mellitus and BMI-matched control subjects.* Diabetologia, 2007. **50**(1): p. 113-20.
18. Distefano, G., et al., *Physical activity unveils the relationship between mitochondrial energetics, muscle quality, and physical function in older adults.* J Cachexia Sarcopenia Muscle, 2018. **9**(2): p. 279-294.
19. Leite, S.A., et al., *Low cardiorespiratory fitness in people at risk for type 2 diabetes: early marker for insulin resistance.* Diabetol Metab Syndr, 2009. **1**(1): p. 8.
20. Lee, D.C., et al., *Associations of cardiorespiratory fitness and obesity with risks of impaired fasting glucose and type 2 diabetes in men.* Diabetes Care, 2009. **32**(2): p. 257-62.

21.   Cummings, S.R., et al., *The Study of Muscle, Mobility and Aging (SOMMA). A Unique Cohort*
Study about the Cellular Biology of Aging and Age-related Loss of Mobility. J Gerontol A Biol Sci
Med Sci, 2023.

22.   Montoye, A.H.K., et al., *Development of cut-points for determining activity intensity from a wrist-*
worn ActiGraph accelerometer in free-living adults. J Sports Sci, 2020. **38**(22): p. 2569-2578.

23.   Balady, G.J., et al., *Clinician's Guide to cardiopulmonary exercise testing in adults: a scientific*
statement from the American Heart Association. Circulation, 2010. **122**(2): p. 191-225.

24.   Mau, T., et al., *Mitochondrial energetics in skeletal muscle are associated with leg power and*
cardiorespiratory fitness in the Study of Muscle, Mobility, and Aging (SOMMA). J Gerontol A Biol
Sci Med Sci, 2022.

25.   Naressi, A., et al., *Java-based graphical user interface for MRUI, a software package for*
quantitation of in vivo/medical magnetic resonance spectroscopy signals. Comput Biol Med,
2001. **31**(4): p. 269-86.

26.   Borga, M., et al., *Reproducibility and repeatability of MRI-based body composition analysis.*
Magn Reson Med, 2020. **84**(6): p. 3146-3156.

27.   Toledo, F.G., et al., *Effects of physical activity and weight loss on skeletal muscle mitochondria*
and relationship with glucose control in type 2 diabetes. Diabetes, 2007. **56**(8): p. 2142-7.

28.   Coen, P.M., et al., *Exercise and Weight Loss Improve Muscle Mitochondrial Respiration, Lipid*
Partitioning, and Insulin Sensitivity After Gastric Bypass Surgery. Diabetes, 2015. **64**(11): p. 3737-
50.

29.   Pruchnic, R., et al., *Exercise training increases intramyocellular lipid and oxidative capacity in*
older adults. Am J Physiol Endocrinol Metab, 2004. **287**(5): p. E857-62.

30.   Kelley, D.E. and J.A. Simoneau, *Impaired free fatty acid utilization by skeletal muscle in non-*
insulin-dependent diabetes mellitus. J Clin Invest, 1994. **94**(6): p. 2349-56.

31.   Wessels, B., et al., *Metformin impairs mitochondrial function in skeletal muscle of both lean and*
diabetic rats in a dose-dependent manner. PLoS One, 2014. **9**(6): p. e100525.

32.   Konopka, A.R., et al., *Metformin inhibits mitochondrial adaptations to aerobic exercise training*
in older adults. Aging Cell, 2019. **18**(1): p. e12880.

33.   de Maranon, A.M., et al., *Metformin modulates mitochondrial function and mitophagy in*
peripheral blood mononuclear cells from type 2 diabetic patients. Redox Biol, 2022. **53**: p.
102342.

34.   McKenzie, A.I., et al., *Short-term exposure to a clinical dose of metformin increases skeletal*
muscle mitochondrial H(2)O(2) emission and production in healthy, older adults: A randomized
controlled trial. Exp Gerontol, 2022. **163**: p. 111804.

35.   Larsen, S., et al., *Metformin-treated patients with type 2 diabetes have normal mitochondrial*
complex I respiration. Diabetologia, 2012. **55**(2): p. 443-9.

36.   Zhao, G., et al., *Physical activity in U.S. older adults with diabetes mellitus: prevalence and*
correlates of meeting physical activity recommendations. J Am Geriatr Soc, 2011. **59**(1): p. 132-7.

37.   Coen, P.M., et al., *Reduced skeletal muscle oxidative capacity and elevated ceramide but not*
diacylglycerol content in severe obesity. Obesity (Silver Spring), 2013. **21**(11): p. 2362-71.

38.   Studenski, S., et al., *Gait speed and survival in older adults.* JAMA, 2011. **305**(1): p. 50-8.

39.   De Rekeneire, N., et al., *Diabetes is associated with subclinical functional limitation in*
nondisabled older individuals: the Health, Aging, and Body Composition study. Diabetes Care,
2003. **26**(12): p. 3257-63.

40.   Okoro, C.A., et al., *Prevalence of Disabilities and Health Care Access by Disability Status and Type*
Among Adults - United States, 2016. MMWR Morb Mortal Wkly Rep, 2018. **67**(32): p. 882-887.

Table 1: Participant Characteristics

Basic Characteristics	Without Diabetes	With Diabetes	p-value
N	717	159	
Age (year)	76.47 ± 5.05	75.79 ± 4.84	0.12
Female	61(438)	51(81)	**0.02**
Race (white)	88(631)	70(111)	**<0.001**
Hispanic ethnicity	1(7)	1(2)	0.76
Weight (kg)	74.74 ± 14.75	82.41 ± 15.58	**<0.001**
Height (m)	1.66 ± 0.10	1.67 ± 0.10	0.16
Waist circumference (cm)	92.69 ± 12.98	100.90 ± 12.57	**<0.001**
BMI (kg/m^2)	27.17 ± 4.44	29.55 ± 4.63	**<0.001**
HbA1c (%)	5.51 ± 0.31	6.73 ± 0.89	**<0.001**
Systolic blood pressure (mmHg)	129.71 ± 15.80	133.72 ± 16.08	**0.004**
Co-morbidities			
Lung disease	13(93)	14(22)	0.77
Arthritis	57(409)	52 (83)	0.25
Stroke	2(16)	3(5)	0.51
Heart disease	6(42)	11(18)	**0.02**
Cancer	43(308)	24(38)	**<0.001**
Hypertension	26(187)	37(59)	**0.006**
Kidney disease/renal failure	3(24)	6(9)	0.19
Peripheral vascular disease	1(4)	1(2)	0.37
Hypothyroidism	19(133)	17(27)	0.64
Glaucoma	9(65)	11(18)	0.39
Macular degeneration	9(61)	5(8)	0.12
Cataracts	66(472)	62(99)	0.37
Hypoglycemic Medications			
All	--	75(118)	
• Insulin	--	19(23)	
• Metformin	--	84(99)	
• Thiazolidinediones	--	3(3)	
Physical Activity			
Total daily steps	7066.68 ± 3233.95	5848.70 ± 2791.12	**<0.001**
Total sedentary time (min)	610.10 ± 112.28	659.40 ± 108.65	**<0.001**
Time in light physical activity (min)	98.20 ± 26.54	92.84 ± 29.16	**0.027**
Time in moderate to vigorous physical	192.95 ± 86.40	155.66 ± 76.74	**<0.001**

activity (min)

Body Adiposity

Abdominal adipose tissue volume (L)	7.50 ± 3.14	8.30 ± 3.31	**0.005**
Visceral adipose tissue volume (L)	3.96 ± 2.21	5.36 ± 2.31	**<0.001**
Quadricep fat infiltration (%)	0.07 ± 0.02	0.07 ± 0.02	**0.002**

Data presented as means ± standard deviation or % (N). Hypoglycemic medications is calculated only for those taking medications. Significance accepted at **p<0.05** and indicated with bold font.

Figure 1. Percent reduction in the association between diabetes and 400-m and 4-m walking speed after adjusting for both individual and combined carbohydrate and fatty acid supported respiration, and cardiorespiratory fitness. Bars depict the reduction in β coefficient from linear regression models (models 2-6 compared to the base model: [1-(β adjusted/β base adjusted)]. Base model: age, gender, and race, model 2: base model + maxOXPHOS$_{CHO}$, model 3: base model + maxOXPHOS$_{FAO}$, model 4: base model + VO$_2$peak, model 5: base model + VO2peak + maxOXPHOS$_{CHO}$, model 6: base model + VO$_2$peak + maxOXPHOS$_{FAO}$.

Table 2. Cardiorespiratory fitness and mitochondrial energetics in older adults with and without diabetes

Model	Diabetes Status	VO$_2$peak	OXPHOS$_{CHO}$	$_{max}$OXPHOS$_{CHO}$	$_{max}$ETS$_{CHO}$	OXPHOS$_{FAO}$	$_{max}$OXPHOS$_{FAO}$
Unadjusted	WOD	1532.10 ± 16.67	**31.08 ± 0.45**	**60.63 ± 0.74**	**81.43 ± 0.97**	12.94 ± 0.22	**58.22 ± 0.84**
	WD	1513.75 ± 36.19	**28.10 ± 0.97#**	**55.11 ± 1.61#**	**73.80 ± 2.11#**	11.93 ± 0.47	**53.31 ± 1.78#**
Model 1	WOD	**1486.68 ± 19.07**	**30.14 ± 0.91**	**59.02 ± 1.45**	**77.53 ± 1.90**	**12.58 ± 0.47**	**59.25 ± 1.64**
	WD	**1386.69 ± 25.66***	**27.88 ± 1.11#**	**54.98 ± 1.80#**	**71.25 ± 2.37#**	**11.49 ± 0.55#**	**54.12 ± 1.93#**
Model 1 + PA	WOD	**1487.30 ± 19.53**	30.15 ± 0.91	58.65 ± 1.47	**77.02 ± 1.98**	12.60 ± 0.50	**59.21 ± 1.61**
	WD	**1412.11 ± 26.03#**	28.51 ± 1.11	55.89 ± 1.78	**77.83 ± 2.44#**	11.70 ± 0.59	**54.29 ± 1.92#**
Model 1 + ASAT	WOD	**1492.37 ± 19.52**	**30.34 ± 0.94**	**59.50 ± 1.49**	**77.64 ± 1.95**	**12.67 ± 0.48**	**59.83 ± 1.67**
	WD	**1376.07 ± 26.56***	**27.83 ± 1.15#**	**55.07 ± 1.82#**	**70.92 ± 2.45#**	**11.37 ± 0.57#**	**53.85 ± 1.98#**
Model 1 + VAT	WOD	**1488.92 ± 19.56**	30.16 ± 0.94	**59.16 ± 1.49**	**77.33 ± 1.96**	12.62 ± 0.49	**59.57 ± 1.68**
	WD	**1386.86 ± 26.60***	28.04 ± 1.15	**55.46 ± 1.81#**	**71.29 ± 2.45#**	**11.50 ± 0.57#**	**54.66 ± 1.98#**
Model 1 + QFI	WOD	**1486.37 ± 19.00**	**30.28 ± 0.93**	**59.38 ± 1.48**	**77.78 ± 1.94**	**12.68 ± 0.48**	**59.74 ± 1.68**
	WD	**1386.10 ± 25.75***	**27.97 ± 1.14#**	**55.33 ± 1.81#**	**71.06 ± 2.43#**	**11.37 ± 0.57#**	**54.22 ± 1.98#**

Mean ± standard error with significance accepted at ***p<0.001, #p<0.05** and represented with bold font. Model 1 is adjusted for age, race, gender, site/technician, BMI, and comorbidities. PA = physical activity, ASAT = abdominal subcutaneous adipose tissue, VAT = visceral adipose tissue, QFI = muscle quadricep fat infiltration, WOD = without diabetes, WD = with diabetes, OXPHOS$_{CHO}$ = carbohydrate supported respiration, maxOXPHOS$_{CHO}$ = carbohydrate supported maximal complex I+II stimulated respiration, maxETS$_{CHO}$ = carbohydrate supported maximal electron transport system, OXPHOS$_{FAO}$ = fatty acid oxidation (FAO) supported respiration, maxOXPHOS$_{FAO}$ = fatty acid oxidation supported maximal complex I+II stimulated respiration.

Table 3: Multivariable linear regression analysis for the association of diabetes with 400m and 4m walk speed

Multivariable model	Without Diabetes Means ± SE	With Diabetes β ± SE	P
400m walk speed (m/sec)			
Base Model: adjusted for gender, age and race	1.06 ± 0.01	-0.05 ± 0.01	**0.00**
Model 2: base model plus maxOXPHOS$_{CHO}$	1.06 ± 0.01	-0.04 ± 0.02	**0.02**
Model 3: base model plus maxOXPHOS$_{FAO}$	1.06 ± 0.01	-0.04 ± 0.02	**0.02**
Model 4: base model plus VO$_2$peak	1.07 ± 0.01	-0.04 ± 0.01	**0.01**
Model 5: base model plus VO$_2$peak and maxOXPHOS$_{CHO}$	1.07 ± 0.01	-0.03 ± 0.02	0.07
Model 6: base model plus VO$_2$peak and maxOXPHOS$_{FAO}$	1.07 ± 0.01	-0.02 ± 0.02	0.20
Fully adjusted: model 5 plus BMI and comorbidities	1.06 ± 0.01	0.003 ± 0.01	0.87
Fully adjusted: model 6 plus BMI and comorbidities	1.06 ± 0.01	-0.005 ± 0.02	0.78
4m walk speed (m/sec)			
Base Model: adjusted for gender, age, and race	1.05 ± 0.01	-0.04 ± 0.02	**0.01**
Model 2: base model plus maxOXPHOS$_{CHO}$	1.04 ± 0.01	-0.03 ± 0.02	0.12
Model 3: base model plus maxOXPHOS$_{FAO}$	1.05 ± 0.01	-0.01 ± 0.02	0.51
Model 4: base model plus VO$_2$peak	1.05 ± 0.01	-0.04 ± 0.02	**0.04**
Model 5: base model plus VO$_2$peak and maxOXPHOS$_{CHO}$	1.05 ± 0.01	-0.02 ± 0.02	0.27
Model 6: base model plus VO$_2$peak and maxOXPHOS$_{FAO}$	1.05 ± 0.01	0.00 ± 0.02	0.99
Fully adjusted: model 5 plus BMI and comorbidities	1.05 ± 0.01	0.003 ± 0.02	0.84
Fully adjusted: model 6 plus BMI and comorbidities	1.05 ± 0.01	0.01 ± 0.02	0.50

Without diabetes, mean ± standard error. With diabetes, β estimate ± standard error with significance accepted at *p<0.05. maxOXPHOS$_{CHO}$ = carbohydrate supported maximal complex I+II stimulated respiration, maxOXPHOS$_{FAO}$ = fatty acid oxidation supported maximal complex I+II stimulated respiration. Models including maxOXPHOS$_{CHO}$ and maxOXPHOS$_{FAO}$ are adjusted for technician, and models with VO$_2$peak are adjusted for site.

Graphical Abstract

Functional Tests

Without Diabetes	With Diabetes
N=717 400m walk speed N=717 4m walk speed	N=159 400m walk speed N=159 4m walk speed

Mitochondrial Energetics

Without Diabetes	With Diabetes
N=663 ATPmax N=614 $OXPHOS_{CHO}$ N=612 $maxOXPHOS_{CHO}$ N=503 maxETS N=446 $OXPHOS_{FAO}$ N=445 $maxOXPHOS_{FAO}$	N=147 ATPmax N=130 $OXPHOS_{CHO}$ N=130 $maxOXPHOS_{CHO}$ N=106 maxETS N=100 $OXPHOS_{FAO}$ N=99 $maxOXPHOS_{FAO}$

Cardiorespiratory Fitness

Without Diabetes	With Diabetes
N=674 have data for VO_2peak	N=143 have data for VO_2peak

Physical Activity

Without Diabetes	With Diabetes
N=614 Total daily steps N=556 Time spent sedentary N=667 Time spent in light Physical activity N=667 Time spent in moderate to vigorous physical activity	N=137 Total daily steps N=127 Time spent sedentary N=153 Time spent in light Physical activity N=153 Time spent in moderate to vigorous physical activity

Body Composition

Without Diabetes	With Diabetes
N=681 Quadricep muscle fat infiltration N=683 Adipose tissue depots	N=149 Quadricep muscle fat infiltration N=149 Adipose Tissue depots

Supplemental Figure 1. SOMMA cohort sample size for analysis of muscle function, mitochondrial energetics, CRF, PA and body composition. Participants missing data for ex-vivo mitochondrial energetics is due exclusions samples passing quality control. Missing ATPmax measures were due to safety reasons or exclusion due to technical conditions during testing.

Supplemental Table 1. Association of hypoglycemic drugs and metformin with cardiorespiratory fitness and mitochondrial energetics

	All hypoglycemic agents		Metformin	
	β estimate	p-value	β estimate	p-value
VO_2peak	-82.27 ± 62.24	0.19	-13.08 ± 55.47	0.81
ATPmax	**-0.05 ± 0.03**	**0.05**	**-0.06 ± 0.02**	**0.01***
$OXPHOS_{CHO}$	-1.03 ± 2.01	0.61	-0.71 ± 1.81	0.70
$maxOXPHOS_{CHO}$	-3.52 ± 3.20	0.27	-3.11 ± 2.89	0.28
$maxETS_{CHO}$	-2.95 ± 4.45	0.51	-4.07 ± 4.04	0.32
$OXPHOS_{FAO}$	-1.35 ± 1.06	0.21	-0.68 ± 1.00	0.50
$maxOXPHOS_{FAO}$	-2.57 ± 3.69	0.49	-5.30 ± 3.43	0.13

β estimate ± standard error with significance accepted at **p<0.05**. Data adjusted for age, race, and site/technician. ATPmax = maximal adenosine triphosphate production, $OXPHOS_{CHO}$ = carbohydrate supported respiration, $maxOXPHOS_{CHO}$ = carbohydrate supported maximal complex I+II stimulated respiration, $maxETS_{CHO}$ = carbohydrate supported maximal electron transport system, $OXPHOS_{FAO}$ = fatty acid oxidation (FAO) supported respiration, $maxOXPHOS_{FAO}$ = fatty acid oxidation supported maximal complex I+II stimulated respiration.