Epistasis regulates genetic control of cardiac hypertrophy

Qianru Wang*[1], Tiffany M. Tang*[2], Nathan Youlton[1], Chad S. Weldy[1], Ana M. Kenney[2], Omer Ronen[2], J. Weston Hughes[1], Elizabeth T. Chin[1], Shirley C. Sutton[1], Abhineet Agarwal[2], Xiao Li[2], Merle Behr[3], Karl Kumbier[4], Christine S. Moravec[5], W. H. Wilson Tang[5,6], Kenneth B. Margulies[7,8], Thomas P. Cappola[7,8], Atul J. Butte[9,10], Rima Arnaout[9,10], James B. Brown[2,10,11], James R. Priest[1,10,12], Victoria N. Parikh[1], Bin Yu[†2,10,13,14], Euan A. Ashley[†1,10]

* Equal contribution

[†] Equal senior authorship

Affiliations
1. Division of Cardiovascular Medicine, Department of Medicine, Stanford University, Stanford, CA, USA
2. Department of Statistics, University of California, Berkeley, Berkeley, CA, USA
3. Faculty of Informatics and Data Science, University of Regensburg, Regensburg, Germany
4. Department of Pharmaceutical Chemistry, University of California, San Francisco, San Francisco, CA, USA
5. Department of Cardiovascular and Metabolic Sciences, Lerner Research Institute, Cleveland Clinic, Cleveland, OH, USA
6. Department of Cardiovascular Medicine, Heart Vascular and Thoracic Institute, Cleveland Clinic, Cleveland, OH, USA
7. Division of Cardiovascular Medicine, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA
8. Hospital of The University of Pennsylvania, Philadelphia, PA, USA
9. Bakar Computational Health Sciences Institute, University of California, San Francisco, San Francisco, CA, USA
10. Chan Zuckerberg Biohub – San Francisco, San Francisco, CA, USA
11. Division of Environmental Genomics and Systems Biology, Lawrence Berkeley National Laboratory, Berkeley, CA, USA
12. Tenaya Therapeutics, San Francisco, CA, USA
13. Department of Electrical Engineering and Computer Science, University of California, Berkeley, Berkeley, CA, USA
14. Center for Computational Biology, University of California, Berkeley, Berkeley, CA, USA

Correspondence
Dr. Euan Ashley
Roger and Joelle Professor of Genomics and Precision Health
Professor of Cardiovascular Medicine, Genetics and Data Science
Division of Cardiovascular Medicine, Department of Medicine
Stanford University
Falk Cardiovascular Research Building, 870 Quarry Road, Stanford, CA 94304
euan@stanford.edu

Dr. Bin Yu
Professor of Statistics
Department of Statistics and Electrical Engineering and Computer Sciences
University of California Berkeley
367 Evans Hall, University Dr, Berkeley, CA 94720
binyu@berkeley.edu

Abstract

The combinatorial effect of genetic variants is often assumed to be additive. Although genetic variation can clearly interact non-additively, methods to uncover epistatic relationships remain in their infancy. We develop low-signal signed iterative random forests to elucidate the complex genetic architecture of cardiac hypertrophy. We derive deep learning-based estimates of left ventricular mass from the cardiac MRI scans of 29,661 individuals enrolled in the UK Biobank. We report epistatic genetic variation including variants close to *CCDC141*, *IGF1R*, *TTN*, and *TNKS.* Several loci not prioritized by univariate genome-wide association analysis are identified. Functional genomic and integrative enrichment analyses reveal a complex gene regulatory network in which genes mapped from these loci share biological processes and myogenic regulatory factors. Through a network analysis of transcriptomic data from 313 explanted human hearts, we show that these interactions are preserved at the level of the cardiac transcriptome. We assess causality of epistatic effects via RNA silencing of gene-gene interactions in human induced pluripotent stem cell-derived cardiomyocytes. Finally, single-cell morphology analysis using a novel high-throughput microfluidic system shows that cardiomyocyte hypertrophy is non-additively modifiable by specific pairwise interactions between *CCDC141* and both *TTN* and *IGF1R*. Our results expand the scope of genetic regulation of cardiac structure to epistasis.

Main

Heart disease is closely tied to the structure of the heart[1]. Heart failure, a syndrome characterized by increased pressure within, or decreased output from, the heart is influenced by structural features including atrial and ventricular chamber size and wall thickness[2–5]. Left ventricular hypertrophy – increased thickness of the left ventricle (LV) – can be the result of mendelian genetic diseases like hypertrophic cardiomyopathy[6] but is also a complex phenotypic trait influenced by multiple factors, genetic and environmental. Progressive LV hypertrophy carries significant independent risk for incident heart failure, atrial arrhythmia, and sudden death[7–10], highlighting the need to understand genetic determinants of cardiac phenotype.

Recent discoveries leveraging cardiac magnetic resonance imaging in the UK Biobank (UKBB) have revealed that cardiac structure is in part determined by complex genetics[11–14]. Common genetic variants, many located near genetic loci associated with dilated cardiomyopathy and heart failure, have been found to influence LV size and systolic function[11]. Further, specific genetic variants that influence LV trabeculation have been shown to impact systolic function and overall risk of cardiomyopathy[13]. However, these variants remain inadequate to explain the total heritable disease risk[15]. Indeed, common genetic variants rarely act independently and additively as modeled by most genome-wide association studies (GWAS)[16]. There is growing evidence to support a disease risk model in which multiple genes interact non-additively with each other through epistasis[17,18]. Recent work has shown that common genetic variation influences susceptibility and expressivity of hypertrophic cardiomyopathy[14]. This raises the possibility that common epistatic interactions drive cardiac phenotype, holding significant potential for uncovering disease mechanisms and developing potential therapeutic strategies.

Several computational and experimental challenges need to be resolved to allow robust identification of epistasis. First, the combinatorial nature of interactions makes an exhaustive search computationally intractable. To reduce the computational burden and ensure stable discoveries, we developed an approach based on signed iterative random forests[19,20] to uncover higher-order (not limited to pairwise)

nonlinear interactions in a computationally-tractable manner. Second, many previously reported

epistatic relationships were not replicated[21,22]. To achieve more trustworthy results, we adhered to a

new framework for veridical data science[23], centered around the principles of predictability,

computability, and stability (PCS) and the need for transparent documentation of decisions made in

data analysis pipelines. A third challenge is the generally small effect size of common genetic

variants[15,24] which impedes both the data-driven discovery and functional validation of epistatic

interactions. In human biobanks, recent advances in deep-learning-enabled phenotyping[25] using

cardiac magnetic resonance images have led to more refined phenotypes at larger scales. At the

cellular level, high-throughput microfluidic technologies[26–28] have been integrated with artificial

intelligence-based image analysis of single cell morphology[29] and human induced pluripotent stem cell-

derived cardiomyocytes[30], opening up new possibilities for rapid, label-free detection of the phenotypic

consequences of genetic perturbation.

Results

Our methodology includes four major stages: derivation of estimates of LV mass (green boxes, Fig. 1);

computational prioritization of epistatic drivers (orange boxes, Fig.1); functional interpretation of the

hypothesized epistatic genetic loci (purple boxes, Fig.1); and experimental confirmation of epistasis

through perturbation (blue boxes, Fig. 1).

Deep learning of UK Biobank cardiac imaging quantifies left ventricular hypertrophy

We accessed all cardiac magnetic resonance images from the UKBB substudy (44,503 people at the

time of this analysis)[31]. We focused on the largest ancestry subset of 29,661 unrelated individuals

(summary characteristics in Supplementary Table 1) and analyzed the most recent image per

individual. We leveraged a recent deep learning model[25] to quantify LV hypertrophy from these 29,661

multislice cine magnetic resonance images (Fig. 2a). A fully convolutional network had been previously

trained for image segmentation and was evaluated on manual pixelwise-annotations of images from

4,875 UKBB participants[25]. This fully convolutional network learns features across five different

resolutions through sequential convolutional layers interspersed with non-linearities, and has displayed

accurate performance compared to cardiac segmentation by human experts[25]. Using this segmentation

model, we extracted areas of the LV chamber wall in each slice of the short axis image at the end of

diastole. Areas extracted from each image slice in the same image stack were then integrated to

calculate the heart muscle volume, which we converted to the LV mass using a standard density of

1.05 g/mL[32]. This was normalized by body surface area, estimated using the Du Bois formula[33], to

obtain the LV mass index (LVMi, Extended Data Fig. 1). Details regarding this analysis can be found in

Methods.

Low-signal signed iterative random forests prioritizes epistatic genetic loci

We developed low-signal signed iterative random forests (lo-siRF, Fig. 2a-2e) to identify stable epistatic

interactions from the extracted LV mass and single-nucleotide variants (SNVs) from UKBB. This

method is guided by the PCS framework[23] and builds upon signed iterative random forests[19,20], a

computationally-tractable algorithm to extract predictive and stable nonlinear higher-order interactions

that frequently co-occur along decision paths in a random forest. More specifically, lo-siRF proceeds

through four steps:

1. *Dimension reduction* (Fig. 2b)*:* we combined the results of two initial genome-wide association

studies, implemented via PLINK[34] and BOLT-LMM[35] (Extended Data Fig. 2, Extended Data 1) to

reduce the interaction search space from 15 million imputed variants down to 1405 variants

(Extended Data 2). Details can be found in the Methods section *Lo-siRF step 1: Dimension*

reduction of variants via genome-wide association studies.

2. *Binarization* (Fig. 2c): we binarized the LV mass measurements into two (high and low)

balanced categories (Supplementary Table 2). This binarization simplified the original low-signal

regression problem for a (continuous) complex trait to a relatively easier classification task,

which denoises and facilitates model checking. The detailed binarization procedure can be

found in the Methods section *Lo-siRF step 2: Binarization of the left ventricular mass phenotype.*

3. *Prediction* (Fig. 2d): we trained a signed iterative random forest using the 1405 GWAS-filtered

SNVs to predict the binarized LV mass measurements. The learnt model yields on average the

highest (balanced) classification accuracy (55%) compared to other common machine learning

prediction algorithms (Supplementary Table 3). Details about the model and prediction check

can be found in the Methods section *Lo-siRF step 3: Prediction.*

4. *Prioritization* (Fig. 2e): we developed a stability-driven feature importance score (Extended Data

Fig. 3), which leveraged the fitted signed iterative random forest and a permutation test, to

aggregate SNVs into genetic loci and prioritize interactions between genetic loci. This

importance score provides the necessary new interpretable machine learning ingredient to

complete the lo-siRF discovery pipeline. Details can be found in the Methods section *Lo-siRF*

step 4: Prioritization.

Extensive documentation and justification of our modeling decisions can be found in Supplementary

Note 1, an interactive HTML webpage hosted at https://yu-group.github.io/epistasis-cardiac-

hypertrophy/.

Lo-siRF identified six genetic risk loci that exhibited stable and reliable associations with LV mass (Fig.

2f). Because these loci are either located within a gene body or in between two genes (Fig. 3a), for

convenience we denote these loci by their nearest genes. Notably, out of the six loci, three (*TTN*,

CCDC141, and *IGF1R*) were prioritized by lo-siRF as epistatic loci. These loci not only interact with

other loci, but also marginally affect LV mass. The other three lo-siRF-prioritized loci are

LOC157273;TNKS, *MIR588;RSPO3*, and *LSP1*. The *LOC157273;TNKS* locus is located within the

intergenic region between genes *LOC157273* and *TNKS* (semicolon indicates intergenic region). This

locus was prioritized by lo-siRF to be hypostatic (i.e., effects are deemed stable by lo-siRF only when

interacting with the *CCDC141* locus). Interestingly, all three identified epistatic interactions involved the

CCDC141 locus (Fig. 3a, green links in circle 1). Furthermore, while the *MIR588;RSPO3* and *LSP1* loci

lacked evidence for epistasis by lo-siRF, they were each identified to be marginally associated with LV

mass. The specific prioritization order of these loci can be found in Supplementary Table 4, and details

regarding the direction or sign of the interactions can be found in Supplementary Note 1. In total, lo-

siRF identified 283 SNVs located within the six loci (Extended Data 3, Extended Data Fig. 4). Ninety

percent of the 283 SNVs have previously been shown to harbor multiple distinct cardiac function

associations[36] in phenome-wide analyses (e.g., pulse rate, Extended Data 3), suggesting a strong

likelihood that these lo-siRF-prioritized loci contribute to determining cardiac structure and function.

Loci associated with left ventricular mass exhibit regulatory enrichment

We performed functional mapping and annotation (FUMA)[37] for the 283 lo-siRF-prioritized SNVs (Fig. 1,

purple and Fig. 3). For linkage disequilibrium (LD), we used a default threshold of $r^2 = 0.6$ and chose

the UKBB release 2b reference panel created for British and European subjects to match the

population group used for lo-siRF prioritization. FUMA identified 572 additional candidate SNVs

(Extended Data 4) in strong LD ($r^2 > 0.6$) with any of the 283 lo-siRF-prioritized SNVs, including 492

SNVs from the input GWAS associations (points in Fig. 3a, circle 8) and 80 non-GWAS-tagged SNVs

extracted from the selected reference panel (heatmap tracks in Fig. 3a, circle 8). We then assigned

these 572 FUMA-extracted candidate SNVs to a lo-siRF-prioritized locus (Fig. 2f) based on the

corresponding lo-siRF-prioritized SNV (out of the 283 SNVs), which has the maximum r^2 value with the

candidate SNV.

The two loci contributing to the top-ranked epistatic interaction by lo-siRF, the *CCDC141* and *IGF1R*

loci (Fig. 2f), both showed a significant enrichment of intronic variants relative to the background

reference panel (Fig. 3b, Extended Data 5). Over 88% of the SNVs in or in LD with these two loci were

mapped to actively transcribed chromatin states (TxWk) or enhancer states (Enh) in left ventricles

based on the ChromHMM Core 15-state model[38] (Fig. 3a, circle 7). More than 47% and 76% of the

identified SNVs in or in LD with the *CCDC141* and *IGF1R* loci, respectively, showed the highest

RegulomeDB[37,39] categorical score (ranked within category 1 from the 7 main categories). The

Combined Annotation-Dependent Depletion (CADD) score[40] was used to judge the deleteriousness of

prioritized variants (Extended Data 4). As expected, GTEx[41] data revealed that 82% of SNVs in or in LD

with the *IGF1R* locus are expression quantitative trait loci (eQTLs) for the gene *IGF1R*. In contrast, of

the SNVs in or in LD with the *CCDC141* locus, only 14% are eQTLs for gene *CCDC141* and 22% are

splicing quantitative trait loci (sQTLs) for gene *FKBP7*. This differential fractionation of eQTLs and

sQTLs suggests distinct epistatic mechanisms: the *IGF1R* locus is likely to affect the expression level of

gene *IGF1R*, whereas the *CCDC141* locus may play a regulatory role by affecting the alternative

splicing of gene *FKBP7*. Furthermore, Hi-C data indicated that all SNVs identified in or in LD with the

IGF1R locus are in 3D chromatin interaction with gene *SYNM* while more than 54% SNVs identified in

or in LD with the *CCDC141* locus are in 3D chromatin interaction with gene *TTN*. These results suggest

a possibility of higher-order interactions between more than two genes.

The *CCDC141* and *TTN* loci exhibit genomic proximity (Fig. 3a). Their interaction, however, does not

appear to stem from this proximity. Indeed, the *CCDC141* and *TTN* genes have been individually

associated with LV mass[42,43]. Due to this proximity, previous studies[44,45] have assumed *CCDC141* as a

secondary gene that affects LV mass through the *TTN* gene expression. However, we found low LD (r^2

< 0.6) between any two of the 283 lo-siRF-prioritized SNVs, suggesting an independent role of each

locus in their epistasis. In addition, we compared all the epistatic SNVs that were aggregated to the

TTN locus, including both lo-siRF-prioritized SNVs and their linked variants in LD extracted by FUMA,

with their counterparts that were filtered out by lo-siRF. We found that the *TTN* locus showed a

significant depletion of SNVs located close to (<10 kb) the gene *CCDC141* (p = 2.38E-9, two-sided

Fisher exact test). Similarly, the *CCDC141* locus showed a substantially decreased enrichment of

SNVs that are close to gene *TTN* (p = 0.02, two-sided Fisher exact test). These results suggest that

although the *CCDC141* and *TTN* loci are located close to each other in the genome, the prioritized

epistatic SNVs are located farther apart relative to randomly selected SNVs from the two loci.

In contrast to the *CCDC141* and *IGF1R* loci, the *TTN* locus showed a significant enrichment of exonic

variants and intronic variants that are transcribed into non-coding RNA (ncRNA_intronic, Fig. 3b). Of

those exonic variants, 62% are nonsynonymous. This suggests that one mechanism of epistatic effect

for the *TTN* locus could be structural alterations in the titin protein. Over 90% of SNVs in or in LD with

the *TTN* locus were mapped to actively transcribed states (Tx, TxWk) in left ventricles (Fig. 3a, circle 7).

Interestingly, these SNVs were associated with a quiescent chromatin state (Quies) in the right atrium,

indicating that the epistatic effects of the *TTN* locus may be specific to ventricular tissues. Nearly half of

SNVs in or in LD with the *TTN* locus are eQTLs for the gene *FKBP7*. In addition, 83% of these SNVs

are sQTLs for gene *FKBP7* or *TTN*, suggesting a regulatory effect of the *TTN* locus on the expression

and splicing of gene *FKBP7*. Moreover, the *TTN* locus was suggested to impact genes *PDE11A*,

RBM45, *PRKRA*, and *DFNB59* through 3D chromatin interactions.

The hypostatic locus *LOC157273;TNKS* showed a significant enrichment of variants within non-coding

RNA regions of exons and introns (Fig. 3b). Over 95% of identified SNVs in or in LD with this locus

were mapped to inactive chromatin states (ReprPCWk, Quies) in left ventricles (Fig. 3a, circle 7). This

suggests that in the absence of an epistatic partner, the *LOC157273;TNKS* locus is epigenetically

quiescent or repressed by polycomb group proteins. In addition, of all the SNVs in or in LD with this

locus, 66% are eQTLs for *MFHAS1* or *CLDN23* and 22% are in 3D chromatin interaction with gene

TNKS.

Functional annotations for the other two lo-siRF-prioritized loci that were marginally associated with LV

mass can be found in Extended Data 4 and 5.

Epistatic loci functionally map to twenty-one protein-coding genes

Three strategies, positional, eQTL, and chromatin interaction, mapped the six LV hypertrophy risk loci

to 21 protein-coding genes (Fig. 4a). Genes prioritized by eQTL and chromatin interaction mapping are

not necessarily located in the corresponding risk locus, but they are linked to SNVs within or in LD with

the locus (Fig. 3a). Among the 21 genes, *CCDC141* and *IGF1R* were prioritized by all the three

mapping strategies (Fig. 4a), suggesting that these two genes are very likely involved in determining LV

mass. Interestingly, none of the SNVs mapped to *IGF1R* were statistically significant in our GWAS

studies using BOLT-LMM and PLINK (Extended Data Fig. 2 and Extended Data 1). This reveals the

potential of lo-siRF to identify risk loci that may be overlooked by GWAS. Based on the expression data

from GTEx V8, *TTN*, *TNNT3*, and *SYNM* are up-regulated while *CLDN23* and *MFHAS1* are down-

regulated in both heart and muscle tissues (Fig. 4b). In contrast, *CCDC141* is up-regulated specifically

in heart tissues whereas *RSPO3* is down-regulated in heart but up-regulated in muscle tissues (Fig.

4b).

Ten of twenty-one genes mapped from epistatic loci interact in network analysis

We performed gene ontology (GO) and pathway enrichment analysis on the 21 genes mapped from lo-siRF loci. We adopted previously established approaches[46–48] and integrated enrichment results across libraries from multiple sources to establish a GO and pathway co-association network (Fig. 4c). To evaluate the interaction strength between any two genes in the network, we calculated a co-association score for every possible gene-gene combination (n = 72,771) from both genes prioritized and deprioritized by lo-siRF. Lo-siRF-prioritized genes are the 21 genes functionally mapped from the 283 lo-siRF-prioritized SNVs and their LD-linked SNVs (Fig. 4a). Lo-siRF-deprioritized genes are those functionally mapped from the SNVs that failed to pass the lo-siRF prioritization threshold. Compared to random gene pairs in the network, 10 genes that were functionally mapped from the lo-siRF-prioritized epistatic and hypostatic loci showed significant co-associations with multiple GO/pathways (Fig. 4c, Extended Data 6). Consistent with our hypothesized epistasis (Fig. 2f), gene *CCDC141* showed a significant co-association to *SYNM* (functionally linked to the *IGF1R* locus) and *PDE11A* and *PLEKHA3* (both functionally linked to the *TTN* locus) through the GO term of hyperactivity (excessive movement), which has been linked to increased risk of cardiac disease[49]. Beyond that, *TTN*, *IGF1R*, and *SYNM* are co-associated with kinase activity and cardiac structure related GO terms, indicating that these genes may interactively affect cardiac structure by regulating the process of kinase activity.

Genes mapped from epistatic loci are co-associated with myogenic regulatory factors

We next performed an integrative enrichment analysis to assess transcriptional regulation of genes prioritized and deprioritized by lo-siRF. Due to assay-specific limitations and biases, we integrated the enrichment results across nine distinct gene set libraries[46,47] (Fig. 4d, Extended Data 7). We found that the lo-siRF-prioritized epistatic genes shared important myogenic regulatory factors, such as MYOD1, MYF6, and MYOG (Fig. 4d, top). These myogenic regulatory factors coordinate to regulate muscle development and differentiation. In contrast, Transcription factors enriched from lo-siRF-deprioritized genes display a less coordinated regulatory pattern (Fig. 4d, bottom). To further evaluate the interaction strength between any two genes that share transcription factors, we calculated a transcription factor co-association score for all the 72,771 possible gene-gene combinations (see Methods). Compared with

random gene pairs,16 gene-gene combinations from the lo-siRF-prioritized genes displayed a

significant co-association (empirical $p < 0.05$, Fig. 4e). These co-associations were found in gene-gene

combinations from both intra- and inter- lo-siRF-prioritized loci (Fig. 4e). In particular, pairwise

combinations among *TTN*, *TNNT3*, *CCDC141*, and *SYNM* share a common splicing regulator, RBM20

(Extended Data Fig. 5). RBM20 has been reported to regulate the alternative splicing of genes

important for cardiac sarcomere organization[50]. This suggests that the splicing patterns of these four

genes are likely to be co-regulated by RBM20, which is consistent with the exhibited enrichment of

sQTLs by the *CCDC141*, *TTN* and *LSP1* lo-siRF loci (Extended Data 4).

The *CCDC141-TTN* interaction is confirmed in human heart failure transcriptomics

We proceeded to the fourth stage for experimental confirmation (Fig. 1, blue) and evaluated how the

identified epistases contribute to the progression of heart failure (Fig. 5). We employed a series of

weighted gene co-expression networks derived from human cardiac transcriptomic data from 177 failing

hearts isolated at the time of heart transplant and 136 non-failing hearts harvested from cardiac

transplant donors whose organs were not able to be placed[51] (Fig. 5a). We compared the molecular

connectivity of genes identified as likely epistatic interactors. We defined connectivity as the edge

weights between two genes normalized to the distribution of all network edge weights, and compared

this to the connectivity of all other available gene-gene combinations in the network. This revealed a

strong connection between genes *CCDC141* and *TTN* in the healthy control network ($p = 0.0026$, Fig.

5b). Although these two genes also displayed a significant connectivity ($p = 0.006$, Fig. 5c) in the heart

failure network, the connectivity is weakened in the heart failure network ($p = 0.002$, from a third

network that measures the connectivity difference between the control and heart failure networks, Fig.

5d). To gain insight into the potential underlying mechanism for this difference, we compared gene

module memberships. *CCDC141* and *TTN* are co-associated with the electron transport

chain/metabolism module in the control network while *IGF1R* is associated with the unfolded protein

response module (Fig. 5e). In the failing hearts, *TTN* and *IGF1R* are co-associated with the muscle

contraction/cardiac remodeling module, whereas *CCDC141* remains associated with the metabolism

module (Fig. 5f).

Perturbation confirms epistatic relationships in cardiomyocyte hypertrophy

We interrogated epistatic associations in a genetic model of cardiac hypertrophy (Fig. 1, blue): induced

pluripotent stem cell cardiomyocytes derived from patients with and without hypertrophic

cardiomyopathy caused by the cardiac myosin heavy chain (*MYH7*) p.R403Q variant[30] (Fig. 6a).

Cardiac myosin heavy chain 7 is a key component of the cardiac sarcomere, and the most common

cause of hypertrophic cardiomyopathy[30]. The patient presented with typical symptoms, and

echocardiography revealed severe LV hypertrophy and a small LV cavity[30]. At the cellular level,

cardiomyocytes exhibit an elevated mean cell size and non-Gaussian size distribution with a long tail

relative to the unaffected control (Fig. 6d).

To determine if *CCDC141* can act both independently and in epistatic interactions with other genes to

attenuate the pathologic cellular hypertrophy caused by *MYH7*-R403Q, we silenced genes *CCDC141*,

IGF1R, *TTN*, and gene pairs *CCDC141-IGF1R* and *CCDC141-TTN* using siRNAs in both diseased and

healthy cardiomyocytes and compared them with cells transfected with scramble siRNAs (control) (Fig.

6a and 6e). Phenotypic consequences of these perturbations on cellular morphology were then

evaluated in high-throughput using a spiral inertial microfluidic device (Fig. 6b) in combination with

automated single-cell image analysis (Fig. 6c). The microfluidic device adopted the Dean flow focusing

principle[28] (details in Extended Data Fig. 6 and Methods) to mitigate the non-uniform cell focusing[52],

thereby enhancing the imaging resolution[53] affected by the large variations in cardiomyocyte diameter

(Fig. 6d).

We first assessed the non-additive effects of the *CCDC141-IGF1R* interaction on cardiomyocyte size

(Fig. 6f). Bootstrapped hypothesis tests were performed, for which the *p*-values are capped below by p

$< 1E-4$ (Extended Data 8). Silencing *IGF1R* alone reduces the median cell size by 5.3% ± 0.4% ($p <$

1E-4) in diseased cells compared to scrambled control and 6.6% ± 0.5% ($p <$ 1E-4) in healthy cells.

Silencing *CCDC141* alone also decreases median cell size by 3.2% ± 0.5% ($p <$ 1E-4) in diseased

cells, but had no impact on healthy cells. Digenic silencing of *CCDC141* and *IGF1R* reveals a

synergistic effect on attenuating pathologic cell hypertrophy in diseased cells, resulting in an 8.5% ±

0.3% (p < 1E-4) decrease in the median cell size. This is consistent in healthy cells, where silencing

CCDC141 alone fails to affect cell size, but digenic silencing of *CCDC141* and *IGF1R* decreases the

median cell size by 9.3% ± 0.5% (p < 1E-4). Moreover, this effect appears to be non-additive for both

healthy and diseased variant cells (p < 1E-4 for non-additivity), consistent with an epistatic mechanism.

These findings serve to confirm the strongest epistatic association identified by lo-siRF (Fig. 2f).

We found a comparable non-additive effect for the *CCDC141-TTN* interaction. Digenic silencing of

CCDC141-TTN leads to a pronounced reduction in median cell size (by 5.8% ± 0.6% for healthy cells

and 3.3% ± 0.4% for diseased cells, p < 1E-4) relative to monogenic silencing, and appears to be non-

additive for both healthy and diseased cells (p < 1E-4 for non-additivity). Additionally, *CCDC141* and

TTN show distinctive independent roles in repressing cardiomyocyte hypertrophy. In healthy cells,

monogenic silencing of *TTN* leads to a larger cell size reduction compared to the case of silencing

CCDC141. In contrast, diseased cells display a larger size reduction in response to monogenic

silencing of *CCDC141*.

Furthermore, both *CCDC141-IGF1R* and *CCDC141-TTN* interactions show a stronger effect on

rescuing larger cardiomyocytes over smaller ones in both cell lines (Extended Data Fig. 7 and 8). In

contrast, monogenic silencing does not exhibit such a non-uniform effect on reshaping the cell size

distribution, which reinforces the hypothesized non-additivity of these two epistatic interactions (details

in Extended Data 8 and Supplementary Note 2).

Recent studies have shown that cellular morphological features, such as cell boundary and textural

irregularities, are informative readouts of cytoskeletal structure, which is highly associated with disease

state in hypertrophic cardiomyopathy[29,54]. We analyzed relative changes in cell shape and texture (Fig.

6g) by measuring the counts of peak intensities normalized to the total number of pixels enclosed by

the cell boundary (Fig. 6h). Cells with a high normalized peak number display a ruffled texture, which

manifests in unevenly distributed 2D intensities (Fig. 6j). Our analysis shows that silencing both

CCDC141 and *IGF1R* (circles in Fig. 6g, left) yields a larger increase in intensity peak number than

silencing *IGF1R* alone (triangles in Fig. 6g, left) for both cell lines, exhibiting a synergistic epistasis

between *CCDC141* and *IGF1R* (p < 1E-4 for non-additivity). We also analyzed cell roundness error, a

measure of how far radii measured on the cell outline deviate from a perfect circle (Fig. 6h). This

parameter increases with an increasing cell boundary waviness or elongation (Fig. 6i). We show that

the silencing of *CCDC141* and *IGF1R* synergistically interact to increase roundness error of diseased

cardiomyocytes (p < 1E-4 for non-additivity, Fig. 6g, left). In addition, *CCDC141* and *TTN* display

antagonistic epistasis and synergistic epistasis in their impact on roundness error for healthy and

diseased cells (p < 1E-4 for non-additivity, Fig. 6g, right), respectively.

Discussion

While computational models[17,18] have supported epistatic contributions to human complex traits and

disease risk, examples in the literature are rare, with even fewer experimentally confirmed. Here, we

developed a veridical machine learning[23] approach to identify epistatic associations with cardiac

hypertrophy derived from a deep learning model that estimates LV mass from cardiac imaging of

almost thirty thousand individuals in the UK Biobank. We report novel epistatic effects on LV mass of

common genetic variants associated with *CCDC141*, *TTN*, and *IGF1R.* We used established tools to

functionally link risk loci to genes, and then confirmed gene level co-associations through network

analyses, including via shared transcription factors and pathways enriched against multiple annotated

gene set libraries and co-expression networks we built using transcriptomic data from over three

hundred healthy and diseased human hearts. Finally, using a cellular disease model incorporating

monogenic and digenic silencing of individual genes, we assessed phenotypic changes in

cardiomyocyte size and morphology using a novel microfluidic system, confirming the non-additive

nature of the interactions.

Our approach advances epistasis discovery in several key ways. First, unlike studies relying on linear-

based models[55–58], we leverage a more realistic, nonlinear tree-based model that mirrors the

thresholding (or switch-like) behavior commonly observed in biomolecular interactions[59]. Second, in

contrast to other tree-based approaches that evaluate interactions on a variant-by-variant basis[60–64], our

novel stability-driven importance score consolidates individual variants into loci for the assessment of

feature importance, allowing for more reliable extraction of epistatic interactions from weak association

signals. This is particularly valuable for evaluating non-coding variants. Moreover, instead of

exhaustively searching all possible interactions, signed iterative random forests internally employ a

computationally-efficient algorithm, which automatically narrows the search space of interactions to only

those that stably appear in the forest and thus achieves a scalability much higher than existing tree-

based approaches[62,65]. This allows lo-siRF to handle larger datasets without the need for LD pruning

before the interaction search, which may inadvertently eliminate important epistatic variants, given that

epistasis between loci in strong LD has been evidenced by a recent study[66]. Furthermore, our

computational prioritization is rigorously validated through multiple functional network analyses and

robust experimental confirmation.

Our results add to a small literature on epistasis in cardiovascular disease. Two recent studies have

found epistasis influencing the risk of coronary artery disease[17,18]. Li et al.[18] identified epistasis between

ANRIL and *TMEM106B* in coronary artery tissues. Although their method predicted functionally

interpretable interactions between risk loci of interest, they relied heavily on prior knowledge and careful

selection of the causal gene pairs,[18] making the approach challenging to scale. Zeng et al.[17] used

population-scale data and performed epistasis scans from regions around 56 known risk loci. This

study identified epistasis between variants in *cis* at the *LPA* locus without experimental confirmation. In

contrast, our approach allows discovery of not only cis-epistasis, but also long-range interactions

between interchromosomal loci (e.g., *CCDC141* and *IGF1R*) and is supported by gene perturbation

experiments. More importantly, both previous studies searched for interactions around known risk loci

identified by genome-wide association, which can be far away from the possible epistatic or hypostatic

loci that are statistically insignificant in linear univariate association studies. In addition, both studies

relied on a logistic regression model, which imposes restrictive assumptions that can be avoided using

a nonlinear machine learning approach as in lo-siRF.

Our study has limitations. We focused this analysis on a single ancestry in order to enhance the

likelihood of finding reliable interactions from weak association signals. These findings cannot be

automatically applied to others. It was not feasible to conduct a formal genetic replication study

because the UK Biobank is the only large-scale population cohort with integrated cardiac magnetic

resonance images and genetic data. However, to help reduce the possibility of overfitting and increase

generalizability, lo-siRF employed numerous stability analyses (see Supplementary Note 1) in addition

to a proper training-validation-test data split. Beyond these computational checks, we also present

functional supporting evidence and experimental validation. Our computational prioritization via lo-siRF

currently groups SNVs based on genomic proximity, without accounting for their functional

interdependencies, but this could be addressed by integrating functional annotation into the lo-siRF

pipeline. Lastly, lo-siRF is not as scalable as linear-based methods, though it is more scalable than

alternative tree-based methods for epistasis detection. It also should be noted that although this study

did not identify stable higher-order (> order-2) interactions due to the weak association signal between

SNVs and LV mass, the method exhibits the capability to detect such interactions for broader

phenotypes and complex traits without incurring additional computational cost.

In summary, our work adds to the discovery toolkit for the genomic architecture of complex traits and

expands the scope of genetic regulation of cardiac structure to epistasis.

Online Methods

Study participants

The use of human subjects (IRB - 4237) and human-derived induced pluripotent stem cells (SCRO - 568) in this study has been approved by the Stanford Research Compliance Office. The UK Biobank received ethical approval from the North West - Haydock Research Ethics Committee (21/NW/0157).

The UK Biobank (UKBB) is a biomedical database with detailed phenotypic and genetic data from over half a million UK individuals between ages 40 and 69 years at recruitment[67]. In this study, we restricted our analysis to the largest ancestry subset (i.e., the White British population) of 29,661 unrelated individuals who have both genetic and cardiac magnetic resonance imaging (MRI) data from the UKBB (Supplementary Table 1). More specifically, we considered only those individuals from the UKBB cohort who self-reported as White British and have similar genotypic backgrounds based on principal components analysis as described in prior work[67]. We also identified related individuals (i.e., third-degree relatives or closer) via genotyping and omitted all but one individual from each related group in the analysis. Details regarding this cohort refinement have been described and implemented previously[67,68]. This refinement resulted in a cohort of 337,535 unrelated White British individuals from the UKBB, of which 29,661 have both genetic and cardiac MRI data. We randomly split this data into training, validation, and test sets of size 15,000, 5,000, and 9,661 individuals, respectively.

Genotyping and quality control

For the study cohort of 29,661 individuals described above, we leveraged genotype data from approximately 15 million imputed autosomal SNVs. These have been imputed from 805,426 directly assayed SNVs (obtained by the UKBB from one of two similar Affymetrix arrays) using the Haplotype Reference Consortium and UK10K reference panels[67]. Imputed variants were subject to several quality-control filters, including outlier-based filtration on effects due to batch, plate, sex, array, and discordance across control replicates. Further, we excluded variants due to extreme heterozygosity, missingness, minor allele frequency ($< 10^{-4}$), Hardy-Weinberg equilibrium ($< 10^{-10}$), and poor imputation quality (< 0.9). Further details can be found in previous studies[67,68].

Quantification of left ventricular hypertrophy

We retrieved cardiac MRI images from 44,503 UKBB participants, taken during their most recent imaging visit, and closely followed the method previously described by Bai et al.[25]. A fully convolutional network[25] was previously trained using a dataset of 4,875 subjects with 93,500 pixelwise segmentations of UKBB short-axis cardiac MRI multi-slice images generated manually with quality control checks for inter-operator consistency[69]. The network architecture employed multiple convolutional layers to learn image features across five resolution scales. Each scale involved two or three convolutions followed by batch normalization and ReLU transformation. Feature maps from the five scales were upsampled back to the original resolution, combined into a multi-scale feature map, and processed through three additional convolutional layers, followed by a softmax function to predict the segmentation label for each pixel. This approach generates a pixelwise segmentation map for each MRI slice which was previously determined to be at a level of accuracy similar to human experts[25]. We applied this trained deep learning model to our entire dataset of 44,503 cardiac MRIs. This resulted in segmentations of the LV cavity and myocardium from each short axis frame, which allowed for both an area calculation of each segment as well as the application of quality control checks[25] based on consistency within and between slices and time steps. There were 44,219 segmentations that passed the quality control. Using the calculated areas, we computed the volume of the LV myocardium through simple integration over slices. This volume was then converted to a left ventricular mass (LVM) using a standard density estimate of 1.05 g/mL[32]. LVMi was computed by dividing LVM by an estimate of body surface area based on height and body weight calculated using the Du Bois formula[33]. From the 44,219 segmentations, we restricted the analysis to LVMi measurements for 29,661 unrelated White British individuals using the measurements from their most recent imaging visit if multiple imaging visits were recorded.

Lo-siRF step 1: Dimension reduction of variants via genome-wide association studies

As the first step in the lo-siRF pipeline, we performed a genome-wide association study (GWAS) on the training data for the rank-based inverse normal-transformed LVMi using two algorithms, PLINK[34] and BOLT-LMM[35], in order to filter the number of features from over 15 million SNVs to a more

computationally-feasible size (Fig. 2b). This step is akin to typical screening phases in fine-mapping[70]

and other tree-based epistasis detection methods[63,71]. Since BOLT-LMM and PLINK rely on different

statistical models, we chose to employ both implementations to mitigate the dependence of

downstream conclusions on this arbitrary choice. Specifically, for the first GWAS run, we fitted a linear

regression model, implemented via `glm` in PLINK[72]. For the second GWAS run, we used BOLT-

LMM[35], a fast Bayesian-based linear mixed model method. Each GWAS was adjusted for the first five

principal components of ancestry, gender, age, height, and body weight. We then ranked the SNVs by

significance (i.e., the GWAS p-value) for each GWAS run separately and took the union of the top 1000

SNVs (without clumping) from each of the two GWAS runs. This resulted in a set of 1405 GWAS-

filtered SNVs that were used in the remainder of the lo-siRF pipeline. Here, we chose to use the top

1000 SNVs per GWAS method (without clumping) as it yielded the highest validation prediction

accuracy compared to choosing other possible thresholds (500 and 2000 SNVs per GWAS with and

without clumping). Though the GWAS is not the focus of this work, we provide a summary of the PLINK

and BOLT-LMM GWAS results for completeness and for comparison in Extended Data 1. We also

provide a list of the 1405 GWAS-filtered SNVs in Extended Data 2. We note that these 1405 GWAS-

filtered SNVs strictly contain the SNVs that passed the genome-wide significance threshold (p = 5E-8).

Lo-siRF step 2: Binarization of the left ventricular mass phenotype

Next, we binarized the raw (continuous) LVMi phenotype into a low and a high LVMi group before fitting

signed iterative random forest (Fig. 2c). That is, for a given threshold x, we binned individuals within the

top and bottom x% of LVMi values into two classes with the high and low LVMi values, respectively,

while omitting the individuals in the middle quantile range. Due to the sex-specific biological variation of

LVMi (Supplementary Note 1), we performed this binarization for males and females separately. For

males, low and high LVMi was considered under 43.8-46.0 g/m^2 and above 55.4-58.5 g/m^2,

respectively, depending on our choice of binarization threshold (Supplementary Table 2). For females,

low and high LVMi was defined as under 35.1-36.8 g/m^2 and above 43.8-46.1 g/m^2, respectively,

depending on our choice of binarization threshold. We performed this binarization step in order to

simplify the original low-signal regression problem into a relatively easier classification task: to predict

whether an individual has a very high or very low LVMi, in essence, denoising the LVMi signal in the

data. Moreover, as we will see in the next section, this binarization helped us more readily interpret and

assess the performance of prediction methods with respect to the prediction screening step of the PCS

framework[23]. Since the specific threshold choice is arbitrary, we ran the remainder of the lo-siRF

pipeline using three different binarization thresholds (15%, 20%, 25%) that balance the amount of

denoising and data lost. In the end, we aggregated the results that were stable across all three

binarization thresholds, described in the Method section *Lo-siRF step 4.4: Ranking genetic loci and*

interactions between loci.

Lo-siRF step 3: Prediction

Lo-siRF step 3.1: Fitting signed iterative random forest on the binarized LV mass index phenotype

For each binarization threshold, we trained a signed iterative random forest (siRF) model[20] using the

1405 GWAS-filtered SNVs to predict the binarized LVMi phenotype and generate candidate interactions

for further investigation (Fig. 2d). siRF first iteratively grows a sequence of feature-weighted random

forests, re-weighting features in each iteration proportional to their feature importance from the previous

iteration in order to stabilize the decision paths. Then, provided that the resulting stabilized forest

provides reasonable prediction performance (see the Methods section *Lo-siRF step 3.2: Prediction*

check), siRF leverages a computationally-efficient algorithm, random intersection trees[73], to identify

nonlinear higher-order interaction candidates based on frequently co-occurring features on a decision

path. Intuitively, sets of features that frequently co-occur along a decision path together are more likely

to interact and are identified by siRF. siRF is particularly attractive for prioritizing epistatic interactions

as (1) it offers an interaction search engine that can automatically search for higher-order interactions

with the same order of computational cost as a traditional random forest, and (2) the thresholding

behavior of its decision trees resembles the thresholding (or switch-like) behavior commonly observed

in biomolecular interactions[59]. Further, siRF improves upon its predecessor, iterative random forests[19],

by not only tracking which sets of features commonly co-occur on decision paths, but also the sign of

the features, i.e., whether low values (denoted X^-) or high values (denoted X^+) of feature X, appear on

the decision path. We refer to Kumbier et al.[20] for details, but in brief, the signed feature X^- (or

respectively, X^+) signifies that a decision rule of the form $X < t$ (or respectively, $X > t$) for some threshold

t appeared on the decision path. siRF hence outputs a list of candidate *signed* interactions, where each

signed interaction consists of two or more signed features that frequently co-occur on the same

decision path. Note when applying siRF to SNV data in practice, the signed feature SNV^+ typically

represents a heterozygous or homozygous mutation while the signed feature SNV^- typically represents

no mutation at the locus. The following hyperparameters were used to train siRF using the iRF2.0 R

package: number of iterations = 3, number of trees = 500, number of bootstrap replicates = 50, depth of

random intersection tree (RIT) = 3, number of RIT = 500, number of children in RIT = 5, and minimum

node size in RIT = 1. We fit siRF using 10,000 training samples (randomly sampled out of the 15,000

total training samples) and reserved the remaining 5,000 training samples for selecting genetic loci for

the permutation test (see the Method section *Lo-siRF step 4.3: Permutation test for difference in local*

stability importance scores).

Lo-siRF step 3.2: Prediction check

Per the PCS framework for veridical data science[23], we next assessed the validation prediction

accuracy of siRF (Fig. 2d) to evaluate whether the learnt model is capturing some biologically-relevant

phenotypic signal, rather than simply noise, before proceeding to interpret this model in step 4 of lo-

siRF. To serve as baseline comparisons, we fit other popular prediction methods, namely, L_1-

regularized (LASSO) logistic regression[74], L_2-regularized (ridge) logistic regression[75], random forests[76],

and support vector machines[77], and we evaluated prediction performance according to multiple metrics,

namely, classification accuracy, area under the receiver operator curve, and area under the precision-

recall curve. We observed that the prediction power of siRF, though weak (~55% balanced

classification accuracy), was greater than these other commonly used prediction methods across all

binarization thresholds and evaluation metrics, except for the 15% binarization threshold where siRF

performed second-best with respect to classification accuracy (Supplementary Table 3). Since siRF

performed better than random guessing (i.e., >50% balanced classification accuracy, which is not

guaranteed given the high phenotypic diversity of the LVMi trait) and demonstrated higher prediction

power than alternative popular prediction methods, we deemed that the siRF fit for LVMi passed the

prediction screening step of the PCS framework. Hence, we proceeded to interpret this siRF model and

prioritize candidate interactions in step 4 of lo-siRF. We note also that this prediction check played a

key role in our choice of phenotypic data. Prior to studying LVMi, we attempted to run a similar analysis

to predict hypertrophic cardiomyopathy (HCM) diagnosis, defined as any ICD10 billing code diagnosis

of I42.1 or I42.2 in the UKBB data. However, neither siRF nor the other aforementioned prediction

methods passed the 50% balanced classification accuracy requirement for predicting HCM diagnosis.

We thus chose not to proceed with the HCM analysis given the poor prediction accuracy and uncertain

relevance between the prediction models and the underlying biological processes. This failed prediction

check motivated the need for a more refined phenotypic measure of cardiac hypertrophy, which

ultimately led to the deep learning extraction of cardiac MRI-derived LVMi. Further discussion of the

HCM analysis can be found in Supplementary Note 1.

Lo-siRF step 4: Prioritization

To lastly interpret the siRF fit for LVMi, we developed a novel stability-driven importance score to

prioritize genetic loci and more interestingly, interactions between loci for follow-up experimental

validation (Fig. 2e). The assessment of importance at the level of genetic loci, rather than for individual

variants, is necessary since variant-level importances here are incredibly unstable (detailed in

Supplementary Note 1). This is due to the high correlation between SNVs in LD and the weak

phenotypic signal. Consequently, our new importance score aims to aggregate weak, unstable variant-

level importances into stronger, more stable locus-level importances by (1) assigning each variant to a

genetic locus, (2) evaluating the local (per-individual) importance of each genetic loci or interaction

between loci from the siRF fit via a stability-driven measure, and (3) conducting a permutation test to

summarize the importance of the genetic locus or interaction between loci across all individuals. We

provide details for each step next.

Lo-siRF step 4.1: Aggregation of SNVs into loci

We aggregated SNVs into a genetic locus based on genomic proximity. Specifically, we used

ANNOVAR[78] to assign each SNV that appears in the siRF fit to a genetic locus according to the hg19

refSeq Gene annotations (i.e., given by the `Gene.refGene` column in the ANNOVAR output).

ANNOVAR uses a default of 1 kb as the maximum distance between SNVs and gene boundaries. Note

that from these annotations, each SNV is assigned to exactly one genetic locus. Thus, herein in the

context of lo-siRF, a genetic locus is a (non-overlapping) group of SNVs, and a signed genetic locus is

a (non-overlapping) group of signed SNVs with the specified sign (i.e., *Locus⁺* consists of SNV_1^+, ...,

SNV_p^+ while *Locus⁻* consists of SNV_1^-, ..., SNV_p^-).

Lo-siRF step 4.2: Local stability importance score

We next measured the importance of a genetic locus or interaction between loci based on their stability,

or frequency of occurrence, within the siRF fit. However, because the number of variants assigned to

each genetic locus can vary, the raw frequency of occurrence will be biased towards larger loci (i.e.,

those with more variants). A more detailed discussion is provided in Supplementary Note 1. To address

this issue, we developed a *local* (or per-individual) *stability importance score*, which quantifies the

importance of a signed locus or interaction between loci for making the prediction for each individual.

Let $G = \{g_1, \ldots, g_K\}$ denote a signed order-K interaction involving the signed genetic loci $g_1, \ldots, g_K$, and

let $v_1^{(j)}, \ldots, v_{p_j}^{(j)}$ denote the signed SNVs belonging to the signed genetic locus g_j. Then given a forest

T, a signed interaction between loci G, and individual i, the *local stability importance score*, $LSI_T(G, i)$, is

defined as $D_T(G, i) / |T|$, where $|T|$ is the number of trees in the forest T, and $D_T(G, i)$ is the number of

decision paths in the forest T for which two criteria are satisfied: (1) individual i appears in its terminal

node and (2) for each $j = 1, \ldots, K$, there exists an $l \in \{1, \ldots, p_j\}$ such that $v_l^{(j)}$ was used in a decision

split along the path (Supplementary Data Fig. 3a). In other words, $LSI_T(G, i)$ is the proportion of trees in

the forest T for which at least one signed variant from each signed locus in the signed interaction G was

used in making the prediction for individual i. A higher score indicates greater importance of the signed

interaction G for individual i's prediction. Note that a genetic locus can be viewed as an order-1

interaction, and thus, this local stability importance score can also be computed to assess the

(marginal) importance of a single genetic locus.

Lo-siRF step 4.3: Permutation test for difference in local stability importance scores

Once we obtained these local stability importance scores for each individual, we performed a two-

sample permutation test to assess whether the local stability importance scores for a given signed

locus or interaction between loci, G, are different between individuals with high and low LVMi

(conditional on the rest of the fitted forest). More formally, the proposed permutation test tests the null

hypothesis $L = H$ versus the alternative hypothesis $L \neq H$, where L and H are the distributions of local

stability importance scores for individuals with low and high LVMi, respectively. If the local stability

importance scores are indeed different between high and low LVMi individuals, the permutation test

results in a small p-value, indicating that G can differentiate between individuals with high versus low

LVMi and hence is an important locus or interaction between loci for LVMi. We performed this

permutation test using 10,000 permutations, the difference in means as the test statistic, and the 5,000

validation samples. To bolster the reliability of our findings, we only tested a conservative subset of

genetic loci and interactions between loci that passed predictive and stability checks in accordance with

the PCS framework. Namely, we tested:

(a) The top 25 genetic loci, ranked by their average local stability importance scores across 5,000

samples. These 5,000 samples were previously set-aside from within the 15,000 training

samples and were not used in fitting the siRF (see the Methods section *Lo-siRF step 3.1: Fitting*

signed iterative random forest on the binarized LV mass index phenotype);

(b) The signed interactions between loci that were stably identified by siRF across 50 bootstrap

replicates. Here, we performed the random intersection trees search within siRF at the locus-

level (i.e., using the variant-to-locus assignment as the hyper-features or `varnames.grp`

argument when running siRF in R), and we defined a "stable" interaction as one that passed the

following siRF stability metric thresholds: stability score > 0.5, stability score for mean increase

in precision > 0, and stability score for independence of feature selection > 0[19,20]

(Supplementary Table 5). Details on the siRF interaction and stability metrics can be found in

previous work[20].

Lo-siRF step 4.4: Ranking genetic loci and interactions between loci

To select the top lo-siRF recommendations for follow-up experimental validation, we incorporated one

final stability check, recommending only those signed loci and interactions between loci that underwent

the permutation test and yielded a *p*-value < 0.1 in all three binarization runs. For these signed loci and

interactions between loci that were stably important in all three binarization runs, we ranked them by

the mean permutation *p*-value, averaged across the three binarization thresholds (Supplementary

Table 4). Because of our emphasis on prioritizing candidates for experimental validation, if both the +

and - version of the signed locus (or interaction) appear, the final prioritized loci (or interaction) are

ranked according to the smaller one of the two *p*-values (Fig. 2f). We note that though the signed

information is not pertinent to our goal of recommending candidates for experiments, the signed

information from siRF provides more granular information that can improve our interpretation of the fit,

and we discuss this further in Supplementary Note 1. We also provide the permutation *p*-values for all

conducted permutation tests (including the loci and interactions between loci that were unstable across

binarization thresholds) in Supplementary Note 1.

lo-siRF: PCS documentation and additional stability analyses

In an effort to facilitate transparency of human judgment calls that are inevitably made here and

throughout our veridical machine learning pipeline, we provide extensive documentation and discussion

of these human judgment calls in Supplementary Note 1. In particular, Supplementary Note 1 includes

a discussion of our reasoning and motivation behind the choice of phenotypic data as well as choices in

the dimension reduction, binarization, prediction, and prioritization steps. We also performed additional

stability analyses in accordance with the PCS framework[23], detailed in Supplementary Note 1, to

ensure that our findings are stable and robust to these human judgment calls (e.g., the choice of GWAS

method and binarization threshold) and to bolster the reproducibility of our findings. Supplementary

Note 1 is an HTML document, which can be downloaded and displayed in a browser or found at

https://yu-group.github.io/epistasis-cardiac-hypertrophy/.

Functional interpretation of lo-siRF-prioritized variants

Functional interpretation step 1: Extraction of candidate SNVs and LD structures

Our lo-siRF approach described above identified a total of 283 SNVs located within 6 LVMi genetic risk

loci (Fig. 2f). In order to explore the functional consequences of these prioritized genetic variants and

identify genes that are potentially involved in the trait of LV hypertrophy, we performed functional

mapping and annotation using a web-based platform, FUMA (v1.5.4)[37]. The SNP2GENE function in

FUMA was used to incorporate LD structure and prioritize candidate genes. Taking the GWAS

summary statistics from PLINK[34] and BOLT-LMM[35] as an input, we submitted the 283 lo-siRF-

prioritized SNVs into SNP2GENE as predefined SNVs. This allows SNP2GENE to define LD blocks for

each of the 283 lo-siRF-prioritized SNV and use the given 283 SNVs and SNVs in LD with them for

further annotations. We adopted the default r^2 threshold (i.e., 0.6) for defining independent significant

SNVs. Because any two of the 283 lo-siRF-prioritized SNVs are in LD with each other at $r^2 < 0.6$, all of

the 283 SNVs were defined as independent significant SNVs by FUMA. In order to match the

population group used for our lo-siRF prioritization, the reference panel from UKBB release 2b that

FUMA created for British and European subjects was chosen for the computation of r^2 and minor allele

frequencies. A total of 572 candidate SNVs in strong LD ($r^2 < 0.6$) with any of the 283 independent

significant SNVs were extracted from both the inputted GWAS (with the maximum p-value threshold

being 0.05) and the reference panel. These 572 candidate SNVs were then assigned to one of the six

lo-siRF-identified locus (Fig. 2f) based on its corresponding independent significant SNV, which showed

the maximum r^2 value in LD with the given candidate SNV. A combination of the 283 independent

significant SNVs and the 572 FUMA-extracted candidate SNVs in LD with the independent significant

SNVs (details in Extended Data 4) was defined as the lo-siRF-prioritized SNV set, which was used to

generate the list of lo-siRF-prioritized genes (Fig. 4a) for the following enrichment analysis (Fig. 4c-4e).

As a comparison to the lo-siRF-prioritized SNV set, we uploaded all 1405 GWAS-filtered SNVs

(Extended Data 2) as the predefined SNVs in a separate SNP2GENE job. Using the same approach

and parameter settings, 929 independent significant SNVs were identified within the given 1405

GWAS-filtered SNVs, and 5771 candidate SNVs in LD with the 929 independent significant SNVs were

extracted by FUMA. A combination of the 929 independent significant SNVs and the 5771 candidate

SNVs were defined as the reference SNV set. This reference SNV set is purely generated from GWAS

prioritization and excludes the evaluation of epistatic effects between genetic variants by lo-siRF.

Genes functionally mapped from the reference SNV set was used as a comparison group for the lo-

siRF-prioritized gene list in the following enrichment analysis to explore the specific contribution of the

identified epistatic genes in the enriched gene ontologies, pathways, and transcription factors (Fig. 4c-

4e).

Functional interpretation step 2: ANNOVAR enrichment test

To evaluate the functional consequences of the lo-siRF-prioritized genetic loci, we performed

ANNOVAR enrichment test of the aforementioned 283 independent significant SNVs and 572

candidate SNVs in LD with them against the selected reference panel in FUMA. The FUMA

SNP2GENE process generated unique ANNOVAR[78] annotations for all the identified SNVs. The

enrichment score for a given annotation in a given lo-siRF-prioritized genetic locus (Fig. 3b) was

computed as the proportion of SNVs associated to that locus with the given annotation divided by the

proportion of SNVs with the same annotation relative to all available SNVs in the reference panel. For

the i^{th} ANNOVAR annotation in the j^{th} lo-siRF-prioritized locus, the enrichment *p*-value was computed

by performing a two-sided Fisher's exact test on the 2-by-2 contingency table containing $n_j(i)$,

$\sum_t n_j(t) - n_j(i)$, $\quad N(i) - n_j(i)$, and $\sum_t N(t) - \sum_t n_j(t) - n_j(i)$. Here, $n_j(i)$ is the number of SNVs

with the i^{th} annotation in the j^{th} lo-siRF-prioritized locus, $N(i)$ is the number of SNVs with the i^{th}

annotation in the reference panel, $\sum_t n_j(t)$ is the summation of $n_j(i)$ for all available annotations in

the j^{th} lo-siRF-prioritized locus, and $\sum_{t} N(t)$ is the summation of $N(i)$ for all available annotations in

the reference panel. Detailed information can be found in Extended Data 5.

Functional interpretation step 3: Functional annotations

In addition to ANNOVAR annotations, FUMA annotated all 283 independent significant SNVs and 572

SNVs in LD with them for functional consequences on potential regulatory functions (core-15 chromatin

state prediction and RegulomeDB score) and deleterious effects (CADD score). In particular, the core-

15 chromatin state was annotated to all SNVs of interest by ChromHMM[38] derived from 5 chromatin

markers (H3K4me3, H3K4me1, H3K36me3, H3K27me3, and H3K9me3) for 127 tissue/cell types, of

which left ventricle (E095), right ventricle (E105), right atrium (E104), and fetal heart (E083) were taken

into consideration in this study (Fig. 3a, circle 7). Data and a corresponding description of the core-15

chromatin state model can be found at

https://egg2.wustl.edu/roadmap/web_portal/chr_state_learning.html. RegulomeDB[37,39] annotations

guide interpretation of regulatory variants through a seven-level categorical score, of which the

category 1 (including 6 subcategories ranging from 1a to 1f) indicates the strongest evidence for a

variant to result in a functional consequence. Because the RegulomeDB database (v1.1) used in FUMA

has not been updated, we queried all SNVs identified by lo-siRF and FUMA in the RegulomeDB

database v2.2 (https://regulomedb.org/regulome-search). Annotations for deleteriousness were

obtained from the CADD database (v1.4)[40] by matching chromosome, position, reference, and

alternative alleles of all SNVs. High CADD scores indicate highly deleterious effects of a given variant.

A minimum threshold CADD score of 12.37 was suggested by Kircher et al.[40]. In addition to the

aforementioned functional annotations, we extracted information of eQTLs and sQTLs for all

independent significant SNVs and SNVs that are in LD with one of the independent significant SNVs

from GTEx v8[41]. The eQTL information was used for eQTL gene mapping as described in the following

section.

Functional interpretation step 4: Functional gene mapping

In SNP2GENE, we performed three functional gene mapping strategies – positional, eQTL, and 3D

chromatin interaction mapping – using the lo-siRF-prioritized SNV set and the reference SNV set

described in the Methods section *Functional interpretation step 1: Extraction of candidate SNVs and LD*

structures. For positional mapping[37,39], a default value of 10 kb was used as the maximum distance

between SNVs and genes. For eQTL mapping, cis-eQTL information of heart left ventricle, heart atrial

appendage, and muscle skeletal tissue types from GTEx v8[41] was used. Only significant SNV-gene

pairs (FDR < 0.05 and p < 1E-3) were used for eQTL mapping. For 3D chromatin interaction mapping,

Hi-C data of left ventricle tissue from GSE87112 was chosen with a default threshold of FDR < 1E-6. A

default promoter region window was defined as 250 bp upstream and 500 bp downstream of TSS[37,39].

Using these three gene mapping strategies, we mapped the lo-siRF-prioritized SNV set to 21 protein-

coding genes (Fig. 4a), of which 20 are HGNC-recognizable. Each of the 21 genes was also

functionally linked to a specific lo-siRF-prioritized LV hypertrophy risk locus (Fig. 2f), to which the

highest proportion of SNVs mapped to the given gene were assigned. A Circos plot (Fig. 3a) showing

comprehensive information of the lo-siRF-prioritized epistatic interactions, FUMA-prioritized eQTL SNV-

to-gene connections and 3D chromatin interactions, as well as LD structures and prioritized genes was

created by TBtools[79]. We then submitted these 21 genes to the GENE2FUNC process in FUMA and

obtained GTEx gene expression data for 19 (out of the 21) genes across multiple tissue types (Fig. 4b).

In addition, we used the same approach and mapped the reference SNV set (mentioned in the Methods

section *Functional interpretation step 1: Extraction of candidate SNVs and LD structures*) to a separate

gene set that contains 382 HGNC-approved genes. The lo-siRF-prioritized gene set and the reference

gene set were used for gene set enrichment analysis that are described in the following sections.

Gene ontology and pathway enrichment analysis

Genes that co-associate to shared gene ontology (GO) and pathway terms are likely to be functionally

related. To assess the differential GO and pathway co-association among the lo-siRF-prioritized genes

relative to their counterparts that were deprioritized by lo-siRF, we performed an integrative GO and

pathway enrichment analysis followed by an exhaustive permutation of co-association scores between

any possible gene-gene combinations found in the aforementioned 382 HGNC-approved genes (see

the Methods section *Functional interpretation step 4: Functional gene mapping*).

In order to improve GO and pathway prioritization, we adopted the concept from Enrichr-KG[48] and

ChEA3[47] to assess enrichment analysis results across libraries and domains of knowledge as an

integrated network of genes and their annotations. We first queried the 382 HGNC-approved genes

from the reference gene set against various prior-knowledge gene set libraries in Enrichr[46]

(https://maayanlab.cloud/Enrichr/). We selected five representative libraries from the GO and pathway

Enrichr categories as follows: GO biological process[80,81], GO molecular function[80,81], MGI Mammalian

Phenotypes[82], Reactome pathways[83], and KEGG pathways[84]. Other FUMA-extracted genes that were

not approved by HGNC using synonyms or aliases were discarded. This enrichment analysis allowed

us to search for a union of enriched GO or pathway terms and their correspondingly annotated gene

sets, from which we built a co-association network. According to the method by Enrichr-KG[48], nodes in

the co-association network are either the enriched GO and pathway terms or genes.

To measure the degree of co-association to specific GO and pathway terms for two given interactor

genes, we computed a co-association score for each of the 72,771 possible gene-gene combinations

(from the 382 queried genes). The co-association score was calculated by $R = N_{(A \cap B)} \big/ N_{(A \cup B)}$. Here,

$N_{(A \cap B)}$ denotes the number of GO or pathway terms that were significantly enriched for both *gene A* and

gene B in the proposed gene-gene combination, and $N_{(A \cup B)}$ is the number of GO or pathway terms that

were enriched for either *gene A* or *gene B*. For cases where $N_{(A \cup B)} = 0$, we defined $R = 0$ to indicate that

no GO or pathway terms were found to be co-associated with the respective gene-gene combinations.

Of the 382 HGNC-approved genes, 20 genes were mapped to lo-siRF-prioritized loci by FUMA

functional gene mapping (one of the 21 lo-siRF-prioritized genes is HGNC-unrecognizable and is

discarded). We compared the co-association scores R for gene-gene combinations in the lo-siRF-

prioritized gene set relative to the full distribution of R provided by an exhaustive permutation of all

possible gene-gene combinations in the set of 382 HGNC-approved genes. The ranking of gene-gene

combinations was determined by the two-sided empirical *p*-values. Fig. 4c displays significant co-

associations (empirical $p < 0.05$) between enriched GO or pathway terms and genes functionally

mapped to lo-siRF-prioritized epistatic and hypostatic loci (Fig. 2f). Further details can be found in

Extended Data 6.

Transcription factor enrichment analysis

Owing to the limitations and biases of various specific assays, we performed an integrative transcription

factor (TF) enrichment analysis against multiple annotated gene set libraries in ChEA3[47] and Enrichr[46].

To preserve the variety of library types, we assembled 9 gene set libraries (Fig. 4d) from distinct

sources as follows:

1. Putative TF target gene sets determined by ChIP-seq experiments from ENCODE[85];

2. Putative TF target gene sets determined by ChIP-seq experiments from ReMap[86];

3. Putative TF target gene sets determined by ChIP-seq experiments from individual

publications[47];

4. TF coexpression with other genes based on RNA-seq data from GTEx[41];

5. TF coexpression with other genes based on RNA-seq data from ARCHS4[87];

6. Single TF perturbations followed by gene differential expression[46];

7. Putative target gene sets determined by scanning PWMs from JASPAR[88] and TRANSFAC[89] at

promoter regions of all human genes;

8. Gene sets predicted by transcriptional regulatory relationships unraveled by sentence-based

text-mining (TRRUST)[90];

9. Top co-occurring genes with TFs in a large number of Enrichr queries[47].

Of the mentioned 9 gene set libraries, libraries 1, 3, and 5 were assembled by combining gene set

libraries downloaded from both ChEA3[47] and Enrichr[46]. Libraries 2, 4, and 9 were downloaded from

ChEA3[47]. Libraries 6, 7, and 8 were downloaded from Enrichr[46]. According to the integration method by

ChEA3[47], for libraries in which multiple gene sets were annotated to the same TF, the unique gene set

with the lowest FET p-value was used. As mentioned in previous sections, we used separate FUMA

SNP2GENE processes and functionally mapped lo-siRF-prioritized SNVs and all GWAS-filtered SNVs

to a lo-siRF-prioritized gene set (20 HGNC-recognizable genes) and a reference gene set (382 HGNC-

recognizable genes), respectively. Because the lo-siRF-prioritized gene set is a subset of the reference

gene set, we considered the 362 genes complementary to the lo-siRF-prioritized gene set as the lo-

siRF-deprioritized gene set. Taking the 20 lo-siRF-prioritized genes and 263 lo-siRF-deprioritized genes

as two separate input gene sets, we performed enrichment analysis against the 9 gene set libraries.

For each of the 9 libraries, we ranked the significance of overlap between the input gene set and the

TF-annotated gene sets in that library by FET p-values. Those TFs with identical FET p-values were

ranked by the same integer number. A scaled rank was then assigned to each TF by dividing the

corresponding integer rank by the maximum integer rank in its respective library. We then integrated

the 9 sets of TF rankings and re-ordered the TFs by two sequential criteria: (1) the number of libraries

that display a significant overlap with the input gene set (FET $p < 0.05$) and (2) the mean scaled rank

across all libraries containing that TF. Using this method, we prioritized two distinct sets of TFs for the

lo-siRF-prioritized genes (Fig. 4d, top) and lo-siRF-deprioritized genes (Fig. 4d, bottom).

We used the same approach described in the Methods section *Gene ontology and pathway enrichment*

analysis to evaluate the differential TF co-association between the lo-siRF-prioritized genes relative to

the lo-siRF-deprioritized genes. A TF co-association score was computed for each of the 72,771

possible gene-gene combinations from the 382 genes. Still, the TF co-association score was computed

by $R = N_{(A \cap B)} \big/ N_{(A \cup B)}$, except that $N_{(A \cap B)}$ and $N_{(A \cup B)}$ denote the number of enriched TF terms instead of

GO or pathway terms. Pairwise interactions between lo-siRF-prioritized genes were extracted and

ranked by the empirical p-values (Fig. 4e) from an exhaustive permutation of TF co-association scores

for the 72,771 possible gene-gene combinations. Details regarding TF enrichment from both lo-siRF-

prioritized and lo-siRF-deprioritized genes and TF co-association strengths can be found in Extended

Data 7.

Disease-state-specific gene co-expression network analysis

In order to evaluate the connectivity between genes and their potential roles in the transition from

healthy to failing myocardium, we compared gene-gene connectivity and changes in the topological

structure between gene co-expression networks for healthy and failing human heart tissues (Fig. 5). To

construct gene co-expression networks, cardiac tissue samples from 177 failing hearts and 136 donor,

non-failing (control) hearts were collected from operating rooms and remote locations for RNA

expression measurements. We performed weighted gene co-expression network analysis (WGCNA) on

the covariate-corrected RNA microarray data for the control and heart failure networks separately (Fig.

5a). Detailed steps for generating these co-expression networks, which included calculating the

correlation matrix, TOM transformation, and Dynamic Tree Cut module finding, are described in our

previous study[51], and data for these networks is available at https://doi.org/10.5281/zenodo.2600420.

To evaluate the degree of connectivity between interacting genes in each of the networks, we

compared the edge weights between the interacting genes demonstrated in this study relative to the

distribution of all possible pairwise combinations of genes (Fig. 5b and 5c). We also evaluated the

difference of edge weights (Z-score normalized) between the control and heart failure networks to

understand how these gene-gene interactions change between non-failing and failing hearts (Fig. 5d).

The two-tailed empirical *p*-value represents the proportion of the absolute difference in edge weights of

all gene pairs that exceed the absolute difference score for gene pairs of interest. We then compared

the structure of modules derived from dendrograms on the WGCNA control and heart failure networks

(Fig. 5e and 5f). Modules were labeled according to Reactome Enrichment analysis of genes within

each module. The full gene module descriptions and Benjamini-Hochberg-adjusted enrichment *p*-

values can be found in the Supplementary Data 5 and 6 in the study by Cordero et al.[51].

Induced pluripotent stem cell cardiomyocytes differentiation

The studied patient-specific human induced pluripotent stem cells (hiPSCs) were derived from a 45-

year-old female proband with a heterozygous *MYH7*-R403Q mutation. Derivation and maintenance of

hiPSC lines were performed following Dainis et al.[30]. Briefly, hiPSCs were maintained in MTeSR

(StemCell Technologies) and split at a low density (1:12) onto fresh 1:200 matrigel-coated 12 well

plates. Following the split, cells were left in MTeSR media supplemented with 1 μM Thiazovivin. The

hiPSCs were maintained in MTeSR until cells reached 90% confluency, which began Day 0 of the

cardiomyocyte differentiation protocol. Cardiomyocytes were differentiated from hiPSCs using small

molecule inhibitors. For Days 0-5, cells were given RPMI 1640 medium + L-glutamine and B27 - insulin.

On Days 0 and 1, the media was supplemented with 6 µM of the GSK3β inhibitor, CHIR99021. On

Days 2 and 3, the media was supplemented with 5 µM of the Wnt inhibitor, IWR-1. Media was switched

to RPMI 1640 medium + L-glutamine and B27 + insulin on Days 6-8. On Days 9-12, cells were

maintained in RPMI 1640 medium + L-glutamine - glucose, B27 + insulin, and sodium lactate. On Day

13, cells were detached using Accutase for 7-10 minutes at 37 ℃ and resuspended in neutralizing

RPMI 1640 medium + L-glutamine and B27 + insulin. This mixture was centrifuged for 5 minutes at

1000 rpm (103 rcf). The cell pellet was resuspended in 1 µM thiazovivin supplemented RPMI 1640

medium + L-glutamine and B27 + insulin. For the rest of the protocol (Days 14-40), cells were exposed

to RPMI 1640 medium + L-glutamine - glucose, B27 + insulin, and sodium lactate. Media changes

occurred every other day on Days 14-19 and every three days for Days 20-40. On Day 40,

cardiomyocytes reached maturity.

RNA silencing in induced pluripotent stem cell-derived cardiomyocytes

Mature hiPSC-derived cardiomyocytes were transfected with Silencer Select siRNAs (Thermofisher)

using TransIT-TKO Transfection reagent (Mirus Bio). Cells were incubated for 48 hours with 75 nM

siRNA treatments. Four wells of cells were transfected with each of the six siRNAs: scramble,

CCDC141 (ID s49797), *IGF1R* (ID s223918), *TTN* (ID s14484), *CCDC141* and *IGF1R*, and *CCDC141*

and *TTN*. After 2 days, hiPSC-CMs were collected for RNA extraction.

RT-qPCR analysis for siRNA gene silencing efficiency

Following cell morphology measurement, all cells for each condition were centrifuged for 5 minutes at

1000 rpm (103 rcf). Cell pellets were frozen at -80 ℃ prior to RNA extraction. RNA was extracted using

Trizol reagent for RT-qPCR to confirm gene knockdown occurred. Reverse Transcription of RNA was

done using High-Capacity cDNA Reverse Transcription Kit (Thermofisher). qPCR of the single stranded

cDNA was performed using TaqMan Fast Advanced MM (Thermofisher) with the following annealing

temperatures: 95°C 20" and 40 cycles of 95°C 1" and 60°C 20". qPCR of the silenced genes was

performed using TaqMan® Gene Expression Assays, including *CCDC141* (Hs00892642_m1), *IGF1R*

(Hs00609566_m1), and *TTN* (Hs00399225_m1). For gene silencing efficiency analysis, gene RPLP0

(Hs00420895_gH) was used as a reference gene. Data were analyzed using the delta-delta Ct method.

Cell sample preparation for cell morphology measurement

Following siRNA treatments, cells were detached for microfluidic single cell imaging using a mixture of

5 parts Accutase and 1 part TrypLE, treated for 6 minutes at 37 ℃. Cells were then added to the

neutralizing RPMI 1640 medium + L-glutamine and B27 + insulin. These mixtures were centrifuged for

5 minutes at 1000 rpm (103 rcf). For each gene silencing condition, the four wells of cells were

resuspended in 4 mL of the MEM medium, which is composed of MEM (HBSS balanced) medium, 10%

FBS, and 1% Pen Strep (Gibco). Cells were filtered with 100 µm strainers (Corning) before adding into

the microfluidic devices.

Microfluidic inertial focusing device

We developed a new spiral inertial microfluidics system on the basis of the study by Guan et al.[28] to

focus randomly suspended cells into separate single streams based on cell size for high-resolution and

high-throughput single cell imaging. The microfluidic device (Extended Data Fig. 6) contains 5 loops of

spiral microchannel with a radius increasing from 3.3 mm to 7.05 mm. The microchannel has a cross-

section with a slanted ceiling, resulting in 80 µm and 150 µm depths at the inner and outer side of the

channel, respectively. The channel width is fixed to 600 µm. The 495 µm wide slanted region of the

channel ceiling is composed of ten 7 µm deep stairs. This particular geometry induces strong Dean

vortices in the outer half of the channel cross-section, leading to high sensitivity of size separation and

cell focusing. The device has two inlets at the spiral center to introduce cell suspensions and sheath

flow of fresh medium. At the outlet region, the channel is expanded in width and split into two outlet

channels with a width of 845 µm for the top outlet and 690 µm for the bottom outlet. Depths of the two

outlet channels are designed to create equal hydraulic resistance. The top and bottom outlet channels

are connected to 80 µm and 50 µm deep straight observation channels for high-throughput cell

imaging.

Microdevice fabrication

The spiral microchannel was fabricated by CNC micromachining a piece of laser-cut poly (methyl

methacrylate) (PMMA) sheet, which was bonded with a PMMA chip machined only with the inlet

channels and another blank PMMA chip using a solvent-assisted thermal binding process to form the

enclosed channel[26]. Before bonding, PMMA chips were cleaned with acetone, methanol, isopropanol

and deionized water in sequence. Droplets of a solvent mixture (47.5% DMSO, 47.5% water, 5%

methanol) were evenly spread over the cleaned chips. The PMMA chips were assembled appropriately

and clamped using a customized aluminum fixture, and then heated in a ThermoScientific Lindberg

Blue M oven at 96 ℃ for 2 hrs. After bonding, fluid reservoirs (McMaster) were then attached to the

chips using a two-part epoxy (McMaster). Microchannels were flushed with 70% ethanol followed by DI

water for sterilization.

High-throughput single cell imaging

Before each experiment, microchannels were flushed with 3 mL of the MEM medium. Prepared cell

samples and fresh MEM medium were loaded into 3 mL syringes, which were connected to the

corresponding microchannel inlets using Tygon PVC tubing (McMaster). Both cells and the fresh MEM

medium were infused into the microchannel using a Pico Plus Elite syringe pump (Harvard Apparatus)

at 1.2 mL/min. Microscope image sequences of cells focused to the top and bottom observation

channels were captured using a VEO 710S high-speed camera (Phantom) with a sampling rate of 700

fps and a 5 µsec light exposure.

Image analysis for cell feature extraction

For each gene silencing condition of each biological repeat, 21,000 images were processed to extract

cell morphology features. To analyze cell size and shape changes induced by gene silencing, we

developed a MATLAB-based image analysis pipeline, which includes three major steps: image

preparation, feature extraction, and image post-processing (Fig. 6c). In step one, image sequences

were fed into the MATLAB program and subtracted from the corresponding background image to

correct any inhomogeneous illumination. The program automatically generates background images, in

which each pixel value is computed as the mode pixel intensity value among the same pixel of the

entire corresponding image sequence. After illumination correction, step two detects cell edges by

looking for the local maxima of the bright field intensity gradient, following which the program closes

edge gaps, removes cells connected to the image borders, cleans small features (noise), and then fills

holes to generate binary images and centroid positions for each single cell. Cell locations were then

traced and stuck cells were removed by a double-counting filter if present. The double-counting filter

excludes measurements collected around the same location with similar cell sizes using a Gaussian

kernel density method (bandwidth = 0.09) when the estimated density for a certain location and size

exceeds a particular threshold. The maximal density value for experimental runs where no repeated

measurements were observed was used as the threshold. This procedure was manually validated

using visual inspection of the removed cells. Binary images passing the double-counting filter were

used to create coordinates (X, Y) of cell outlines, which leads to a range of cell size and shape

parameters, including cell diameter and area, solidity, roundness error, circularity, and intensity spatial

relationship enclosed within the cell boundary. Cell area was computed as the 2D integration of the cell

outline, and the cell diameter was computed as $2\sqrt{Area/\pi}$. Solidity is the ratio of cell area to the area of

the smallest convex polygon that contains the cell region. Roundness error was computed as the ratio

between the standard deviation and mean of radii on the cell outline measured from the centroid.

Circularity was calculated as $4 Area\pi/Perimeter^2$. The 2D intensity distributions within cell outlines

were used to derive peak locations and count peak numbers, which is a measure of intensity spatial

relationship and a gating parameter to remove clumped cells. In the post-processing step, data were

cleaned using three filters with the following gating threshold. To remove large clumps, the peak-solidity

filter removes data outside of the polygonal region defined by {(0.9, 0), (0.9, 3.2), (0.934, 8.26), (1, 28),

(1, 0)} in the (solidity, peak No.) space. Then, the roundness filter removes cells with weird shapes by

excluding data with a roundness error higher than 0.3 or a circularity lower than 0.6. Finally, the small

size filter removes cell debris whose major diameter is lower than 15 μm (12 μm) or minor diameter is

lower than 12 μm (10 μm) for images photographed at the top (bottom) outlet microchannels.

Statistical analysis of experimental results

To analyze the experimental results, we compared differences in cell sizes and morphological features between cells with silenced genes or gene pairs and their scramble controls using various summary statistics. To study the centers of the distributions, we compared the medians using a traditional Wilcoxon signed rank test (Fig. 6f) and a bootstrap quantile test at the 0.5 quantile level. In accordance with the PCS framework, we used two different tests here to ensure that our findings are robust to this arbitrary modeling choice and that the underlying assumptions do not drive our results. We further performed a bootstrap-t for trimmed means of cell size distributions and evaluated the stability of results by varying the amount of trimming (ranging from 0-0.3) to verify that outliers (e.g., from large clumps of cells) do not drive the conclusions (data not shown). In addition to comparisons of central behavior, we performed the bootstrap quantile test across increasing quantile levels (ranging from 0.5-0.9) to study cell size changes (Extended Data Fig. 7) that favor hypertrophic cells, which is more relevant to the pathologic phenotype of cardiac hypertrophy. All tests are performed on each experimental batch, and the maximum *p*-value across batches are reported in the main text. To formally quantify whether interactions are non-additive, for each interaction (i.e., *CCDC141-TTN* and *CCDC141-IGF1R*), we tested whether the interaction effect is significant when regressing cell size on the interaction and main effects for each relevant gene. Again in accordance with the PCS framework, we performed a stability analysis by comparing results across different regression parameters – namely, quantile regression with increasing quantile levels (ranging from 0.5-0.9). In each regression, percentile bootstrap t-tests were used to compute *p*-values. Traditional t-test *p*-values supporting these results are also provided in Extended Data 8 as a stability check. Since these regressions require comparisons between gene silencing experiments (e.g., silencing *CCDC141* and *TTN* vs only silencing *CCDC141*), and gene silencing efficiency varied across experiments, batches were merged based on those with the highest efficiency for each regression.

Fig. 1: Schematic of the study workflow.

The study workflow includes four major stages: (a) derivation of left ventricular mass from cardiac magnetic resonance imaging (green boxes); (b) computational prioritization of epistatic drivers (orange); (c) functional interpretation of the hypothesized epistatic genetic loci (purple); and (d) experimental confirmation of epistasis in cardiac tissues and cells (blue). Abbreviations: MRI, magnetic resonance imaging; LV, left ventricle; LVM, left ventricular mass; LVMi, left ventricular mass indexed by body surface area; SNV, single-nucleotide variant; GWAS, genome-wide association study; BOLT-LMM[35] and PLINK[34], two different GWAS software packages; lo-siRF, low-signal signed iterative random forest; ANNOVAR[78], a software for functional annotation of genetic variants; CADD[40], combined annotation dependent depletion, which scores the deleteriousness of variants;

RegulomeDB[39], a database that scores functional regulatory variants; ChromHMM[38], a multivariate Hidden Markov Model for chromatin state annotation; eQTL, expression quantitative trait locus; sQTL, splicing quantitative trait locus; Hi-C, high-throughput chromosome conformation capture; PheWAS, phenome-wide association study; siRNA, small interfering RNA; hiPSC-CM, human induced pluripotent stem cell-derived cardiomyocyte; HCM, hypertrophic cardiomyopathy.

Fig. 2: Low-signal signed iterative random forest (lo-siRF) prioritizes risk loci and epistatic interactions for left ventricular hypertrophy.

a-e, Workflow of low-signal signed iterative random forest (lo-siRF). **a**, Lo-siRF took in as input single-nucleotide variant (SNV) data and cardiac MRI-derived left ventricular mass indexed by body surface area (LVMi) from 29,661 UK Biobank participants. **b**, Dimension reduction was performed via a genome-wide association study (GWAS) to concentrate the analysis on a smaller set of SNVs. **c**, LVMi was binarized into high and low LVMi categories according to three different binarization thresholds (represented by the stacked boxes). **d**, For each of the three binarization thresholds, a signed iterative random forest was fitted using the GWAS-filtered SNVs to predict the binarized LVMi phenotype. The validation prediction accuracy was assessed prior to interpreting the model fit. **e**, SNVs used in the fitted signed iterative random forest were aggregated into genetic loci based on annotations using

ANNOVAR[78]. Genetic loci and pairwise interactions between loci were finally ranked according to their importance across the three signed iterative random forest fits, as measured by our proposed stability-driven importance score. **f**, Lo-siRF-prioritized risk loci and epistatic interactions. (1) Loci stably prioritized by lo-siRF as epistasis participants are highlighted in green. (2) nIndSigSNVs, the number of independent significant SNVs that are stably prioritized by lo-siRF across the three different LVMi binarization thresholds (panel **c**). (3) nSNVs, the number of candidate SNVs extracted by FUMA[37] (v1.5.4) in strong LD ($r^2 > 0.6$) with any of the lo-siRF-prioritized independent significant SNVs. (4) Lo-siRF p-value, the mean p value from lo-siRF, averaged across the three LVMi binarization thresholds. (5) Max CADD, the maximum CADD[40] score of SNVs within or in LD with the specific locus. A high CADD score indicates a strong deleterious effect of the variant. A threshold of 12.37 has been suggested by Kircher et al.[40]. (6) Min RDB, the minimum RegulomeDB[39] score of SNVs within or in LD with the specific locus. RDB is a categorical score to guide interpretation of regulatory variants (from 1a to 7, with 1a being the most biological evidence for an SNV to be a regulatory element)[37,39]. (7) The top-ranked SNV or SNV-SNV pair showing the highest occurrence frequency (Extended Data Fig. 4) averaged across lo-siRF fits from the three LVMi binarization thresholds. A full list of lo-siRF-prioritized SNVs and SNV-SNV pairs can be found in Extended Data 3. (8) Genomic location (hg38) and GWAS statistics information (using PLINK[34]) of the top SNV for each lo-siRF-prioritized locus. Abbreviations: MAF, minor allele frequency; NEA/EA, non-effect-allele/effect-allele; SE, standard error. (9) nPartnerSNVs, number of partner SNVs that interact with the given SNV in lo-siRF. These SNV-SNV pairs interacted in at least one lo-siRF decision path across every LVMi binarization threshold (details in Methods).

Fig. 3: lo-siRF finds epistatic interactions between genetic risk loci for left ventricular hypertrophy.

a, Circos plot showing the genetic risk loci identified by lo-siRF (green, circle 2) and regions after clumping FUMA-extracted SNVs in LD ($r^2 > 0.6$) with lo-siRF-prioritized SNVs (black, circle 3). Circle 1 shows the top 300 epistatic SNV-SNV pairs with the highest frequency of occurrence in lo-siRF (green), SNV-gene linkages (FDR < 0.5) based on GTEx[41] V8 cis-eQTL information from heart and skeletal muscle tissues (purple), and 3D chromatin interactions[37] based on Hi-C data of left ventricular tissue obtained from GSE87112. Circle 5 and 6 show bar plots of the occurrence frequency and number of partner SNVs in epistasis (normalized by the maximum value of the corresponding locus) identified by lo-siRF, respectively. Circle 7 shows the ChromHMM[38] core-15 chromatin state for left ventricle (LV), right ventricle (RV), right atrium (RA), and fetal heart (Fetal). Circle 8 shows the GWAS Manhattan plot from PLINK[34] (circles) where only SNVs with $p < 0.05$ are displayed. The 283 lo-siRF-prioritized SNPs and their LD-linked ($r^2 > 0.6$) SNVs are color-coded as a function of their maximum r^2 value. A portion of these LD-linked SNPs (the outer heatmap layer in circle 8) are extracted from the selected FUMA reference panel (thereby with no GWAS p-values). SNVs that are not in LD ($r^2 \le 0.6$) with any of the 283 lo-siRF-prioritized SNVs are gray. Dashed line indicates GWAS $p = 5E-8$. Circle 9 shows the 21 protein-coding genes mapped by FUMA. **b**, Pie charts showing ANNOVAR enrichment performed for each of the 6 lo-siRF loci (circle 2 in Fig. 3a and Fig. 2f). The arc length of each slice indicates the proportion of SNVs with a specific functional annotation. The radius of each slice indicates $\log_2(E + 1)$, where E is the enrichment score computed as (proportion of SNVs with an annotation for a given locus)/(proportion of SNVs with an annotation relative to all available SNVs in the FUMA reference panel). The dashed circle indicates E = 1 (no enrichment). Asterisks indicate two-sided p-values of Fisher's exact tests for the enrichment of each annotation. Details can be found in Extended Data 4 and 5.

Fig. 4: Genes mapped from epistatic loci interact in multiple functional co-association networks.

a, UpSet plot showing the number of lo-siRF-prioritized SNVs (dark blue) and their LD-linked ($r^2 > 0.6$)

SNVs (light blue, circle 8 in Fig. 3a) that are functionally mapped to each of the 21 protein-coding genes

by positional, eQTL, and/or chromatin interaction (CI) mapping using FUMA[37]. *CCDC141* and *IGF1R*

(highlighted in red) are prioritized by all the three types of SNV-to-gene mapping. Details can be found

in Extended Data 4. **b**, Heatmap of averaged expression (from GTEx) per tissue type per gene (50%

winsorization, $\log_2(\text{TPM} + 1)$) for these functionally mapped genes. **c**, Co-association network built from an enrichment analysis integrating multiple annotated gene set libraries for gene ontology (GO) and pathway terms from Enrichr[46]. The co-association network connects top enriched GO and pathway terms with genes (green nodes in the network) functionally linked from lo-siRF-prioritized epistatic and hypostatic loci (Fig. 2f). Strengths (indicated by the edge width in the network) of the co-association between enriched terms and genes were measured and ranked by the empirical *p*-value from an exhaustive permutation of the co-association score for all possible gene-gene combinations in the network (Details in Methods and Extended Data 6). **d,** A comparison between the top 10 transcription factors (TFs) enriched from genes prioritized (top) and deprioritized (bottom) by lo-siRF. The lo-siRF-prioritized genes are genes functionally linked from lo-siRF-prioritized SNVs (panel **a**). The lo-siRF-deprioritized genes are genes functionally linked from SNVs that failed to pass the lo-siRF prioritization thresholds. For each of the two gene groups, enrichment results against nine TF-annotated gene set libraries from ChEA3[47] and Enrichr[46] were integrated and ranked by the number of significantly (FET $p <$ 0.05) overlapped libraries (numbers in the *nLibraries* column) and the mean scaled rank across all libraries containing that TF (colored boxes in the *nLibraries* column). The balloon plot shows the lowest FET *p*-values for each TF (horizontal axis) and the proportion of overlapped genes (balloon size) between the input gene set and the corresponding TF-annotated gene set. **e**, Heatmap showing the TF co-association strength of gene-gene combinations among lo-siRF-prioritized genes relative to randomly selected gene pairs in the co-association network. More details are available in Extended Data Fig. 5 and Extended Data 7.

Fig. 5: Network analysis using transcriptomic data from 313 human hearts confirms the *CCDC141-TTN* interaction.

a, Control (blue) and heart failure (red) gene co-expression networks were established from a weighted gene co-expression network analysis (WGCNA) on transcriptomic data obtained from 313 non-failing and failing human heart tissues[51]. **b-c**, The connectivity between lo-siRF-prioritized genes in this study was compared against the full connectivity distributions for all possible gene-gene combinations in the control (**b**) and heart failure (**c**) networks. *CCDC141* and *TTN* show significant connectivity in both networks. **d**, Comparing the difference in connectivities between the control and heart failure networks indicate a change in the connectivity of the hypothesized *CCDC141-TTN* interaction during the progression of failing hearts. **e-f**, Dendrograms from WGCNA control (**g**) and heart failure (**h**) networks show distinctive gene module structures.

Fig. 6: *CCDC141* non-additively interacts with *TTN* and *IGF1R* to modify cardiomyocyte morphology.

a, Human induced pluripotent stem cell (iPSC)-derived cardiomyocytes with and without hypertrophic cardiomyopathy (carrying an *MYH7*-R403Q mutation) were transfected with scramble siRNA or siRNAs specifically targeting single (*CCDC141*, *IGF1R*, and *TTN*) or combined (*CCDC141-IGF1R* and *CCDC141-TTN*) genetic loci prioritized by lo-siRF. **b**, Gene-silenced cardiomyocytes were bifurcated into two focused streams of large and small cells using a spiral microfluidic device (cell focusing mechanism illustrated in Extended Data Fig. 6) to allow high-resolution single cell imaging. **c**, Workflow of the image analysis process. Time-lapse image sequences of single cells passing through the top and bottom microchannel outlets (panel **b**) were fed into a customized MATLAB-based program that extracts cell size/shape features via a sequential process of bright field background correction, cell boundary detection, cell tracking and stuck cell removal, cell feature extraction, data quality control and postprocessing, and morphological feature analysis. Extracted single cell features for each gene-silencing condition were compared with their scrambled control values to validate the potential role of epistasis in the genetic regulation of cardiomyocyte hypertrophy (**d-j**). **d**, Violin plots of cell diameters of unaffected (blue) and *MYH7*-R403Q variant (red) cardiomyocytes. Solid and dashed lines in box plots represent median and mean values, respectively. Asterisks indicate significant difference (***p < 1E-36, Wilcoxon signed rank test). **e**, Gene-silencing efficiency in unaffected (blue, n = 5 to 9) and *MYH7*-R403Q variant (red, n = 3) cells based on RT-qPCR analysis (details in Methods). Error bars indicate standard deviations. **f**, Percent change (relative size difference) in median cell diameter of gene-silenced cardiomyocytes relative to scramble control values indicates that *CCDC141* interacts with *IGF1R* to rescue cardiomyocyte hypertrophy. Relative size differences were averaged across data from two to four independent batches of cells. Error bars indicate standard deviations computed on 1000 bootstrap samples of these batches with the following sample size: n = 13147 (*TTN*), 19460 (*IGF1R*), 45304 (*CCDC141*), 19979 (*CCDC141-TTN*), and 26135 (*CCDC141-IGF1R*) for unaffected cells and n = 22134 (*TTN*), 33801 (*IGF1R*), 21158 (*CCDC141*), 39515 (*CCDC141-TTN*), and 52049 (*CCDC141-IGF1R*) for *MYH7*-R403Q variant cells. Asterisks indicate significant difference between gene-silencing and scramble control conditions based on the maximum *p*-values of Wilcoxon signed rank test across all batches of cells (*p < 0.05, **p < 0.001, and ***p < 1E-4). **g**, *CCDC141* non-additively interacts with

IGF1R (left) and *TTN* (right) to modify boundary and texture features of unaffected (blue) and *MYH7-R403Q* variant (red) cells. Cell boundary waveness and texture irregularity were measured by the roundness error (**h**, top) and normalized peak number (**h**, bottom), respectively. **i**, Representative single-cell images overlapped with detected cell boundaries (red lines) show that a higher roundness error indicates increased irregularity of the cell boundary. **j**, Representative single-cell images with detected peaks (blue plus signs) of the brightfield intensity distribution enclosed within the cell boundaries (red lines) indicate a varying level of cell textural irregularity. Scale bars: 10 µm. Detailed statistical information of cell morphology measurements and non-additivity analysis for the studied gene pairs can be found in Extended Data 8.

Extended Data Figures

Extended Data Fig. 1: Distribution of LVM and LVMi measurements for 29,661 UK Biobank participants.

Left ventricular mass (LVM, **a**) and LVM indexed to body surface area (LVMi, **b**) measurements were extracted from cardiac magnetic resonance imaging for 29,661 unrelated White British individuals via deep learning[25]. **c,** A high Pearson correlation of 0.92 was observed between these LVM and LVMi measurements.

Extended Data Fig. 2: LVMi GWAS using BOLT-LMM and PLINK.

GWAS using BOLT-LMM (**a**) and PLINK (**b**) identified associations with LVMi, of which the lead SNV rs3045696 showed the highest significance. This SNV rs3045696 was also identified as the top lead SNV by both BOLT-LMM and PLINK while other lead SNVs (labeled) were significant in either the BOLT-LMM GWAS or the PLINK GWAS but not both. The red dashed line denotes the genome-wide significance threshold ($p < 5E-8$). The two SNVs, rs3045696 and rs67172995, are also stably prioritized by lo-siRF as epistasis interactor variants. Details can be found in Extended Data 1.

Extended Data Fig. 3: Differences in local stability scores between high and low LV mass highlight the importance of the lo-siRF-prioritized interactions between genetic loci.

a, Schematic of local stability importance score computation. Given a locus (light blue transcript), the local stability importance score for an individual is defined as the proportion of trees for which at least one SNV (shaded black region) in the locus is used in the individual's decision path. This computation (top) was performed for each individual (denoted by the stacked boxes). Then, a permutation test was

conducted to assess the difference in these local stability importance scores between the low and high LVMi individuals (bottom). **b**, Differences in the distribution of local stability importance scores suggest that the identified interactions between genetic loci are important for differentiating individuals with high (dark gray) and low (orange) LVMi in the siRF fit. This result, evaluated on the validation data, is stable across the three binarization thresholds and is quantified by a permutation *p*-value given in the top right corner of each subplot.

Extended Data Fig. 4: Top SNVs from lo-siRF-prioritized loci and interactions between loci.

The most important SNVs and SNV-SNV pairs, as measured by their proportion of occurrence in the siRF fit are annotated for the top lo-siRF-prioritized interactions between loci in **a** and top genetic loci in **b**. The y-axis shows the proportion of decision paths in siRF, for which the SNV or SNV-SNV pair occurs, averaged across all three binarization thresholds. In each of the interactions between genetic loci, the SNV rs7591091 in the *CCDC141* locus appears most frequently, suggesting a key role in cardiac hypertrophy.

Extended Data Fig. 5: Genes mapped from epistatic loci share transcription factors and splicing regulators.

For the gene pairs exhibiting a strong co-association to transcription factors (TFs) and RNA-binding regulators ($p < 0.05$, Fig. 4e), the horizontal bars indicate the number of shared TFs or RNA-binding regulators (bottom axis), of which the top-ranked one with the lowest enrichment p-value (dots, top axis) are labeled on the bars, which are colored by the corresponding gene set library. Detailed information can be found in Extended Data 7.

Extended Data Fig. 6: Spiral-shaped inertial microfluidic channel for cell focusing and imaging.

a, Schematic of an inertial microfluidic cell focusing device. Cell suspensions and fresh medium were introduced into the microfluidic device through the cell and sheath flow inlets, respectively, using a syringe pump and flowed down the 5-loop spiral microchannel with the same flow rate (1.2 mL/min). Inserted microscopy images show that randomly dispersed cells were separated by size and bifurcated into the top (large cells) or bottom (small cells) outlets. Scale bar, 10 μm. Outlet channels are connected to straight observation channels where flowing cells were further focused in the channel height direction and imaged using a high-speed camera for morphological feature extraction (Fig. 6c).

b, Schematic of the cell focusing principle. The spiral microchannel has a cross-section with a slanted ceiling, resulting in different depths at the inner and outer side of the microchannel. This geometry

induces strong Dean vortices (counter rotating vortices in the plane perpendicular to the main flow direction) in the outer half of the microchannel cross-section. The interplay between drag forces (F_D) induced by Dean vortices and lift forces (F_L) due to shear gradient and the channel wall drives cell transverse migration towards equilibrium positions where the net force is zero. As a result, large cells in a heterogeneous population progressively migrate closer to the inner channel wall, while smaller cells move towards the outer channel wall. Details about microchannel dimensions can be found in Methods.

Extended Data Fig. 7: Epistatic genes non-uniformly reshape cardiomyocyte size distributions.

a, A heatmap of relative differences of cell sizes at various quantile levels between gene-silencing and scramble control conditions for unaffected and *MYH7*-R403Q variant cardiomyocytes. Larger quantiles correspond to larger cells in the cell size distribution. Dark red indicates strong reduction of cell sizes at the specified quantile level in gene-silenced cells relative to the scramble control. The corresponding statistical differences (**b**) were evaluated by the maximum *p*-values across all batches of cells using a bootstrap quantile test (with 10,000 bootstrapped samples). **c**, Representative QQ-plots of cell size quantiles comparing between gene-silenced cells and scramble controls for both unaffected (top row) and *MYH7*-R403Q variant (bottom row) cardiomyocytes indicate a clear size-bias in the effect of silencing *CCDC141-IGF1R* on correcting cardiomyocyte hypertrophy.

Extended Data Fig. 8: Effects of lo-siRF-prioritized genes and gene-gene interactions on hypertrophic and non-hypertrophic cell morphology.

Relative differences in median cell size (**a**), normalized peak number (**b**), and roundness error (**c**) of gene-silenced human induced pluripotent stem cells-derived cardiomyocytes compared to scramble controls computed for hypertrophic cells (size-sorted by microfluidic channel illustrated in Extended Data Fig. 6, gray bars), non-hypertrophic cells (white bars), or both (black bars). Error bars indicate standard deviation calculated from bootstrapping samples of 2 to 4 batches of cells. Asterisks indicate significant differences compared to the scramble control based on the maximum p-values of Wilcoxon signed rank test across all batches of cells (*$p < 0.05$, **$p < 0.001$, and ***$p < 1E-4$).

Supplementary Tables

Supplementary Table 1: Characteristics of 29,661 analyzed participants in the UK Biobank.

	Female	Male
N	15189	14472
LVMi (g/m^2)	40.7 (5.7)	51.3 (7.8)
LVM (g)	70.7 (12.3)	102.7 (18.6)
Age (y)	63.4 (7.5)	64.9 (7.7)
Height (cm)	162.9 (6.2)	176.2 (6.6)
Weight (kg)	69.0 (13.0)	83.7 (13.3)
Hypertensive Diseases	20.1%	29.8%
Aortic Stenosis	0.1%	0.2%
Heart Failure	0.1%	0.6%
Type II Diabetes	1.6%	3.2%
Blood Pressure Medication	12.1%	20.0%

Summary statistics of the 29,661 unrelated White British individuals analyzed in this study. Means and standard deviations (in parentheses) are reported for continuous measurements (LVMi, LVM, age, height, and weight) alongside the number of individuals (N) and the percentage of individuals with various cardiac hypertrophy-related diseases or on blood pressure medication. We define hypertensive diseases as anyone with self-reported hypertension, high blood pressure as diagnosed by a doctor, or any ICD10 billing code diagnosis in I10-I16; aortic stenosis as self-reported aortic stenosis or an ICD10 billing code diagnosis of I35; heart failure as self-reported heart failure or an ICD10 billing code diagnosis of I50; and type II diabetes as self-reported type II diabetes or an ICD10 billing code diagnosis of E11.

Supplementary Table 2: Thresholds defining low and high LVMi groups used in siRF fit

Binarization Threshold	Male		Female	
	Low LVMi Threshold	High LVMi Threshold	Low LVMi Threshold	High LVMi Threshold
0.15	43.8	58.5	35.1	46.1
0.20	45.1	56.8	36.0	44.9
0.25	46.0	55.4	36.8	43.8

For each of the three binarization thresholds used in lo-siRF (corresponding to the bottom/top 15[th], 20[th], and 25[th] quantiles), we provide the gender-specific LVMi cutoffs for the low and high LVMi groups. All thresholds were measured in g/m^2.

Supplementary Table 3: Prediction accuracies of methods across different LVMi binarization thresholds.

Method	Binarization Threshold = 0.15			Binarization Threshold = 0.20			Binarization Threshold = 0.25		
	Accuracy	AUROC	AUPRC	Accuracy	AUROC	AUPRC	Accuracy	AUROC	AUPRC
siRF	0.554	**0.585**	**0.579**	**0.557**	**0.583**	**0.562**	**0.563**	**0.582**	**0.556**
RF	0.546	0.572	0.571	0.553	0.569	0.549	0.554	0.569	0.555
Lasso	0.547	0.559	0.541	0.545	0.556	0.529	0.534	0.550	0.526
Ridge	**0.559**	0.567	0.550	0.539	0.563	0.539	0.541	0.555	0.536
SVM	0.553	0.566	0.552	0.551	0.565	0.541	0.544	0.558	0.541

Maximum prediction accuracies highlighted in bold. The siRF model performs better or on par with other commonly used machine learning methods when predicting the binarized LVMi phenotype. This result holds across all three binarization thresholds and three different classification metrics, i.e., classification accuracy, area under the receiver operator curve (AUROC), and area under the precision-recall curve (AUPRC). In accordance with the prediction check component of the PCS framework, siRF is an appropriate fit for the given data.

Supplementary Table 4: Top signed loci and interactions between loci, prioritized by lo-siRF across LVMi binarization thresholds.

Loci / Interaction	Binarization Threshold			Mean p-value
	0.15	0.20	0.25	
$CCDC141^- - IGF1R^-$	$< 10^{-3}$	$< 10^{-4}$	$< 10^{-4}$	$< 10^{-3}$
$IGF1R^-$	$< 10^{-3}$	$< 10^{-3}$	$< 10^{-3}$	$< 10^{-3}$
$MIR588;RSPO3^+$	0.002	0.004	$< 10^{-4}$	0.002
TTN^-	0.022	0.006	$< 10^{-3}$	0.009
$CCDC141^- - TTN^-$	0.030	0.002	$< 10^{-3}$	0.011
TTN^+	0.030	0.005	$< 10^{-3}$	0.012
$MIR588;RSPO3^-$	0.016	0.014	0.009	0.013
$LSP1^-$	0.029	0.019	0.002	0.017
$CCDC141^-$	0.033	0.007	0.015	0.018
$CCDC141^- - LOC157273;TNKS^-$	0.099	0.024	0.044	0.056

A list of the top signed loci and interactions between loci, prioritized by lo-siRF, that were stably important across all three LVMi binarization thresholds (Supplementary Table 2). These loci and interactions between loci are ranked by the lo-siRF *p*-value, averaged across the three binarization thresholds.

Supplementary Table 5: Summary of siRF evaluation metrics for top interactions between loci.

Threshold	Prevalence	Precision	Class Difference in Prevalence	Stability of Class Difference in Prevalence	Independence of Feature Selection	Stability of Independence of Feature Selection	Increase in Precision	Stability of Increase in Precision	Stability
CCDC141⁻–IGF1R⁻									
Binarization Threshold = 0.15	0.064	0.56	0.012	1.0	0.00094	0.78	0.017	1.0	0.68
Binarization Threshold = 0.2	0.062	0.54	0.0095	1.0	0.0044	1.0	0.014	1.0	0.64
Binarization Threshold = 0.25	0.077	0.54	0.012	1.0	0.011	1.0	0.019	1.0	0.82
CCDC141⁻–LOC157273;TNKS⁻									
Binarization Threshold = 0.15	0.20	0.56	0.040	1.0	0.012	1.0	0.020	1.0	1.0
Binarization Threshold = 0.2	0.19	0.56	0.037	1.0	0.020	1.0	0.028	1.0	1.0
Binarization Threshold = 0.25	0.23	0.55	0.043	1.0	0.0068	1.0	0.022	1.0	1.0
CCDC141⁻–TTN⁻									
Binarization Threshold = 0.15	0.12	0.56	0.023	1.0	0.0066	1.0	0.016	1.0	1.0
Binarization Threshold = 0.2	0.14	0.55	0.024	1.0	0.0042	0.96	0.017	1.0	1.0
Binarization Threshold = 0.25	0.085	0.52	0.0076	1.0	-0.0036	0.020	0.0024	0.88	0.92

Though prediction accuracy is weak (indicated by precision scores close to 0.5), the lo-siRF-prioritized interactions are stable across binarization thresholds and across bootstrap replicates (indicated by all types of stability scores being close or equal to 1). Here, prevalence measures the proportion of high LVMi individuals for which the interaction appears. Precision measures the probability of having high LVMi given that the interaction is active. The class difference in prevalence is the prevalence of the interaction in high LVMi individuals minus the prevalence in low LVMi individuals. Independence of feature selection evaluates whether the interaction is collectively or individually associated with the responses. The stability of each of these metrics evaluates how stable the respective scores are across 50 bootstrap replicates. The overall stability score (last column) is the proportion of times that the interaction is identified by siRF across 50 bootstrapped replicates. Higher scores for each listed metric indicate greater importance.

Data availability

All genotype and cardiac MRI data used as input to the lo-siRF pipeline are available from the UK Biobank (https://www.ukbiobank.ac.uk/). This work was conducted under the UK Biobank application 22282. GWAS-filtered SNVs using PLINK[34] and BOLT-LMM[35] are summarized in Extended Data 2. Data for the gene co-expression networks from 313 explanted human hearts is available at https://doi.org/10.5281/zenodo.2600420.

Code availability

All code for running the lo-siRF analysis and analyzing the experimental results can be found on GitHub (https://github.com/Yu-Group/epistasis-cardiac-hypertrophy). This lo-siRF analysis was conducted using R version 3.6.1, Python 3.6.1, and iRF2.0 (https://github.com/karlkumbier/iRF2.0). The LVMi derivation from cardiac MRI images and corresponding deep learning model have been published elsewere[25] (https://github.com/baiwenjia/ukbb_cardiac). PLINK[34] (https://www.cog-genomics.org/plink/) and BOLT-LMM[35] (https://alkesgroup.broadinstitute.org/BOLT-LMM/BOLT-LMM_manual.html) were used to perform the GWAS dimension reduction. ANNOVAR[78] (https://annovar.openbioinformatics.org/en/latest/) was used to map each SNV to a genetic locus within lo-siRF. FUMA GWAS[37] (https://fuma.ctglab.nl/) version 1.5.4 was used to functionally annotate SNVs and map to genes.

Acknowledgements

The authors would like to acknowledge Dr. David Amar, Dr. Srigokul Upadhyayula, Dr. Haiyan Huang, and Dr. Ziad Obermeyer for their critical comments and discussions on this work, Stanford nanofabrication facility and Elmer Enriquez for the technical support of microfluidic device fabrication, and Dr. Anna Shcherbina and Dr. Manuel Rivas for their technical help with the UK Biobank data. This work was supported by the Chan Zuckerberg Biohub – San Francisco through the Intercampus Research Awards (2019 - 2022) to R.A., J.R.P., J.B.B., A.J.B., E.A.A., and B.Y. E.A.A. received funding from National Institutes of Health (NIH) through grant number 1R01HL144843. B.Y. received support from National Science Foundation (NSF) through grants DMS-1613002 and IIS 1741340, and a Weill

Neurohub grant. V.N.P. received funding from K08HL143185. T.M.T. was supported by the National Science Foundation (NSF) Graduate Research Fellowship Program DGE-2146752. Q.W. received funding from American Heart Association Postdoctoral Fellowship through grant number 23POST1023278. C.S.W. received support from NIH through grants F32HL160067 and L30HL159413.

Author Information

These authors contributed equally: Qianru Wang and Tiffany M. Tang
Equal senior authors: Bin Yu and Euan A. Ashley

Authors and Affiliations

Department of Medicine, Division of Cardiovascular Medicine, Stanford University, Stanford, CA, USA
Qianru Wang, Nathan Youlton, Chad S. Weldy, J. Weston Hughes, Elizabeth T. Chin, Shirley C. Sutton, James R. Priest, Victoria N. Parikh, and Euan A. Ashley

Department of Statistics, University of California, Berkeley, Berkeley, CA, USA
Tiffany M. Tang, Ana M. Kenney, Omer Ronen, Abhineet Agarwal, Xiao Li, James B. Brown, and Bin Yu

Faculty of Informatics and Data Science, University of Regensburg, Regensburg, Germany
Merle Behr

Department of Pharmaceutical Chemistry, University of California, San Francisco, San Francisco, CA, USA
Karl Kumbier

Department of Cardiovascular and Metabolic Sciences, Lerner Research Institute, Cleveland Clinic, Cleveland, OH, USA
Christine S. Moravec and W. H. Wilson Tang

Department of Cardiovascular Medicine, Heart Vascular and Thoracic Institute, Cleveland Clinic, Cleveland, OH, USA

W. H. Wilson Tang

Division of Cardiovascular Medicine, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

Kenneth B. Margulies and Thomas P. Cappola

Hospital of The University of Pennsylvania, Philadelphia, PA, USA

Kenneth B. Margulies and Thomas P. Cappola

Bakar Computational Health Sciences Institute, University of California, San Francisco, San Francisco, CA, USA

Atul J. Butte and Rima Arnaout

Chan Zuckerberg Biohub – San Francisco, San Francisco, CA 94158, USA

Atul J. Butte, Rima A. Arnaout, James B. Brown, James R. Priest, Bin Yu, and Euan A. Ashley

Division of Environmental Genomics and Systems Biology, Lawrence Berkeley National Laboratory, Berkeley, CA, USA

James B. Brown

Tenaya Therapeutics, San Francisco, CA, USA

James R. Priest

Department of Electrical Engineering and Computer Science, University of California, Berkeley, Berkeley, CA, USA

Bin Yu

Center for Computational Biology, University of California, Berkeley, Berkeley, CA, USA

Bin Yu

Current addresses

Department of Statistics, University of Michigan, Ann Arbor, MI, USA

Tiffany M. Tang

Department of Molecular Cell and Developmental Biology, University of California Santa Cruz, Santa Cruz, CA, USA

Nathan Youlton

Department of Statistics, University of California Irvine, Irvine, CA, USA

Ana M. Kenney

Department of Biostatistics, Johns Hopkins University, Baltimore, MD, USA

Elizabeth T. Chin

Contributions

Q.W., T.M.T., C.S.W., J.W.H., A.J.B., R.A., J.B.B., J.R.P., V.N.P., B.Y., and E.A.A. conceived and designed research. J.W.H. performed LVMi extraction from cardiac MRI images. T.M.T., A.A., X.L., M.B., and K.K. performed exploratory data investigations leading to development of lo-siRF. T.M.T., A.A., and B.Y. developed the lo-siRF pipeline; T.M.T. performed the lo-siRF analysis. Q.W. and T.M.T. performed the FUMA SNP2GENE process using lo-siRF-prioritized SNVs and GWAS-filtered SNVs. Q.W. evaluated functional annotation results from FUMA, performed ANNOVAR enrichment test for each lo-siRF loci and functional gene mapping. Q.W. performed integrative biological enrichment analyses and evaluated the co-associations between lo-siRF hypothesized gene-gene interactions and the enriched GOs, pathways, and TFs. E.T.C., C.S.M., W.H.W.T., K.B.M., T.P.C., V.N.P., and E.A.A. contributed to the data collection and construction of WGCNA healthy and heart failure coexpression networks. E.T.C. evaluated connectivity differences of epistatic genes hypothesized by lo-siRF between cardiac co-expression networks of failing and non-failing hearts. Q.W. designed and created microfluidic devices; N.Y., S.C.S. and V.N.P. created gene-silenced hiPSC-CM lines; Q.W. and N.Y. performed the microfluidic single cell imaging experiments. Q.W., O.R., and A.M.K. performed single cell image analysis and morphological feature extraction. Q.W., T.M.T., A.M.K., O.R., C.S.W., V.N.P., J.W.H., B.Y., and E.A.A. interpreted results of experiments; Q.W., T.M.T., and A.M.K. prepared figures;

Q.W., T.M.T., C.S.W. and J.W.H. drafted manuscript; All authors contributed in editing and revising manuscript.

Corresponding authors

Correspondence to Bin Yu (binyu@berkeley.edu) and Euan A. Ashley (euan@stanford.edu).

Competing interests

E.A.A. is a Founder of Personalis, Deepcell, Svexa, RCD Co, and Parameter Health; Advisor to Oxford Nanopore, SequenceBio, and Pacific Biosciences; and a non-executive director for AstraZeneca. C.S.W. is a consultant for Tensixteen Bio and Renovacor. V.N.P. is an SAB member for and receives research support from BioMarin, Inc, and is a consultant for Constantiam, Inc. and viz.ai. The remaining authors declare no competing interests.

Reference

1. Weldy, C. S. & Ashley, E. A. Towards precision medicine in heart failure. *Nat. Rev. Cardiol.* 1–18 (2021).

2. Sharir, T. *et al.* Ventricular systolic assessment in patients with dilated cardiomyopathy by preload-adjusted maximal power. Validation and noninvasive application. *Circulation* **89**, 2045–2053 (1994).

3. Bastos, M. B. *et al.* Invasive left ventricle pressure–volume analysis: overview and practical clinical implications. *Eur. Heart J.* **41**, 1286–1297 (2019).

4. Udelson, J. E., 3rd, R. O. C., Bacharach, S. L., Rumble, T. F. & Bonow, R. O. Beta-adrenergic stimulation with isoproterenol enhances left ventricular diastolic performance in hypertrophic cardiomyopathy despite potentiation of myocardial ischemia. Comparison to rapid atrial pacing. *Circulation* **79**, 371–382 (1988).

5. Burkhoff, D., Mirsky, I. & Suga, H. Assessment of systolic and diastolic ventricular properties via pressure-volume analysis: a guide for clinical, translational, and basic researchers. *American Journal of Physiology-Heart and Circulatory Physiology* **289**, H501–H512 (2005).

6. Marian, A. J. & Braunwald, E. Hypertrophic Cardiomyopathy: Genetics, Pathogenesis, Clinical Manifestations, Diagnosis, and Therapy. *Circ. Res.* **121**, 749–770 (2017).

7. Haider, A. W., Larson, M. G., Benjamin, E. J. & Levy, D. Increased left ventricular mass and hypertrophy are associated with increased risk for sudden death. *J. Am. Coll. Cardiol.* **32**, 1454–1459 (1998).

8. Chrispin, J. *et al.* Association of Electrocardiographic and Imaging Surrogates of Left Ventricular Hypertrophy With Incident Atrial Fibrillation. *Journal of the American College of Cardiology* vol. 63 2007–2013 Preprint at https://doi.org/10.1016/j.jacc.2014.01.066 (2014).

9. Kawel-Boehm, N. *et al.* Left Ventricular Mass at MRI and Long-term Risk of Cardiovascular Events: The Multi-Ethnic Study of Atherosclerosis (MESA). *Radiology* **293**, 107–114 (2019).

10. Bluemke, D. A. *et al.* The Relationship of Left Ventricular Mass and Geometry to Incident Cardiovascular Events The MESA (Multi-Ethnic Study of Atherosclerosis) Study. *J. Am. Coll.*

Cardiol. **52**, 2148–2155 (2008).

11. Pirruccello, J. P. *et al.* Analysis of cardiac magnetic resonance imaging in 36,000 individuals yields genetic insights into dilated cardiomyopathy. *Nat. Commun.* **11**, 2254 (2020).

12. Bai, W. *et al.* A population-based phenome-wide association study of cardiac and aortic structure and function. *Nat. Med.* **26**, 1654–1662 (2020).

13. Meyer, H. V. *et al.* Genetic and functional insights into the fractal structure of the heart. *Nature* **584**, 589–594 (2020).

14. Harper, A. R. *et al.* Common genetic variants and modifiable risk factors underpin hypertrophic cardiomyopathy susceptibility and expressivity. *Nat. Genet.* **53**, 135–142 (2021).

15. O'Sullivan, J. W. *et al.* Polygenic Risk Scores for Cardiovascular Disease: A Scientific Statement From the American Heart Association. *Circulation* **146**, e93–e118 (2022).

16. Guindo-Martínez, M. *et al.* The impact of non-additive genetic associations on age-related complex diseases. *Nat. Commun.* **12**, 2436 (2021).

17. Zeng, L. *et al.* Cis-epistasis at the LPA locus and risk of cardiovascular diseases. *Cardiovasc. Res.* **118**, 1088–1102 (2022).

18. Li, Y. *et al.* Statistical and Functional Studies Identify Epistasis of Cardiovascular Risk Genomic Variants From Genome-Wide Association Studies. *J. Am. Heart Assoc.* **9**, e014146 (2020).

19. Basu, S., Kumbier, K., Brown, J. B. & Yu, B. Iterative random forests to discover predictive and stable high-order interactions. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 1943–1948 (2018).

20. Kumbier, K. *et al.* Signed iterative random forests to identify enhancer-associated transcription factor binding. *arXiv [stat.ML]* (2023).

21. Reimherr, M. & Nicolae, D. L. You've gotta be lucky: Coverage and the elusive gene-gene interaction. *Ann. Hum. Genet.* **75**, 105–111 (2011).

22. Murk, W., Bracken, M. B. & DeWan, A. T. Confronting the missing epistasis problem: on the reproducibility of gene-gene interactions. *Hum. Genet.* **134**, 837–849 (2015).

23. Yu, B. & Kumbier, K. Veridical data science. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 3920–3929 (2020).

24. Koch, E. M. & Sunyaev, S. R. Maintenance of Complex Trait Variation: Classic Theory and Modern Data. *Front. Genet.* **12**, 763363 (2021).

25. Bai, W. *et al.* Automated cardiovascular magnetic resonance image analysis with fully convolutional networks. *J. Cardiovasc. Magn. Reson.* **20**, 65 (2018).

26. Wang, Q., Jones, A.-A. D., 3rd, Gralnick, J. A., Lin, L. & Buie, C. R. Microfluidic dielectrophoresis illuminates the relationship between microbial cell envelope polarizability and electrochemical activity. *Sci Adv* **5**, eaat5664 (2019).

27. Di Carlo, D. Inertial microfluidics. *Lab Chip* **9**, 3038–3046 (2009).

28. Guan, G. *et al.* Spiral microchannel with rectangular and trapezoidal cross-sections for size based particle separation. *Sci. Rep.* **3**, 1475 (2013).

29. Wu, P.-H. *et al.* Single-cell morphology encodes metastatic potential. *Sci Adv* **6**, eaaw6938 (2020).

30. Dainis, A. *et al.* Silencing of MYH7 ameliorates disease phenotypes in human iPSC-cardiomyocytes. *Physiol. Genomics* **52**, 293–303 (2020).

31. Littlejohns, T. J. *et al.* The UK Biobank imaging enhancement of 100,000 participants: rationale, data collection, management and future directions. *Nat. Commun.* **11**, 2624 (2020).

32. Grothues, F. *et al.* Comparison of interstudy reproducibility of cardiovascular magnetic resonance with two-dimensional echocardiography in normal subjects and in patients with heart failure or left ventricular hypertrophy. *Am. J. Cardiol.* **90**, 29–34 (2002).

33. Du Bois, D. & Du Bois, E. F. Clinical calorimetry: tenth paper a formula to estimate the approximate surface area if height and weight be known. *JAMA Intern. Med.* **17**, 863–871 (1916).

34. Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet.* **81**, 559–575 (2007).

35. Loh, P.-R. *et al.* Efficient Bayesian mixed-model analysis increases association power in large cohorts. *Nat. Genet.* **47**, 284–290 (2015).

36. McInnes, G. *et al.* Global Biobank Engine: enabling genotype-phenotype browsing for biobank summary statistics. *Bioinformatics* **35**, 2495–2497 (2019).

37. Watanabe, K., Taskesen, E., van Bochoven, A. & Posthuma, D. Functional mapping and

annotation of genetic associations with FUMA. *Nat. Commun.* **8**, 1826 (2017).

38. Roadmap Epigenomics Consortium *et al.* Integrative analysis of 111 reference human epigenomes. *Nature* **518**, 317–330 (2015).

39. Boyle, A. P. *et al.* Annotation of functional variation in personal genomes using RegulomeDB. *Genome Res.* **22**, 1790–1797 (2012).

40. Kircher, M. *et al.* A general framework for estimating the relative pathogenicity of human genetic variants. *Nat. Genet.* **46**, 310–315 (2014).

41. GTEx Consortium. The Genotype-Tissue Expression (GTEx) project. *Nat. Genet.* **45**, 580–585 (2013).

42. Khurshid, S. *et al.* Clinical and genetic associations of deep learning-derived cardiac magnetic resonance-based left ventricular mass. *Nat. Commun.* **14**, 1558 (2023).

43. Aung, N. *et al.* Genome-Wide Analysis of Left Ventricular Image-Derived Phenotypes Identifies Fourteen Loci Associated With Cardiac Morphogenesis and Heart Failure Development. *Circulation* **140**, 1318–1330 (2019).

44. Verweij, N., van de Vegte, Y. J. & van der Harst, P. Genetic study links components of the autonomous nervous system to heart-rate profile during exercise. *Nat. Commun.* **9**, 898 (2018).

45. Thorolfsdottir, R. B. *et al.* Genetic insight into sick sinus syndrome. *Eur. Heart J.* **42**, 1959–1971 (2021).

46. Kuleshov, M. V. *et al.* Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. *Nucleic Acids Res.* **44**, W90–7 (2016).

47. Keenan, A. B. *et al.* ChEA3: transcription factor enrichment analysis by orthogonal omics integration. *Nucleic Acids Res.* **47**, W212–W224 (2019).

48. Evangelista, J. E. *et al.* Enrichr-KG: bridging enrichment analysis across multiple libraries. *Nucleic Acids Res.* **51**, W168–W179 (2023).

49. Li, L. *et al.* Attention-deficit/hyperactivity disorder as a risk factor for cardiovascular diseases: a nationwide population-based cohort study. *World Psychiatry* **21**, 452–459 (2022).

50. Parikh, V. N. *et al.* Regional Variation in RBM20 Causes a Highly Penetrant Arrhythmogenic

Cardiomyopathy. *Circ. Heart Fail.* **12**, e005371 (2019).

51. Cordero, P. *et al.* Pathologic gene network rewiring implicates PPP1R3A as a central regulator in pressure overload heart failure. *Nat. Commun.* **10**, 2760 (2019).

52. Hood, K., Kahkeshani, S., Di Carlo, D. & Roper, M. Direct measurement of particle inertial migration in rectangular microchannels. *Lab Chip* **16**, 2840–2850 (2016).

53. Stavrakis, S., Holzner, G., Choo, J. & deMello, A. High-throughput microfluidic imaging flow cytometry. *Curr. Opin. Biotechnol.* **55**, 36–43 (2019).

54. Alizadeh, E., Xu, W., Castle, J., Foss, J. & Prasad, A. TISMorph: A tool to quantify texture, irregularity and spreading of single cells. *PLoS One* **14**, e0217346 (2019).

55. Wu, X. *et al.* A novel statistic for genome-wide interaction analysis. *PLoS Genet.* **6**, e1001131 (2010).

56. Wan, X. *et al.* BOOST: A fast approach to detecting gene-gene interactions in genome-wide case-control studies. *Am. J. Hum. Genet.* **87**, 325–340 (2010).

57. Kam-Thong, T. *et al.* EPIBLASTER-fast exhaustive two-locus epistasis detection strategy using graphical processing units. *Eur. J. Hum. Genet.* **19**, 465–471 (2011).

58. Ueki, M. & Cordell, H. J. Improved statistics for genome-wide interaction analysis. *PLoS genetics* vol. 8 e1002625 (2012).

59. Nelson, D. L., Lehninger, A. L. & Cox, M. M. *Lehninger Principles of Biochemistry.* (Macmillan, 2008).

60. Stephan, J., Stegle, O. & Beyer, A. A random forest approach to capture genetic effects in the presence of population structure. *Nat. Commun.* **6**, 7432 (2015).

61. Li, J., Malley, J. D., Andrew, A. S., Karagas, M. R. & Moore, J. H. Detecting gene-gene interactions using a permutation-based random forest method. *BioData Min.* **9**, 14 (2016).

62. Adams, S. M. *et al.* Genome Wide Epistasis Study of On-Statin Cardiovascular Events with Iterative Feature Reduction and Selection. *J Pers Med* **10**, (2020).

63. Saha, S., Perrin, L., Röder, L., Brun, C. & Spinelli, L. Epi-MEIF: detecting higher order epistatic interactions for complex traits using mixed effect conditional inference forests. *Nucleic Acids Res.*

50, e114 (2022).

64. Hornung, R. & Boulesteix, A.-L. Interaction forests: Identifying and exploiting interpretable quantitative and qualitative interaction effects. *Comput. Stat. Data Anal.* **171**, 107460 (2022).

65. Jiang, R., Tang, W., Wu, X. & Fu, W. A random forest approach to the detection of epistatic interactions in case-control studies. *BMC Bioinformatics* **10 Suppl 1**, S65 (2009).

66. Singhal, P. *et al.* Evidence of epistasis in regions of long-range linkage disequilibrium across five complex diseases in the UK Biobank and eMERGE datasets. *Am. J. Hum. Genet.* **110**, 575–591 (2023).

67. Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data. *Nature* **562**, 203–209 (2018).

68. Morgan, M. D. *et al.* Genome-wide study of hair colour in UK Biobank explains most of the SNP heritability. *Nat. Commun.* **9**, 5271 (2018).

69. Petersen, S. E. *et al.* Reference ranges for cardiac structure and function using cardiovascular magnetic resonance (CMR) in Caucasians from the UK Biobank population cohort. *J. Cardiovasc. Magn. Reson.* **19**, 18 (2017).

70. Schaid, D. J., Chen, W. & Larson, N. B. From genome-wide associations to candidate causal variants by statistical fine-mapping. *Nat. Rev. Genet.* **19**, 491–504 (2018).

71. Yoshida, M. & Koike, A. SNPInterForest: a new method for detecting epistatic interactions. *BMC Bioinformatics* **12**, 469 (2011).

72. Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer datasets. *Gigascience* **4**, 7 (2015).

73. Shah, R. D. & Meinshausen, N. Random Intersection Trees. *The Journal of Machine Learning Research* **15**, 629–654 (2014).

74. Tibshirani, R. Regression shrinkage and selection via the lasso. *J. R. Stat. Soc. Series B Stat. Methodol.* **58**, 267–288 (1996).

75. Hoerl, A. E. & Kennard, R. W. Ridge Regression: Biased Estimation for Nonorthogonal Problems. *Technometrics* **12**, 55–67 (1970).

76. Breiman, L. Random Forests. *Mach. Learn.* **45**, 5–32 (2001).

77. Cortes, C. & Vapnik, V. Support-vector networks. *Mach. Learn.* **20**, 273–297 (1995).

78. Wang, K., Li, M. & Hakonarson, H. ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic Acids Res.* **38**, e164 (2010).

79. Chen, C. *et al.* TBtools: An Integrative Toolkit Developed for Interactive Analyses of Big Biological Data. *Mol. Plant* **13**, 1194–1202 (2020).

80. Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat. Genet.* **25**, 25–29 (2000).

81. Gene Ontology Consortium *et al.* The Gene Ontology knowledgebase in 2023. *Genetics* **224**, (2023).

82. Smith, C. L. & Eppig, J. T. The mammalian phenotype ontology: enabling robust annotation and comparative analysis. *Wiley Interdiscip. Rev. Syst. Biol. Med.* **1**, 390–399 (2009).

83. Gillespie, M. *et al.* The reactome pathway knowledgebase 2022. *Nucleic Acids Res.* **50**, D687–D692 (2022).

84. Kanehisa, M. & Goto, S. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res.* **28**, 27–30 (2000).

85. Davis, C. A. *et al.* The Encyclopedia of DNA elements (ENCODE): data portal update. *Nucleic Acids Res.* **46**, D794–D801 (2018).

86. Chèneby, J., Gheorghe, M., Artufel, M., Mathelier, A. & Ballester, B. ReMap 2018: an updated atlas of regulatory regions from an integrative analysis of DNA-binding ChIP-seq experiments. *Nucleic Acids Res.* **46**, D267–D275 (2018).

87. Lachmann, A. *et al.* Massive mining of publicly available RNA-seq data from human and mouse. *Nat. Commun.* **9**, 1366 (2018).

88. Portales-Casamar, E. *et al.* JASPAR 2010: the greatly expanded open-access database of transcription factor binding profiles. *Nucleic Acids Res.* **38**, D105–10 (2010).

89. Matys, V. *et al.* TRANSFAC and its module TRANSCompel: transcriptional gene regulation in eukaryotes. *Nucleic Acids Res.* **34**, D108–10 (2006).

90. Han, H. *et al.* TRRUST v2: an expanded reference database of human and mouse transcriptional regulatory interactions. *Nucleic Acids Res.* **46**, D380–D386 (2018).

a UK Biobank Data
❶ SNV Data
Individuals (29,661)
SNVs (~15 million)

❷ Cardiac MRI-derived Phenotype
Cardiac MRIs → deep learning → LVMi
Individuals

b Dimension Reduction of SNVs
LVMi GWAS using BOLT-LMM
LVMi GWAS using PLINK
Select union of top 1000 SNVs from each GWAS

c Binarization of LVMi Phenotype
Frequency
LVMi → Binarized LVMi

d Prediction
GWAS-filtered SNV Data
Binarized LVMi phenotype using x% threshold
Fit signed iterative random forest
Check prediction accuracy

e Prioritization
Fitted signed iterative random forest
Aggregate SNV-level features into genetic loci
Compute local stability importance scores
Conduct permutation test

Rank loci and interactions between loci by permutation p-values, averaged across binarization thresholds

f

Lo-siRF-prioritized LV hypertrophy risk loci

Lo-siRF locus[1]	nIndSigSNVs[2]	nSNVs[3]	Lo-siRF p-value[4]	Max CADD[5]	Min RDB[6]	Top SNV[7]	Chr:Pos[8]	MAF[8]	NEA/EA[8]	Function	GWAS p-value[8]	GWAS Beta[8]	GWAS SE[8]
TTN	52	148	0.009	25.5	1b	rs66733621	2:178799323	0.290	A/G	intronic	8.78E-6	0.0437	0.00983
CCDC141	65	85	0.018	27.3	1a	rs7591091	2:178889467	0.298	C/T	intronic	3.67E-11	-0.0646	0.00975
MIR588;RSPO3	75	218	0.002	20.2	1b	rs9401921	6:126925592	0.379	A/G	intergenic	3.14E-7	0.0474	0.00926
LOC157273;TNKS	76	50	0.142	16.88	1b	rs6999852	8:9478458	0.301	A/G	intergenic	1.15E-5	-0.0426	0.00970
LSP1	9	11	0.017	10.91	1b	rs569550	11:1865838	0.386	T/G	ncRNA_intronic	3.85E-6	0.0423	0.00915
IGF1R	6	60	< 1E-3	17.46	1a	rs62024491	15:98733068	0.312	G/A	intronic	9.43E-6	-0.0426	0.00962

Lo-siRF-prioritized epistatic interactions between LV hypertrophy risk loci

Epistatic interaction[1]	Lo-siRF p-value[4]	Top SNV-SNV interaction[7]	Chr:Pos[8]	nPartnerSNVs[9]
CCDC141 - IGF1R	< 1E-3	rs7591091	2:178889467	133
		rs62024491	15:98733068	62
CCDC141 - TTN	0.011	rs7591091	2:178889467	133
		rs66733621	2:178799323	61
CCDC141 - LOC157273;TNKS	0.056	rs7591091	2:178889467	133
		rs6999852	8:9478458	59

a

MFHAS1
CLDN23
ECHDC1
RNF146
RSPO3
TNKS
LSP1
C11orf89
TNNT3
Chr8
Chr6
Chr11
Chr2
Chr15
IGF1R

Interactions
Epistasis
eQTLs
Chromatin interactions

①

② Lo-siRF loci
③ LD clumping
④ Chromosome
⑤ Occurrence frequency in lo-siRF
⑥ N lo-siRF interactors
⑦ ChromHMM 15-state

LV
RV
RA
Fetal

⑧ GWAS -log₁₀(p-val)

CCDC141
TTN
PLEKHA3
FKBP7
DFNB59
PRKRA
RBM45
PDE11A
⑨ Functionally mapped genes

SYNM
DKFZP779J2370
LYSMD4

TssA	ZNF/Rpts
TssAFlnk	Het
TxFlnk	TssBiv
Tx	BivFlnk
TxWk	EnhBiv
EnhG	ReprPC
Enh	ReprPCWk
	Quies

GWAS -log₁₀(p-val)
0.6 0.7 0.8 0.9 1   r²

b

TTN locus   *CCDC141* locus   *MIR588;RSPO3* locus   *LOC157273;TNKS* locus   *LSP1* locus   *IGF1R* locus

exonic	upstream
intronic	downstream
intergenic	UTR3
ncRNA_exonic	UTR5
ncRNA_intronic	

** p < 1E-5
* p < 0.002

Functional gene mapping from lo-siRF-prioritized genetic variants

a

SNVs

■ Lo-siRF-prioritized SNVs
■ LD-linked SNVs ($r^2 > 0.6$)

b

Functionally mapped genes

Gene expression log₂(TPM+1)

Lo-siRF locus
- TTN
- CCDC141
- MIR588;RSPO3
- LOC157273;TNKS
- LSP1
- IGF1R

c Gene ontology (GO) and pathway coassociation

- MGI Mammalian Phenotype Level 4 2021
- GO Biological Process 2023
- GO Molecular Function 2023
- Reactome 2022
- Gene

-log₁₀(P)

d Transcription factor (TF) enrichment

from genes prioritized by lo-siRF

TF	nLibraries p < 0.05	Mean scaled rank	-LOG₁₀(p)
MYOD1	4	0.298	
HOXA4	3	0.057	
MYF6	3	0.063	
MYOG	3	0.287	
TBX15	2	0.011	
YBX3	2	0.011	
ZNF493	2	0.046	
NFE2L1	2	0.075	
LBX1	2	0.079	
SIX4	2	0.079	

Overlap: 10%, 30%

from genes deprioritized by lo-siRF

TF	nLibraries p < 0.05	Mean scaled rank	-LOG₁₀(p)
TBX20	3	0.135	
MYOG	3	0.188	
IRF2	3	0.289	
PGR	3	0.352	
ARID5A	2	0.029	
DPF3	2	0.043	
SATB2	2	0.055	
GCM1	2	0.077	
ZNF704	2	0.078	
PHF21A	2	0.088	

Overlap: 10%, 30%

- ■ ENCODE ChIP-seq
- ■ ReMap ChIP-seq
- ■ Literature ChIP-seq
- ■ ARCHS4 Coexpression
- ■ GTEx Coexpression
- ■ Enrichr Queries
- ■ Gene DE following TF perturbations
- ■ TRRUST text-mining
- ■ JASPAR and TRANSFAC PWMs

e Transcription factor coassociation

Lo-siRF locus
- TTN
- CCDC141
- MIR588;RSPO3
- LOC157273;TNKS
- LSP1
- IGF1R

-LOG₁₀(p)

a Cardiac transplant tissue from:
136 donor (control) hearts

WGCNA on covariate-corrected RNA expression data

Control network
connectivity

177 failing hearts

Heart failure network

Differential network
connectivity difference

b Control network

CCDC141–IGF1R
empirical *p* = 0.521

CCDC141–TTN
empirical *p* = 0.0026

Probability

Connectivity score in control network

c Heart failure network

CCDC141–IGF1R
empirical *p* = 0.321

CCDC141–TTN
empirical *p* = 0.006

Probability

Connectivity score in heart failure network

d Differential network

CCDC141–TTN
two-tailed empirical
p = 0.002

CCDC141–IGF1R
two-tailed empirical
p = 0.372

Probability

Connectivity difference score

e Control

Height

IGF1R   *TTN*   *CCDC141*

Dynamic Tree Cut

Merged dynamic

f Heart failure

IGF1R   *TTN*   *CCDC141*

a siRNA gene silencing in iPSC derived cardiomyocytes

differentiation & maturation
40 days

donor/patient → skin fibroblasts → induced pluripotent stem cells (iPSCs) → iPSC derived cardiomyocytes

siRNAs
2 days

RISC assembly → RISC activation → Target mRNA recognetion → Target mRNA cleavage

b High-throughput cell focusing and imaging

Io-siRF-prioritized genes & gene-gene interactions

cell lines

Inlet
Sheath flow
High-speed time lapse imaging

Top
Bottom

Bottom outlet   Top outlet

c Image analysis workflow

Load image sequence

Generate cell free background
Frame No.   Brightness
(X,Y)
Background   mode

Background substraction

Image preparation

Detect cell boundary

Trace cell centroid
Centroid X (μm), Y (μm)

Double counts filter → FALSE
TRUE
Remove stuck cells

Feature extraction

Extract cell features
Diameter (μm),
Solidity
Peaks of 2D brightness field
Roundness
Circularity

Peaks - Solidity filter → TRUE → Remove large clumps
FALSE
Roundness filter → TRUE → Remove weird shapes
FALSE
Small size filter → TRUE → Remove debris
FALSE

Changes in cell size/shape
Silenced genes

Gene silencing quantiles
Control quantiles

Postprocessing

Analysis

Epistatic effects on cardiomyocyte morphology

d

Cell size (μm)
***

Unaffected (n = 12455)   MYH7-R403Q (n = 21086)

e

Gene silencing efficiency (%)

Unaffected
MYH7-R403Q

TTN, IGF1R, CCDC141, CCDC141-TTN (TTN), CCDC141-TTN (CCDC141), CCDC141-IGF1R (IGF1R), CCDC141-IGF1R (CCDC141)

f

	Unaffected	MYH7-R403Q
TTN	**	*
IGF1R	***	***
CCDC141		***
CCDC141-TTN	***	***
CCDC141-IGF1R	***	***

Relative size difference (%)

g

Relative difference of roundness error (%)

IGF1R
CCDC141
CCDC141-IGF1R

TTN
CCDC141
CCDC141-TTN

Relative difference of norm. peak No. (%)

h

r_i

$$\text{Roundness error} = \frac{\text{STD}[r_i]}{\text{Mean}[r_i]}$$

n_{pk}: No. of peaks (+)
N_{tot}: No. of pixels

$$\text{Norm. peak No.} = n_{pk}/N_{tot}$$

i

Roundness error < 0.08

Roundness error > 0.23

j

Norm. Pk No. < 1%   Norm. Pk No. > 1%