1   **Title.** Derivation and Validation of a Machine Learning Approach to Detect and Mitigate

2   Biases in Healthcare Data

3
Running title. AI for mitigating dataset bias.

Authors. Faris F. Gulamali[1], Ashwin S. Sawant[1,2], Lora Liharska[1], Carol R. Horowitz[3],

Lili Chan[1,2], Patricia H. Kovatch[4], Ira Hofer[1,2,5], Karandeep Singh[6], Lynne D.

Richardson[3], Emmanuel Mensah[7], Alexander W Charney[1], David L. Reich[5], Jianying

Hu[2,8], Girish N. Nadkarni[1,2*]

Affiliations.

1. The Charles Bronfman Institute of Personalized Medicine, Icahn School of Medicine

at Mount Sinai; New York, 10029, USA.

2. The Division of Data Driven and Digital Medicine, The Samuel L Bronfman

Department of Medicine, Icahn School of Medicine at Mount Sinai; New York, 10029,

USA.

3. Institute for Health Equity Research, Icahn School of Medicine at Mount Sinai; New

York, 10029, USA.

4. Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount

Sinai; New York, 10029, USA.

5. Department of Anesthesiology, Perioperative and Pain Medicine, Icahn School of

Medicine at Mount Sinai; New York, 10029, USA.

6. Department of Learning Health Sciences, University of Michigan Medical School, Ann

Arbor, MI

7. Department of Medicine, Beth Israel Deaconess Medical Center / Harvard Medical

School; Boston, 02215, USA.

8. IBM Research, Yorktown Heights, NY

*Corresponding author.

Girish N Nadkarni, MD, MPH

One Gustave L Levy Place, Box 1003,

New York, NY-10029

girish.nadkarni@mountsinai.org

Word count (main text): 5,650

Word count (abstract): 301

Abstract

Background
Broad adoption of artificial intelligence (AI) algorithms in healthcare has led to perpetuation of bias found in datasets used for algorithm training. Methods to mitigate bias involve approaches after training leading to tradeoffs between sensitivity and specificity. There have been limited efforts to address bias at the level of the data for algorithm generation.

Methods
We generate a data-centric, but algorithm-agnostic approach to evaluate dataset bias by investigating how the relationships between different groups are learned at different sample sizes. We name this method AEquity and define a metric AEq. We then apply a systematic analysis of AEq values across subpopulations to identify and mitigate manifestations of racial bias.

Findings
We demonstrate that AEquity helps mitigate different biases in three different chest radiograph datasets, a healthcare costs dataset, and when using tabularized electronic health record data for mortality prediction. In the healthcare costs dataset, we show that AEquity is a more sensitive metric of label bias than model performance. AEquity can be utilized for label selection when standard fairness metrics fail. In the chest radiographs dataset, we show that AEquity can help optimize dataset selection to mitigate bias, as measured by nine different fairness metrics across nine of the most frequent diagnoses and four different protected categories (race, sex, insurance status, age) and the intersections of race and sex. We benchmark against approaches currently used after algorithm training including recalibration and balanced empirical risk minimization. Finally, we utilize AEquity to characterize and mitigate a previously unreported bias in mortality prediction with the widely used National Health and Nutrition Examination Survey (NHANES) dataset, showing that AEquity outperforms currently used approaches, and is effective at both small and large sample sizes.

Interpretation
AEquity can identify and mitigate bias in known biased datasets through different strategies and an unreported bias in a widely used dataset.

Summary
AEquity, a machine learning approach can identify and mitigate bias the level of datasets used to train algorithms. We demonstrate it can mitigate known cases of bias

75   better than existing methods, and detect and mitigate bias that was previously

76   unreported.

EVIDENCE IN CONTEXT

Evidence before this study

Methods to mitigate algorithmic bias typically involve adjustments made after training, leading to a tradeoff between sensitivity and specificity. There have been limited efforts to mitigate bias at the level of the data.

Added value of this study

This study introduces a machine learning based method, AEquity, which analyzes the learnability of data from subpopulations at different sample sizes, which can then be used to intervene on the larger dataset to mitigate bias. The study demonstrates the detection and mitigation of bias in two scenarios where bias had been previously reported. It also demonstrates the detection and mitigation of bias the widely used National Health and Nutrition Examination Survey (NHANES) dataset, which was previously unknown.

Implications of all available evidence

AEquity is a complementary approach that can be used early in the algorithm lifecycle to characterize and mitigate bias and thus prevent perpetuation of algorithmic disparities.

98   **MAIN**

99   **INTRODUCTION**
100
The adoption of diagnostic and prognostic algorithms in healthcare has led to

perpetuation of bias against underrepresented groups. These biases exist even in

algorithms trained using diverse populations, because longstanding societal biases and

disparities are reflected in the data used to develop algorithms [1].

Attempts to mitigate bias have focused on fairness-metric based algorithm

optimization after training on the data [2–4]. Recent developments in generative algorithms

have focused on amplifying representation [5]. However, if the underlying dataset is

biased, algorithms will exhibit residual unfairness or performance tradeoffs even after

algorithm optimization or synthetic data generation [6–9]. Second, many forms of implicit

bias are not captured by fairness metrics or algorithm performance, and detecting these

biases is difficult [6,10]. There is increasing recognition of the need to address bias earlier

in the algorithm development pipeline, when data is collected and curated or when

outcomes are chosen [11–13].

We present AEquity, a machine learning-based method, which is used to detect

and evaluate bias and perform an appropriate intervention to mitigate it using a small

number of training samples. Since this technique is agnostic of the machine learning

algorithm and architecture, it can be applied in a variety of settings. We first

demonstrate its application to detect and mitigate bias in two scenarios wherein

healthcare algorithms have been shown to be biased: 1) an algorithm used to predict

healthcare costs which exhibits bias against Black patients [10] (healthcare costs

scenario), and 2) a standard computer vision algorithm applied to chest radiographs

results in selective underdiagnosis in people who are Black, Non-White Hispanic/Latino,

or Medicaid [14] (chest radiographs scenario). Finally, we utilize AEquity to detect,

characterize and mitigate a previously unreported, undetected bias in prediction of

mortality with the National Health and Nutrition Examination Survey dataset (NHANES

scenario)[15]. We provide an easy-to-use implementation and release this for use without

any restrictions (**Supplementary Methods**).

Generally, the task can be described as: We have a subset of data containing

two distinct groups distinguished by a characteristic (like race), with features x and

potential outcomes y. The first task is to determine which outcome (or label) should be

considered to minimize residual bias towards a group, where residual bias is after using

methods after training[15]. We demonstrate optimal label selection to minimize bias using

the healthcare costs scenario. The next task is to determine which samples from the

dataset need to be prioritized to reduce differences in performances against, or in favor

of, a group based on a characteristic. We demonstrate guided sample selection using

the chest radiographs scenario and demonstrate that this works for algorithms based on

both convolutional neural network and transformer architectures, which form the

foundation of computer vision. Finally, we demonstrate the use of AEquity to

sequentially perform label selection and guided sample selection in the NHANES

scenario, where prior biases have not been reported. Thus, we demonstrate a task-,

algorithm- and architecture-agnostic method for data-centric debiasing, potentially

usable beyond healthcare tasks. We make it open source with documentation and a

video tutorial to increase accessibility.

METHODS

For clarity, we use "groups" to refer to various subpopulations, for example by race, gender, or insurance status. We will use "outcome" to refer to diagnostic findings, for example "pneumonia" and "edema". A "protected group" refers to categories of people who are legally protected from discrimination under US law. It includes, among other things, age, sex, and race. Following an accepted framework [1], we characterize dataset biases into main types: sampling bias; complexity bias; and label bias (Figure 1). Sampling bias arises when there is a low frequency of patients with a given diagnosis from a protected group. In complexity bias, a protected group presents more heterogeneously with a diagnosis, and consequently, one group exhibits a class label with greater complexity than another group. Subsequently, even if groups are drawn with equivalent frequencies, generalization performance on the more complex group is worse than other groups for the same class. For example, this could cover a group for whom the diagnostic criteria aren't precise due to historical reasons, a group with large translational distribution shift due to geographic differences, or a group for which there are differences in data quality. In each of these three cases, the complexity of the underlying data-distribution is expected to increase. Third, label bias occurs when outcome labels are placed incorrectly at different rates for different groups and can lead to increased misclassification errors for the affected group. For example, in published work, Black patients are more frequently labeled as requiring fewer healthcare resources despite having the same number of comorbidities because of lack of access to care [10].

We develop and validate AEquity, a machine learning-based method, that can be

used for the characterization of these three biases (sampling, complexity, and label),

followed by guided mitigation, in healthcare datasets. AEquity works by appending a

simple compressive network to a dataset or the latent space of existing machine

learning algorithms to generate a single value, AEq, characterizing an outcome with

respect to a group and the priors established by the algorithm. In general, a higher AEq

value for a group with respect to an outcome and a given machine learning algorithm

suggests that generalization is more difficult and more or higher quality samples are

needed for learning [19,20]. Thus, AEq is a metric that is calculated at the intersection of

an outcome like "pneumonia", a group label like "Black", and a machine learning

algorithm like "ResNet-50." AEq is reported as $\log_2$(sample size estimate). AEq values

are only meaningful when constrained to comparing groups with the same outcome in

the dataset, and interpretation of the metric is invariant to architecture of the machine

learning algorithm, the compressive network or the dataset type or characteristics [17–19].

Related mathematical work has sought to characterize the capacity of neural networks

with topological data analysis and approximations of inductive bias complexity, which

we do by utilizing a small number of samples to establish empirical approximations of

relationships between groups [20,21].

To mitigate a specific type of bias, the AEq values can be used to evaluate the

right composition of the dataset for training a machine learning algorithm, or for label

selection without changing the dataset composition. To evaluate bias mitigation, we

evaluated an extensive variety of fairness metrics (area-under-the-curve, true positive

rate, false negative rate, true negative rate, positive predictive value, false positive rate,

false discovery rate, and precision) and showed that AEquity can help improve these

metrics over just algorithm optimization. While previous work has sought to optimize for

these metrics, these works are algorithm-centric and optimization based, rather than

modifications of the underlying dataset. Previous methods that have utilized information

bottleneck theory make no changes to the dataset nor identify which label to best select,

and therefore lead to a tradeoff between overall accuracy and fairness, which is true of

any algorithm-centered approach [22,23]. Our approach is fundamentally different –

engineering the dataset used for training and selecting appropriate labels so that

fairness is improved without sacrificing accuracy. While data-centric approaches in the

literature have been discussed [7,24], state-of-the-art remains balanced empirical risk

minimization [12], which performs at the same level as or worse than AEquity. Moreover,

AEquity functions at a significantly smaller sample size (3-10% of the data). As a result,

AEquity can be used early in the algorithm development pipeline and can be

complementary to downstream optimizations.

<u>Description of AEquity based data curation</u>

AEquity-based data curation is achieved through a multi-step pipeline (Figure 2).

First, a group-balanced subset of the patient data is collected, and this data is labeled.

Next, AEq values by group for each outcome are calculated. The difference in the AEq

values between groups for each outcome can help us determine if label bias exists in

the dataset. Fairness metrics are not sensitive measures of label bias because

algorithms can learn the relationship between specific patient populations and a given

outcome, even if the outcome is systemically distorted [6,10]. However, if AEq measures

are significantly different between groups for the same label, then the label may not

represent the same distribution for different groups. The label with least distributional

difference (smallest difference in AEq values between groups) should be selected to

mitigate the bias. Next, AEq is calculated with respect to each outcome when groups

are combined into a joint dataset. The relationship between individual group and joint

AEq values allows us to determine if sampling or complexity bias exists, and the

differences in AEq values for different patient groups for the same label are an indicator

of label bias. When predictive bias is driven primarily by sampling bias in the dataset,

combining groups drives the AEq value at or below the over-sampled data

(**Supplementary Methods**). In sampling bias, balanced sampling from each group is

sufficient to mitigate the bias because the data is equitably represented within the

algorithm. When complexity bias was the only type of dataset bias, combining the

groups results either in an increase in the value of AEq or an AEq closer to value that

had been higher prior to the combination. In complexity bias, collecting data exclusively

from the protected population, due to its relative heterogeneity compared to the over-

represented group, is necessary to mitigate bias. If the algorithm exhibits residual

unfairness, then the AEq will be different for two groups for the same outcome. Thus,

AEquity can help with quantifying biases within outcome metrics and subsequently

selecting appropriate outcomes to minimize AEq values between groups.

<u>Machine Learning and Statistical Analysis of AEquity based data curation</u>

For all machine learning tasks, the training, validation and test set are kept independent

with no overlapping patients. The training and validation sets were seeded and

bootstrapped fifty times. AEquity was applied to a small subset of the training data for all

three tasks (< 10%). The mean and standard errors are calculated across all fifty

bootstraps and 95% confidence intervals are reported. P-values are calculated via t-

tests for two-group quantitative analyses and ANOVA for multi-group quantitative

variant analyses. All reported P-values are adjusted with Bonferroni correction.

When training machine learning algorithms, all algorithms are trained on an

independent training set, validated on a validation set, and performances are reported

on an independent test set (10%). The test set consists of held-out (non-overlapping)

samples prior to any dataset modification or intervention. Training and validation sets

are bootstrapped fifty times, and performance is reported on the held-out test set.

Means and 95% confidence interval are reported.

RESULTS

<u>Detection and Mitigation of Racial Bias from Systematic Label Distortion in Electronic</u>

<u>Healthcare Record Datasets at Small Sample Sizes (healthcare costs scenario)</u>

In this example, we highlight the role of AEquity when labels for a particular

group are systematically distorted for a given patient population. Obermeyer et al.

showed that a algorithm's predictive power is not a sensitive metric for bias. In other

words, an algorithm trained on labels that are significantly biased against Black patients

has an accuracy comparable to an algorithm trained with more equitable labels.

Moreover, despite equivalent predictive power, the algorithm trained on biased labels

produces biased outcomes. We examined whether AEquity can detect the biased label

and help select a more equitable label.

We examined a publicly available tabularized EHR dataset of 49,618 primary

care patients enrolled in risk-based contracts from 2013 to 2015, from a large academic

hospital, described in Obermeyer et al[1]. Data included demographics and visit

information. In that paper, the authors examined an algorithm which predicted total

healthcare costs for a patient as a proxy for their predicted healthcare needs. They

categorized individuals at or above the 50th percentile of predicted outcome as 'high

risk' and these patients would be screened for a care management program, whereas

patients at the 95th percentile would be automatically enrolled. They found that Black

patients screened for the high-risk category were significantly sicker compared to White

patients in that category. This is a form of bias because it resulted in a reduced

allocation of resources to Black patients in comparison with their healthcare needs.

They found that this discrepancy persisted when patients were categorized based on

another predicted cost-based metric-avoidable costs. However, the discrepancy

disappeared if predicted active chronic conditions were used for categorization. In short,

they discuss in their paper that using a better label can mitigate bias. With AEquity, we

can empirically replicate these conclusions quantitatively at small sample sizes, and

subsequently guide the development of minimally biased algorithms prior to complete

data collection, algorithm training and validation.

AEquity-based data curation was achieved through a multi-step pipeline. First, a

race-balanced subset of the patient data was collected (n=1,024, 3% of the total

available data) and this data was labeled according to each of the available labels

(active chronic conditions, avoidable costs, and total costs). We calculated AEq by race

(Black, White), for each risk category (high risk, low risk), for each of the three

outcomes (total costs, avoidable costs, active chronic conditions) (Table 1). To

determine if Black and White patients in each risk category were similar, we then

calculated the difference in AEq between races, for each risk category, for each

outcome.

First, the difference in AEq was statistically significant across outcome measures

for the population designated as high-risk (ANOVA, $P = 1.59 \times 10^{-8}$). When predicted

cost-based metrics (total costs, avoidable costs) were used to designate patients at

highest risk, there was a significant difference in AEq between Black and White patients

(Figure 3A, 3B; Table 1). For example, with total costs, the difference was 0.47 (95% CI,

0.36-0.52). Thus, Black patients predicted to be at high risk were significantly different

from their White counterparts which led to inequitable resource allocation.

However, the difference between Black and White patients categorized to be at

high risk disappeared when using predicted active chronic conditions (instead of

predicted costs) as the outcome measure; the difference in AEq was not statistically

significantly different from 0 [0.06 (95% CI, -0.04 to 0.16)], indicating that Black and

White patients in that risk category were similar. Thus, AEq values could *a priori* guide

the choice of a better outcome measure to identify patients at high risk and thus mitigate

bias.

We conducted additional analyses in patients at lower risk and found that the

differences in AEq across races were all statistically significant (Figure 3B). In the low-

risk group, Black patients display more heterogeneous underlying characteristics than

White patients. We also observed that the differences in the AEq metric were smaller

when using active chronic conditions than avoidable costs or total costs, indicating

mitigation of bias for this lower-risk group despite the increased heterogeneity (Table 1).

Nevertheless, the presence of differences in AEquity-score in the low-risk group may

indicate that there may be some residual complexity or sampling bias even after

changing the outcome metric.

Detection at Small Sample Sizes and Mitigation of Under-diagnosis Bias in Chest X-ray

(chest radiographs scenario)

We then evaluated AEquity on MIMIC-CXR, a publicly available dataset

containing 377,110 chest radiographs of 65,379 patients from the Beth Israel

Deaconess Medical Center between 2011 and 2016 [25]. We also used two additional

publicly available chest radiograph datasets to validate our findings – CheXpert

containing 224,316 chest radiographs from 65,240 patients, and ChestX-ray14

containing 108,948 chest radiographs from 32,717 patients [26,27] . MIMIC-CXR and

CheXpert each contain 14 binary diagnostic findings, ChestX-ray14 contains 15, and all

three contain an additional binary label for radiographs with no positive diagnostic

finding. (Supplementary Table 1) [26,27]. We generated a subset from MIMIC-CXR by

including only postero-anterior radiographs of patients who self-identified as either Black

or White, and with either one of the nine most frequently encountered diagnostic

findings (atelectasis, cardiomegaly, consolidation, edema, effusion, enlarged

cardiomediastinum, opacity, pneumonia, pneumothorax) or 'no finding'. In Seyyed-

Kalantari et al. [27–29], a convolutional neural network was trained on Chest X-rays

belonging to the MIMIC-CXR dataset and evaluated on protected populations. The

algorithms demonstrated significant differences in performance on protected

populations, indicating bias. Subsequent work has shown that fairness methods which

strive to achieve better worst-group performance (GroupDRO, JTT, CRT, DFR, etc.) do

not outperform balanced empirical risk minimization (balanced ERM) [2,12,28,29]. Thus, we

compared AEquity to balanced ERM.

We followed a multi-step pipeline for AEquity-based data curation. First, we

collected a balanced subset of the patient data (n=1,024, 3% of the total available data,

512 Black patients, 512 white patients) and labeled it according to each of the available

labels. For each diagnostic finding, we calculated AEq by race (Black, White) and also

for the joint dataset containing samples from both Black and White patients. We then

used these AEq values to guide data curation - choosing either a) to group-balance

with an equal number of additional samples from both Black and White patients, or b) to

prioritize data collection of a single group to a total of 30,000 samples, a sample size

sufficient to produce performance comparable to state-of-the-art classification

algorithms (Table S2). We then demonstrated that AEquity can help analyze causes of

bias at both small and large dataset sizes but becomes more precise at larger sample

sizes (**Supplementary Figure S7**). We split the subset 60%-20%-20% into training

samples, validation samples, and test samples, ensuring no patients overlapped, and

fine-tuned two different pre-trained machine learning algorithms (ResNet-50, VIT-B-16)

with PyTorch Lightning [30–32]. First, we compared these algorithms to state-of-the-art

Chest X-ray classification algorithms (Table S2). Next, we showed how AEquity can

improve performance over naive data collection (Figure 4, Table 2), and benchmarked

against Balanced ERM, which has previously been shown to outperform other variants

of subgroup characterization such as Just-Think-Twice, GroupDRO [30–32] (Table 3,

Figure S6). Balanced empirical risk minimization counterbalances varying population

frequencies with over-weighing or over-sampling under-represented groups, and while

this is a valid strategy for sampling bias, AEquity demonstrates that in many instances,

groups exhibit complexity bias or label bias, which require alternative strategies. We

validated against threshold-independent metrics such as area-under-the-curve as well

as fairness metrics such as sensitivity, specificity, FPR, FDR, FNR in intersectional

populations (Table S3). We bootstrapped each experiment 50 times, in keeping with

available computational resources [33–35]. We defined bootstrapping as sampling with

replacement, but avoided sampling patients and their chest X-rays from the test dataset.

First, we observed that a higher frequency of occurrence for a diagnosis

corresponded to a lower AEq value across Black and White patients (Figure 4c). This

indicates that labels with more samples, like lung opacity (AEq ≈ 7.5), generally trend

towards lower AEqs, whereas labels with fewer samples such as consolidation (AEq ≈

9.8), enlarged cardio-mediastinum (AEq ≈ 9.9) or pneumothorax (AEq ≈ 9.95) trend

towards having higher AEqs. We posit that the latter group of diagnoses may have the

highest potential for bias because of difficulty in generalization samples from patients

not present in the training dataset. Second, we noticed that six out of the nine

diagnoses, accounting for 71% of the positive samples, demonstrate higher AEq values

in Black patients than in White patients. Higher AEq values indicate that, in general,

algorithms built on chest X-rays from Black patients for Black patients generalize more

poorly than algorithms trained on chest X-rays from White patients for the purpose of

diagnosing White patients. In other words, more samples from Black patients than

White patients are required to mitigate bias.

Next, we examined pneumothorax. Chest X-rays from Black individuals

diagnosed with pneumothorax have significantly larger AEq values than chest X-rays

belonging to White individuals with the same diagnosis ($AEq_{Black, Pneumothorax}$ = 10.10;

95% CI: (10.01, 10.18), $AEq_{White, Pneumothorax}$ = 9.50; 95% CI: (9.44, 9.58); $P < 0.05$). For

the diagnosis of pneumothorax, the AEq of the joint dataset is smaller than the AEq of

chest X-rays belonging to each subgroup, which implies that ($AEq_{Joint, Pneumothorax}$ = 9.44,

95% CI: (9.35, 9.52)). Subsequently, we see that adding dataset diversity by equitably

sampling from each race was associated with an improvement in classifier performance

for Black individuals when compared to the naïve approach (Bias Reduction (Absolute)

= 2.43×10^{-2}; 95% CI: (2.22×10^{-2}, 2.59×10^{-2}); Bias Reduction (%) = 49.37%, 95% CI:

(49.3, 49.5)). Second, we examined edema, where the joint AEq was higher than the

AEq for either race, consistent with complexity bias. For edema, AEq_{Joint} is larger than

both AEq_{Black} and AEq_{White} ($AEq_{Joint, Edema}$ = 9.48; 95% CI: (9.37, 9.58), $AEq_{Black, Edema}$ =

8.78; 95% CI: (8.69, 8.86); $AEq_{White, Edema}$ = 9.12; 95% CI: (9.08, 9.20), $P < 0.05$).

Prioritizing data collection from Black patients better captured the group-specific

presentations, and this strategy was associated with an algorithm that generalized

better to Black patients when compared to a naïve approach (Bias Reduction (Absolute)

= 1.95×10^{-2}; 95% CI: (1.90×10^{-2}, 2.01×10^{-2}); Bias Reduction (%) = 60.02%, 95% CI:

(59.98, 60.06)). Bias for each diagnostic finding in the chest radiograph dataset

decreased by between 29% and 96.5% following intervention (Table 2; Figure 4D).

Finally, we demonstrated that 1) AEquity remains valid for mitigation of bias by

sex, 2) AEquity-guided data curation is effective for individuals at the intersection of

more than one disadvantaged group, 3) targeted data collection and curation is robust

to choice of fairness metric, and 4) these results are robust when trained on transformer

architectures with 86 million parameters. When evaluating sex, the joint AEq score for

all 9 diagnostic findings was not significantly different from, or lower than the AEq score

for females, indicating that predictive biases are primarily driven by sampling bias

(Figure S1). Larrazabal et al [12] has previously demonstrated that group balancing is the

optimal approach to mitigate gender biases in this setting. Second, we demonstrate the

effectiveness of AEquity on intersectional populations, who are at greater risk of being

affected by systemic biases [36,37], and we show that this technique is robust to the choice

of fairness metrics. When we examined Black patients on Medicaid, at the intersection

of race and socioeconomic status, we found that AEquity-based interventions reduced

overall false negative rate by 33.3% (Bias Reduction Absolute = 1.88×10^{-1}; 95% CI

(1.4×10^{-1}, 2.5×10^{-1}); Bias Reduction (%) 33.3% (95% CI, 26.6-40.0)), Precision Bias by

7.50×10^{-2}; 95% CI (7.48×10^{-2}, 7.51×10^{-2}); Bias Reduction (%) 94.6% (95% CI, 94.5-

94.7%); False Discovery Rate by 94.5% (Absolute Bias Reduction = 3.50×10^{-2}; 95% CI:

(3.49×10^{-2}, 3.50×10^{-2}) (Figure S4). Finally, we demonstrated AEquity works on

VIT_B_16, an 86 million parameter Vision Transformer ([Figure S6](#)). AEquity

outperformed balanced ERM across every diagnostic finding when measuring bias and

resulted in reductions of up to 55% in bias compared to the originaldataset ([Table S5](#)).

Notably, in four of the nine diagnostic findings (opacity, effusion, edema and enlarged

cardiomediastinum), analysis of AEquity values resulted in the curation of a dataset

reflecting the underlying empirical group frequencies. However, in the remaining five

(cardiomegaly, pneumonia, pneumothorax, and consolidation), AEquity led to a dataset

that prioritized one group, outperforming balanced ERM for the mitigation of bias[38].

We do not examine label bias due to unavailability of gold-standard labels

besides those already present in publicly available dataset. However, Zhang et al have

previously shown that, at least for the 'No Finding' label in the chest radiograph dataset,

the prevalence of incorrect labeling did not vary significantly by race and sex, but that it

was more prevalent in the subgroup of patients over age 80 [38–40].

Detection of Previously Unknown Bias in the National Health and Nutrition Examination

Survey for predicting all-cause mortality (NHANES scenario)

The National Health and Nutrition Examination Survey (NHANES) data was

collected between 1999 and 2014, and includes demographic, laboratory, examination

and questionnaire features. The NHANES dataset has been used for a wide range of

applications including the measurement of the population prevalence of metabolic

syndrome, demographic trends in major depressive disorder, and associations between

factors such as inflammation and autoimmune diseases [39–41]. Machine learning has

been used on the NHANES dataset to predict coronary heart disease, hypertension,

and all-cause mortality [42–44].

The prediction of all-cause mortality is an important task because all-cause

mortality is a commonly chosen outcome in epidemiological studies [45]. All-cause

mortality can be used as a metric for health disparities, to guide public health

interventions, and to affect health resource allocation [46–48]. Therefore, detecting,

characterizing, and mitigating biases in the prediction of all-cause mortality can

potentially affect large scale epidemiological interventions. To evaluate biases in the

prediction of all-mortality in the NHANES dataset, we pre-processed the data

(N=47,261) and selected variables as described in past work by Qiu et al [44]. To detect

potential biases, we calculate AEq values at small sample sizes (N=400) for Black and

White individuals, and all-cause mortality for different follow up periods in the NHANES

dataset (Figure 5A). Next, we qualitatively and quantitatively analyzed the AEquity

values to identify the types of biases in the dataset. Finally, we used AEquity to guide

data-centric intervention, and compared this intervention with simple algorithmic

calibration and balanced ERM. All experiments were bootstrapped 50 times with the

Pytorch library.

First, we evaluated for label bias by calculating the difference in AEquity score

between Black and White patients for all-cause mortality at different follow-up periods

(Figure 5B). AEquity scores for Black and White patients are least significantly different

at 5-year mortality ($\text{AEq}_{\text{Difference, 5-year-mortality}}$ = 0.60; 95% CI: (0.52, 0.68); $\text{AEq}_{\text{Difference, 1-}}$

$_{\text{year-mortality}}$ = 1.00; 95% CI: (0.83, 1.17); $P < 0.05$). Next, we evaluated for complexity and

sampling bias based on 5-year-mortality and race (Figure 5C). The AEq value for the

joint distribution was significantly larger than the AEq values for either Black or White

patients, signaling complexity bias ($\text{AEq}_{\text{Joint, 5-year-mortality}}$ = 7.00; 95% CI: (6.93, 7.07);

AEq$_{White, 5\text{-}year\text{-}mortality}$ = 7.00; 95% CI: (6.93, 7.07); AEq$_{Black, 5\text{-}year\text{-}mortality}$ = 6.6; 95% CI:

(6.53, 6.66); *P < 0.05*).

Based on the results from AEquity, we curated a dataset with prioritized data

from Black patients, trained a gradient-boosted tree to predict 5-year-mortality, and

compared its fairness metrics to that of a gradient boosted tree trained to predict 1-year-

mortality with a dataset sampled based on population prevalence. First, AEquity

significantly reduced bias as measured by the difference in area-under-the-curve (AUC)

between Black and White patients, and outperformed both calibration and balanced

ERM (Figure 5D). Second, label selection and dataset prioritization had synergistic bias

mitigation effects (Figure 5D). Third, AEquity's bias mitigation relative to training with

random sampling from the original dataset is invariant to sample size (Figure 5E).

DISCUSSION

Algorithmic bias is one of the major challenges to the adoption of AI in healthcare

and is a focus for regulation. For example, the Good Machine Learning Practice

guidance from the FDA and EMA emphasizes the importance of ensuring that datasets

are representative of the intended patient population [49]. However, this is not sufficient

because significant bias can result from the choice of outcome or differences in group

complexity even when datasets are diverse. AEquity, the machine-learning based

technique described in this paper, can help characterize and mitigate bias early in the

algorithm development lifecycle.

The responsibility for ensuring algorithmic equity is being placed on algorithm

developers and institutions. The US Department of Health and Human Services has

proposed a rule under Section 1557 of the Affordable Care Act to ensure that "a

covered entity must not discriminate against any individual on the basis of race, color,

national origin, sex, age, or disability through the use of clinical algorithms in decision-

making". More recently, an executive order has placed at least some of the liability for

algorithmic bias on developers and deployers of these algorithms [50]. Therefore, the

need to develop tools to identify and remediate potential bias in healthcare datasets is

urgent from both ethical and regulatory perspectives.

The algorithm development pipeline involves task identification, data collection,

data curation, training, validation, and l deployment, each of which plays an important

role in whether it adds value to the healthcare system or propagates disparities that it

seeks to address [51–54]. The ability of this approach to quantify and address bias at small

sample sizes makes it an attractive addition to the toolset, akin to sample size

determination but for bias [54]. Nevertheless, data-centric curation is only one of many

approaches in which bias can be mitigated [55].

Currently, the most prevalent solution for bias mitigation with data-centric

approaches is the comparison of predictive power across protected groups and

balanced ERM. We have demonstrated the application of AEquity to datasets where

biases have been confirmed and described in high-quality peer reviewed studies, but

application of the current state-of-the-art method falls short. Algorithms predicting all

three labels in the healthcare costs scenario of Obermeyer et al have similar predictive

power, but two of the three are biased, demonstrating that equal predictive power is not

a reliable indicator of the absence of bias. AEquity can identify the biased labels by

examining a small subset of the data. Second, we demonstrated in the chest radiograph

scenario that the optimal mitigation of bias sometimes requires prioritization of data from

the minority population and at other times, it requires a larger, more balanced dataset.

AEquity can help distinguish these two scenarios early in the data collection process,

because it only requires a small amount of data for analysis. Third, we use AEquity to

describe and mitigate a previously undetected and unreported bias in all-cause mortality

prediction based on data from the National Health and Nutrition Examination Survey

(NHANES). Applying AEquity to the NHANES dataset can detect the bias at small

sample sizes and help optimize label selection and dataset curation to mitigate bias

across a range of sample sizes.

Data-guided strategies face challenges due to the lack of representation in many

datasets [56,57]. Moreover, once a gap has been identified, targeted data collection of

protected populations raises ethical concerns given the long history of mistrust [58–63].

Nevertheless, acknowledging the existence of these biases through quantitative

measures can build trust, a crucial step towards mitigating some of these biases at a

grassroots level [64,65]. Investigating the reason why AEq scores are different between

different populations may provide key insights into underlying mechanisms of racial

inequities [66,67]. More formally, US regulatory guidelines require that AI algorithms in

healthcare not be biased against protected classes [68]. The collection of appropriate data

to test this against protected classes is obligatory even if it is difficult. AEq can help

guide sampling strategies early in the data collection process because it needs

relatively few samples.

Our examination of AEquity builds on previous work examining bias in

healthcare. As with many machine learning approaches and clinical studies, AEquity is

empirical, and no theoretical proof exists. In past work, we highlighted that the sample

size at which the training loss curve of an autoencoder converges corresponds to the

sample size at which a training loss curve for a neural network classifier trained on the

same data converges. We showed that this holds in a variety of computer vision

datasets and on latent spaces belonging to a variety of structures, which included

varying the number of classes, number of informative features, number of total features,

number of class-specific subgroups, and the relationships between different variables

within the dataset [68,69]. In this work, we build on that approach to the problem of bias

mitigation in healthcare. Here, we do not use it to predict sample size for training a

neural network. Rather, we show that it can be used as a simple marker of the

learnability of data subsets and to further guide dataset intervention which can reduce

bias of classifiers trained on the data. However, because no theoretical proof exists, its

performance should be verified when applying it to a new setting. Nevertheless, the

demonstration of AEquity in multiple state-of-the-art algorithms, multiple tasks and

multiple metrics provides strong validation for its use in practice.

While our results are promising, there are some limitations to AEquity and

directions for future inquiry. First, we primarily focus on three case examples –

underdiagnosis, under-allocation of resources and unreported bias for outcome

prediction. However, validation of AEquity on a broader scale across different

phenotypes, protected groups, data modalities and algorithms types are essential.

Second, AEq in its current form focuses primarily on classification tasks. A growing

number of algorithms, however, are: a) using various forms of regularization and

attention to produce more informative latent spaces [69,70] ; b) being used for generation

rather than simply classification or regression [71] and c) are multi-modal [72]. Extending

AEq to these latent spaces and generative algorithms may help investigate and mitigate

bias in those settings. However, generative algorithms are not substitutes for

appropriate data curation, [5] and generative algorithms themselves can contain biases

[73–75]. For example, the use of diffusion algorithms in conjunction with data-centric

strategies may yield compounding effects [5]. Third, the labels for the MIMIC-CXR

dataset were obtained using the CheXpert system of natural language processing of

radiology reports, which is known to have a substantial and varying amount of

mislabeling for different diagnoses [69]. Results from Zhang et al suggest there is no

statistically detectable label bias by race in MIMIC-CXR for the 'no finding' label [72].

Fourth, we cannot exclude the possibility of label bias related to the rate of mislabeling

being different for different groups for other diagnoses. AEquity can help choose a less

biased label, but the cost of labeling healthcare data often precludes the availability of

alternative, high-quality labels. The application of frameworks for dataset development

which include expert labeling of the most informative samples may help overcome this

problem [76]. We also posit that identification of the type of bias can assist in identifying

the other origins of bias. For example, AEquity can help characterize the bias as

primarily complexity-related, which may lead an investigator to hypothesize that it may

be due to more advanced disease due to lack of access to care. However, we concede

that AEquity cannot confirm that lack of access to care is indeed the cause. Similarly, in

the case of label bias, one may hypothesize that structural racism may contribute to the

decision to focus on labels which perpetuate inequities. However, a method like

AEquity, which looks at patient-level predictors and outcomes, is not expected to be

able to confirm such a hypothesis.

In summary, we present AEquity, a task-, algorithm-, and architecture-agnostic

machine learning based metric that may be valuable for disentangling and quantifying

various types of biases at the dataset level. We demonstrate the application of AEquity

to guide specific, effective mitigation measures in datasets where biases have been

confirmed and described in high-quality peer-reviewed studies. We demonstrate its

robustness by applying it to different datasets and algorithms, intersectional analyses

and measuring its effectiveness with respect to a range of traditional fairness metrics.

Finally, we use it to detect and mitigate label and complexity bias in a dataset where

bias had not been described before. Although we demonstrate its utility in the urgent

use case of algorithmic bias in healthcare, this approach can be used to address biases

across the spectrum of algorithmic applications.

Acknowledgements

The authors would like to acknowledge Jill Gregory for support in illustrating the figures

and the staff at the Minerva High Performance Computing Cluster for maintaining the

servers used to run the experiments.

Funding:

The study was funded by the National Center for Advancing Translational Sciences

(NCATS) and the National Institutes of Health (NIH). NCATS and NIH had no role in

study design, data collection, data analysis, preparation of the manuscript, or the

decision to submit for publication.

Data and materials availability:

The publicly available images can be access via the PyTorch library ([torchvision — Torchvision main documentation (pytorch.org)](#)). The MIMIC-CXR version 2.0 dataset is publicly available by registration at https://physionet.org/content/mimic-cxr/2.0.0/. The CheXpert dataset is publicly available by registration at https://stanfordmlgroup.github.io/competitions/chexpert/. The NIH-CXR dataset is publicly available at https://nihcc.app.box.com/v/ChestXray-NIHCC. The dataset used in the Obermeyer analysis is available in the supplement ([Dissecting racial bias in an algorithm used to manage the health of populations | Science](#)). The code for the analyses are available here ([Nadkarni-Lab/AEquitas: Deep Learning Based Metric to Mitigate Dataset Bias (github.com)](#)), and the instructions for usage are included in the supplement.

Author contributions:

Conceptualization: FFG, ASS, GNN

Methodology: FFG, ASS, GNN

Investigation: FFG, ASS, GNN

Visualization: FFG

Funding acquisition: GNN

Project administration: ASS, GNN

Supervision: DLR, GNN, AWC

Writing – original draft: FFG, ASS, GNN

Writing – review & editing: LL, CRH, LC, PHK, IH, KS, AWC, JH

References

1. Mehrabi, N., Morstatter, F., Saxena, N., Lerman, K. & Galstyan, A. A survey on bias and fairness in machine learning. *ACM Comput. Surv.* **54**, 1–35 (2021).

2. Zhang, H. *et al.* Improving the Fairness of Chest X-ray Classifiers. *arXiv [cs.LG]* (2022).

3. Bellamy, R. K. E. *et al.* AI Fairness 360: An extensible toolkit for detecting and mitigating algorithmic bias. *IBM J. Res. Dev.* **63**, 4:1-4:15 (01 July-Sep 2019).

4. Rajkomar, A., Hardt, M., Howell, M. D., Corrado, G. & Chin, M. H. Ensuring Fairness in Machine Learning to Advance Health Equity. *Ann. Intern. Med.* **169**, 866–872 (2018).

5. Ktena, I. *et al.* Generative models improve fairness of medical classifiers under distribution shifts. *Nat. Med.* 1–8 (2024).

6. Kallus, N. & Zhou, A. Residual Unfairness in Fair Machine Learning from Prejudiced Data. in *Proceedings of the 35th International Conference on Machine Learning* (eds. Dy, J. & Krause, A.) vol. 80 2439–2448 (PMLR, 10--15 Jul 2018).

7. Petersen, E., Holm, S., Ganz, M. & Feragen, A. The path toward equal performance in medical machine learning. *Patterns (N Y)* **4**, 100790 (2023).

8. Petersen, E., Ganz, M., Holm, S. & Feragen, A. On (assessing) the fairness of risk score models. in *Proceedings of the 2023 ACM Conference on Fairness, Accountability, and Transparency* 817–829 (Association for Computing Machinery, New York, NY, USA, 2023).

9. Dolata, M., Feuerriegel, S. & Schwabe, G. A sociotechnical view of algorithmic fairness. *Inf. Syst. J.* **32**, 754–818 (2022).

10. Obermeyer, Z., Powers, B., Vogeli, C. & Mullainathan, S. Dissecting racial bias in

an algorithm used to manage the health of populations. *Science* **366**, 447–453

(2019).

11. Rich, A. S. & Gureckis, T. M. Lessons for artificial intelligence from the study of

natural stupidity. *Nature Machine Intelligence* **1**, 174–180 (2019).

12. Larrazabal, A. J., Nieto, N., Peterson, V., Milone, D. H. & Ferrante, E. Gender

imbalance in medical imaging datasets produces biased classifiers for computer-

aided diagnosis. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 12592–12594 (2020).

13. Jain, S., Smit, A., Ng, A. Y. & Rajpurkar, P. Effect of Radiology Report Labeler

Quality on Deep Learning Models for Chest X-Ray Interpretation. *arXiv [eess.IV]*

(2021).

14. Seyyed-Kalantari, L., Zhang, H., McDermott, M. B. A., Chen, I. Y. & Ghassemi, M.

Underdiagnosis bias of artificial intelligence algorithms applied to chest radiographs

in under-served patient populations. *Nat. Med.* **27**, 2176–2182 (2021).

15. National Center for Health Statistics (NCHS). National Health and Nutrition

Examination Survey Data. *Centers for Disease Control and Prevention (CDC)*.

16. Gulamali, F. F. *et al.* Autoencoders for sample size estimation for fully connected

neural network classifiers. *npj Digital Medicine* vol. 5 Preprint at

https://doi.org/10.1038/s41746-022-00728-0 (2022).

17. Du, X., Farrahi, K. & Niranjan, M. Information bottleneck theory based exploration of

Cascade Learning. *Entropy (Basel)* **23**, 1360 (2021).

18. Tishby, N. & Zaslavsky, N. Deep learning and the information bottleneck principle.

in *2015 IEEE Information Theory Workshop (ITW)* (IEEE, 2015).

doi:10.1109/itw.2015.7133169.

19. Shwartz-Ziv, R. & Tishby, N. Opening the Black Box of Deep Neural Networks via
    Information. *arXiv [cs.LG]* (2017).

20. Goldfarb, D. Understanding Deep Neural Networks Using Topological Data
    Analysis. *arXiv [cs.LG]* (2018).

21. Boopathy, A. *et al.* Model-agnostic Measure of Generalization Difficulty. in
    Proceedings of the 40th International Conference on Machine Learning (eds.
    Krause, A. et al.) vol. 202 2857–2884 (PMLR, 23--29 Jul 2023).

22. Gronowski, A., Paul, W., Alajaji, F., Gharesifard, B. & Burlina, P. Classification
    Utility, Fairness, and Compactness via Tunable Information Bottleneck and Rényi
    Measures. *IEEE Trans. Inf. Forensics Secur.* **19**, 1630–1645 (2024).

23. Liu, D. *et al.* Mitigating Confounding Bias in Recommendation via Information
    Bottleneck. in *Proceedings of the 15th ACM Conference on Recommender Systems*
    351–360 (Association for Computing Machinery, New York, NY, USA, 2021).

24. Li, N., Goel, N. & Ash, E. Data-Centric Factors in Algorithmic Fairness. in
    Proceedings of the 2022 AAAI/ACM Conference on AI, Ethics, and Society 396–
    410 (Association for Computing Machinery, New York, NY, USA, 2022).

25. Johnson, A. E. W. *et al.* MIMIC-CXR, a de-identified publicly available database of
    chest radiographs with free-text reports. *Sci Data* **6**, 317 (2019).

26. Wang, Peng, Lu & Lu. Chestx-ray8: Hospital-scale chest x-ray database and
    benchmarks on weakly-supervised classification and localization of common thorax
    diseases. *Proc. Estonian Acad. Sci. Biol. Ecol.*

27. Irvin, J. *et al.* CheXpert: A large chest radiograph dataset with uncertainty labels

and expert comparison. *Proc. Conf. AAAI Artif. Intell.* **33**, 590–597 (2019).

28. Idrissi, B. Y., Arjovsky, M., Pezeshki, M. & Lopez-Paz, D. Simple data balancing

achieves competitive worst-group-accuracy. in *Proceedings of the First Conference*

on Causal Learning and Reasoning (eds. Schölkopf, B., Uhler, C. & Zhang, K.) vol.

177 336–351 (PMLR, 11--13 Apr 2022).

29. Pfohl, S. R. *et al.* A comparison of approaches to improve worst-case predictive

model performance over patient subpopulations. *Sci. Rep.* **12**, 3254 (2022).

30. Falcon, W. A. Pytorch lightning. *GitHub* **3**, (2019).

31. Mei, X. *et al.* RadImageNet: An open radiologic deep learning research dataset for

effective transfer learning. *Radiol. Artif. Intell.* **4**, e210315 (2022).

32. Dosovitskiy, A. *et al.* An Image is Worth 16x16 Words: Transformers for Image

Recognition at Scale. *arXiv [cs.CV]* (2020).

33. García-Martín, E., Rodrigues, C. F., Riley, G. & Grahn, H. Estimation of energy

consumption in machine learning. *J. Parallel Distrib. Comput.* **134**, 75–88 (2019).

34. Lacoste, A., Luccioni, A., Schmidt, V. & Dandres, T. Quantifying the Carbon

Emissions of Machine Learning. *arXiv [cs.CY]* (2019).

35. Henderson, P. *et al.* Towards the systematic reporting of the energy and carbon

footprints of machine learning. *J. Mach. Learn. Res.* **21**, 10039–10081 (2020).

36. Gottfried, H. Reflections on intersectionality: Gender, class, race and nation. *Indian*

J. Gend. Stud. **11**, 23–40 (2008).

37. Hooks, B. *Feminist Theory: From Margin to Center*. (Pluto Press, 2000).

38. Paszke, A. *et al.* PyTorch: An imperative style, high-performance deep learning

library. *arXiv [cs.LG]* (2019).

39. Mozumdar, A. & Liguori, G. Persistent increase of prevalence of metabolic

syndrome among U.S. adults: NHANES III to NHANES 1999-2006. *Diabetes Care*

34, 216–219 (2011).

40. Yu, B. *et al.* Trends in depression among Adults in the United States, NHANES

2005–2016. *J. Affect. Disord.* **263**, 609–620 (2020).

41. Liu, B., Wang, J., Li, Y.-Y., Li, K.-P. & Zhang, Q. The association between systemic

immune-inflammation index and rheumatoid arthritis: evidence from NHANES

1999–2018. *Arthritis Res. Ther.* **25**, 34 (2023).

42. Li, X. *et al.* Development of an interpretable machine learning model associated

with heavy metals' exposure to identify coronary heart disease among US adults via

SHAP: Findings of the US NHANES from 2003 to 2018. *Chemosphere* **311**, 137039

(2023).

43. López-Martínez, F., Núñez-Valdez, E. R., Crespo, R. G. & García-Díaz, V. An

artificial neural network approach for predicting hypertension using NHANES data.

Sci. Rep. **10**, 10620 (2020).

44. Qiu, W. *et al.* Interpretable machine learning prediction of all-cause mortality.

Commun. Med. **2**, 125 (2022).

45. Weiss, N. S. All-cause mortality as an outcome in epidemiologic studies: proceed

with caution. *Eur. J. Epidemiol.* **29**, 147–149 (2014).

46. Singh, G. K. & Siahpush, M. Widening rural-urban disparities in all-cause mortality

and mortality from major causes of death in the USA, 1969-2009. *J. Urban Health*

91, 272–292 (2014).

47. Harel, Z. *et al.* Nephrologist follow-up improves all-cause mortality of severe acute

kidney injury survivors. *Kidney Int.* **83**, 901–908 (2013).

48. Agrawal, V., Cantor, J. H., Sood, N. & Whaley, C. *The Impact of the COVID-19*

Pandemic and Policy Responses on Excess Mortality. (National Bureau of

Economic Research, 2021).

49. Food and Drug Administration, U. S. Good machine learning practice for medical

device development: guiding principles. *FDA Med. Bull.*

50. The White House. Executive order on the safe, secure, and trustworthy

development and use of artificial intelligence. *The White House*

https://www.whitehouse.gov/briefing-room/presidential-

actions/2023/10/30/executive-order-on-the-safe-secure-and-trustworthy-

development-and-use-of-artificial-intelligence/ (2023).

51. Wiens, J. *et al.* Do no harm: a roadmap for responsible machine learning for health

care. *Nat. Med.* **25**, 1337–1340 (2019).

52. Chen, I. Y., Joshi, S. & Ghassemi, M. Treating health disparities with artificial

intelligence. *Nat. Med.* **26**, 16–17 (2020).

53. Arora, A. *et al.* The value of standards for health datasets in artificial intelligence-

based applications. *Nat. Med.* (2023) doi:10.1038/s41591-023-02608-w.

54. Chen, R. J. *et al.* Algorithmic fairness in artificial intelligence for medicine and

healthcare. *Nat Biomed Eng* **7**, 719–742 (2023).

55. Hooker, S. Moving beyond "algorithmic bias is a data problem." *Patterns Prejudice*

2, 100241 (2021).

56. Khan, S. M. *et al.* A global review of publicly available datasets for ophthalmological

imaging: barriers to access, usability, and generalisability. *Lancet Digit Health* **3**,

e51–e66 (2021).

57. Wen, D. *et al.* Characteristics of publicly available skin cancer image datasets: a

systematic review. *Lancet Digit Health* **4**, e64–e74 (2022).

58. Kalivas, J. H. & Palmer, J. Characterizing multivariate calibration tradeoffs (bias,

variance, selectivity, and sensitivity) to select model tuning parameters. *J. Chemom.*

28, 347–357 (2014).

59. Bridges, C. R., Edwards, F. H., Peterson, E. D. & Coombs, L. P. The effect of race

on coronary bypass operative mortality. *J. Am. Coll. Cardiol.* **36**, 1870–1876 (2000).

60. Ng, D. K., Moxey-Mims, M., Warady, B. A., Furth, S. L. & Muñoz, A. Racial

differences in renal replacement therapy initiation among children with a

nonglomerular cause of chronic kidney disease. *Ann. Epidemiol.* **26**, 780-787.e1

(2016).

61. Hung, M.-C., Ekwueme, D. U., Rim, S. H. & White, A. Racial/ethnicity disparities in

invasive breast cancer among younger and older women: An analysis using

multiple measures of population health. *Cancer Epidemiol.* **45**, 112–118 (2016).

62. Patzer, R. E. *et al.* Racial disparities in pediatric access to kidney transplantation:

does socioeconomic status play a role? *Am. J. Transplant* **12**, 369–378 (2012).

63. Jucker, A. H., Schneider, K. P. & Bublitz, W. *Methods in Pragmatics*. (Walter de

Gruyter GmbH & Co KG, 2018).

64. Wesson, D. E., Lucey, C. R. & Cooper, L. A. Building Trust in Health Systems to

Eliminate Health Disparities. *JAMA* **322**, 111–112 (2019).

65. Gille, F., Jobin, A. & Ienca, M. What we talk about when we talk about trust: Theory

of trust for AI in healthcare. *Intell. Based Med.* **1–2**, 100001 (2020).

66. Ferryman, K., Mackintosh, M. & Ghassemi, M. Considering Biased Data as

Informative Artifacts in AI-Assisted Health Care. *N. Engl. J. Med.* **389**, 833–838

(2023).

67. Hurston, Z. N., Plant, D. G. & Walker, A. Barracoon: the story of the last" black

cargo". (2018).

68. Shachar, C. & Gerke, S. Prevention of Bias and Discrimination in Clinical Practice

Algorithms. *JAMA* **329**, 283–284 (2023).

69. Oikonomou, E. K., Spatz, E. S., Suchard, M. A. & Khera, R. Individualising intensive

systolic blood pressure reduction in hypertension using computational trial

phenomaps and machine learning: a post-hoc analysis of randomised clinical trials.

Lancet Digit Health **4**, e796–e805 (2022).

70. Freitas, J. K. D. *et al.* Phe2vec: Automated disease phenotyping based on

unsupervised embeddings from electronic health records. *Patterns* vol. 2 100337

Preprint at https://doi.org/10.1016/j.patter.2021.100337 (2021).

71. Singhal, K. *et al.* Large language models encode clinical knowledge. *Nature* **620**,

172–180 (2023).

72. Belyaeva, A. *et al.* Multimodal LLMs for health grounded in individual-specific data.

arXiv [q-bio.QM] (2023).

73. Schwarz, K., Liao, Y. & Geiger, A. On the frequency bias of generative models.

arXiv [cs.CV] 18126–18136 (2021).

74. Zhao, S. *et al.* Bias and generalization in deep generative models: An empirical

study. *arXiv [cs.LG]* (2018).

75. Kirk, H. *et al.* Bias out-of-the-box: An empirical analysis of Intersectional

occupational biases in popular generative language models. *arXiv [cs.CL]* (2021).

76. Hao, Moon, Didari, Woo & Bangert. Highly Efficient Representation and Active

Learning Framework and Its Application to Imbalanced Medical Image

Classification. *datacentricai.org*.

77. He, K., Zhang, X., Ren, S. & Sun, J. Deep residual learning for image recognition. in

Proceedings of the IEEE conference on computer vision and pattern recognition

770–778 (2016).

827
828

Figures

Figure 1. Description of Different Biases and Proposed Role of AEquity in Identifying

and Mitigating Bias

Figure 1. Description of Different Biases and Proposed Role of AEquity in

Identifying and Mitigating Bias. A. A prototypical dataset containing samples from two

groups each of which has four informative features. B. When trained in a similar manner

on a more diverse dataset, the resulting model is biased because it performs worse for

a particular group. C. Sampling bias occurs when one group is dramatically under-

represented compared to a second group. D. Complexity bias results when one group

presents more heterogeneously than a second group; for example patients with edema

on chest radiographs present more heterogeneously in the general population than in

each subgroup. E. Label bias occurs when mislabeling burdens one group more than

another; Light green represents a higher burden of mislabeling.

Figure 2. Application of AEquity to detect and mitigate diagnostic biases in MIMIC-CXR.

Figure 2. Schema Of AEquity Workflow to detect and mitigate diagnostic biases in

MIMIC-CXR.

A. Red – White individuals; Blue – Black individuals. B. Frequency of diagnostic findings

by race. Purple – average of White and Black. C. AEquity values (log(N)) by diagnosis

and race. D. Effect of data-centric mitigation intervention on bias against Black patients.

E. Decrease in bias as measured by different fairness metrics in Black patients on

Medicaid.

Figure 3. Detecting bias in prediction of healthcare needs with the AEquity metric.

Figure 3. Detecting bias in prediction of healthcare needs with the AEquity metric.

A. A dataset with claims data used to calculate a score based on number of active

chronic conditions, and two cost-based metrics. B. Difference in AEquity values

between Black and White patients for comorbidity- and cost-based outcomes, stratified

by risk level.

Figure 4. Application of AEquity to detect and mitigate diagnostic biases in

MIMIC-CXR.

Figure 4. Application of AEquity to detect and mitigate diagnostic biases in

MIMIC-CXR.

A. Red – White individuals; Blue – Black individuals. B. Frequency of diagnostic findings

by race. Purple – average of White and Black. C. AEquity values (log(N)) by diagnosis

and race. D. Effect of data-centric mitigation intervention on bias against Black patients.

E. Decrease in bias as measured by different fairness metrics in Black patients on

Medicaid.

Figure 5. Application of AEquity to Detect, Characterize, and Mitigate a Previously
Undetected Racial Bias in Mortality Prediction on a Commonly Used Dataset

Figure 5. Application of AEquity to Detect, Characterize, and Mitigate a Previously

Undetected Racial Bias in Mortality Prediction on a Commonly Used Dataset

A. Schematic representations of AEquity applied to label selection and demographic

prioritization of bias in the National Health and Nutrition Examination Survey (NHANES)

B. AEquity reveals that the least biased label for mortality prediction occurs at 5-years

because the groups have similar levels of complexity.

C. AEquity determines that 5-year mortality has complexity bias because the joint AEq

exceeds the AEq value of both black and white patients. D. AEquity outperforms other

modalities of bias reduction, namely Balanced Empirical Risk Minimization (BERM), and

Calibration. E. AEquity demonstrates that bias reduction is invariant to sample size.

Tables

Table 1. Measurement of the discrepancy between Black and White patients for each

risk score in the EHR dataset using the AEquity metric.

Risk Metric	Risk Score	AEquity Metric
Avoidable Costs	Low Risk	0.81 (0.71, 0.92)
	High Risk	0.41 (0.32, 0.49)
Total Cost	Low Risk	0.81 (0.73, 0.90)
	High Risk	0.47 (0.36, 0.57)
Active Chronic Conditions	Low Risk	0.34 (0.26, 0.43)
	High Risk	0.06 (-0.04, 0.16)

893
We show the AEquity metric calculated with respect to each risk metric in both the
predicted low risk and high-risk population. When stratifying via avoidable costs and
total costs, the AEquity metric is elevated, but when stratifying by active chronic
conditions, the AEquity metric is not elevated indicating that active chronic conditions is
a better choice of outcome to mitigate bias. We report 95% confidence intervals in the
parenthesis.

Table 2. Measurement of bias and intervention effect by diagnosis in the radiograph

dataset with RadImageNet (ResNet-50 with pre-training on Medical Images)

Data Curation	Pre-Intervention			AEquity-Guided			
Diagnosis	White Patients	Black Patients	Bias	White Patients	Black Patients	Bias	Bias Reduction
Pneumothorax*	0.73 (0.72,0.74)	0.68 (0.67,0.69)	7.5	0.73 (0.72,0.74)	0.70 (0.68,0.72)	3.8	49.4 (49.3, 49.5)
Atelectasis*	0.69 (0.68, 0.69)	0.64 (0.64,0.65)	6.4	0.70 (0.70,0.70)	0.69 (0.69,0.69)	0.6	90.6 (90.5, 90.7)
Edema*	0.75 (0.75,0.75)	0.72 (0.71,0.72)	5.0	0.78 (0.78,0.78)	0.79 (0.79,0.80)	2.0	60.0 (59.9, 60.1)
Consolidation*	0.67 (0.67,0.67)	0.65 (0.64,0.66)	3.8	0.63 (0.63,0.63)	0.61 (0.60,0.63)	2.8	29.0 (28.8,29.2)
Pneumonia*	0.67 (0.66,0.67)	0.64 (0.64,0.65)	3.8	0.65 (0.65,0.65)	0.64 (0.64,0.65)	1.5	59.5 (59.4, 59.6)
Enlarged CM*	0.62 (0.61,0.63)	0.60 (0.59,0.60)	3.6	0.61 (0.61,0.62)	0.61 (0.60,0.61	1.1	69.9 (69.6, 70.1)
Cardiomegaly*	0.76 (0.76,0.76)	0.75 (0.75,0.75)	1.9	0.76 (0.76,0.76)	0.76 (0.76,0.76)	0.1	96.5 (96.4,96.6)
Pleural Effusion	0.64 (0.64, 0.64)	0.64 (0.64,0.64)	0.3	0.65 (0.65,0.65)	0.65 (0.65,0.65)	0.2	35.0 (33.7,36.2)
Opacity	0.69 (0.69, 0.69)	0.71 (0.70,0.71)	-2.0	0.69 (0.69,0.69)	0.70 (0.69,0.70)	-0.8	61.2 (61.0,61.4)

We show AUROC before and after intervention with 95% Confidence Intervals, for Black

and White patients. Bias is the difference in the AUROC between Black and White

patients calculated in percentages. With a dataset intervention, we reduce the

discrepancy between the two groups and calculate percentage change in the

discrepancy. Percentages have been rounded to one decimal place, and all other

numbers have been rounded to two decimal places. * - diagnostic findings with bias

against Black patients.

Table 3. Percentage Bias against Black patients by method in a ResNet-50 naive to

medical images as measured by Area-Under-the-Curve.

Label	Balanced ERM[1] (%)	AEquity (%)	% Improvement over sub-optimal policy
Atelectasis	0.85 (0.76, 0.94)	**0.51 (0.42, 0.60)**	40.07 (39.09, 41.05)
Pneumonia	1.70 (1.64, 1.76)	**1.39 (1.33, 1.45)**	24.3 (24.26, 24.34)
Pneumothorax	7.11 (7.00, 7.22)	**6.42 (6.31, 6.53)**	24.79 (24.61, 24.79)
Consolidation	**2.36 (2.24, 2.48)**	**2.36 (2.24, 2.48)**	0.00 (-0.7, 0.7)
Edema	2.11 (2.06, 2.16)	2.03 (1.98, 2.08)	3.77 (3.73, 3.81)
Effusion	0.41 (0.35, 0.46)	**0.35 (0.30, 0.41)**	12.9 (10.61, 15.21)
Cardiomegaly	**0.18 (-0.13, 0.49)**	**0.18 (-0.13, 0.49)**	0.66 (0.54, 0.78)
Enlarged CM	3.59 (3.42, 3.76)	**3.12 (2.95, 3.29)**	13.1 (12.83, 13.35)
Opacity	0.49 (0.38, 0.60)	**0.24 (0.13, 0.36)**	50.3 (50.10, 50.34)

917 Balanced sampling of groups has been previously determined to be state-of-the-art

918 across different fairness-optimizing procedures ([70]). We benchmark against empirical

919 risk minimization procedures, showing the advantages of using AEquity over ERM.

Supplement

Materials and Methods

Background on the AEquity metric

AEquity is rooted in information bottleneck theory. The core idea of information

bottleneck theory is that a deep learning classifier mapping input X to output y does so

by retaining the relevant information and eliminating the irrelevant information through a

hidden layer h[1]. In contrast, an autoencoder reconstructs X, i.e., maps input X to input X

through a hidden layer h and must retain all meaningful information. If y semantically

encodes all the meaningful aspects of X, then the information loss between X and y

through h in the classifier is minimal. The information, therefore $I(X;y)$ is approximately

equal to $I(X;X)$ at all sample sizes, and rate of learning for the autoencoder and the

generalizability of a deep learning model is similar[2]. Information flow is invariant to

autoencoder architecture and loss, which makes this technique generalizable across

different hyper-parameters[3].

This approximation can be utilized for sample size determination for deep

learning models. A standard practice in traditional experimental design, sample size

estimation takes place prior to data generation, and describes the minimum number of

samples required to observe an effect of an independent variable on an outcome. The

deep learning equivalent is the minimum convergence sample, defined by [73], which

describes the minimum number of samples needed for an algorithm trained on that

dataset to start generalizing to unseen data. If f_n is a deep learning algorithm that maps

an input X to h to y trained on n samples where h is the latent representation, then the

minimum convergence sample n is the number of training samples such that E[AUC

$(f_n(X_{test}, y_{test})] > 0.5$. In practice, this number cannot be directly observed without

prospective validation because the test set, by definition, is not available until training is

complete. It can, however, be approximated using an autoencoder, which we define as

g_n: X to h to X, where h is the latent representation. The minimum number of samples n

such that $E[g_n(X_{train}, X_{train})] < 1.0$ is defined as the minimum convergence sample

estimate. Above the minimum convergence sample, Gulamali et al have shown that the

decrease in autoencoder loss is proportional to the gain in generalizability as measured

by area-under-the-curve metric[4]. The minimum convergence sample estimate remains a

valid approximation of generalizability across model architecture, hyperparameter

selections, dataset dimensions and dataset complexity.

Bias is marked by differences in generalization performance by group or class.

Consequently, we examined the MCSE metric on subsets of the data consisting of

individual groups and classes. The resulting *AEquity* metric is defined as

$$AEq_i := arg\ min_n \frac{d^2 L_{2_i}}{dn^2}$$　　　　　　　　　　**Equation 1**

where L_{2_i} is the i^{th} class/group-specific loss curve for a network trained on

samples drawn from the entire dataset. A smaller n tends to indicate a better

generalization performance, whereas a larger n indicates poor generalization

performance when compared to other classes.

Code Availability

The code for the analyses is available at (removed for double-blind peer review)

The steps to use the repository are listed below.

1. Install requirements.

These requirements are publicly sourced such as

sklearn torch scipy torchvision pandas

pip install requirements.txt

2. Setup configuration in a config.yaml file.

data_path: ./data/custom_data.tsv # Contains path to data. Contains independent

variables, demographics, and outcome variables.

demographics_col: demographics # Name of demographics variable in data_path.

outcome_cols: outcome_1 # Name of outcome variable in data_path

exclude_cols: None # name of columns to exclude if there are extraneous columns.

out_data: ./output/data.tsv # Output directory for AEq analyses.

bootstraps: 10 # Number of bootstraps. 30-50 is typically recommended for resolution at

5000 samples.

start_seed: 42 # Seed experiments.

input_dim: 149 # Number of independent columns in data_path

max_sample_size: 5000 # Max sample size to calculate from. Usually only require 128-

512 samples.

root_dir: ./weights # Root directory to output weights, and other output files.

3. Run the measure and mitigate experiments.

python measure_disparity.py --config config.yaml

python mitigate_disparity.py --config config.yaml

Additional configurations are built for more complex models (AlexNet, ResNet,

EfficientNet), but require a custom dataloader. Custom dataloders can be built by

modifying the cnnMCSE repository. See the cnnMCSE repository for more details.

Materials

Algorithm Implementation for Medical Imaging

We have utilized three large public chest radiograph datasets in this study:

MIMIC-CXR, CheXpert, and ChestX-ray14 (**Table 1**). The MIMIC-CXR dataset was

collected from Beth Israel Deaconess Medical Center (Boston, MA, United States)

between 2011 and 2016, the CheXpert dataset was collected from Stanford Hospital

(Stanford, CA, United States) between October 2002 and July 2017, and the NIH

ChestX-ray14 dataset was collected from the NIH Clinical Center (Bethesda, MD,

United States) between 1992 and 2015. We focus on diagnostic labels with sufficiently

high frequencies.

1006

A standard practice in deep learning on medical imaging classification tasks is

the use of transfer learning – utilization of a pre-trained model such as AlexNet which is

then fine-tuned on another dataset for a given task. We employ transfer learning

baseline architectures and fine-tune the classification block to generate our classifiers,

and the encoder-decoder blocks to generate our autoencoders. We seed each

experiment with PyTorch, bootstrap experiments at 50x and report the Area Under the

Curve Metric on a held-out test dataset (AUC) for our classifiers and the L_2-norm for the

autoencoders. Code is provided as an attachment.

1015

To compute the AEq values, we utilized standard machine learning practices to

generate an autoencoder with tunable weights and biases. We generated joint datasets

by merging a balanced sample of chest X-rays belonging to each race, controlling for

sampling bias across different groups. We adapted a pre-trained ResNet-50[77], which

has 23 million parameters for the convolutional and max pooling layers. We replaced

the linear layers with an autoencoder with a latent space h = 2, selected via a grid-

search based method. The autoencoder loss was trained via mean-squared error or L_2-

norm. The network was trained via Adam optimization, with a learning rate determined

by a simple grid search. All experiments were bootstrapped fifty times, and error bars

represent 95% Confidence Intervals. The data-loaders and networks classes were

generated with PyTorch. All experiments were trained on a single NVIDIA A100 GPU

using the CUDA toolkit backend. All values and 95% confidence intervals are reported.

P values are unpaired t-tests for two groups, and ANOVA for multiple groups.

1029

We used a pretrained ResNet-50 model to generate encodings for the Chest X-

ray. We added three dense layers and a SoftMax layer to the top of the model to

generate a single-class classifier. We used a simple grid search with values ranging

from 10^{-5} to 10^{-1} for the learning rate and 4 to 256 for the batch size. After the

hyperparameter optimization, we fine-tuned the dense layers and classification layers to

generate classifications on a training set that was guided by AEquity values in

comparison to naïve dataset collection. For example, in the case of the pneumothorax,

the AEquity value dictated a diversification of the dataset, which was a 50-50 split

between Blackand white patients, whereas in the case of edema, the AEquity value

required population prioritization, which meant collecting samples belonging to Black

patients exclusively.

1041

The train-validation-test split was 60-20-20, with the size of the training dataset

being 30,000 samples from the MIMIC Chest-X-ray dataset. We trained for a maximum

of 30 epochs and employed early stopping on validation area-under-the-curve metrics.

For each experiment, we bootstrap the train and validation datasets with different

seeds, and show that our results for generalizability are robust to sampling. The train,

validation and test set, which consist of mutually exclusive samples, are bootstrapped

fifty times and 95% confidence intervals are reported.

We take a data-centric approach with simple training of a ResNet-50 pretrained

model and achieve similar performance to state-of-the-art approaches which use self-

supervision and knowledge distillation, label dependencies, hand-crafted features,

location aware dense networks. We report the results from the knowledge distillation

paper below, which was benchmarked against the latter methods (**Supplementary**

Table S2).

1055

We define bias as the discrepancy between generalization performance (as measured

by the area under the receiver operating characteristic curve) for two groups, for

example, White and Black patients. We choose this over traditional fairness metrics like

precision or false negative rates because AEquity works directly with the dataset and

doesn't concern itself with model calibration. We later show empirically that bias

mitigation with AEquity is robust to the choice of fairness metric (i.e., bias as measured

by traditional fairness metrics is mitigated as well).

1063

1064 We calculate % reduction in bias with the following steps.

1065 First, we calculate the bias without a data-centric intervention.

1066

1067 $$Bias_{pre} = \left(Test\ AUC_{pre,white} - Test\ AUC_{pre,black}\right)$$
1068

1069 Second, we calculate the bias with a data-centric intervention.

1070

1071 $$Bias_{post} = \left(Test\ AUC_{post,white} - Test\ AUC_{post,black}\right)$$
1072

1073 Third, we calculate the reduction in bias as a percentage of the original bias.

1074

1075 $$Bias\ Reduction = 100 \times \frac{Bias_{Pre} - Bias_{post}}{Bias_{Pre}}$$

1076

1077 We report the reduction in bias for both diversification and population prioritization.

1078

1079

1080   Algorithm Implementation for Cost-Utilization Analysis

1081

1082   In the data provided by (*6*), the data frame has two elements.

1083   1. Y_{it}: Outcome – Avoidable Cost, Total Cost or Comorbidity Score.

1084   2. $X_{i,t-1}$: The data collected by the patients insurer over the year *t-1*.

1085   a. Demographics (age, sex excluding race)

1086   b. Insurance type

1087   c. ICD-9 diagnosis and procedure codes

1088   d. Prescribed medications.

1089   e. Encounters

1090   f. Billed Amounts.

From these, 149 features which include biomarkers, demographics and individual

datasets are generated. The output feature is either the total cost, avoidable costs due

to emergency visits and hospitalizations, and health which measures how many chronic

conditions are flaring up and driving utilization. Those whose metrics exceed a lower

threshold, the 50th percentile, are typically referred to their primary care physicians,

who are provided with additional metrics about the patients and prompted to consider

whether they would benefit from enrollment. We assign them to the high-risk category.

1098

To compute the AEquity values, we utilized standard machine learning practices

to generate an autoencoder with tunable weights and biases. We generated combined

datasets by merging a balanced sample of electronic health record-derived data

belonging to each race, controlling for sampling bias across different groups. The

tabular nature of the dataset requires a fully connected network encoder and decoder,

which was trained via Adam optimization, with a learning rate determined by a simple

grid search. All experiments were bootstrapped fifty times with different seeds that

initialized different weights with Xavier initialization. The error bars represent 95%

confidence intervals. All experiments were trained on a single NVIDIA A100 GPU using

the CUDA toolkit backend. All values and 95% confidence intervals are reported. P

values are unpaired t-tests for two groups, and ANOVA for multiple groups.

1110

1111   We provide a notebook demonstration and YouTube tutorial on how to use this

1112   method for any tabularized dataset, using the EHR dataset as the example (removed for

1113   double-blind peer review). Selecting an informative label on the electronic health record

1114   dataset can improve test AUC by 0.11 (ΔAUC = 0.109; 95% CI: (0.106 – 0.112); $P <$

1115   0.05) on Black Patients by decreasing false negative rate and increasing precision.

1116

1117

1118

1119   Supplementary Results

1120

1121   Replication on Sex

1122

In this section, we demonstrate the potential of using the AEquity metric to

understand biases based on sex (Figure S1). For the MIMIC-CXR dataset, we notice

that while most diagnoses (5/9) have a higher AEquity value for females over males, the

labels that appear at higher frequencies such as opacity and effusion tend to have a

higher AEquity value for males rather than females. However the joint AEquity values

for these labels – opacity and effusion – tend to be lower than the AEq on individual

groups, which may mitigate the biases of these labels in practice. In the CheXPert

dataset, we see that only three labels appear to have statistically significant differences

in AEquity values between males and females – cardiomegaly, enlarged cardio-

mediastinum, and consolidation. Two of these labels have higher AEquity values for

females (cardiomegaly and enlarged cardio-mediastinum), and one has a higher

AEquity value for males (consolidation). In the NIH dataset, we see that while females

have a lower AEquity value across the in five out of the seven diagnoses, the joint

AEquity values are significantly larger in four of these, which means that training on a

joint dataset may perpetuate some of the biases, and it makes more sense to further

prioritize collection of data on female individuals.

1139

1140

Replication on Age

Next, we look at the effect of age on AEquity values (Figure S2). In the MIMIC-CXR, we see that an older age corresponds to a higher AEquity value in three of the diagnoses – enlarged cardiomediastinum, pneumonia and pneumothorax, and a younger age corresponds to a higher AEquity value for edema and cardiomegaly. In the ChestX-ray14, we see that generally AEquity is generally balanced with a higher AEquity value for younger individuals in two out of the seven labels (opacity and pneumothorax). In the CheXPert data, we see that age is relatively balanced with a higher AEquity value in two out of the nine diagnoses for the younger age – pneumonia and effusion.

Investigations and Interventions based on Medicare Status

We sought to validate our interventions in comparisons of insurance status (Figure S3). First, however, we investigated the coherence between age-related biases, as measured by Medicare status, and the race-related biases that we previously measured. We wanted to see if the impacts of age and race on biases, measured by AEquity, are coherent or dis-coherent.

We draw data from individuals with Medicare and those individuals with private insurance in Whites only, to control for the effects of race on data distribution observed above. We see some differences in distribution between the privately and Medicare-insured individuals. For example, chest radiographs labeled with pneumothorax in

individuals with Medicare have a significantly higher AEquity value than those in

individuals that are privately insured ($AEq_{Medicare, Pneumothorax}$ = 9.89; 95% CI: (9.80, 9.90),

$AEquity_{Private, Pneumothorax}$ = 8.71; 95% CI: (8.64, 8.78); *P < 0.01*) (**Fig S3(a), 3(b)**).

Moreover, the joint AEquity value is strictly in between the two different insurance

status, which may suggest that there is some but not all overlap between the two

distributions ($AEquity_{Joint, Pneumothorax}$ = 8.95; 95% CI: (8.87, 9.04)). This difference was

observed in racial bias experiments, which may suggest that the differences in

distribution are somewhat captured by inequities in care that are overlapping between

Black individuals and White individuals that are insured on Medicare. However, the joint

AEquity values of the datasets merged by racewere slightly lower than either of the

individual group's AEquity values, which indicates that Blacks and Whites had

overlapping distributions ($AEquity_{Joint, Pneumothorax}$ = 9.44; 95% CI: (9.35, 9.52),

$AEquity_{Black, Pneumothorax}$ = 10.1; *95% CI*: (10.01, 10.18), $AEquity_{White, Pneumothorax}$ = 9.50;

95% CI: (9.41, 9.57), *P > 0.05*). When stratifying by insurance status, the joint AEquity is

between the two individual insurance status, which indicates that there is a significant

complexity bias in the distribution of pneumothorax when stratifying by privately and

Medicare-insured individuals.

1181

We see the opposite trend in chest radiographs labeled with a diagnosis of

atelectasis where privately insured individuals have a significantly larger AEquity value

than individuals on Medicare ($AEquity_{Private, Atelectasis}$ = 8.82; *95% CI:* (8.72, 8.90),

$AEquity_{Medicare, Atelectasis}$ = 8.08; 95% CI: (8.00, 8.15); *P < 0.05*). The joint AEquity value is

between the two distributions, which indicates a non-overlapping data distribution

(AEquity$_{\text{Joint, Atelectasis}}$ = 8.59; 95% CI: (8.50, 8.67)). In contrast, Black individuals have an

AEquity value higher than White individuals for this specific diagnosis, which indicates

that this trend cannot be solely explained by systemic disparities in healthcare

(AEquity$_{\text{Black,Atelectasis}}$ = 9.12; 95% CI: (9.01, 9.22), AEquity$_{\text{White, Atelectasis}}$ = 8.22; 95% CI:

(8.13, 8.30), *P < 0.05*).

1192

Next, we highlighted that our interventions, which initially were conducted on

race, are valid to mitigate bias as measured by Insurance status (**Fig S3(c), S3(d)**). or

effusion, AEquity$_{\text{joint}}$ is smaller than AEquity$_{\text{private}}$, AEquity$_{\text{Medicare}}$. (AEquity$_{\text{Private, Effusion}}$ =

8.08; 95% CI : (8.00, 8.16), AEquity$_{\text{Medicare, Effusion}}$ = 7.81; 95% CI: (7.73, 7.89),

AEquity$_{\text{Joint, Effusion}}$ = 7.65; 95% CI: (7.56, 7.73)). In the case where the AEquity value

requires us to increase dataset diversity, we add samples to the training dataset from

each group equally – drawing equally from patients with and without private insurance.

In the case where the AEquity value recommends population prioritization, we add

samples exclusively from patients not on private insurance. We compare both to the

naïve approach of data collection. For effusion at the largest evaluated sample size,

there was a 0.02 increase in test AUC on chest X-rays belonging to individuals with

Medicare following the intervention, (AUC$_{\text{pre}}$ = 0.79; *95% CI*: (0.79, 0.80) vs. AUC$_{\text{post}}$ =

0.83; *95% CI*: (0.82, 0.84) n = 8,192). For pneumothorax, AEq$_{\text{joint}}$ is larger than

AEquity$_{\text{private}}$ but smaller than AEquity$_{\text{Medicare}}$ (AEquity$_{\text{Joint, Pneumothorax}}$ = 8.96; 95% CI:

(8.87, 9.04), AEquity$_{\text{Private, Pneumothorax}}$ = 8.71; 95% CI: (8.64, 8.78) , AEquity$_{\text{Medicare,}}$

$_{\text{Pneumothorax}}$ = 9.89; 95% CI: (9.80, 9.99), P < 0.05). Population prioritization leads to a

0.06 increase in test AUC for pneumothorax on Chest X-rays belonging to individuals

with Medicare (AUC_{Pre} = 0.59; 95% CI: (0.59, 0.60) vs AUC_{Post} = 0.65; 95% CI: (0.65,

0.66), $P < 0.05$ n = 8,192)

Intersectional Analysis on Black Patients and Age, Socioeconomic Status, and Sex

In this analysis, we investigate the intersection of race and age, socioeconomic

status and sex (**Fig S4**). We run AEquity analysis on datasets that intersect between

different under-represented groups including age, socioeconomic status and sex. First,

we notice that for age, we compared patients that were between the ages of 20-40 and

the ages of 40-60. For the analysis on the intersection of age and race, we see that AEq

value is significantly higher for the older group in 5 out the 9 positive labels, versus

higher for the younger group in 2 out of the 9 positive labels. The "No finding" label is

higher for patients in the younger demographic, which may be explained by significant

more variability in patient presentation where X-ray may have been ordered as a rule-

out diagnosis rather than to specifically attribute a diagnostic cause. However, we notice

that the joint is lower than that for older individuals for the labels Opacity, None,

Pneumothorax, Edema and Atelectasis, which may mean that training on a joint dataset

may mask some of potential biases we could have seen between the groups.

Second, we investigate the role of the intersection between sex and race. We

analyze the chest X-rays in Blackmale patients and Blackfemale patients. We previously

saw that white female patients generally have a higher AEquity value than their male

counterparts in the MIMIC Chest X-ray dataset This trend also holds true in Black

patients. We see that in four of the 9 labels, AEquity values are significantly higher for

females than males, whereas we see that Blackmale patients have a significantly higher

AEquity in three of the 9 labels. However, in these three labels, we see that the joint

AEquity is larger than AEquity values of the individual subgroups (Opacity,

Cardiomegaly and Effusion), which means that these populations have different

distributions for these labels. Training on a balanced dataset or one that may have been

sampled more Blackmale patients, for example, would have still induced generalizability

biases on the under-represented population.

In our third analysis, we looked at the intersection between Black patients on

private insurance and Black patients on Medicaid. Here, we notice the biggest potential

biases occur when socioeconomic status is stacked on race. Looking simply at 'no

diagnosis', we see that the AEquity for Black patients on Medicaid is 10.8, whereas the

AEquity for Black patients that are privately insured is 9.2. This difference is both

statistically significant ($P < 0.05$) and clinically significant because it implies there is a

significant amount of complexity that occurs at baseline when comparing Black patients

on Medicare to Black patients on private insurance. Second, we notice that the AEquity

values are significantly higher for Black patients on Medicaid than Black patients that

are privately insured in five of the class labels, but the opposite is only true for a single

label, while the remaining labels are neutral. The intersection between race and

socioeconomic status potentiates the largest potential for generalizability error, when

compared to the intersectionality of race and sex or race and age.

Finally, we apply the relevant dataset intervention to Black patients on Medicaid

with respect to the 'No Finding' or 'None' diagnosis. The AUC increases significantly,

and we see that this change in the AUC increases by approximately 0.03 and is

statistically significant ($P < 1\times10^{-5}$). To further dissect where this increase in AUC is

coming from, we sought to further investigate the different fairness metrics including

True Negative Rate, True Positive Rate, False Positive Rate, as well as further derived

fairness metrics such as Precision, False Discovery Rate, and Predicted Prevalence.

We see that there is a clinically and statistically significant decrease in false positive

rate for the diagnosis of "No Finding" (ΔFPR = 0.10) and a corresponding increase in

true negative rate (ΔTNR = 0.10) , with a clinically insignificant difference in false

negative rate (ΔFNR = -0.01) and true positive rate (ΔTPR = -0.01). A lower false

positive rate for the diagnosis of "No Finding" is equivalent to having a lower under-

diagnosis bias rate. Furthermore, this translates into a higher precision, lower false

discovery rate, and lower predicted prevalence. Changing the dataset gets at the root of

the model's problem described previously, which was under-diagnosis of patients in

under-represented models. By taking the relevant mitigation measures, AEquity can

reduce the false discovery rate and predicted prevalence of "No Finding", thereby

reducing the under-diagnosis bias in intersectional under-represented minorities.

1269

Analysis for Asian patients

In the MIMIC-CXR dataset, there are multiple races represented at lower

sampling frequencies than White including Black and Asian. However, models trained

on chest X-rays perform worse on Black patients compared to Asian patients. According

to an analysis of AEquity values, this is because the sampling bias is compounded by

complexity bias in the case of Black patients, whereas in Asian patients, the complexity

bias may be less severe if present at all. When we calculate AEquity values with respect

to Asian patients, we notice two things – first that the AEquity values for Asian patients

tend to be lower than those of White patients. Second, the joint AEquity values between

the Asian patients and White patients are also lower, indicating sampling bias. This is

because the chest X-rays of Asian and White patients are also relatively similar

compared to the difference between X-rays for Black and White patients. We have

computed the AEquity for these patients below and reported them in Supplemental

Figure 5.

1284

1285

1286

1287

1288

1289

1290

1291

1292

1293

1294

1295 Supplementary Figures

1296 Figure S1. Replication on sex demographic

1297

1298

Fig. S1. **Replication on sex demographic**. Red is female, orange is male and blue is

the combined dataset. **a)** Histogram by class label of MIMIC-CXR and **b)** AEquity values

for MIMIC-CXR. (R = -0.80, P = 4.94 x 10^{-7}) **c)** Histogram by class label for CheXPert

and **d)** AEquity values by class label for CheXPert. (R = -0.41, P = 8 x 10^{-3}). **e)**

Histogram by class label for ChestX-ray14, and **f)** Corresponding AEquity values for

ChestX-ray14 (R = -0.41, P = 7 x 10^{-3}).

 Figure S2. Replication on Age Demographic

1307

Fig S2. **Replication on age demographic**. Dark blue consists of individuals age (20-

40) and yellow consists of individuals from 40-60. Light blue represents the joint dataset

a) Histogram for individuals by age on MIMIC-CXR, **b)** Corresponding AEquity values

for age on MIMIC-CXR (R =-0.88, P = 2 x 10^{-8}). **c)** Histogram for individuals by age for

ChestX-ray14 and **d)** corresponding AEquity values (R=-0.42, P = 3.4 x 10^{-5}). **e)**

Histogram for individuals by age on CheXPert and **f)** corresponding AEquity values (R =

-0.32, P = 9.2 x 10^{-3}).

1315

1316

1317

1318

1319

1320

1321   Figure S3. AEquity applied to the MIMIC-CXR dataset stratified by Insurance Status.

1322

Fig S3. **AEquity applied to the MIMIC-CXR dataset stratified by Insurance Status.**

Dark green represents privately insured, brown represents insured by Medicare, and

beige represents joint data. **a)** Histogram of diagnostic frequencies by Insurance Status

b) AEquity values by Insurance status. **c)** Intervention to increase dataset diversity

improves performance in individuals on Medicare with effusion. **d)** Intervention to

prioritize populations improves performance significantly in individuals on Medicare.

1329

1330

1331

1332

1333

1334

1335

1336

1337

1338

1339

1340   Figure S4. AEquity Applied to Intersectional Populations.

1341

Fig S4. **AEquity applied to intersectional patient populations.** (a) Age – Black

patients aged 20-40 are colored red, Black patients aged 40-60 are colored green, and

the combined dataset is colored gold. Sex – Male Black patients are colored lighter

turquoise, Female Black patients are colored light green, and the combined dataset is

colored dark blue. Socioeconomic Status – Black patients on private insurance are

colored pink, Black patients on Medicaid are colored dark purple, and the combined

dataset is colored dark green. (b) AEq values calculated for each group across each

label. (c) Interventions applied to Black patients on Medicaid and the resulting effects on

different fairness metrics.

Figure S5. AEquity applied to patient population identifying as Asian.

1352

Fig S5. AEquity applied to patient population identifying as Asian. The bright red

represents AEquity values calculated for individuals identifying as Asian. The green

represents AEquity values calculated for individuals identifying as White, and the blue

represents AEquity values calculated for a combined subset of individuals identifying as

either Asian or White individuals. The AEquity of the combined subset is smaller than

the individual subsets in eight of the 10 labels.

1359

1360

1361 Figure S6. Benchmarking AEquity against Balanced ERM in Vision Transformers as

1362 measured by Area-Under-the-Curve

1363

1364

1365 **Figure S6.** Measurement of bias by diagnosis in the radiograph dataset with

1366 VisionTransformers (86 M parameter VIT_B_16 with pre-training on IMAGENET) via

1367 area-under-the-curve. Light-blue is AEquity, Dark Blue is Balanced Empirical Risk

1368 Minimization.

1369

1370 Figure S7. AEquity remains functional at smaller sample sizes.

1371

1372

1373 **Figure S7.** AEquity becomes increasingly precise as the number of samples increases

1374 and generalizes across different sample sizes when using small convolutional neural

1375 networks.

1376

1378 Table S1. Description of Datasets Utilized in Paper.

Dataset	Description	Label/Outcome	Groups examined
MIMIC-CXR	377,110 chest radiographs of 65,379 patients from the Beth Israel Deaconess Medical Center between 2011 and 2016.	Opacity, cardiomegaly, effusion, pneumonia, atelectasis, edema, enlarged cardiomediastinum, consolidation, pneumothorax, no finding.	Race, sex, age, insurance status; Sex, age and insurance in Black patients.
Healthcare costs	All primary care patients ($N = 49,618$) enrolled in risk-based contracts from 2013 to 2015, from an unspecified large academic hospital. Data included demographics, visit information, costs, comorbidities and biomarkers.	Total costs, avoidable costs, comorbidity score.	Race.
ChestX-ray14	108,948 frontal chest radiographs from 32,717 patients from the NIH Clinical Center between 1992 and 2015.	Atelectasis, cardiomegaly, effusion, infiltration, mass, nodule, pneumonia, pneumothorax.	Sex, age.
CheXpert	224,316 chest radiographs of 65,240 patients performed between October 2002 and July 2017 in both inpatient and outpatient centers affiliated with Stanford Hospital.	Opacity, atelectasis, effusion, pneumothorax, edema, cardiomegaly, pneumonia.	Sex, age.
NHANES	47,261 samples (2000-2014) with demographic, laboratory, examination, and questionnaire features that could be automatically matched across different NHANES cycles. Follow-up mortality data is provided from the date of survey participation through December 31, 2015.	All-cause mortality for follow-up times of 1-year, 2-year, 3-year, 4-year, and 5-year.	Race.

1379

1380 Description of datasets examined in this paper.

1381 **Table S2**. Comparing AEquity to SOTA methods on MIMIC Chest X-ray classification

1382 tasks.

Method	Test Area-Under-Curve
RadImageNet (2020)[5]	0.71
Wang et al. (2020)[7]	0.74
Ho et al. (2020)[8]	0.74
Yao et al (2018)[9]	0.78
Ours (RadImageNet)	0.74
Ours (Vit_B_16)	0.83

1383

1384 **Comparing AEquity to SOTA methods on MIMIC Chest X-ray classification tasks.**

1385 We take a data-centric approach with simple training of a ResNet-50 pretrained model

1386 and achieve similar performance to state-of-the-art approaches that utilize a ResNet-50

1387 backbone.

1388

1389

1390 **Table S3.** Measurement of bias and intervention effect by metric on an Intersectional

1391 Population

Metric	Bias Reduction (%)
True Positive Rate	14.4 (14.3, 14.5)
True Negative Rate	56.6 (51.9, 61.3)
False Positive Rate	68.7 (68.4, 69.0)
False Negative Rate	33.3 (26.6, 40.04)
Precision	94.6 (94.5, 94.7)
False Discovery Rate	94.5 (94.3, 94.7)

1392

1393 We show the effect of the interventions on various metrics used to measure fairness for

1394 a patient population that is both Black and on Medicaid compared to White patients.

1395 With a dataset intervention, we have reduced the discrepancy between the two groups.

1396

1397

1398

1399 Table S4. AEquity metric by race and predicted risk level, for each outcome

Metric	Category	Black	White	Difference
Total Costs	high risk	7.13 (7.05, 7.21)	7.59 (7.47, 7.71)	0.47 (0.36, 0.57)
	low risk	7.03 (6.97, 7.09)	7.84 (7.72, 7.96)	0.81 (0.73, 0.90)
Avoidable Costs	high risk	7.09 (7.01, 7.17)	7.50 (7.42, 7.58)	0.41 (0.32, 0.49)
	low risk	7.91 (7.81, 8.01)	7.09 (6.99, 7.19)	0.81 (0.71, 0.92)
Active chronic conditions	high risk	7.22 (7.12, 7.32)	7.28 (7.18, 7.38)	0.06 (-0.04, 0.16)
	low risk	6.97 (6.89, 7.05)	7.31 (7.23, 7.39)	0.34 (0.26, 0.43)

1400

1401 **AEquity metric by race and predicted risk level, for each outcome**. We show the

1402 calculated AEquity metric for highest risk and lower risk individuals for Black and White

1403 races, for each of the three outcomes (total costs, avoidable costs and active chronic

1404 conditions).

1405

1406

1407

     Table S5. AEquity-guided interventions for Vision Transformers

Method	Naive Data Collection		Balanced Empirical Risk Minimization		AEquity-Guided	
Label	White	Black	White	Black	White	Black
Opacity	0.8126 (0.8102, 0.8149)	0.8154 (0.8131, 0.8178)	0.8139 (0.8119, 0.8159)	0.8169 (0.8149, 0.8188)	0.8126 (0.8102, 0.8149)	0.8154 (0.8131, 0.8178)
Effusion	0.925 (0.9242, 0.9257)	0.9195 (0.917, 0.9221)	0.9249 (0.9239, 0.9259)	0.9192 (0.9164, 0.922)	0.925 (0.9242, 0.9257)	0.9195 (0.917, 0.9221)
Cardiomegaly	0.8613 (0.8603, 0.8624)	0.8611 (0.8596, 0.8625)	0.864 (0.8625, 0.8654)	0.8647 (0.8632, 0.8662)	0.8633 (0.8622, 0.8644)	0.8632 (0.8619, 0.8645)
Atelectasis	0.8512 (0.8497, 0.8527)	0.8463 (0.8431, 0.8495)	0.8508 (0.8496, 0.852)	0.8457 (0.8432, 0.8481)	0.8426 (0.8413, 0.8439)	0.8396 (0.8375, 0.8418)
Pneumonia	0.7781 (0.7769, 0.7792)	0.771 (0.7677, 0.7743)	0.7772 (0.7754, 0.7791)	0.7706 (0.7671, 0.7741)	0.7782 (0.7769, 0.7794)	0.7836 (0.7803, 0.7869)
Edema	0.9276 (0.9263, 0.9289)	0.9303 (0.9278, 0.9327)	0.925 (0.9226, 0.9274)	0.9286 (0.9254, 0.9318)	0.9276 (0.9263, 0.9289)	0.9303 (0.9278, 0.9327)
Enlarged CM	0.6948 (0.687, 0.7025)	0.6591 (0.6422, 0.6761)	0.6974 (0.6898, 0.7049)	0.6588 (0.6412, 0.6763)	0.6948 (0.687, 0.7025)	0.6591 (0.6422, 0.6761)
Pneumothorax	0.818 (0.8114, 0.8247)	0.7439 (0.7123, 0.7755)	0.8139 (0.8063, 0.8215)	0.7408 (0.7103, 0.7713)	0.8075 (0.7995, 0.8155)	0.7374 (0.708, 0.7668)
Consolidation	0.7903 (0.7847, 0.7958)	0.7372 (0.7145, 0.76)	0.7897 (0.7847, 0.7947)	0.7361 (0.7136, 0.7587)	0.7948 (0.7896, 0.7999)	0.7443 (0.7213, 0.7674)

1409

Table S5. Measurement of bias and intervention effect by diagnosis in the radiograph

dataset with VisionTransformers (86 M parameter VIT_B_16 with pre-training on

IMAGENET)

Supplement References

1.   Tishby N, Zaslavsky N. Deep Learning and the Information Bottleneck Principle.

Published online 2015. doi:10.48550/ARXIV.1503.02406

2.   Du X, Farrahi K, Niranjan M. Information Bottleneck Theory Based Exploration of

Cascade Learning. *Entropy*. 2021;23(10):1360. doi:10.3390/e23101360

3.   Lee S, Jo J. Information Flows of Diverse Autoencoders. *Entropy*.

2021;23(7):862. doi:10.3390/e23070862

4.   Gulamali FF, Sawant AS, Kovatch P, et al. Autoencoders for sample size

estimation of fully connected neural network classifiers. *npj Digital Medicine*.

2022;5(1):180. doi:10.1038/s41746-022-00728-0

5.   Mei X, Lee HC, Diao K yue, et al. Artificial intelligence–enabled rapid diagnosis of

patients with COVID-19. *Nat Med*. 2020;26(8):1224-1228. doi:10.1038/s41591-020-

0931-3

6.   Obermeyer Z, Powers B, Vogeli C, Mullainathan S. Dissecting racial bias in an

algorithm used to manage the health of populations. *Science*. 2019;366(6464):447-

453. doi:10.1126/science.aax2342

7.   Wang H, Xia Y. ChestNet: A Deep Neural Network for Classification of Thoracic

Diseases on Chest Radiography. Published online 2018.

doi:10.48550/ARXIV.1807.03058

8.   Ho TKK, Gwak J. Utilizing Knowledge Distillation in Deep Learning for

Classification of Chest X-Ray Abnormalities. *IEEE Access*. 2020;8:160749-160761.

doi:10.1109/ACCESS.2020.3020802

9.   Yao L, Poblenz E, Dagunts D, Covington B, Bernard D, Lyman K. Learning to

diagnose from scratch by exploiting dependencies among labels. Published online

2017. doi:10.48550/ARXIV.1710.10501

1438