Evolutionary dynamics of 1,976 lymphoid malignancies predict clinical outcome

Calum Gabbutt[1,2,13*], Martí Duran-Ferrer[3,13], Heather Grant[1], Diego Mallo[4], Ferran Nadeu[3,5], Jacob Househam[1,2], Neus Villamor[3,5,6], Olga Krali[7], Jessica Nordlund[7], Thorsten Zenz[8], Elias Campo[3,5,6,9], Armando Lopez-Guillermo[3,5,6], Jude Fitzgibbon[2], Chris P Barnes[10], Darryl Shibata[11], José I Martin-Subero[2,9,12,14*], Trevor A Graham[1,2,14*]

1. Centre for Evolution and Cancer, Institute of Cancer Research, London, UK
2. Centre for Genomics and Computational Biology, Barts Cancer Institute, Queen Mary University of London, London, UK
3. Fundació de Recerca Clínic Barcelona-Institut d'Investigacions Biomèdiques August Pi i Sunyer (FRCB-IDIBAPS), Barcelona, Spain
4. Arizona Cancer Evolution Center, Biodesign Institute and School of Life Sciences, Arizona State University, Tempe, AZ, United States
5. Centro de Investigación Biomédica en Red de Cáncer (CIBERONC), Madrid, Spain
6. Hospital Clínic de Barcelona, Barcelona, Spain
7. Department of Medical Sciences, Molecular Precision Medicine and Science for Life Laboratory, Uppsala University, Uppsala, Sweden
8. Department of Medical Oncology and Hematology, University Hospital and University of Zürich, Zurich
9. Departament de Fonaments Clínics, Universitat de Barcelona, Barcelona, Spain
10. Department of Cell & Developmental Biology, University College London, London, UK
11. University of Southern California, Keck School of Medicine, Department of Pathology, Los Angeles, CA, USA
12. Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain
13. These authors contributed equally to this work
14. These authors jointly supervised this work

* Correspondence should be addressed to Calum Gabbutt (calum.gabbutt@icr.ac.uk), José I Martin-Subero (imartins@recerca.clinic.cat) or Trevor A Graham (trevor.graham@icr.ac.uk)

Abstract

Cancer development, progression, and response to treatment are evolutionary processes, but characterising the evolutionary dynamics at sufficient scale to be clinically-meaningful has remained challenging. Here, we develop a new methodology called EVOFLUx, based upon natural DNA methylation barcodes fluctuating over time, that quantitatively infers evolutionary dynamics using only a bulk tumour methylation profile as input. We apply EVOFLUx to 1,976 well-characterised lymphoid cancer samples spanning a broad spectrum of diseases and show that tumour growth rates, malignancy age and epimutation rates vary by orders of magnitude across disease types. We measure that subclonal selection occurs only infrequently within bulk samples and detect occasional examples of multiple independent primary tumours. Clinically, we observe that tumour growth rates are higher in more aggressive disease subtypes, and in two series of chronic lymphocytic leukaemia patients, evolutionary histories are independent prognostic factors. Phylogenetic analyses of longitudinal CLL samples using EVOFLUx detect the seeds of future Richter transformation many decades prior to presentation. We provide orthogonal verification of EVOFLUx inferences using additional genetic and clinical data. Collectively, we show how widely-available, low-cost bulk DNA methylation data precisely measures cancer evolutionary dynamics, and provides new insights into cancer biology and clinical behaviour.

Introduction

The growth and dissemination of cancer is fundamentally an evolutionary process (Nowell 1976; Merlo et al. 2006); consequently, the past evolutionary trajectory of a cancer should strongly influence its future trajectory and allow inference of the clinical path of a patient (Greaves and Maley 2012). However, testing this hypothesis directly is challenging as a precise characterization of cancer evolutionary dynamics would require the analysis of multiple sequential patient samples. Instead, evolutionary trajectories must be inferred by means of a proxy; for example, somatic (epi)mutations are patterned in distinctive ways by differing evolutionary dynamics (Turajlic et al. 2019). In the haematological system, genome sequencing of single cells or single cell colonies are employed to infer the phylogenetic relationships among cells (Gaiti et al. 2019; Lee-Six et al. 2018; N. Williams et al. 2022). The expense of this approach has restricted analyses to small numbers of cases, which limits its scaling for clinical translation.

DNA methylation can also be employed as a heritable lineage marker, recording the clonal architecture of a population of cells (Yatabe, Tavaré, and Shibata 2001; Hong et al. 2010; Brocks et al. 2014; Hao et al. 2016; Siegmund et al. 2009) or the total cell proliferative history (Duran-Ferrer et al. 2020; Endicott et al. 2022). We have recently identified CpGs whose methylation state in each allele is heritable, but stochastically fluctuate over time. These fluctuating CpGs (fCpGs) function as a methylation barcode and so offer a low-cost strategy to provide high temporal resolution lineage tracing in patient samples (Gabbutt et al. 2022).

In a single diploid region, an fCpG can take one of three states: neither allele methylated (0% methylated), one allele methylated (50% methylated) or both alleles methylated (100% methylated). If multiple fCpGs are considered together, collectively they represent a vast

number of possibilities (i.e. 3 to the power of the number of fCpGs) where distantly related cells carry distinct fCpG barcodes. These barcodes are not stable, but stochastically methylate and demethylate over time at a timescale measured in years (Gabbutt et al. 2022). Therefore, they not only provide a means to trace the ancestor of a clonal expansion, but also an "evolving barcode" that tracks clonal evolution: two somatic cells with close ancestry will share a near-identical pattern of fCpG methylation, whereas distantly related cells will have divergent fCpG methylation patterns. In bulk populations of somatic cells, the dominant fCpG pattern in the sample represents the fCpG state of the founder cell of the population and intermediate methylation values that deviate from 0, 0.5 and 1 are caused by subclonal expansions in the population (fig 1A). Consequently, the full bulk distribution of fCpG methylation is determined by the evolution of the population and so provides a precise proxy-measurement of the evolutionary dynamics.

Here, we use fCpGs to infer the evolutionary dynamics of cancer cells from clinical specimens, at scale. We focus on lymphoid neoplasms, which cover a broad spectrum of diseases and subtypes with highly variable biological features and clinical manifestations (de Leval et al. 2022; Arber et al. 2022). These tumours have been extensively profiled by DNA methylation arrays, which have provided important insights into the cellular origin, pathogenesis, and clinical behaviour of these tumours (Oakes and Martin-Subero 2018). While their temporal clonal dynamics has been analysed in part (Gruber et al. 2019; Gutierrez et al. 2021), their precise evolutionary dynamics remains poorly characterised.

We construct a quantitative modelling framework called "EVOlutionary inference using FLUctuating methylation" (EVOFLUx) that enables precise quantitative estimation of cancer cell dynamics in lymphoid tumours from individual patients. In 1,976 individual lymphoid malignancies, we precisely measure evolutionary history and show that the patient-specific evolutionary dynamics is strongly associated with disease outcomes.

Results

Bulk Methylation Array Samples from 2,430 samples

We assembled and processed with a harmonized pipeline 2,430 bulk sample Illumina methylation array data of normal and neoplastic lymphoid cells (Kulis et al. 2015; Nordlund et al. 2013; Reinius et al. 2012; S. T. Lee et al. 2015; Queirós et al. 2016; Nadeu et al. 2020; Duran-Ferrer et al. 2020; Nadeu et al. 2022; Oakes et al. 2016; Dietrich et al. 2018; Agirre et al. 2015). (Methods). After quality control, we retained 2,204 samples from 2,054 patients (including 22 technical replicates, 3 synchronic and 125 longitudinal samples from the same patients; tables 1, S1-2) and 389,180 CpGs for downstream analyses. As healthy control samples, this dataset contained sorted $CD19^+$ B cells (n=40), $CD3^+$ T cells (n=35), peripheral blood mononuclear cells (PBMCs, n=6) and whole blood samples (n=6). As tumor samples, we included precursor 797 B- and 90 T-acute lymphoblastic leukemias (B- and T-ALL, respectively) at diagnosis, 28 B and 2 T-ALL at relapse as well as 74 B and 12 T-ALL at complete remission (i.e. normal blood); 149 mantle cell lymphomas (MCL); 722 chronic lymphocytic leukemias (CLL), 55 of its precursor condition monoclonal B-cell lymphocytosis (MBL), and 6 samples from CLL patients undergoing a diffuse large B-cell lymphoma (DLBCL)

transformation called Richter transformation (RT); 62 primary DLBCL, not otherwise specified (NOS); and 104 multiple myeloma (MM) and 16 of its precursor condition monoclonal gammopathy of undetermined significance (MGUS). In addition, clinical follow-up was available for 1,667 patients and matched DNA methylation with genetic (whole genome/exome sequencing (WGS/WES)) or with RNA-seq data was available for 505 and 293 CLL/RT samples, respectively (tables 1, S1). Therefore, this exquisite dataset was unique from multiple perspectives, including tumors derived from lymphocytes at various maturation stages, tumors with a highly proliferative acute presentation and more indolent chronic leukemias, tumors from patients ranging from infants to old adults, and tumor samples from different niches and disease stages with available genetic, transcriptomic, and clinical data.

		Sex		Tissue source					Age at sampling			Purity			Clinical follow-up	fCpG identification	EVOFLUx inference	
	N	F	M	BM	PB	LN	Other	Unknown	Distribution	Mean	Range	Distribution	Mean	Range			Discovery	Validation
Bcell	40	20	14	-	26	-	14	-		28	2-65		95	95-95	-	-	-	-
Tcell	35	-	-	35	-	-	-	-		-	-		90	90-90	-	-	-	-
PBMCs	6	-	-	-	6	-	-	-		-	-		90	90-90	-	-	-	-
Whole blood	6	-	-	-	6	-	-	-		-	-		90	90-90	-	-	-	-
T-ALL	90	-	-	70	20	-	-	-		9	1-18		79	60-93	90	90	90	-
T-ALL-relapse	2	-	-	2	-	-	-	-		6	5-6		62	60-63	2	2	2	-
T-ALL-remission	12	-	-	-	1	-	-	11		12	4-18		-	-	12	-	-	-
B-ALL	797	81	90	507	60	-	214	16		6	0-19		88	60-100	608	608	608	170
B-ALL-relapse	28	-	-	27	-	-	-	1		10	3-18		86	60-98	28	28	28	-
B-ALL-remission	74	-	-	16	2	-	-	56		6	1-20		-	-	74	-	-	-
MCL	149	38	111	4	86	52	6	1		65	34-89		86	60-100	117	77	59	71
MBL	55	32	23	-	55	-	-	-		71	39-89		95	85-100	55	53	53	2
CLL	722	256	399	2	718	2	-	-		66	19-94		94	65-100	675	437	437	281
RT	6	4	2	-	3	3	-	-		71	45-84		84	75-93	6	-	-	6
DLBCL-NOS	62	31	31	-	-	30	19	13		64	18-90		80	60-99	-	56	52	6
MGUS	16	3	13	16	-	-	-	-		72	56-83		91	84-97	-	16	9	-
MM	104	62	42	104	-	-	-	-		69	38-88		95	81-100	-	104	102	-
Total N	2204	527	725	748	1018	87	253	98	-	-	-	-	-	-	1667	1471	1440	536

Table 1: A summary of methylation array samples
Abbreviations: Peripheral blood mononuclear cells (PBMCs); precursor T-acute lymphoblastic leukemias (T-ALL); precursor B-acute lymphoblastic leukemias (B-ALL); mantle cell lymphoma; monoclonal B-cell lymphocytosis (MBL); chronic lymphocytic leukemias (CLL); Richter transformation (RT); diffuse large B-cell lymphoma, not otherwise specified (DLBCL-NOS); monoclonal gammopathy of undetermined significance (MGUS); and multiple myeloma (MM). Female (F); Male (M); bone marrow (BM); peripheral blood (PB); lymph node (LN).

Identification of fCpG Loci in Blood Malignancies

We have previously observed that fCpG loci are tissue-specific (Gabbutt et al. 2022). To identify fCpGs in the lymphoid system we first partitioned our dataset into discovery (n=1,471) and validation sets (n=733, tumour samples n=536; tables 1, S2). Using the discovery set, we selected for CpG loci that fulfilled the following criteria:

142   (i)   Heterogeneous across different subjects with the same disease (by accepting
143   CpG loci with top 5% of standard deviation of methylation value within a cancer
144   type)

146   (ii)   Equally likely to be methylated or unmethylated (by averaging across all cases in
147   a cancer type and accepting only CpG sites with average methylation of
148   approximately 0.5 in the cancer type)

149   (iii)   Unlikely to be associated with specific cell or cancer types. This was done using
150   an unsupervised Laplacian Score (LS) feature selection metric (He, Cai, and Niyogi
151   2005) that ranked CpG loci by their tendency to preserve the nearest-neighbour
152   graph, where we accepted the 5% least informative CpG loci.

To visualise how the LS quantified the informativeness of individual CpG loci, we performed
a principal component analysis (PCA) on methylation values (fig. 1B). The first 2 principal
components revealed disease specific clustering in methylation. The CpG loci with the most
informative features for clustering (lowest LS) showed disease-specific methylation. In
contrast, the CpG loci that were least informative for clustering (highest LS) tended to have
either low variability or high variability that was well-dispersed across the PCA. Combining
these filters yielded 978 pan-lymphoid cancer fCpGs (table S3). These fCpGs did not cluster
the samples by either disease (fig. 1C) or disease-subtype in either discovery or validation
samples (fig. S1A-D), nor with the version of methylation array employed.

We then examined the distribution of these 978 fCpG methylation values in individual
samples (table S4). As we previously predicted (Gabbutt et al. 2022), in each cancer sample
the fCpG loci followed a characteristic "W-shaped" distribution that depicts the fCpG
methylation patterns of the founder cell of the cancer sample (fig 1D). In contrast, the
healthy B cell subpopulations, which were not included in the discovery set, had unimodal
distributions with intermediate methylation consistent with these being polyclonal
populations (fig. 1D). The fCpG methylation distributions of samples from ALL patients in
remission (i.e. contain mostly healthy cells) similarly had unimodal distributions and
clustered with the normal B cell subpopulations (fig 1C-D).

Employing the standard deviation of each sample's fCpG methylation distribution as a proxy
for its "W-shape", we observed that lymphoid cancers had significantly higher standard
deviation than both peripheral blood cells and sorted B and T lymphocytes, as expected (fig
1D).

Figure 1 – Selection and properties of fCpG loci

A: Illustration of how fCpGs enable tracking of lineage dynamics, describing three limiting cases of a population evolutionary structure: (top) polyclonal, in which the most recent common ancestor (MRCA) exists far in the past; (middle) clonal, in which the MRCA is very recent; (bottom) ongoing evolution following a population bottleneck. In a polyclonal population, the different fCpGs (white – unmethylated, grey – heterozygous methylated, black – methylated) are all different due to fluctuations that occurred since the MRCA. Thus, the average methylation of the population as each fCpG is roughly the steady state methylation value. Following a clonal expansion, all the fCpGs in the population are

the same, so the average methylation reflects that of the methylation of the founder cell. However, as the population continues to grow ongoing fluctuations alter the barcodes in individual cells from the ancestral barcode, changing the distribution of bulk methylation values. **B**: fCpGs were selected by combining three filters: (i) CpGs with low intra-disease heterogeneity were removed as sites that were likely not fluctuating, (ii) CpGs with a mean methylation far from 0.5 were removed as likely belonging to CpG sites with skewed methylation/demethylation rates, and (iii) CpGs which preserved the nearest-neighbour graph (quantified using the Laplacian Score) were removed as sites that were likely under selection/strict regulation. **C**: A hierarchically-clustered heatmap of the 978 fCpGs. **D**: (top) Example methylation distributions of the fCpGs from healthy B cells, B-ALL cells and whole blood from a patient in remission from B-ALL. (bottom) Boxplots (whiskers extending to ±1.5×IQR) displaying the standard deviation of the methylation distributions separated by cell/disease type. **E**: A heatmap showing the log2 fold change in chromatin status at the site of fCpG loci compared to other CpG loci on the 450K Illumina array in healthy lymphoid cells, MCL, CLL and MM samples. White stars signify the degree of statistical significance (χ^2 tests, hs correction). **F**: Expression analysis on the genes associated with fCpG and non-fCpG loci: (left) a Q-Q comparing non-fCpG and fCpG associated genes, demonstrating fCpG associated genes have a lower expression (in transcripts per million), (middle) a comparison between fCpG and non-fCpG associated genes in a single CLL sample (P=3.96e10^{-11}, Wilcoxon test), (right) the expression of fCpG associated genes separated by discretised methylation status (all p>0.05, Wilcoxon test).

fCpG loci are enriched in silent regions of the genome

fCpGs represented less than 1% of measured CpGs and were generally evenly dispersed across all chromosomes, with the exception of an enrichment on chromosome 19 (P=2.2e-5, χ^2 test, holm-sidak (hs) correction, fig. S2A). Despite the similar distribution across chromosomes, fCpG loci were enriched on the shores of CpG islands, whilst being depleted within CpG islands themselves and the open sea (all significant p values < 10^{-7}, χ^2 test, hs correction, fig. S2B). Remarkably, fCpG were poorly overlapped with CpG previously used in a variety of "epigenetic clocks", including mitotic age, chronological age, trait predictors and biological and mortality clocks (fig S2C, table S5, Methods).

Next, to extend our analyses beyond Illumina array data, we looked at the methylation patterns of sorted bulk B-cell and T-cell populations using publicly available whole genome bisulfite sequencing data (Loyfer et al. 2023). Methylation at fCpG loci in these polyclonal normal samples was largely intermediate (fig. S3A), consistent with our previous findings in array data (fig. 1C-D). Over a small window of 100bp, as the distance from the fCpG increased, an increasing fraction of the neighbouring CpG loci were either hyper- or hypo-methylated (fig. S3A-B), suggesting that a particular 3D DNA structure may allow the fluctuation of methylation at the specific sites where fCpGs are located.

The chromatin configuration of fCpG loci in normal and neoplastic B cells showed that fCpG loci were enriched in weak promoters and enhancers as well as H3K27me3-marked regions, whereas in active promoters and H3K36me3-marked regions they were significantly underrepresented (fig 1E). In line with this, RNA-seq analysis of 294 CLL samples showed that the expression of genes associated with fCpGs was highly skewed towards lower expression compared to genes exclusively associated with non-fCpG loci (fig 1F). Using a subset of CLL samples (n=224) having matched RNA-seq and DNA methylation data, we observed similarly low gene expression levels between fCpG-associated genes regardless of the methylation allele states (fig. 1F, Methods). These results were further supported by a variety of statistical analyses and database annotations, including a significantly lower likelihood of fCpGs being annotated as associated with a gene (fig. S2DC, p<0.001) or annotated as promoters by the ENCODE Methylation Consortium (Dunham et al. 2012) (fig. S2E p<10^{-10}, χ^2 test, hs correction). fCpG associated genes were underrepresented in pathways ubiquitously expressed across multiple tissue types (e.g. lymph node, tonsil and

213 testis) by gene-set enrichment analysis (fig. S4A, table S6). Conversely, fCpGs were enriched
214 in CpGs unclassified by ENCODE (p < 10^{-10}, χ^2 test, hs correction, fig. S2E) and enriched in
215 developmental pathways (fig. S4B, sup table S7).
216
217 Together, these results indicate that fCpG loci are located in particular regions of the
218 genome where it is likely that the DNA structure enables fluctuation, and further that fCpGs
219 do not regulate transcription and thus are unlikely to experience clonal selection.
220

EVOFLUx measures clonal evolution

222 We developed a stochastic modelling and inference framework called EVOFLUx to simulate
223 how clonal evolution quantitatively determines fluctuating methylation values and enables
224 inference of evolutionary dynamics on a case-by-case basis from patient data.
225
226 The model simulates the ongoing gain and loss of methylation at fCpGs within a lineage
227 from a patient's birth until the beginning of a cancer-associated clonal expansion at some
228 specified time, and then continues to simulate methylation fluctuations within the growing
229 population of cancer cells until the cancer sample is collected at time T (fig 2A; Methods
230 and Supplementary Information). The key parameters in the model are:
231
232 • Cancer growth rate per year (θ), assuming an exponentially growing population.
233 • Patient age at cancer emergence (τ), measured in years.
234 • fCpG switching rates per allele per year. Four parameters corresponding to the four
235   possible transitions between homozygous unmethylated, heterozygous methylated
236   and homozygous methylated (μ, υ, γ and ζ).
237
Importantly, the time of cancer emergence should be thought of as the time of occurrence
of the most recent common ancestor (MRCA) of the sampled extant cancer population,
rather than the cancer age *per se*, as there may have been previous clonal expansions prior
to the "final" clonal sweep that produced a clinically-detectable cancer, and the MRCA of
the sample is inevitably the same or more recent than the MRCA of the whole cancer
population. By combining the cancer growth rate and the time that has passed since the
MRCA, the cancer effective population size (N$_e$) present within the sample can be calculated
as $N_e = e^{\theta(T-\tau)}$. Note that this is the N_e of the sample, rather than the cancer as a whole,
and under the assumption the cancer is growing exponentially.
247
EVOFLUx accounts for sample impurity (contaminating non-cancer cells in the sample) by
modifying the methylation fraction of the cancer cells by the average methylation of the
contaminating cells (assumed to be random), weighted by sample purity (Methods). The
framework also mimics the technical noise introduced by the methylation array, by adding
beta-distributed noise to the methylation values centred upon the "true" methylation value
in the population.
254
Simulations showed that the model qualitatively recapitulates the patterns observed in
patient data, including the characteristic W-shape (fig. 2B). The specific shape of the
resulting W-distribution is determined by the model parameters. For example, as the cancer
age is increased (i.e. the time τ occurs earlier in a patient's life, all other parameters held

constant), there is more time for fluctuations to "desynchronise" the methylation pattern of
the founder cell, causing the flanking peaks of the W-shaped distribution to move towards
the central peak (fig. 2C). Similarly, if we keep the time since the MRCA and the epimutation
rate (per year) fixed but decrease the growth rate of the tumour population, each
subsequent division has more time to accumulate epimutations, magnifying the
stochasticity of the system and broadening the width of the peaks in the W-distribution (fig.
2C).
266
EVOFLUx utilises an extensive Bayesian inference method to learn model parameters from
input fCpG methylation distribution data (Methods). To evaluate the accuracy of the
inference, we simulated fCpG data and tested the ability of EVOFLUx to recover the known
input parameters (fig. 2C). These values were inferred with good confidence, with the 95%
credible intervals encompassing the ground truth values (fig. 2D), significant narrowing of
the posterior compared to the prior and with the posterior predictive distribution well-
matching that of the original simulated dataset (fig. 2E). Hence, these simulation
experiments demonstrate that the parameters defining the evolutionary dynamics of
growing cancers can be measured using bulk methylation arrays.

Figure 2 – A stochastic mathematical model well-describes the observed patterns in cancer fCpG data.
A: An illustration of the mathematical model describing how the fCpG distributions vary with the evolutionary parameters of a cancer. (top) The model is split into 2 phases, prior to the MRCA (τ) whereby methylation changes occur in a single cell, and following the MRCA where the population grows exponentially (θ). (bottom) At each time step, epigenetic switching is

allowed to occur between the 3 possible states (rates μ, υ, ζ and γ). **B**: An example simulation run, which has the characteristic "W-shape" observed in lymphoid cancer data (e.g. fig. 1D). **C**: Illustrative limiting cases of: (top) a very old cancer in which the fCpGs have had a long time to desynchronise vs a young cancer in which the fCpGs are largely synchronised, (bottom) a rapidly growing cancer in which the rate at which cells divide is high compared to the switching rate vs a slowly growing cancer in which the effect of stochasticity is amplified. **D**: The resulting posterior distributions after fitting the simulated data in fig. 2B with the Bayesian inference method. The posterior (orange) displays tightening around the ground truth parameter values (red) compared to the prior (grey). **E**: The posterior predictive distribution (orange) superimposed upon the simulated data (blue), showing an excellent fit.

The evolutionary dynamics of lymphoid malignancies

We applied EVOFLUx to 1,976 samples from lymphoid cancer (including T-ALL, B-ALL, CLL,
MCL, DLBCL and MM) and premalignant conditions (i.e. MBL and MGUS) where we had age
and tumour cell purity information, and inferred the cancer growth rate, time to the MRCA
and epigenetic switching rates of each cancer independently (table S8). Posterior
distributions were well-formed (fig S5A), and posterior predictive distributions recapitulated
the input data well (fig S5B), emphasising the excellent fit of the model to data.

Acute paediatric leukemias derived from precursor B/T cells and adult lymphoid neoplasms
derived from mature B cells were observed to have markedly different evolutionary
histories (taking the median as a point estimate; fig. 3A). In particular, EVOFLUx inferred
ALLs growing at a significantly higher rate (fig. S6A, P= 9.3e-306, MW-U, hs correction),
having a smaller cancer effective population size (N_e, fig. S6B, P= 8.1e-25) and shorter time
since the MRCA (fig. S6C, P= 6.0e-306) than all other lymphoid malignancies. T-ALL were
found to grow faster than B-ALL (P=0.0017, hs correction), although the variance on the B-
ALL growth rates was larger than for T-ALL (P=0.00044, Levene test). In adult cancers, MBL,
an asymptomatic precursor condition to CLL, was found to have lower growth rates than CLL
(fig. S6A, P= 9.7e-10) and longer time since the MRCA (P=9.9e-13), although there was no
statistically significant difference in the inferred N_e (P= 0.53). DLBCL was found to have the
largest N_e, despite not having a particularly elevated growth rate compared to the other
adult cancers (potentially due to the lower purity of DLBCL samples).

There was also significant variation in the epigenetic switching rates (epimutation rates)
between diseases, with acute paediatric leukemias having an order of magnitude faster
switching than adult mature B cell neoplasms (fig. 3B; fig S6D, P=5.6e-301). Further, we
observed a positive correlation between growth rate and methylation switching rate in both
B-ALL and T-ALL (fig. S6E, P= 2.4e-98, R^2=0.44 & P=5.9e-06, R^2=0.22 respectively), a weak
negative correlation in MM (P=1.6e-05, R^2=0.18) and no correlation in the other adult
malignancies, suggesting that fCpG switching rates are elevated during childhood, but fall to
a steady rate during adulthood. Notably, when comparing the four epigenetic switching
parameters (μ, υ, γ and ζ), the different cancer types occupied different regions of the
epimutation parameter space (fig. S6F).

Figure 3 – EVOFLUx reveals the evolutionary dynamics of lymphoid cancers

A: A scatter plot of the inferred growth rate (θ) vs effective population size (N_e, $e^{\theta(T-\tau)}$) per sample, using the posterior median as a descriptive summary statistic. Coloured according to cancer type. **B:** A scatter plot of the inferred time since the MRCA (τ) vs the mean epigenetic switching rate (averaged across different rate parameters) per sample. **C-E:** 3 example histograms samples in which the neutral (**C**), subclonal selection (**D**) and independent clonal origins (**E**) models are preferred. **F:** A histogram showing the variant allele frequency spectrum of matched whole exome sequencing (WES) data, with the results of subclonal deconvolution (performed using Mobster (Caravagna et al. 2020)) overlaid. In this sample, there is strong evidence of an emerging subclone (C2), in line with the inference from EVOFLUx. **G:** Boxplots comparing the distribution of subclonal weightings inferred by EVOFLUx in samples called as neutral vs under subclonal selection via the WES data.

The Majority of CLLs Grow Neutrally

312 Previous work has proposed that many cancers grow effectively neutrally with no subclonal
313
selection detectable (Sottoriva et al. 2015; M. J. Williams et al. 2016; 2018; Ling et al. 2015).
In CLL, two major molecular subtypes are defined based on the extent of somatic
hypermutation in the heavy-chain variable region of the IG gene, namely unmutated (U-

CLL), and mutated (M-CLL). Nonetheless, a small fraction of CLL patients present with
multiple IGHV rearrangements (Plevova et al. 2014), and likely consistent of two cancers
with independent clonal origins. Subclonal, independent and neutral growth dynamics are
distinguishable by characteristic patterns of mutations across the cancer genome
(Caravagna et al. 2020; M. J. Williams et al. 2018).

Simulations showed that in fCpG data the signature of subclonal selection is the presence of
intermediate peaks between the peaks corresponding to fCpG loci that were in the
homozygous unmethylated, heterozygous, and homozygous methylated states in the MRCA
cell (fig. S7A). These additional intermediate peaks are formed by fCpG loci that have
switched methylation state in the lineage that formed the subclone within the intervening
time between the MRCA of the sample population and the emergence of the outgrowing
subclone. However, direct detection of these fCpG methylation patterns in the data (e.g. via
a peak detection method) was generally impractical due to the relatively small number of
fCpG loci that are expected to change methylation status in the time between the MRCA
and the emergence of the subclone (e.g. assuming $\mu \approx 0.01$ per year, as in CLL, $1 -$
$e{-}2\mu\tau2{-}\tau1{\approx}2\%$ of fCpGs in the homozygous unmethylated state will have undergone 1
fluctuation in a single year) and technical noise of fCpG methylation measured by the
methylation array (fig. S7B).

Therefore, we re-fitted each of our patient's fCpG methylation distributions with modified
EVOFLUx models that included subclonal selection and independent clonal origins, and used
Bayesian model selection and a leave-one-out approach to determine which model best
represented the data (fig. 3C-E, S7C, Methods). We found that in the majority of samples
(1610/1976), there was no evidence of either subclonal or two independent cancers at 95%
confidence. There was also a high variance in the fraction of cancers growing under
subclonal selection between different cancer types (fig. S8A, from over 30% (232/718) in CLL
to less than 5% in DLBCL (1/57). However, these results need to be interpreted cautiously
given the heterogeneity in purity across the sample groups, which likely limits the power to
detect subclonal selection in lower purity samples.

We note that only strong selection occurring neither "too early" or "too late" is detectable
in the data (fig. S7D), analogously to detecting selection using point mutations in WGS
(Heide et al. 2018). This is because late arising or weakly selected subclones will only be
present at very low frequencies, whereas very strongly selected early subclones will tend to
have swept prior to sampling, and very early arising subclones have not had sufficient time
to evolve fCpG patterns dissimilar enough to be distinguishable from the ancestor clone.
Furthermore, whilst our simulations demonstrated that EVOFLUx can correctly identify
subclonal section, the absolute value of the inferred selection coefficients are not well
informed due to the noisiness of the methylation array data (fig. S7E).

To verify EVOFLUx based inferences of subclonal selection we used matched WES data
available for 425 CLL cases. Using MOBSTER, our subclonal deconvolution tool (Caravagna et
al. 2020), to detect emerging subclone(s) from the variant allele frequency (VAF)
distributions of somatic mutations (fig. 3F, table S9) we found 78 /425 samples contained
evidence of subclonality by WES (table S10). Samples which were classified as subclonal by

MOBSTER had significantly higher subclonality weights inferred from our EVOFLUx inference
(P= 2.0e-4, MW-U test, fig. 3G).
365
The paucity of point mutations in WES data limits the power to detect subclonality.
MOBSTER was more likely to assign a sample as containing a subclone when the sample
contained more mutations (p= 0.0023, logistic regression against log number of mutations,
fig. S8B). Hence, we suggest that those samples with high subclonality weights from our
EVOFLUx inference, despite being labelled as "neutral" by MOBSTER, may truly contain
subclones. Conversely, in cases assigned as subclonal by MOBSTER, there was no association
between the mutation burden and EVOFLUx subclonality weighting, suggesting these may
be genuinely subclonal samples that EVOFLUx has missed (P=0.24, logistic regression, fig.
S8C).
375
Cancers with two independent clonal origins were detected in 22/718 CLL cases. We
attempted to validate these inferences by comparing them to orthogonal measurement of
the number of IG gene rearrangements based on WES/WGS and RNA-seq (Knisbacher 2022)
(table S11). Those patients with multiple IG gene rearrangements had elevated independent
origins model weightings compared to those without (fig S8D, P= 0.028), although we note
that EVOFLUx was not as accurate at detecting independent cancers as detecting
subclonality.
383
EVOFLUx detects early subclonal origins of Richter transformed cells in CLL
A small fraction of CLL patients will undergo RT, the emergence of an aggressive phenotype
with dismal prognosis. Recently, using single cell genomics and transcriptomics, we found
that RT cells are present at low frequencies within the diagnostic CLL sample, preceding the
clinical emergence of RT in some cases by two decades (Nadeu et al. 2022). For 2 of the
same patients analysed previously by WGS and single cell approaches (Nadeu et al. 2022),
we also had DNA methylation arrays of longitudinal samples from diagnosis to RT (fig 4A,
table S12). We used EVOFLUX to construct phylogenetic relationships between samples
from differences in fCpG methylation (Methods), extracting valuable temporal information
regarding the divergence of RT cells from the ancestral clone.
394
We found the inferred phylogeny well matched the known clinical relationships between
the different samples (fig. 4B). Notably, the evolutionary distance between two technical
replicates of untreated samples (450k and EPIC data) was small in both cases (fig. 4B). In
both phylogenies, the RT clone diverged exceptionally early, roughly a decade before the
MRCA of the non-transformed samples (9 and 12 years for case 12 and 19, respectively).
This is consistent with our previous work (Nadeu et al. 2022) in which we detected RT cells
within the diagnostic sample, but suggests that the initial RT divergence occurred well
before diagnosis, over 30 years before the clinical presentation of RT.
403
To validate the fCpG phylogenetic inference, we built phylogenetic trees on SNV data from
matched WGS (Nadeu et al. 2022) (fig. 4A and S9, table S12). We restricted our analysis to
clock-like SBS1 mutations (Nik-Zainal et al. 2012) which were clonal within the sample
(Caravagna et al. 2020). The topology of the SNV phylogenies matched that of the fCpG
phylogenies and the 95% credible intervals of the estimated MRCA overlapped. Hence,

fCpGs have the potential to serve as a high temporal resolution phylogenetic character,
enabling multi-sample phylogenetic reconstruction at a fraction of the cost of WGS.
411
In addition, we applied our phylogenetic analysis to 3 longitudinal B-ALL samples, tracking
the disease course from initial diagnosis to relapse using the same methodology (fig. S10). In
each of the patients, the subsequent relapse samples formed a separate clade from the
initial diagnostic samples, suggesting the cancer population has undergone a significant
bottleneck through treatment. The branch lengths of the diagnostic samples were all very
short, suggesting there was initially little evolutionary distance between the surviving
resistant population and the sensitive cells.
419

Figure 4 – fCpGs allow for phylogenetic reconstruction of longitudinal lymphoid cancer samples.
A: Timeline of two CLL patients with samples collected longitudinally, annotated with treatment received. Circles represent methylation array (blue: 450K, orange EPIC), squares represent whole genome sequencing, treatment is represented with a green vertical line and Richter transformation (RT) is represented with a vertical pink line. **B:** (left) The reconstructed phylogenies of the relationship between samples, annotated with the clinical classification of each sample. The black triangles represent the time that occurred since the most recent common ancestor, taken as the posterior median of $T - \tau$ from the single-sample EVOFLUx inferences. (right) A heatmap representing the 978 fCpG loci, with the colour a representing the fraction methylated (0% blue, 100% red).

Evolutionary growth dynamics of CLL predicts clinical outcomes

Since cancer development and progression is fundamentally an evolutionary process, the evolutionary trajectory of the disease should predict the future course of the disease and clinical outcome of the patients. To evaluate this hypothesis, we first compared the inferred growth rate and N_e between different molecular subtypes of the same cancer type.

In B-ALL, cases having the translocation 11q23/MLL had a significantly higher growth rate (fig. 5A, P=1.3e-13, MW-U, 44.3±6.1 vs 11.7±0.2 per year [mean ± standard error]), but lower N_e (P=3.7e-07, 1.8e5±0.2e5 vs 3.0e5±0.06e5 cells) than the other subtypes, consistent with its distinct clinical behaviour (Rice and Roy 2020). In MCL, the generally more indolent nnMCL (Nadeu et al. 2020) had a lower growth rate (fig. 5B, P=1.1e-03, 1.7±0.1 vs 2.1±0.1 per year) and N$_e$ (P=7.4e-05, 4.7e5±1.4e5 vs 1.5e6±0.2e6 cells) than the more aggressive conventional MCL (cMCL). In DLBCL transcriptomic subtypes (Alizadeh et al. 2000) there was no significant differences, likely due to the smaller number of cases and the lower sample purity (fig. S11A&B).

In CLL, the more aggressive U-CLL subtype (Hamblin et al. 1999; Damle et al. 1999) showed a significantly higher growth rates (fig. 5C, P=1.3e-32, 2.3±0.04 vs 1.8±0.02 per year) and N_e (P=2.1e-22, 7.2e5±0.3e5 vs 4.1e5±0.3e5 cells) compared to M-CLL. In its precursor condition MBL, although we had little power to detect a difference between U-CLL and M-CLL subtypes, we could observe a non-significant increase in growth rate in U-CLL cases mirroring the findings in CLL (fig. S11C&D, P=0.093, 1.7±0.1 vs 1.5±0.06 per year). This result is consistent with the known enrichment of MBLs in IGHV mutated cases, and suggestive that MBL patients with an unmutated IGHV gene generally progress more rapidly towards CLL and therefore the MBL phase is less frequently observed in the clinical setting.

To further measure the influence the past evolutionary history of a cancer imparts upon its future clinical course, we exploited our access to a large clinically well-annotated series of CLL cases to evaluate the prognostic value of the inferred evolutionary parameters. We used univariate Cox models to assess the impact of each parameter on time to first treatment (TTFT), a commonly used variable that describes the natural history (biological behaviour) of the CLL cells, as well as overall survival (OS) which is a more complex endpoint as it convolves disease biology with heterogeneous treatments administered to the patients.

The univariate prognostic impact of the inferred growth rate of the cancer on both TTFT (fig. 5D, P=1.4e-30, hazard ratio (HR) HR=3.95) and OS (P=0.0053, HR=1.51) was stark. The N_e of a cancer did not have a strong impact on TTFT (P=0.058, HR=1.17), but its effect on OS was clear (P=1.3e-4, HR=1.41). The patient age at the time of the MRCA of the CLL population is highly corelated with the age of the patient (fig S12A), so unsurprisingly older patients had worse OS (P=2.3e-11, HR=1.79). The decrease in risk of progression with cancer age (P=3.8e-17, HR=0.65), measured by the time since the MRCA, was likely due to confounding with the growth rate, as these parameters were negatively correlated (fig. S12A). The epigenetic switching rate parameters were largely uninformative of prognosis.

As the growth rate was different between U-CLL and M-CLL (fig. 5C), we next analysed its prognostic impact within each of them. Indeed, we observed that the estimated growth rate has a profound impact on TTFT in both CLL subtypes separately (fig. 5E, P=1.4e-5 for M-CLL,

P=1.56e-7 for U-CLL, overall *P*=2.1e-53). Cases with higher growth rate consistently showed
a shorter TTFT. Considering the growth rate as continuous variable in a multivariate Cox
regression model, it maintained a strong independent prognostic impact (*P*=2.2e-10,
HR=2.28) in the context of other well-established clinical variables, including the IGHV
mutational status and *TP53* aberrations (including both mutations and deletions) and age at
sampling (fig. 5F). The cancer N_e was more significantly correlated with OS than the growth
rate, and this effect was preserved in the U-CLL subtype (fig S12B, *P*=0.55 for M-CLL, *P*=1.0e-
6 for U-CLL, overall *P*=1.7e-9), and in multivariate setting with IGHV status, *TP53* aberrations
and age at sampling (fig S12C, *P*=0.025, HR=1.33).
477
The inference of the evolutionary parameters on our initial cohort was wholly blinded to the
clinical outcomes. Nevertheless, we validated the prognostic value of the inferred
evolutionary parameters using a second independent cohort of 210 CLL patients (136
untreated at sampling; table S2) (Oakes et al. 2016; Dietrich et al. 2018). We found that the
growth rate was prognostically relevant as a continuous (fig. S13A, *P*=7.4e-4, HR=2.39) and
dichotomic variable (fig. S13B, *P*=5.1e-5). In a multivariate Cox model with IGHV, *TP53* status
and age the growth rate approached significance (fig. S13C, *P*=0.11, HR=1.62). We note that
the power of the study in this cohort was limited, as even the well-validated IGHV status had
borderline significance (*P*=0.053). In this cohort, the predictive impact of N_e on OS was not
found to be significant (fig. S13A, *P*=0.25, HR=1.39), but we stress that TTFT is a better
reflection of the natural history of disease, not confounded by heterogenous treatment
decisions.
490
491 These results demonstrate that the evolutionary parameters inferred from a cost-effective
492 methylation array could have a direct clinical application by contributing to predict the
493 clinical behaviour of CLL patients independently from well-established prognostic variables.
494

Figure 5 – The evolutionary history of a tumour varies between molecular subtype and is predictive of clinical outcome.
A-C: The inferred growth rate (top) and effective population size (N_e, bottom) of individual cancer samples separated by molecular subtype in B-acute lymphoblastic leukaemia (**A**, BCP-ALL), mantle cell lymphoma (**B**, MCL) and chronic lymphocytic leukaemia (**C**, CLL). Differences between subtypes were tested using Mann-Whitney U tests, with Holm-Bonferroni corrections applied. In **A**, the 11q23/MLL subtype had significantly higher growth rate than each of the other B-ALL subtypes (all $P<0.005$), but for ease of presentation, only the comparison with t(1;19) is shown here. **D:** Univariate survival analysis of the time to first treatment (TTFT, blue) and overall survival (OS, red) in the discovery CLL cohort for evolutionary variables inferred via EVOFLUx. **E:** Kaplan-Meier curves comparing the TTFT between patients with high vs low inferred cancer growth rates, separated by IGHV mutational status. **F:** Multivariate Cox regression of the TTFT shows the cancer growth rate is significant when controlling for IGHV status, TP53 status and age.

Discussion

498 Our study establishes a framework, called EVOFLUx, that enables quantitative measurement
499 of the evolutionary dynamics of human malignancies. These dynamics represent the long-

term evolutionary history of the neoplasm and are fundamentally distinct from characterisations of the contemporary phenotype of cancer cells, such as the fraction of proliferating cells. The framework only requires data obtained from single-timepoint bulk tumour samples that have been processed with widely-available and low-cost DNA methylation arrays as input. The methodology should also work identically for sequencing-based measurement of DNA methylation, should generalise across cancer types, and be applicable to proxy tumour measures such as tumour-derived cell free DNA (cfDNA) extracted from blood. Consequently, this strategy enables measurement of evolutionary dynamics at massive scale, which paves the way towards clinical translation.

EVOFLUx utilises the clonal lineage identity signal that derives from stochastic fluctuations in methylated and unmethylated states at a subset of CpG sites across the genome, and outputs the evolutionary parameters defining the longitudinal dynamics of cancer growth, alongside precise measurement of DNA methylation and demethylation rates at these sites. Epimutation rates were found to differ by more than an order of magnitude between paediatric and adult malignancies. Thus, we suggest that future analyses of these quantitative data are likely to reveal new mechanistic insight into the somatic maintenance of DNA methylation.

Applying our framework in lymphoid malignancies reveals that cancer evolutionary dynamics are strongly associated with patient clinical outcomes across disease types, and further that evolutionary dynamics add significant new prognostic value to state-of-the-art approaches currently applied to the management of CLL in the clinical setting. We consider this strong evidence that clonal evolution, the fundamental cellular process of disease development, underlies the clinical course of the disease. Further, as cancer development and progression are in essence evolutionary processes, we expect these results to generalise across all cancer types. We note that genome-wide DNA methylation analyses also measure other important biological features of a cancer (for example in CLL: (Duran-Ferrer et al. 2020)), that could be combined with EVOFLUx-based inference of evolutionary dynamics to further improve the prognostic value of DNA methylation data.

In summary, we present a cost-effective high-throughput platform for measuring cancer evolutionary dynamics in patient samples. These fundamental measurements of the disease biology hold significant prognostic value and represent an innovative asset in the field of precision oncology.

Author Contributions

TAG and JIMS conceived, funded, and supervised the study. CG and TAG conceived and designed the modelling scheme and the Bayesian inference framework, with support from CPB, and CG implemented and ran this. CG, MDF, HG, JH and FN performed bioinformatic analysis. MDF and JIMS performed survival analysis. CG, HG and DM designed and ran the phylogenetic analysis. MDF, FN, NV, OK, JN, TZ, EC and ALG provided data and contributed to sample biological and/or clinical annotation. MDF, NV, JF, DS and JIMS provided clinical insight. Our study builds upon DS's earlier conception of the fluctuating methylation phenomenon. CG, MDF, JIMS and TAG wrote the manuscript, and all authors approved the final version.

Acknowledgements:

This research was primarily funded by an Accelerator award Cancer Research UK/AIRC/AECC joint funder-partnership (to JF, JIMS and TAG) and the US National Institutes of Health National Cancer Institute (U54 CA217376, to DS and TAG). Additional funding: Cancer Research UK (A19771 and DRCNPG-May21_100001) to TAG, the European Research Council under the European Union's Horizon 2020 Research and Innovation Program (810287, BCLLatlas, to EC and JIMS), La Caixa Foundation (CLLEvolution LCF/PR/HR17/52150017 [HR17- 00221LCF] and CLLSYSTEMS - LCF/PR/HR22/52420015 [HR22-00172] Health Research 2017 and 2022 Programs, to EC), Generalitat de Catalunya Suport Grups de Recerca AGAUR (2021-SGR-01343 to JIMS and 2021-SGR-01172 to EC). JN and OK were supported by the Swedish Research Council (2019-01976), Swedish Childhood Cancer Fund (PR2019-0046/PR2022-0082), and Swedish Cancer Society (CAN2022-2395). CG was further supported by the BBSRC London Interdisciplinary Doctoral Programme (LIDo) and Thorton Foundation funding to the Institute of Cancer Research. MDF was supported by a postdoctoral fellowship of the AECC Scientific Foundation. This research utilized the Cancer Research UK City of London High Performance Computing (HPC) facility, and was partially developed at the Centro Esther Koplowitz (CEK, Barcelona, Spain). We acknowledge the SciLifeLab National Genomics Infrastructure, SNP&SEQ Technology Platform in Upppsala, Sweden, funded by the Swedish Research Council and the Knut and Alice Wallenberg Foundation, for assistance with DNA methylation analyses.

Methods

Assembly and quality control of DNA methylation data

We assembled and processed with a harmonized pipeline (Duran-Ferrer et al. 2020) (version 4.1, see code availability section) 2,430 bulk sample Illumina methylation array data of normal and neoplastic lymphoid cells from previous publications (Kulis et al. 2015; Nordlund et al. 2013; Reinius et al. 2012; Lee et al. 2015; Queirós et al. 2016; Nadeu et al. 2020; Duran-Ferrer et al. 2020; Nadeu et al. 2022; Oakes et al. 2016; Dietrich et al. 2018; Agirre et al. 2015). Briefly, raw idat files were loaded and processed with R (version 4.3.1) using *minfi* package (Aryee et al. 2014; Fortin, Triche Jr, and Hansen 2017) (version 1.46.0) in batches as specified in the column "SSNOB_NORMALIZATION_BATCH" of table S2. Briefly, the data was processed for each batch as follows. First, idats files were loaded into a *RGChannelSet* object, and *minfi* quality metrics using the *qcReport* function were performed, controlling for biased distributions of methylation values and low signal intensities of internal control probes for each sample. Next, further quality metrics were derived using the function *minfiQC* on unnormalized *RGChannelSet* obejct. Those samples with median signal intensities of unmethylated and methylated channels of at least 10.5 in log2 scale were considered as having good signal intensities (default value in *minfi*). Subsequently, detection *P*-values were calculated across all CpGs and samples using the *detectionP* function for the unnormalized *RGChannelSet* object. Samples were considered as good if having a mean detection P-value across all CpGs pf $P \leq 0.01$. On a CpG level, we retained CpGs with a detection *P*-value $\leq 1e^{-16}$ in $\geq 90\%$ of the samples. The *RGChannelSet* object was normalized with the *preprocessNoob* function. We next retained only CpGs (excluding CH probes) that did not contain any single nucleotide polymorphism neither in the interrogated CpGs nor in the probe using the *dropMethylationLoci* and *dropLociWithSnps* functions. Furthermore,

CpGs with any previous evidence of potential cross-hybridization were excluded (Chen et al.
2013) and only CpGs mapping to autosomal chromosomes were subsequently retained for
downstream analyses. Finally, we checked the distribution of normalized methylation values
and performed principal component analyses separately for samples passing all quality
checks as well as those considered to as bad samples. The final DNA methylation matrix
contained 2,204 samples and 389,180 CpGs passing all the aforementioned quality controls,
and included 2,054 patients (22 technical replicates, 3 synchronic and 125 longitudinal
samples from the same patients).
599
To determine to purity of samples, we used our previously deconvolution strategy to infer
tumor cell content by DNA methylation (Duran-Ferrer et al. 2020), which was used as a
consensus purity in all the tumor samples except for DLBCL and MM. In these two tumor
entities, we have previously identified a DNA methylation signature loss causing inaccurate
tumor purity predictions using DNA methylation data, and therefore we used available
genetic or flow cytometry data for DLBCL and MM, respectively.
606

Statistical analysis

607
608 Statistical tests performed throughout the study were performed as two-sided. Appropriate
609 multiple test correction, such as the holm-sidak (hs) correction, is noted when applied.
610

Characterisation of fCpGs

611
To characterise the genomic and regulatory context of fCpGs we employed a series of
statistical analyses and database annotations. We annotated fCpGs using Illumina manifest
and other genomic annotationa available at Bioconductor packages
IlluminaHumanMethylation450kanno.ilmn12.hg19 (version 0.6.1) and
IlluminaHumanMethylationEPICanno.ilm10b2.hg19 (version 0.6.0). We used Chi-squared
tests χ^2 to assess the enrichment of fCpGs in distinct genomic regions or elements. We also
used CpG annotations of the regulatory features provided by the ENCODE Methylation
Consortium (Dunham et al. 2012). We performed gene-set enrichment analysis upon the
fCpG-associated genes using gProfiler (Raudvere et al. 2019), specifically focusing upon the
Gene Ontology biological processes (Ashburner et al. 2000) and the Human Protein Atlas
(Uhlén et al. 2015). The statistical domain space was limited to genes targeted by at least
one CpG in the 389,180 candidate CpG set and significance was determined using the g:SCS
algorithm (Reimand et al. 2007). Previous chromatin segmentation of normal and neoplastic
B cells was used to assess the chromatin state enrichment of fCpG (Duran-Ferrer et al. 2020;
Beekman et al. 2018).
627
fCpG were checked for their overlap with previous "epigenetic clocks", including mitotic
(Duran-Ferrer et al. 2020; Yang et al. 2016; Teschendorff 2020; Youn and Wang 2018; Zhou
et al. 2018), chronological age (Bocklandt et al. 2011; Garagnani et al. 2012; Hannum et al.
2013; Horvath 2013; Lin et al. 2016; Vidal-Bralo, Lopez-Golan, and Gonzalez 2016; Weidner
et al. 2014; Zhang et al. 2019; Horvath et al. 2018; Shireby et al. 2020), gestational age
(Bohlin et al. 2016; Knight et al. 2016; Y. Lee et al. 2019; Mayne et al. 2017; McEwen et al.
2020), biological age and mortality (Belsky et al. 2022; Levine et al. 2018; Lu et al. 2019), and
trait predictors (McCartney et al. 2018; Liang et al. 2020). The package *methylCIPHER*

(https://github.com/MorganLevineLab/methylCIPHER) was used to integrate the majority of
the epigenetic clocks.
638

CLL RNA-seq data

Previously available RNA-seq data for 294 CLL patients was obtained (Puente et al. 2015)
and processed as previously described (Nadeu et al. 2020). Matched RNA-seq data and DNA
methylation data for the same patients at the same timepoint was available for 224 CLL
patients. Transcript per million counts (tpms) were used to represent differential gene
expression values across genes and samples. We used the gene annotation provided in the R
Bioconductor package *IlluminaHumanMethylationEPICanno.ilm10b2.hg19* to classify genes
associated with fCpGs. Genes targeted by any fCpG were considered as "fCpG genes".
647

In each methylation sample, the 978 fCpG were discretised as homozygous demethylated,
heterozygous methylated or homozygous methylated (coded as [0,1,2] respectively). This
was done by separately fitting a beta mixture model with 3 components to each sample
using Stan (Carpenter et al. 2017) and extracting the component mixture probability. The
gene expression value for genes classified as having and fCpG with 0, 1 or 2 alleles
methylated were plotted as previously described.
654

A Stochastic Model of how fCpGs vary in large, neutrally growing populations of cells

657 We built a generative computational model of how the patterns of fCpGs vary over time (t)
658 according to the evolutionary history of a cancer. Initially, our model focused on neutral
659 evolution, before expanding to non-neutral modes of tumour evolution below. For the full
660 explanation of the model, please see the supplementary information.
661

Our model was parameterised in terms of the age of the patient at which the most recent
common ancestor (MRCA) emerged (τ), the exponential growth rate of the cancer (θ), and
the epigenetic switching rates of the fCpGs (μ, γ, ν, ζ). The model was partitioned into 2
phases, prior and post the emergence of the MRCA. At time $t = 0$, the fCpGs were assumed
to be equally likely to be homozygously methylated or demethylated. The fCpG status of the
MRCA at time $t = \tau$ was calculated by applying matrix exponentiation.
668

The second phase of the model consisted of a discrete time Markov process. The effective
population size of the growing cancer was modelled as growing according to a deterministic
exponential growth equation, $N_e = e^{\theta(t-\tau)}$. Each fCpG was considered independently – at
each time step, $t \rightarrow t + \delta t$, the number of homozygous methylated (m), heterozygous
methylated (k) and homozygous demethylated cells (w) at a specific fCpG was updated
according to the epigenetic switching rates.
675

676 At the time of sample, T, the fraction methylation of each simulated fCpG was calculated by
677 summing the number of methylated alleles and normalising by the total number of alleles in
678 the   population:

$$\beta_c = \frac{k + 2m}{2N_e}$$

We further accounted for contaminating normal cells and the technical noise introduced by
the methylation bead array. The methylation of the contaminated samples was assumed to
be an average of the cancer methylation, $\beta_c(t)$, weighted by the tumour purity ρ, and the
average of the normal population, β_n, weighted by $1 - \rho$. Following our previous work, the
bead array was assumed to saturate at extreme methylation values, shifting the minimum
and maximum methylation by δ and ε respectively. The noise of the bead array was
assumed to be beta distributed, with precision parameter κ.
686

Non-neutral models of tumour evolution

688   Alongside our model of neutral exponentially growing cancer populations, we devised two
689   alternative models of cancer growth:
690   1. A subclonal selection model in which a single cell within the cancer develops a
691   selective advantage and begins to grow at an increased growth rate.
692   2. An independent clonal origins model, whereby a patient has developed two distinct
693   cancers concurrently.
694

For the subclonal selection model, we replaced the growth rate (θ) and the time of the
MRCA (τ) with the growth rates and time of the MRCA of the initial, slower-growing
population $(\theta_1$ and τ_1 respectively), and that of the more recently emerging, faster-growing
population $(\theta_2$ and $\tau_2)$, constraining $\tau_1 < \tau_2$ and $\theta_1 < \theta_2$ (fig. S7C). We assumed that the
initial cancer population begins exponentially growing at τ_1 as above, but at time $t = \tau_2$ we
select a single cell with a set of fCpG states drawn according to the cancer population and
allow this second population to grow concurrently with a growth rate θ_2.
702

The independent-cancer model followed the same scheme as the nested subclonal selection
model, except the methylation status of the emerging cancer was that of an independent
cell which experienced random fluctuations between $t = 0$ and $t = \tau_2$.
If we let the number of cells in the less fit subclone in each methylation state be
$\{m_1, k_1, w_1\}$ and in the fitter subclone be $\{m_2, k_2, w_2\}$, following the convention above,
then in both cases the measured methylation patterns at the time of sample are:

$$\beta_c(t) = \frac{k_1(t) + 2m_1(t) + k_2(t) + 2m_2(t)}{2N_e(t)}$$

709   Where $N_e(t) = e^{\theta_1(t-\tau_1)} + e^{\theta_2(t-\tau_2)}$.
710

Prior functions

For each methylation array blood sample, we had matched age (T) and purity (ρ)
information. Hence, the parameters to be inferred are the growth rate (θ), the age of the
patient when the MRCA emerged (τ), the epigenetic switching rates $(\mu, \upsilon, \gamma, \zeta$; defined in fig
2A), the average fraction methylated of contaminating normal cells (β_n), the beta offsets
from 0 and 1 due to the background noise on the methylation array (Δ and ϵ, respectively),
and the precision of the beta distributed noise (κ).
718

These parameters are constrained either to be positive $(\theta, \mu, \upsilon, \gamma, \zeta, \kappa > 0)$ or to lie within a
specified range $(0 < \frac{\tau}{T}, \beta_n, \Delta, \epsilon < 1)$, which we achieved using appropriate prior
distributions. To better allow for priors to be set on a biologically meaningful scale, the
priors for the lognormal distribution were set in terms of the real scale mean and standard

deviation, rather than the standard log-scale. To reduce correlations in the posterior and
make sampling more efficient, the variables v and ζ were normalised by μ and γ
respectively.

The priors are as follows:

$$\theta \sim \text{lognormal}(3, 2)$$
$$\frac{\tau}{T} \sim \text{beta}(2, 2)$$
$$\mu \sim \text{halfnormal}(0, 0.05)$$
$$\gamma \sim \text{halfnormal}(0, 0.05)$$
$$\frac{v}{\mu} \sim \text{lognormal}(1, 0.7)$$
$$\frac{\zeta}{\gamma} \sim \text{lognormal}(1, 0.7)$$
$$\beta_n \sim \text{beta}(2, 2)$$
$$\Delta \sim \text{beta}(5, 95)$$
$$\epsilon \sim \text{beta}(95, 5)$$
$$\kappa \sim \text{halfnormal}(100, 30)$$

729   When fitting non-neutral models of tumour growth, the inference is parameterised in terms
730   of the relative growth of the fitter subclone, $\tilde{\theta}_2 = \frac{\theta_2}{\theta_1}$, and the fraction of the population
731   consisting of the fitter subclone, $f = \frac{e^{\theta_2(t-\tau_2)}}{e^{\theta_1(t-\tau_1)}+e^{\theta_2(t-\tau_2)}}$. The age at which the second clone
732   emerges is then:

$$\tau_2 = T - \frac{(T-\tau_1)\theta_1}{\theta_2} - \frac{\text{logit}(f)}{\theta_2}$$

733   This parameterisation induces less correlations in the resulting posterior, which significantly
734   improves the sampling efficiency. The priors on these additional parameters are:

$$\frac{\tau_1}{T} \sim \text{beta}(2, 2)$$
$$\tilde{\theta}_2 \sim \text{lognormal}(1, 0.7)$$
$$f \sim \text{beta}(2, 2)$$

All the other priors were the same as in the neutral case.

Bayesian Inference
We developed a stochastic estimator of the loglikelihood function at a given set of
parameters by simulating the fCpG methylation distribution a large number of times,
correcting for the bias inherent with using a finite number of simulations and penalising the
loglikelihood for extreme values of the N_e (see supplementary for details).

The standard Bayesian algorithms developed to infer the posterior for a given set of data
(e.g. MCMC, nested sampling) are typically used when the log-likelihood is analytically
tractable and can be calculated exactly. Remarkably, it has been shown that, as long as the
stochastic approximation of the log-likelihood is unbiased, MCMC methods can obtain an
exact Bayesian inference of the true posterior, as in pseudo-marginal Metropolis–Hastings
(Andrieu and Roberts 2009).

Here, we employed a nested sampling approach using the dynesty package (Skilling 2004;
2006; Speagle 2019). Unlike pseudo-marginal Metropolis–Hastings, nested sampling is able
to efficiently explore multi-modal posterior landscapes (which can occur under the
subclonal and independent cancer models).

Model comparison between non-neutral models of tumour evolution
756
We employed an expected log pointwise predictive density (ELPD) (Vehtari, Gelman, and
Gabry 2015) approach to compare our competing models of evolution for each sample using
the arviz Python package (Kumar et al. 2019), which uses PSIS-LOO-CV to compare the out-
of-sample prediction accuracy between models whilst naturally penalising more complex
models. This required the loglikelihood per datapoint and the posterior predictive for every
point in the posterior. The weights of the respective models were calculated using pseudo-
Bayesian Model averaging using Akaike-type weighting, stabilized using the Bayesian
bootstrap (Yao et al. 2017).

CLL and RT genomic analyses
766
767 Previous mutated annotation files (MAF) from WES data (Knisbacher et al. 2022) and WGS
768 (Nadeu et al. 2022) were used to further validate our distinct EVOFLUx evolutionary modes
769 (i.e., neutral, subclonal and independent) and RT phylogenies.

Subclonal deconvolution of WES and WGS data
771
To detect subclones in bulk WES and WGS data, we employed MOBSTER (Caravagna et al.
2020), which fits the variant allele frequency (VAF) spectrum with a mixture model
containing a Pareto distribution to account for the neutral tail (M. J. Williams et al. 2016)
and a variable number of beta distributions to account for the clonal and subclonal peaks.

We ran MOBSTER using the default parameters, except employing a minimum 5% VAF
threshold and lowering the minimum number of mutations to compose a cluster to 5 in WES
samples due to the low number of mutations. We then manually QC'd all 377 WES samples
and 10 WGS, tuning the fitting parameters to better represent the data (for instance, when
the clonal peak had been called at a low frequency despite the median tumour purity being
95%).

Phylogenetic inference of longitudinal methylation data
784
A novel Bayesian phylogenetic method was used to reconstruct the evolutionary
relationships and the time to most recent common ancestor (tMRCA) of longitudinal
samples from the same patients. This was carried out in the BEAST v.1.8.4 framework
(Drummond and Rambaut 2007; Drummond et al. 2012) using custom models implemented
in PISCA v1.1 (available from https://github.com/adamallo/PISCA) (Martinez et al. 2018).

EVOFLUx provided an estimate of the age of the patient when the MRCA of each bulk
sample emerged. To estimate the methylation status of each fCpG at the sample's MRCA in
each of our longitudinal samples, we discretised the fCpGs as described above (RNAseq
Methods).

We implemented a 4-parameter biallelic binary substitution model analogous to the pre-
growth EVOFLUx model in PISCA. This plugin contains all the required statistical machinery
to use this model for somatic phylogenetic estimation. The biallelic binary substitution
model has three relative rate parameters 1) heterozygous methylation $\tilde{\upsilon}$, 2) homozygous
demethylation $\tilde{\gamma}$, and 3) heterozygous demethylation $\tilde{\zeta}$, where homozygous methylation $\tilde{\mu}$
was normalised to 1. For all relative transition rate parameters, a lognormal prior with mean
1 and standard deviation of 0.6 was used, with a half normal prior with mean 0 and
standard deviation 0.13 for the molecular clock rate, using a strict clock model for the rate
of evolution across the tree. Two demographic tree models, constant population size
(Kingman 1982) and exponential growth (Griffiths and Tavaré 1994), were compared by
marginal likelihood estimation using path-sampling (Baele et al. 2012) and a constant
population model was deemed more appropriate.
808
809 MCMC chains were run for 100 million generations sampled every 100,000 generations and
810 convergence was assessed using Tracer v.1.7 (Rambaut et al. 2018), ensuring effective
811 sample sizes greater than 500 for all parameters. Maximum clade credibility trees were then
812 made using 10% burn-in and medium node heights. The resulting trees were plotted using
813 ggtree (Yu et al. 2017).
814

Phylogenetic inference of SNVs from WGS data

Each bulk sample is represented by a set of clonal mutations found during the
deconvolution of WGS data (see above). Where a mutation was deemed absent in the clonal
peak, the reference nucleotide was used. Mutational signature assignment (Díaz-Gay et al.
2023) was used to select mutations in the clock-like SBS1 channel (Tate et al. 2019). BEAST
v.1.10 (Suchard et al. 2018) was then used with the simple binary substitution model (since
SBS1 effectively represents just C to T substitutions), a strict clock model, a constant
population size prior (Kingman 1982), and a flat prior on the age of most recent common
ancestor (from zero to earliest patient sample), with ancestral state estimation at the root.
Chains were run and ESS values assessed as described above. The distance between the
ancestral state of the root at each MCMC state, and the clock rate were used to calculate
the expected evolution distance between the root and the known germline. This was used
to inform the length of the branch between germline (at birth) and the MRCA of the
samples.
829

Survival Analysis

Clinical analyses were performed in CLL for time to first treatment (TTFT) and overall
survival (OS) from the time of sampling. Tumour growth rate (θ), effective population size
(N_e) and epigenetic switching rates were analysed as continuous variables in univariate Cox
regression models for both TTFT and OS. The effect size of hazard ratios (HRs) for each
evolutionary variable were analysed considering different scaling factors. In particular, the
growth rate was analysed in exponential growth per year (i.e., for $\theta = 1$, the population per
year is e=2.71 times bigger) ; the N_e was considered per million cells; the cancer age or time
from the MRCA was analysed for each 10 years; the homozygous to heterozygous
(de)methylation rates (γ and μ, respectively) and heterozygous to homozygous methylation
rate (υ) were analysed multiplied by a factor of 100 (per allele per year); and heterozygous
to homozygous demethylation rate (ζ) was multiplied by a factor of 1,000. In addition,

growth rate and effective population were analysed as continuous variables in multivariate
Cox regression models together with *TP53* aberrations (considering mutations and deletions
together), IGHV gene mutational status and the age of patients at sampling. Kaplan-Meier
(KM) curves were generated for low and high growth rate and effective population size
within IGHV subtypes using maximally selected log-rank statistic using maxstats package
(version 0.7-25). *P* values from KM curves were derived using the Log-rank statistic. Survival
(version 3.5-7), survminer (version 0.4.9) and ggsurvfit (version 0.3.1) packages were used
under R version 4.3.1.

Data availability

851　No new data was generated in the course of this study. The harmonised and filtered
852　methylation matrix is available (preferred browser Firefox) at: http://inb-
853　cg.bsc.es/hcli_priv/publications_data/fluctuating_cpgs_gabbutt_duran_ferrer_2023/QC_22
854　04_ssNob.tsv.7z - the password for this file is available upon request to the authors.
855
Previously published DNA methylation data re-analysed in this study can be found under
accession codes: B cells, EGAS00001001196;
ALL, GSE56602, GSE49032, GSE76585, GSE69229;
MCL, EGAS00001001637, EGAS00001004165; CLL, EGAD00010000871, EGAD00010000948,
EGAD00010001975; MM, EGAS00001000841; DLBCL, EGAD00010001974.

CLL gene expression data is available EGAS00001000374 and EGAS00001001306. ChIP-seq
datasets are available from Blueprint https://www.blueprint-epigenome.eu/ under the
accession EGAS00001000326.

Matched WES and WGS are available under accessions EGAS00000000092 and
EGAD00001008954 respectively.

Code Availability

867　The EVOFLUx code employed to infer the evolutionary history of cancer samples from
868　methylation array data is attached as a bundled Zip file and will be made publicly available
869　via Github prior to publication at https://github.com/CalumGabbutt. Further code will also
870　be made available at https://github.com/Duran-FerrerM and
871　https://github.com/adamallo/PISCA.

Ethics statement

873　The study was approved by the ethics committees of the institutions that generated the
874　data used in the present study.

Reference

Agirre, Xabier, Giancarlo Castellano, Marien Pascual, Simon Heath, Marta Kulis, Victor
Segura, Anke Bergmann, et al. 2015. 'Whole-Epigenome Analysis in Multiple Myeloma
Reveals DNA Hypermethylation of B Cell-Specific Enhancers'. *Genome Research* 25 (4):
478–87. https://doi.org/10.1101/GR.180240.114.
Alizadeh, Ash A., Michael B. Elsen, R. Eric Davis, Ch L. Ma, Izidore S. Lossos, Andreas
Rosenwald, Jennifer C. Boldrick, et al. 2000. 'Distinct Types of Diffuse Large B-Cell

Lymphoma Identified by Gene Expression Profiling'. *Nature* 403 (6769): 503–11.
https://doi.org/10.1038/35000501.

Andrieu, Christophe, and Gareth O. Roberts. 2009. 'The Pseudo-Marginal Approach for
Efficient Monte Carlo Computations'. *Https://Doi.Org/10.1214/07-AOS574* 37 (2): 697–
725. https://doi.org/10.1214/07-AOS574.

Arber, Daniel A., Attilio Orazi, Robert P. Hasserjian, Michael J. Borowitz, Katherine R. Calvo,
Hans Michael Kvasnicka, Sa A. Wang, et al. 2022. 'International Consensus Classification
of Myeloid Neoplasms and Acute Leukemias: Integrating Morphologic, Clinical, and
Genomic Data'. *Blood* 140 (11): 1200–1228.
https://doi.org/10.1182/BLOOD.2022015850.

Aryee, Martin J, Andrew E Jaffe, Hector Corrada-Bravo, Christine Ladd-Acosta, Andrew P
Feinberg, Kasper D Hansen, and Rafael A Irizarry. 2014. 'Minfi: A Flexible and
Comprehensive Bioconductor Package for the Analysis of Infinium DNA Methylation
Microarrays'. *Bioinformatics* 30 (10): 1363–69.
https://doi.org/10.1093/bioinformatics/btu049.

Ashburner, Michael, Catherine A. Ball, Judith A. Blake, David Botstein, Heather Butler, J.
Michael Cherry, Allan P. Davis, et al. 2000. 'Gene Ontology: Tool for the Unification of
Biology'. *Nature Genetics 2000 25:1* 25 (1): 25–29. https://doi.org/10.1038/75556.

Baele, Guy, Philippe Lemey, Trevor Bedford, Andrew Rambaut, Marc A. Suchard, and
Alexander V. Alekseyenko. 2012. 'Improving the Accuracy of Demographic and
Molecular Clock Model Comparison While Accommodating Phylogenetic Uncertainty'.
Molecular Biology and Evolution 29 (9): 2157–67.
https://doi.org/10.1093/MOLBEV/MSS084.

Beekman, Renée, Vicente Chapaprieta, Núria Russiñol, Roser Vilarrasa-Blasi, Núria
Verdaguer-Dot, Joost H.A. Martens, Martí Duran-Ferrer, et al. 2018. 'The Reference
Epigenome and Regulatory Chromatin Landscape of Chronic Lymphocytic Leukemia'.
Nature Medicine 2018 24:6 24 (6): 868–80. https://doi.org/10.1038/s41591-018-0028-
4.

Belsky, D. W., A. Caspi, D. L. Corcoran, K. Sugden, R. Poulton, L. Arseneault, A. Baccarelli, et
al. 2022. 'DunedinPACE, A DNA Methylation Biomarker of the Pace of Aging'. *ELife* 11
(January). https://doi.org/10.7554/ELIFE.73420.

Bocklandt, Sven, Wen Lin, Mary E. Sehl, Francisco J. Sánchez, Janet S. Sinsheimer, Steve
Horvath, and Eric Vilain. 2011. 'Epigenetic Predictor of Age'. *PLOS ONE* 6 (6): e14821.
https://doi.org/10.1371/JOURNAL.PONE.0014821.

Bohlin, J., S. E. Håberg, P. Magnus, S. E. Reese, H. K. Gjessing, M. C. Magnus, C. L. Parr, C. M.
Page, S. J. London, and W. Nystad. 2016. 'Prediction of Gestational Age Based on
Genome-Wide Differentially Methylated Regions'. *Genome Biology* 17 (1): 1–9.
https://doi.org/10.1186/S13059-016-1063-4/TABLES/5.

Brocks, David, Yassen Assenov, Sarah Minner, Olga Bogatyrova, Ronald Simon, Christina
Koop, Christopher Oakes, et al. 2014. 'Intratumor DNA Methylation Heterogeneity
Reflects Clonal Evolution in Aggressive Prostate Cancer'. *Cell Reports* 8 (3): 798–806.
https://doi.org/10.1016/J.CELREP.2014.06.053/ATTACHMENT/0EB6C7A6-FCFD-4977-
A0EC-D8324921D744/MMC1.PDF.

Caravagna, Giulio, Timon Heide, Marc J. Williams, Luis Zapata, Daniel Nichol, Ketevan
Chkhaidze, William Cross, et al. 2020. 'Subclonal Reconstruction of Tumors by Using
Machine Learning and Population Genetics'. *Nature Genetics 2020 52:9* 52 (9): 898–
907. https://doi.org/10.1038/s41588-020-0675-5.

Carpenter, Bob, Andrew Gelman, Matthew D Hoffman, Daniel Lee, Ben Goodrich, Michael
Betancourt, Marcus Brubaker, Jiqiang Guo, Peter Li, and Allen Riddell. 2017. 'Stan: A
Probabilistic Programming Language'. *Journal of Statistical Software; Vol 1, Issue 1*
(2017), January. https://www.jstatsoft.org/v076/i01.
Chen, Yi An, Mathieu Lemire, Sanaa Choufani, Darci T. Butcher, Daria Grafodatskaya, Brent
W. Zanke, Steven Gallinger, Thomas J. Hudson, and Rosanna Weksberg. 2013.
'Discovery of Cross-Reactive Probes and Polymorphic CpGs in the Illumina Infinium
HumanMethylation450 Microarray'. *Epigenetics* 8 (2): 203–9.
https://doi.org/10.4161/epi.23470.
Damle, Rajendra N., Tarun Wasil, Franco Fais, Fabio Ghiotto, Angelo Valetto, Steven L. Allen,
Aby Buchbinder, et al. 1999. 'Ig V Gene Mutation Status and CD38 Expression As Novel
Prognostic Indicators in Chronic Lymphocytic Leukemia'. *Blood* 94 (6): 1840–47.
https://doi.org/10.1182/BLOOD.V94.6.1840.
Díaz-Gay, Marcos, Raviteja Vangara, Mark Barnes, Xi Wang, S M Ashiqul Islam, Ian Vermes,
Nithish Bharadhwaj Narasimman, et al. 2023. 'Assigning Mutational Signatures to
Individual Samples and Individual Somatic Mutations with SigProfilerAssignment'.
BioRxiv, July, 2023.07.10.548264. https://doi.org/10.1101/2023.07.10.548264.
Dietrich, Sascha, Małgorzata Oleś, Junyan Lu, Leopold Sellner, Simon Anders, Britta Velten,
Bian Wu, et al. 2018. 'Drug-Perturbation-Based Stratification of Blood Cancer'. *The*
Journal of Clinical Investigation 128 (1): 427–45. https://doi.org/10.1172/JCI93801.
Drummond, Alexei J., and Andrew Rambaut. 2007. 'BEAST: Bayesian Evolutionary Analysis
by Sampling Trees'. *BMC Evolutionary Biology* 7 (1). https://doi.org/10.1186/1471-
2148-7-214.
Drummond, Alexei J., Marc A. Suchard, Dong Xie, and Andrew Rambaut. 2012. 'Bayesian
Phylogenetics with BEAUti and the BEAST 1.7'. *Molecular Biology and Evolution* 29 (8):
1969–73. https://doi.org/10.1093/MOLBEV/MSS075.
Dunham, Ian, Anshul Kundaje, Shelley F. Aldred, Patrick J. Collins, Carrie A. Davis, Francis
Doyle, Charles B. Epstein, et al. 2012. 'An Integrated Encyclopedia of DNA Elements in
the Human Genome'. *Nature* 489 (7414): 57–74.
https://doi.org/10.1038/NATURE11247.
Duran-Ferrer, Martí, Guillem Clot, Ferran Nadeu, Renée Beekman, Tycho Baumann, Jessica
Nordlund, Yanara Marincevic-Zuniga, et al. 2020. 'The Proliferative History Shapes the
DNA Methylome of B-Cell Tumors and Predicts Clinical Outcome'. *Nature Cancer* 1 (11):
1066–81. https://doi.org/10.1038/s43018-020-00131-2.
Endicott, Jamie L., Paula A. Nolte, Hui Shen, and Peter W. Laird. 2022. 'Cell Division Drives
DNA Methylation Loss in Late-Replicating Domains in Primary Human Cells'. *Nature*
Communications 2022 13:1 13 (1): 1–12. https://doi.org/10.1038/s41467-022-34268-8.
Fortin, Jean-Philippe, Timothy J Triche Jr, and Kasper D Hansen. 2017. 'Preprocessing,
Normalization and Integration of the Illumina HumanMethylationEPIC Array with
Minfi'. *Bioinformatics* 33 (4): 558–60. https://doi.org/10.1093/bioinformatics/btw691.
Gabbutt, Calum, Ryan O. Schenck, Daniel J. Weisenberger, Christopher Kimberley, Alison
Berner, Jacob Househam, Eszter Lakatos, et al. 2022. 'Fluctuating Methylation Clocks
for Cell Lineage Tracing at High Temporal Resolution in Human Tissues'. *Nature*
Biotechnology 2022, January, 1–11. https://doi.org/10.1038/s41587-021-01109-w.
Gaiti, Federico, Ronan Chaligne, Hongcang Gu, Ryan M Brand, Steven Kothen-Hill, Rafael C
Schulman, Kirill Grigorev, et al. 2019. 'Epigenetic Evolution and Lineage Histories of

Chronic Lymphocytic Leukaemia'. *Nature* 569 (7757): 576–80. https://doi.org/10.1038/s41586-019-1198-z.

Garagnani, Paolo, Maria G. Bacalini, Chiara Pirazzini, Davide Gori, Cristina Giuliani, Daniela Mari, Anna M. Di Blasio, et al. 2012. 'Methylation of ELOVL2 Gene as a New Epigenetic Marker of Age'. *Aging Cell* 11 (6): 1132–34. https://doi.org/10.1111/ACEL.12005.

Greaves, Mel, and Carlo C. Maley. 2012. 'Clonal Evolution in Cancer'. *Nature 2012 481:7381* 481 (7381): 306–13. https://doi.org/10.1038/nature10762.

Griffiths, R. C., and S. Tavaré. 1994. 'Sampling Theory for Neutral Alleles in a Varying Environment'. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* 344 (1310): 403–10. https://doi.org/10.1098/RSTB.1994.0079.

Gruber, Michaela, Ivana Bozic, Ignaty Leshchiner, Dimitri Livitz, Kristen Stevenson, Laura Rassenti, Daniel Rosebrock, et al. 2019. 'Growth Dynamics in Naturally Progressing Chronic Lymphocytic Leukaemia'. *Nature* 570 (7762): 474–79. https://doi.org/10.1038/S41586-019-1252-X.

Gutierrez, Catherine, Aziz M. Al'Khafaji, Eric Brenner, Kaitlyn E. Johnson, Satyen H. Gohil, Ziao Lin, Binyamin A. Knisbacher, et al. 2021. 'Multifunctional Barcoding with ClonMapper Enables High-Resolution Study of Clonal Dynamics during Tumor Evolution and Treatment'. *Nature Cancer* 2 (7): 758–72. https://doi.org/10.1038/S43018-021-00222-8.

Hamblin, Terry J., Zadie Davis, Anne Gardiner, David G. Oscier, and Freda K. Stevenson. 1999. 'Unmutated Ig VH Genes Are Associated With a More Aggressive Form of Chronic Lymphocytic Leukemia'. *Blood* 94 (6): 1848–54. https://doi.org/10.1182/BLOOD.V94.6.1848.

Hannum, Gregory, Justin Guinney, Ling Zhao, Li Zhang, Guy Hughes, Srini Vas Sadda, Brandy Klotzle, et al. 2013. 'Genome-Wide Methylation Profiles Reveal Quantitative Views of Human Aging Rates'. *Molecular Cell* 49 (2): 359–67. https://doi.org/10.1016/J.MOLCEL.2012.10.016.

Hao, Jia Jie, De Chen Lin, Huy Q. Dinh, Anand Mayakonda, Yan Yi Jiang, Chen Chang, Ye Jiang, et al. 2016. 'Spatial Intratumoral Heterogeneity and Temporal Clonal Evolution in Esophageal Squamous Cell Carcinoma'. *Nature Genetics 2016 48:12* 48 (12): 1500–1507. https://doi.org/10.1038/ng.3683.

He, Xiaofei, Deng Cai, and Partha Niyogi. 2005. 'Laplacian Score for Feature Selection'. In *Advances in Neural Information Processing Systems*, edited by Y Weiss, B Schölkopf, and J Platt. Vol. 18. MIT Press. https://proceedings.neurips.cc/paper/2005/file/b5b03f06271f8917685d14cea7c6c50a-Paper.pdf.

Heide, Timon, Luis Zapata, Marc J. Williams, Benjamin Werner, Giulio Caravagna, Chris P. Barnes, Trevor A. Graham, and Andrea Sottoriva. 2018. 'Reply to "Neutral Tumor Evolution?"' *Nature Genetics 2018 50:12* 50 (12): 1633–37. https://doi.org/10.1038/s41588-018-0256-z.

Hong, You Jin, Paul Marjoram, Darryl Shibata, and Kimberly D. Siegmund. 2010. 'Using DNA Methylation Patterns to Infer Tumor Ancestry'. *PloS One* 5 (8). https://doi.org/10.1371/JOURNAL.PONE.0012002.

Horvath, Steve. 2013. 'DNA Methylation Age of Human Tissues and Cell Types'. *Genome Biology* 14 (10): R115. https://doi.org/10.1186/gb-2013-14-10-r115.

Horvath, Steve, Junko Oshima, George M. Martin, Ake T. Lu, Austin Quach, Howard Cohen, Sarah Felton, et al. 2018. 'Epigenetic Clock for Skin and Blood Cells Applied to

Hutchinson Gilford Progeria Syndrome and *Ex Vivo* Studies'. *Aging* 10 (7): 1758–75. https://doi.org/10.18632/AGING.101508.

Kingman, J. F.C. 1982. 'The Coalescent'. *Stochastic Processes and Their Applications* 13 (3): 235–48. https://doi.org/10.1016/0304-4149(82)90011-4.

Knight, Anna K., Jeffrey M. Craig, Christiane Theda, Marie Bækvad-Hansen, Jonas Bybjerg-Grauholm, Christine S. Hansen, Mads V. Hollegaard, et al. 2016. 'An Epigenetic Clock for Gestational Age at Birth Based on Blood Methylation Data'. *Genome Biology* 17 (1): 1–11. https://doi.org/10.1186/S13059-016-1068-Z/FIGURES/4.

Knisbacher, Binyamin A., Ziao Lin, Cynthia K. Hahn, Ferran Nadeu, Martí Duran-Ferrer, Kristen E. Stevenson, Eugen Tausch, et al. 2022. 'Molecular Map of Chronic Lymphocytic Leukemia and Its Impact on Outcome.' *Nature Genetics* 54 (11): 1664–74. https://doi.org/10.1038/S41588-022-01140-W.

Kulis, Marta, Angelika Merkel, Simon Heath, Ana C. Queirós, Ronald P. Schuyler, Giancarlo Castellano, Renée Beekman, et al. 2015. 'Whole-Genome Fingerprint of the DNA Methylome during Human B Cell Differentiation'. *Nature Genetics 2015 47:7* 47 (7): 746–56. https://doi.org/10.1038/ng.3291.

Kumar, Ravin, Colin Carroll, Ari Hartikainen, and Osvaldo Martin. 2019. 'ArviZ a Unified Library for Exploratory Analysis of Bayesian Models in Python'. *Journal of Open Source Software* 4 (33): 1143. https://doi.org/10.21105/JOSS.01143.

Lee, Seung Tae, Marcus O. Muench, Marina E. Fomin, Jianqiao Xiao, Mi Zhou, Adam De Smith, José I. Martín-Subero, et al. 2015. 'Epigenetic Remodeling in B-Cell Acute Lymphoblastic Leukemia Occurs in Two Tracks and Employs Embryonic Stem Cell-like Signatures'. *Nucleic Acids Research* 43 (5): 2590–2602. https://doi.org/10.1093/NAR/GKV103.

Lee, Yunsung, Sanaa Choufani, Rosanna Weksberg, Samantha L. Wilson, Victor Yuan, Amber Burt, Carmen Marsit, et al. 2019. 'Placental Epigenetic Clocks: Estimating Gestational Age Using Placental DNA Methylation Levels'. *Aging* 11 (12): 4238–53. https://doi.org/10.18632/AGING.102049.

Lee-Six, Henry, Nina Friesgaard Øbro, Mairi S Shepherd, Sebastian Grossmann, Kevin Dawson, Miriam Belmonte, Robert J Osborne, et al. 2018. 'Population Dynamics of Normal Human Blood Inferred from Somatic Mutations'. *Nature* 561 (7724): 473–78. https://doi.org/10.1038/s41586-018-0497-0.

Leval, Laurence de, Ash A. Alizadeh, P. Leif Bergsagel, Elias Campo, Andrew Davies, Ahmet Dogan, Jude Fitzgibbon, et al. 2022. 'Genomic Profiling for Clinical Decision Making in Lymphoid Neoplasms'. *Blood* 140 (21): 2193–2227. https://doi.org/10.1182/BLOOD.2022015854.

Levine, Morgan E., Ake T. Lu, Austin Quach, Brian H. Chen, Themistocles L. Assimes, Stefania Bandinelli, Lifang Hou, et al. 2018. 'An Epigenetic Biomarker of Aging for Lifespan and Healthspan'. *Aging* 10 (4): 573–91. https://doi.org/10.18632/AGING.101414.

Liang, Xiaoyu, Amy C. Justice, Kaku So-Armah, John H. Krystal, Rajita Sinha, and Ke Xu. 2020. 'DNA Methylation Signature on Phosphatidylethanol, Not on Self-Reported Alcohol Consumption, Predicts Hazardous Alcohol Consumption in Two Distinct Populations'. *Molecular Psychiatry 2020 26:6* 26 (6): 2238–53. https://doi.org/10.1038/s41380-020-0668-x.

Lin, Qiong, Carola I. Weidner, Ivan G. Costa, Riccardo E. Marioni, Marcelo R.P. Ferreira, Ian J. Deary, and Wolfgang Wagner. 2016. 'DNA Methylation Levels at Individual Age-

Associated CpG Sites Can Be Indicative for Life Expectancy'. *Aging* 8 (2): 394–401.
https://doi.org/10.18632/AGING.100908.
Ling, Shaoping, Zheng Hu, Zuyu Yang, Fang Yang, Yawei Li, Pei Lin, Ke Chen, et al. 2015.
'Extremely High Genetic Diversity in a Single Tumor Points to Prevalence of Non-
Darwinian Cell Evolution'. *Proceedings of the National Academy of Sciences of the*
United States of America 112 (47): E6496–6505.
https://doi.org/10.1073/PNAS.1519556112/SUPPL_FILE/PNAS.1519556112.SD01.XLSX.
Loyfer, Netanel, Judith Magenheim, Ayelet Peretz, Gordon Cann, Joerg Bredno, Agnes
Klochendler, Ilana Fox-Fisher, et al. 2023. 'A DNA Methylation Atlas of Normal Human
Cell Types'. *Nature 2023 613:7943* 613 (7943): 355–64.
https://doi.org/10.1038/s41586-022-05580-6.
Lu, Ake T., Austin Quach, James G. Wilson, Alex P. Reiner, Abraham Aviv, Kenneth Raj, Lifang
Hou, et al. 2019. 'DNA Methylation GrimAge Strongly Predicts Lifespan and
Healthspan'. *Aging* 11 (2): 303–27. https://doi.org/10.18632/AGING.101684.
Martinez, Pierre, Diego Mallo, Thomas G. Paulson, Xiaohong Li, Carissa A. Sanchez, Brian J.
Reid, Trevor A. Graham, Mary K. Kuhner, and Carlo C. Maley. 2018. 'Evolution of
Barrett's Esophagus through Space and Time at Single-Crypt and Whole-Biopsy Levels'.
Nature Communications 2018 9:1 9 (1): 1–12. https://doi.org/10.1038/s41467-017-
02621-x.
Mayne, Benjamin T., Shalem Y. Leemaqz, Alicia K. Smith, James Breen, Claire T. Roberts, and
Tina Bianco-Miotto. 2017. 'Accelerated Placental Aging in Early Onset Preeclampsia
Pregnancies Identified by DNA Methylation'. *Epigenomics* 9 (3): 279–89.
https://doi.org/10.2217/EPI-2016-0103/SUPPL_FILE/SUPPLEMENTARY_FIGURE_1.PDF.
McCartney, Daniel L., Robert F. Hillary, Anna J. Stevenson, Stuart J. Ritchie, Rosie M. Walker,
Qian Zhang, Stewart W. Morris, et al. 2018. 'Epigenetic Prediction of Complex Traits
and Death'. *Genome Biology* 19 (1): 136. https://doi.org/10.1186/S13059-018-1514-
1/TABLES/3.
McEwen, Lisa M., Kieran J. O'Donnell, Megan G. McGill, Rachel D. Edgar, Meaghan J. Jones,
Julia L. MacIsaac, David Tse Shen Lin, et al. 2020. 'The PedBE Clock Accurately Estimates
DNA Methylation Age in Pediatric Buccal Cells'. *Proceedings of the National Academy of*
Sciences of the United States of America 117 (38): 23329–35.
https://doi.org/10.1073/PNAS.1820843116/SUPPL_FILE/PNAS.1820843116.SAPP.PDF.
Merlo, Lauren M.F., John W. Pepper, Brian J. Reid, and Carlo C. Maley. 2006. 'Cancer as an
Evolutionary and Ecological Process'. *Nature Reviews Cancer*. Nature Publishing Group.
https://doi.org/10.1038/nrc2013.
Nadeu, Ferran, David Martin-Garcia, Guillem Clot, Ander Díaz-Navarro, Martí Duran-Ferrer,
Alba Navarro, Roser Vilarrasa-Blasi, et al. 2020. 'Genomic and Epigenomic Insights into
the Origin, Pathogenesis, and Clinical Behavior of Mantle Cell Lymphoma Subtypes'.
Blood 136 (12): 1419–32. https://doi.org/10.1182/BLOOD.2020005289.
Nadeu, Ferran, Romina Royo, Ramon Massoni-Badosa, Heribert Playa-Albinyana, Beatriz
Garcia-Torre, Martí Duran-Ferrer, Kevin J. Dawson, et al. 2022. 'Detection of Early
Seeding of Richter Transformation in Chronic Lymphocytic Leukemia'. *Nature Medicine*
2022 28:8 28 (8): 1662–71. https://doi.org/10.1038/s41591-022-01927-8.
Nik-Zainal, Serena, Ludmil B. Alexandrov, David C. Wedge, Peter Van Loo, Christopher D.
Greenman, Keiran Raine, David Jones, et al. 2012. 'Mutational Processes Molding the
Genomes of 21 Breast Cancers'. *Cell* 149 (5): 979–93.
https://doi.org/10.1016/j.cell.2012.04.024.

Nordlund, Jessica, Christofer L. Bäcklin, Per Wahlberg, Stephan Busche, Eva C. Berglund, Maija Leena Eloranta, Trond Flaegstad, et al. 2013. 'Genome-Wide Signatures of Differential DNA Methylation in Pediatric Acute Lymphoblastic Leukemia'. *Genome Biology* 14 (9): 1–15. https://doi.org/10.1186/GB-2013-14-9-R105/TABLES/3.

Nowell, Peter C. 1976. 'The Clonal Evolution of Tumor Cell Populations'. *Science* 194 (4260): 23–28. https://doi.org/10.1126/science.959840.

Oakes, Christopher C., and Jose I. Martin-Subero. 2018. 'Insight into Origins, Mechanisms, and Utility of DNA Methylation in B-Cell Malignancies'. *Blood* 132 (10): 999–1006. https://doi.org/10.1182/BLOOD-2018-02-692970.

Oakes, Christopher C., Marc Seifert, Yassen Assenov, Lei Gu, Martina Przekopowitz, Amy S. Ruppert, Qi Wang, et al. 2016. 'DNA Methylation Dynamics during B Cell Maturation Underlie a Continuum of Disease Phenotypes in Chronic Lymphocytic Leukemia'. *Nature Genetics* 48 (3): 253–64. https://doi.org/10.1038/NG.3488.

Plevova, Karla, Hana Skuhrova Francova, Katerina Burckova, Yvona Brychtova, Michael Doubek, Sarka Pavlova, Jitka Malcikova, Jiri Mayer, Boris Tichy, and Sarka Pospisilova. 2014. 'Multiple Productive Immunoglobulin Heavy Chain Gene Rearrangements in Chronic Lymphocytic Leukemia Are Mostly Derived from Independent Clones'. *Haematologica* 99 (2): 329–38. https://doi.org/10.3324/HAEMATOL.2013.087593.

Puente, Xose S., Silvia Beà, Rafael Valdés-Mas, Neus Villamor, Jesús Gutiérrez-Abril, José I. Martín-Subero, Marta Munar, et al. 2015. 'Non-Coding Recurrent Mutations in Chronic Lymphocytic Leukaemia'. *Nature 2015 526:7574* 526 (7574): 519–24. https://doi.org/10.1038/nature14666.

Queirós, Ana C., Renée Beekman, Roser Vilarrasa-Blasi, Martí Duran-Ferrer, Guillem Clot, Angelika Merkel, Emanuele Raineri, et al. 2016. 'Decoding the DNA Methylome of Mantle Cell Lymphoma in the Light of the Entire B Cell Lineage'. *Cancer Cell* 30 (5): 806–21. https://doi.org/10.1016/J.CCELL.2016.09.014.

Rambaut, Andrew, Alexei J. Drummond, Dong Xie, Guy Baele, and Marc A. Suchard. 2018. 'Posterior Summarization in Bayesian Phylogenetics Using Tracer 1.7'. *Systematic Biology* 67 (5): 901–4. https://doi.org/10.1093/SYSBIO/SYY032.

Raudvere, Uku, Liis Kolberg, Ivan Kuzmin, Tambet Arak, Priit Adler, Hedi Peterson, and Jaak Vilo. 2019. 'G:Profiler: A Web Server for Functional Enrichment Analysis and Conversions of Gene Lists (2019 Update)'. *Nucleic Acids Research* 47 (W1): W191–98. https://doi.org/10.1093/NAR/GKZ369.

Reimand, Jüri, Meelis Kull, Hedi Peterson, Jaanus Hansen, and Jaak Vilo. 2007. 'G:Profiler--a Web-Based Toolset for Functional Profiling of Gene Lists from Large-Scale Experiments'. *Nucleic Acids Research* 35 (Web Server issue). https://doi.org/10.1093/NAR/GKM226.

Reinius, Lovisa E., Nathalie Acevedo, Maaike Joerink, Göran Pershagen, Sven Erik Dahlén, Dario Greco, Cilla Söderhäll, Annika Scheynius, and Juha Kere. 2012. 'Differential DNA Methylation in Purified Human Blood Cells: Implications for Cell Lineage and Studies on Disease Susceptibility'. *PLOS ONE* 7 (7): e41361. https://doi.org/10.1371/JOURNAL.PONE.0041361.

Rice, Siobhan, and Anindita Roy. 2020. 'MLL-Rearranged Infant Leukaemia: A "thorn in the Side" of a Remarkable Success Story'. *Biochimica et Biophysica Acta. Gene Regulatory Mechanisms* 1863 (8). https://doi.org/10.1016/J.BBAGRM.2020.194564.

Shireby, Gemma L., Jonathan P. Davies, Paul T. Francis, Joe Burrage, Emma M. Walker, Grant W.A. Neilson, Aisha Dahir, et al. 2020. 'Recalibrating the Epigenetic Clock: Implications

for Assessing Biological Age in the Human Cortex'. *Brain* 143 (12): 3763–75. https://doi.org/10.1093/BRAIN/AWAA334.

Siegmund, Kimberly D., Paul Marjoram, Yen Jung Woo, Simon Tavaré, and Darryl Shibata. 2009. 'Inferring Clonal Expansion and Cancer Stem Cell Dynamics from DNA Methylation Patterns in Colorectal Cancers'. *Proceedings of the National Academy of Sciences of the United States of America* 106 (12): 4828. https://doi.org/10.1073/PNAS.0810276106.

Skilling, John. 2004. 'Nested Sampling'. In *AIP Conference Proceedings*, 735:395–405. AIP Publishing. https://doi.org/10.1063/1.1835238.

———. 2006. 'Nested Sampling for General Bayesian Computation'. *Bayesian Analysis* 1 (4): 833–60. https://doi.org/10.1214/06-BA127.

Sottoriva, Andrea, Haeyoun Kang, Zhicheng Ma, Trevor A. Graham, Matthew P. Salomon, Junsong Zhao, Paul Marjoram, et al. 2015. 'A Big Bang Model of Human Colorectal Tumor Growth'. *Nature Genetics 2015 47:3* 47 (3): 209–16. https://doi.org/10.1038/ng.3214.

Speagle, Joshua S. 2019. 'Dynesty: A Dynamic Nested Sampling Package for Estimating Bayesian Posteriors and Evidences'. *Monthly Notices of the Royal Astronomical Society* 493 (3): 3132–58. https://doi.org/10.1093/mnras/staa278.

Suchard, Marc A, Philippe Lemey, Guy Baele, Daniel L Ayres, Alexei J Drummond, and Andrew Rambaut. 2018. 'Bayesian Phylogenetic and Phylodynamic Data Integration Using BEAST 1.10'. *Virus Evolution* 4 (1). https://doi.org/10.1093/VE/VEY016.

Tate, John G., Sally Bamford, Harry C. Jubb, Zbyslaw Sondka, David M. Beare, Nidhi Bindal, Harry Boutselakis, et al. 2019. 'COSMIC: The Catalogue Of Somatic Mutations In Cancer'. *Nucleic Acids Research* 47 (D1): D941–47. https://doi.org/10.1093/NAR/GKY1015.

Teschendorff, Andrew E. 2020. 'A Comparison of Epigenetic Mitotic-like Clocks for Cancer Risk Prediction'. *Genome Medicine* 12 (1): 1–17. https://doi.org/10.1186/S13073-020-00752-3/FIGURES/4.

Turajlic, Samra, Andrea Sottoriva, Trevor Graham, and Charles Swanton. 2019. 'Resolving Genetic Heterogeneity in Cancer'. *Nature Reviews Genetics 2019 20:7* 20 (7): 404–16. https://doi.org/10.1038/s41576-019-0114-6.

Uhlén, Mathias, Linn Fagerberg, Bjö M. Hallström, Cecilia Lindskog, Per Oksvold, Adil Mardinoglu, Åsa Sivertsson, et al. 2015. 'Tissue-Based Map of the Human Proteome'. *Science* 347 (6220). https://doi.org/10.1126/SCIENCE.1260419/SUPPL_FILE/1260419_UHLEN.SM.PDF.

Vehtari, Aki, Andrew Gelman, and Jonah Gabry. 2015. 'Practical Bayesian Model Evaluation Using Leave-One-out Cross-Validation and WAIC'. *Statistics and Computing* 27 (5): 1413–32. https://doi.org/10.1007/s11222-016-9696-4.

Vidal-Bralo, Laura, Yolanda Lopez-Golan, and Antonio Gonzalez. 2016. 'Simplified Assay for Epigenetic Age Estimation in Whole Blood of Adults'. *Frontiers in Genetics* 7 (JUL): 209192. https://doi.org/10.3389/FGENE.2016.00126/BIBTEX.

Weidner, Carola I., Qiong Lin, Carmen M. Koch, Lewin Eisele, Fabian Beier, Patrick Ziegler, Dirk O. Bauerschlag, et al. 2014. 'Aging of Blood Can Be Tracked by DNA Methylation Changes at Just Three CpG Sites'. *Genome Biology* 15 (2): 1–12. https://doi.org/10.1186/GB-2014-15-2-R24/COMMENTS.

Williams, Marc J., Benjamin Werner, Chris P. Barnes, Trevor A. Graham, and Andrea Sottoriva. 2016. 'Identification of Neutral Tumor Evolution across Cancer Types'. *Nature Genetics* 48 (3): 238–44. https://doi.org/10.1038/ng.3489.

Williams, Marc J., Benjamin Werner, Timon Heide, Christina Curtis, Chris P. Barnes, Andrea Sottoriva, and Trevor A. Graham. 2018. 'Quantification of Subclonal Selection in Cancer from Bulk Sequencing Data'. *Nature Genetics* 50 (6): 895–903. https://doi.org/10.1038/s41588-018-0128-6.

Williams, Nicholas, Joe Lee, Emily Mitchell, Luiza Moore, E. Joanna Baxter, James Hewinson, Kevin J. Dawson, et al. 2022. 'Life Histories of Myeloproliferative Neoplasms Inferred from Phylogenies'. *Nature 2022 602:7895* 602 (7895): 162–68. https://doi.org/10.1038/s41586-021-04312-6.

Yang, Zhen, Andrew Wong, Diana Kuh, Dirk S. Paul, Vardhman K. Rakyan, R. David Leslie, Shijie C. Zheng, Martin Widschwendter, Stephan Beck, and Andrew E. Teschendorff. 2016. 'Correlation of an Epigenetic Mitotic Clock with Cancer Risk'. *Genome Biology* 17 (1): 1–18. https://doi.org/10.1186/S13059-016-1064-3/FIGURES/6.

Yao, Yuling, Aki Vehtari, Daniel Simpson, and Andrew Gelman. 2017. 'Using Stacking to Average Bayesian Predictive Distributions'. *Bayesian Analysis* 13 (3). https://doi.org/10.1214/17-BA1091.

Yatabe, Yasushi, Simon Tavaré, and Darryl Shibata. 2001. 'Investigating Stem Cells in Human Colon by Using Methylation Patterns'. *Proceedings of the National Academy of Sciences* 98 (19): 10839–44. https://doi.org/10.1073/pnas.191225998.

Youn, Ahrim, and Shuang Wang. 2018. 'The MiAge Calculator: A DNA Methylation-Based Mitotic Age Calculator of Human Tissue Types'. *Epigenetics* 13 (2): 192–206. https://doi.org/10.1080/15592294.2017.1389361.

Yu, Guangchuang, David K. Smith, Huachen Zhu, Yi Guan, and Tommy Tsan Yuk Lam. 2017. 'Ggtree: An r Package for Visualization and Annotation of Phylogenetic Trees with Their Covariates and Other Associated Data'. *Methods in Ecology and Evolution* 8 (1): 28–36. https://doi.org/10.1111/2041-210X.12628.

Zhang, Qian, Costanza L. Vallerga, Rosie M. Walker, Tian Lin, Anjali K. Henders, Grant W. Montgomery, Ji He, et al. 2019. 'Improved Precision of Epigenetic Clock Estimates across Tissues and Its Implication for Biological Ageing'. *Genome Medicine* 11 (1): 1–11. https://doi.org/10.1186/S13073-019-0667-1/FIGURES/4.

Zhou, Wanding, Huy Q. Dinh, Zachary Ramjan, Daniel J. Weisenberger, Charles M. Nicolet, Hui Shen, Peter W. Laird, and Benjamin P. Berman. 2018. 'DNA Methylation Loss in Late-Replicating Domains Is Linked to Mitotic Cell Division'. *Nature Genetics 2018 50:4* 50 (4): 591–602. https://doi.org/10.1038/s41588-018-0073-4.