TITLE

Improving Diagnostic Sensitivity for Imbalanced Musculoskeletal Disorder Data: A Sensitivity-Based

Multi-Sampling Technique for Osteoarthritis Prediction

Name of Author (Academic degree)

Jun-hee Kim (PhD, PT)

Affiliation

1, Yeonsedae-gil, Maeji-ri, Heungeop-myeon, Wonju-si, Gangwon-do, 26493, Department of Physical

Therapy, College of Software and Digital Healthcare Convergence, Yonsei University, Wonju, South

Korea.

Corresponding Author

Jun-hee Kim, PhD, PT

move@yonsei.ac.kr / +82-10-9636-3638

ABSTRACT

Background: Medical datasets containing musculoskeletal disorders may have data imbalances due to the incidence of the disease, which may limit the predictive ability, such as the sensitivity, of musculoskeletal diagnostic prediction models built from these data. This study aimed to increase the sensitivity performance of osteoarthritis (OA) prediction when building a model by adjusting an OA imbalanced dataset using a sensitivity-based multi-sampling (SMS) technique.

Methods: OA Data were obtained from the Korea National Health and Nutrition Examination Survey (KNHANES). SMS technique combining oversampling and undersampling was applied to the imbalanced OA data, and the RandomForest algorithm was used for machine learning modeling. Model performance was evaluated based on accuracy, sensitivity, and specificity and compared with other hybrid sampling techniques.

Result: In the SMS technique, ADASYN, Borderline-SMOTE, SMOTE oversampling and ENN undersampling techniques were combined and applied. The OA prediction model using the SMS technique showed the highest sensitivity (82.20) but the lowest specificity (82.26) and accuracy (82.26) compared to other hybrid models.

Conclusion: SMS technique offers a potential solution for improving sensitivity performance for prediction models built on medical data imbalances due to low-incidence diseases. Nonetheless, caution is warranted due to the concern that while improving sensitivity, it may decrease specificity with a trade-off.

Key words: Machine learning, sample size, algorithm, sampling, musculoskeletal disorder, osteoarthritis

INTRODUCTION

Musculoskeletal diagnosis refers to the process of determining and diagnosing diseases, damage and abnormalities related to the human musculoskeletal system, which consists of structures such as muscles, bones, joints, and ligaments [1,2]. The musculoskeletal diagnosis provides medical professionals with essential information to understand the patient's condition and to develop an appropriate treatment and care plan [1,3]. In addition, accurate musculoskeletal diagnosis allows medical personnel and related professionals to identify the extent of a patient's illness or damage [1–3]. This allows experts to take proper care and take precautions before the disease becomes serious [3]. This accurate diagnosis can help improve the quality of life of patients who can experience pain, discomfort, and functional limitations, and speed up the recovery of daily activity [1]. Therefore, accurate diagnosis can help improve the quality of life of patients, who can experience pain, discomfort, and functional limitations, and speed up the recovery of daily activity [1,3].

The onset of musculoskeletal disorder is affected by several factors. These conditions have been reported to be primarily related to age, gender, genetic factors, lifestyle, and environmental factors [4]. Typically, lack of exercise, excessive weight, smoking, alcohol consumption, insufficient nutrient intake, excessive stress, and poor working environment can act as factors that can affect the onset of musculoskeletal disorder. Depending on these factors, the incidence of musculoskeletal disorders varies [4,5]. In 2017, the highest incidence of musculoskeletal disorders worldwide was 36.8% due to back pain, followed by osteoarthritis (OA) with 19.3%, neck pain with 18.4%, gout with 2.6%, and rheumatoid arthritis with 1.3% [5].

However, medical data, including musculoskeletal disorder data, have varying incidence rates, resulting in a data imbalance problem [6]. This data imbalance occurs when the occurrence of a particular disease is rare or uncommon [7]. Since the incidence of musculoskeletal disorders is generally lower than in healthy people, there may be fewer data points for disease classes, which may cause machine learning models to distort to primarily train normal classes, limiting their predictive ability for minority disease classes [7,8]. For this reason, data imbalances primarily seen in medical data negatively affect the performance of machine learning models [7,8]. It has been a particularly important issue in the healthcare and medical fields, and when predicting whether to diagnose a disorder with a small incidence, the sensitivity of the model may be limited due to data, which may result in missing actual patients [7,8].

Various sampling techniques are used to address data imbalance. One sampling technique is oversampling, which creates a more balanced set of data by replicating or synthesizing instances of a minority class [9,10].

Typically, oversampling technique such as SMOTE (synthetic prime oversampling technique) and ADASYN

(Adaptive Synthetic Sampling) exist [9,10]. Undersampling, on the other hand, involves reducing the number of

instances in multiple classes, such as Tomek link or Neighbourhood-based undersampling [11,12]. Also,

Combinations of oversampling and undersampling techniques can also be used to maximize the benefits of both

approaches. SMOTE-Tomek, which combines SMOTE and Tomek, or SMOTE-ENN, which combines SMOTE

and Edited Nearest Neighbors, can be seen as examples of hybrid sampling [13,14]. These techniques are

effective at mitigating data imbalances but can lead to overfitting or potentially losing valuable information

when training machine learning models [9–11,13,15].

The purpose of this study was to develop and apply sampling techniques to improve the low sensitivity

performance of models built with medical datasets such as musculoskeletal disorder that indicate data

imbalances according to low incidence rates. Therefore, in this study, a sensitivity-based multiple sampling

(SMS) technique was developed, and the SMS technique was applied to construct a dataset with an emphasis on

improving sensitivity to predict OA, a musculoskeletal disorder with a low incidence. And this technique was

applied to machine learning model training and compared to the performance of machine learning models of

other sampling techniques.

MATERIALS AND METHODS

Data Sources

The flowchart of this study is shown in Figure 1. The study samples were obtained from Korea National Health and Nutrition Examination Survey (KNHANES), a national periodic survey conducted by the Korea Disease Control and Prevention Agency. This study used 8110 datasets, the first-year data of the 8th KNHANES (2019–2020). Among the datasets, data that meet the following criteria were selected. 1) Participants over the age of 19 2) Data with diagnosis as OA 3) Data with no missing values in variables to be used as predictors. Of the 8110 samples without data missing values, 621 samples with only osteoarthritis without rheumatoid arthritis and 4988 data samples without both OA and rheumatoid arthritis were finally selected for this study. Variables representing individual characteristics, physical activity, lifestyle, and descriptions used in this study are shown in Table 1.

Figure 1. Flowchart of this study

Table 1. List of variables and variable descriptions (mean $\pm$ SD)

	OA (621)	Normal (4988)	Description
Sex	M:116 / F:505	M:2403 / F:2585	Gender (M: male / F: female)
Age	67.84 $\pm$ 9.95	49.04 $\pm$ 16.21	Age (year)
Height	156.72 $\pm$ 8.27	164.74 $\pm$ 9.06	Height (cm)
Weight	61.35 $\pm$ 10.70	64.95 $\pm$ 12.97	Weight (kg)
BMI	24.92 $\pm$ 3.46	23.82 $\pm$ 3.61	Weight/Height^2
BO1_1	1:426 / 2:90 / 3:105	1:3122 / 2:671 / 3:1195	Weight change (1: none / 2: increase / 3: decrease)
HE_wc	87.69 $\pm$ 9.46	83.51 $\pm$ 10.48	Waist circumference (cm)
GS	21.97 $\pm$ 7.68	29.53 $\pm$ 9.91	Grip strength (kg)
LQ4_00	1:131 / 2:490	1:278 / 2:4710	Activity limitation (1: yes / 2: no)
EQ5D	0.86 $\pm$ 0.15	0.96 $\pm$ 0.09	Quality of life (EQ5D index)
EC1_1	1:245 / 2:376	1:3199 / 2:1789	Work (1: yes / 2: no)
EC_wht_23	14.91 $\pm$ 21.17	27.91 $\pm$ 21.92	Work time
BE3_71	1:2 / 2:619	1:82 / 2:4906	Work activity (1: yes / 2: no)
BE3_75	1:11 / 2:610	1:558 / 2:4430	Leisure activity (1: yes / 2: no)
BD3_2	1.97 $\pm$ 1.78	3.00 $\pm$ 1.74	Smoking (per day)
BD1_11	0.72 $\pm$ 3.44	2.42 $\pm$ 5.89	Drinking alcohol (per week)
BP16_1, BP16_2	6.39 $\pm$ 1.56	6.88 $\pm$ 1.28	Sleep time (hour)

Data Pre-processing

Inter Quarter Range (IQR) and Boxplot were utilized to detect and eliminate outliers. Outliers in numerical data were identified and processed based on IQR, and scaling was performed using the StandardScaler function for numerical variables. Categorical variables were converted into dummy data using the LabelEncoder function. This pre-processed data was divided into training and test data at an 8:2 ratio and utilized for model learning and performance evaluation.

Sensitivity-based Multi-sampling and Hybrid Sampling

SMS, which was designed in this study, is a technique that is a combination of oversampling and undersampling techniques (Figure 2). According to the results of the test performance evaluation of each model learned with data sampled by five oversampling techniques, the top three oversampling techniques were selected based on sensitivity score. In addition, the undersampling technique with the highest sensitivity score was selected according to the test performance evaluation results of each model learned with data sampled using the four undersampling techniques. In the original data, the data was augmented by dividing half the sample size difference between classes into three selected oversampling techniques. And finally, a selected undersampling technique reduced the data by half of the sample size difference between classes. Hybrid sampling techniques, SMOTE-Tomek and SMOTE-ENN, were employed for comparison with the SMS technique in this study. Hybrid sampling involves a combination of oversampling and undersampling techniques to harness the benefits of both approaches.

Figure 2. Sensitivity-based Multi-sampling algorithm

Machine-learning modeling

In this study, the RandomForest machine learning algorithm was used to compare the differences in OA prediction performance between sampling techniques. Hyperparameters were not fine-tuned and default values were used, and the default values were as follows: n_estimators (decision-making number of trees): 100, criterion (splitting criterion): "gini", max_depth (maximum tree depth): no limit, min_samples_split (minimum number of samples for node splitting): 2, min_samples_leaf (minimum number of samples required from a terminal node) : 1, max_features (maximum number of features to use for splitting at each node): auto. The default values of hyperparameters were used to create a RandomForest machine-learning model for OA diagnosis prediction.

Model evaluation

In applying the SMS technique, the performance of machine learning models built by each oversampling and

undersampling technique evaluated accuracy, sensitivity, and specificity. In addition, the performance of

machine learning models built with respective training datasets sampled by SMS, SMOTE-Tomek, and

SMOTE-ENN was also evaluated and compared for accuracy, sensitivity, and specificity.

RESULTS

Oversampling and Undersampling model performance

The accuracy of RandomForrest models built with training data applied with oversampling techniques was

found to be 89.61 high from 87.90 (Table 2). In terms of sensitivity, ADASYN, Borderline-SMOTE, and

SMOTE were found to have high performance in order.

Table 2. Oversampling techniques application and model evaluation scores

	Accuracy	**Sensitivity**	**Specificity**
SMOTE	88.71	59.32	92.18
ADASYN	87.90	60.17	91.18
Borderline-SMOTE	88.62	60.17	91.98
SVMSMOTE	89.61	57.63	93.39
Synthetic Over-sampling	88.08	57.63	91.68

152   The accuracy of RandomForrest models built with training data applied with this downsampling technique

153 was found to be 89.52 high at 86.74 (Table 3). In terms of sensitivity, the model built by the sampling technique

154 of Repeated Edited Nearest Neighbours showed the highest performance at 69.49.

Table 3. Undersampling techniques application and model evaluation scores

	Accuracy	**Sensitivity**	**Specificity**
Tomek Links	89.52	33.90	96.09
Edited Nearest Neighbours	88.98	59.32	92.48
Repeated Edited Nearest Neighbours	86.74	69.49	88.78
Neighbourhood Cleaning Rule	89.25	60.17	92.69

Sensitivity-based Multi-sampling

The three oversampling techniques chosen by SMS were ADASYN, Borderline-SMOTE, and SMOTE, and

the undersampling technique was Repeated Edited Nearest Neighbours (Figure 3). A total of 1758 OA data were

added to the original train data, 586 each by three oversampling techniques according to the SMS algorithm.

Then, 1247 normal data were removed by Repeated Edited Nearest Neighbours, selected by the under-sampling

technique, leaving only 2743 normal data.

163
Figure 3. Train set structure after Sensitivity-based Multi-sampling technique

Sensitivity-based Multi-sampling and Hybrid Sampling Model Performance

Table 4 shows the performance of the RandomForrest model for OA prediction built with train datasets

sampled by the Hybrid sampling techniques. The accuracy performance of the model built using the SMS

technique was 82.26, the lowest among the three models, but the sensitivity performance was the highest at

82.20. On the other hand, the accuracy performance of the model built using the SMOTE-Tomek technique was

found to be the highest at 88.35, but the lowest sensitivity performance of 56.78. SMOTE-ENN showed the

second highest performance with accuracy performance of 84.05 and sensitivity of 77.12, respectively. In terms

of specificity, SMOTE-TOMEK was the highest at 92.08, SMOTE-ENN was second at 84.87, and SMS was the

lowest at 82.26.

Table 4. Hybrid sampling techniques application and model evaluation score

	Accuracy	**Sensitivity**	**Specificity**
Sensitivity-based Multi-sampling	82.26	82.20	82.26
SMOTE-Tomek	88.35	56.78	92.08
SMOTE-ENN	84.05	77.12	84.87

DISCUSSION

In this study, a SMS technique was introduced and applied to address the data imbalance problem in musculoskeletal disorder prediction, especially focusing on OA. An OA prediction model using the SMS technique was constructed, and sensitivity performance was compared with other prediction models. The model was constructed by applying a combination of oversampling (ADASYN, Borderline-SMOTE, SMOTE) and undersampling (iteratively edited nearest neighbors) using a SMS technique. The accuracy and specificity of the built model decreased compared to other models, but in terms of sensitivity, it showed higher performance than other models.

In medical data, where data imbalance inevitably occurs due to the rate of disease occurrence, model performance often deteriorates due to data imbalance [16,17]. If a prediction model is created with imbalanced medical data, sensitivity performance may decrease [16,17]. Shim et al. (2020) built an osteoporosis risk prediction model using osteoporosis patients and normal subjects with a sample size that was approximately three times that of the patients [17]. The sample ratio differed by only about 3 times, but when the model was built using the RandomForest algorithm, the sensitivity was only 59 [17]. In the case of medical data, various sampling techniques have been proposed to improve the poor performance caused by data imbalance [7,18]. Sampling techniques to resolve data imbalance are largely divided into undersampling and oversampling [15,19]. Oversampling involves creating additional instances of minority classes to balance the class distribution [15,19]. This helps the model learn minority classes better, but generating too many synthetic examples can lead to overfitting. This means that the model may become too tailored to the training data and perform poorly on new data [15,19]. Undersampling involves reducing the number of instances of a majority class, which can lead to information loss issues [15,19]. This results in the loss of valuable information present in the majority class, potentially resulting in a less representative model [15,19]. If many classes contain important patterns or variations, removing instances may reduce the model's ability to generalize to unseen data [15,19].

To address the problems associated with oversampling and undersampling applied to building models containing such imbalanced medical data, various hybrid sampling techniques have been proposed and used. Zeng et al. (2016) proposed a preprocessing technique combining SMOTE with Tomek link and applied it to imbalanced medical data sets of three diseases [20]. The combined technique of SMOTE and Tomek Link showed marked improvement in 31, 27, and 30 out of 32 evaluation indicators for diabetes, Parkinson's disease, and spinal disease compared to the oversampling using SMOTE alone [20]. Lamari et al. (2021) used an

ensemble technique that combines SMOTE, which synthesizes samples belonging to minority classes, and

Edited Nearest Neighbors (ENN), which examines the neighbors of a given sample and removes unnecessary

samples [14]. Using ensemble techniques, they improved the sensitivity performance of appendicitis, diabetes,

and Parkinson's disease classification models from an average of 78 to 82 [14]. Khushi et al. (2021) built a

model by applying both hybrid sampling techniques (SMOTE-ENN, SMOTE-Tomek) to the lung cancer dataset,

and the AUC performance of this model increased from 50 to 65.50 [16]. Our study also showed performance

improvement in terms of sensitivity. With only oversampling or undersampling, sensitivity performance ranged

from 33.90 to 69.49. In the case of hybrid techniques combining oversampling and undersampling, sensitivity

performance ranged from 56.78 to 77.12, except for SMS. In comparison, SMS showed a high sensitivity

performance of 82.20.

Sensitivity is a measure of how accurately true positives can be detected among positive cases and is very

important in diagnosing a disease [21–23]. In other words, the higher the sensitivity, the fewer false negatives,

allowing more effective identification of actual positive cases [21]. Especially in disease diagnosis, missing a

positive test can be a big problem, and if a positive test is not accurately detected, the patient may not receive

appropriate treatment or measures, which can lead to serious problems due to delayed diagnosis or inaccurate

results [22]. Considering the aspect of sensitivity in medical diagnosis, this study designed a sampling technique

to improve low sensitivity performance due to data imbalance. The reason why SMS had the highest sensitivity

in this study may be because oversampling and undersampling techniques were selected in order of high

sensitivity. However, there is a trade-off between sensitivity and specificity [24,25]. When sensitivity is high, the

model is good at finding positive samples without missing them, but this makes it easier for the model to predict

positive ones and may mispredict negative ones as positive [24]. This may result in an increase in the false

positive rate and a decrease in specificity [24]. In other words, sensitivity and specificity are complementary

indicators, and improving one may decrease the other, and in medical diagnosis, sensitivity may be more

important than specificity [26–28]. For this reason, sensitivity-based multi-sampling technique to improve

sensitivity would have reduced specificity and further reduced accuracy.

This study has some limitations. First, the effectiveness of the SMS technique may vary depending on the

characteristics of the disease dataset used. Different disease datasets may have different patterns of imbalance

depending on the incidence of the disease, so sampling rates may need to be adjusted. Second, the

hyperparameters for the RandomForest machine learning algorithm were not tuned and default values were used.

Since hyperparameter tuning can improve model performance, it will be necessary to investigate the impact of

hyperparameter tuning on sensitivity-based multi-sampling technique. Third, since the variables used to

construct this OA prediction model were not medical measurement variables used to diagnose OA, it is unclear

how reliably they can be applied in actual medical settings. Therefore, although this model can assist in

diagnosing OA, it cannot replace medical diagnostic criteria.

CONCLUSION

This study attempted to solve the decrease in sensitivity performance caused by data imbalance in predicting

musculoskeletal disorder, focusing on OA, by introducing and applying the SMS technique. In this study, the

RandomForest machine learning algorithm was used, and the performance of the SMS technique was compared

with other techniques combining oversampling and undersampling. According to the study results, the SMS

technique, which was designed to improve the sensitivity of predicting OA, had the highest sensitivity, but

showed lower specificity, which is a trade-off, compared to other models. In conclusion, SMS technique

provides valuable insight into the challenge of addressing data imbalances in musculoskeletal disorder

prediction. SMS technique has shown potential to improve sensitivity, an important indicator of disease

diagnosis, but their use should be approached cautiously, considering the trade-offs and limitations observed in

specificity and overall model performance.

REFERENCES

1. Shultz SJ, Houglum PA, Perrin DH. Examination of musculoskeletal injuries. Human Kinetics; 2015.

2. Yelin E, Weinstein S, King T. The burden of musculoskeletal diseases in the United States. In: Seminars in arthritis and rheumatism. 2016. p. 259–60.

3. Braddom RL. Physical medicine and rehabilitation e-book. Elsevier Health Sciences; 2010.

4. Woolf AD, Pfleger B. Burden of major musculoskeletal conditions. Bull World Health Organ. 2003;81(9):646–56.

5. Safiri S, Kolahi A, Cross M, Hill C, Smith E, Carson☐Chahhoud K, et al. Prevalence, deaths, and disability☐adjusted life years due to musculoskeletal disorders for 195 countries and territories 1990–2017. Arthritis & rheumatology. 2021;73(4):702–14.

6. Ali H, Salleh MNM, Saedudin R, Hussain K, Mushtaq MF. Imbalance class problems in data mining: A review. Indonesian Journal of Electrical Engineering and Computer Science. 2019;14(3):1560–71.

7. Khalilia M, Chakraborty S, Popescu M. Predicting disease risks from highly imbalanced data using random forest. BMC Med Inform Decis Mak. 2011;11:1–13.

8. Thabtah F, Hammoud S, Kamalov F, Gonsalves A. Data imbalance in classification: Experimental evaluation. Inf Sci (N Y). 2020;513:429–41.

9. He H, Bai Y, Garcia EA, Li S. ADASYN: Adaptive synthetic sampling approach for imbalanced learning. In: 2008 IEEE international joint conference on neural networks (IEEE world congress on computational intelligence). Ieee; 2008. p. 1322–8.

10. Chawla N V, Bowyer KW, Hall LO, Kegelmeyer WP. SMOTE: synthetic minority over-sampling technique. Journal of artificial intelligence research. 2002;16:321–57.

11. Swana EF, Doorsamy W, Bokoro P. Tomek link and SMOTE approaches for machine fault classification with an imbalanced dataset. Sensors. 2022;22(9):3246.

12. Vuttipittayamongkol P, Elyan E. Neighbourhood-based undersampling approach for handling imbalanced and overlapped data. Inf Sci (N Y). 2020;509:47–70.

13. Wang ZHE, Wu C, Zheng K, Niu X, Wang X. SMOTETomek-based resampling for personality recognition. IEEE access. 2019;7:129678–89.

14. Lamari M, Azizi N, Hammami NE, Boukhamla A, Cheriguene S, Dendani N, et al. SMOTE–ENN-based data sampling and improved dynamic ensemble selection for imbalanced medical data

classification. In: Advances on Smart and Soft Computing: Proceedings of ICACIn 2020. Springer;

2021. p. 37–49.

15.   Mishra S. Handling imbalanced data: SMOTE vs. random undersampling. Int Res J Eng Technol.

2017;4(8):317–20.

16.   Khushi M, Shaukat K, Alam TM, Hameed IA, Uddin S, Luo S, et al. A comparative performance

analysis of data resampling methods on imbalance medical data. IEEE Access. 2021;9:109960–75.

17.   Shim JG, Kim DW, Ryu KH, Cho EA, Ahn JH, Kim JI, et al. Application of machine learning

approaches for osteoporosis risk prediction in postmenopausal women. Arch Osteoporos. 2020;15:1–9.

18.   Zhang PI, Hsu CC, Kao Y, Chen CJ, Kuo YW, Hsu SL, et al. Real-time AI prediction for major adverse

cardiac events in emergency department patients with chest pain. Scand J Trauma Resusc Emerg Med.

2020;28(1):1–7.

19.   Shelke MS, Deshmukh PR, Shandilya VK. A review on imbalanced data handling using undersampling

and oversampling technique. Int J Recent Trends Eng Res. 2017;3(4):444–9.

20.   Zeng M, Zou B, Wei F, Liu X, Wang L. Effective prediction of three common diseases by combining

SMOTE with Tomek links technique for imbalanced medical data. In: 2016 IEEE International

Conference of Online Analysis and Computing Science (ICOACS). IEEE; 2016. p. 225–8.

21.   Van Stralen KJ, Stel VS, Reitsma JB, Dekker FW, Zoccali C, Jager KJ. Diagnostic methods I: sensitivity,

specificity, and other measures of accuracy. Kidney Int. 2009;75(12):1257–63.

22.   Akobeng AK. Understanding diagnostic tests 1: sensitivity, specificity and predictive values. Acta

Paediatr. 2007;96(3):338–41.

23.   Enøe C, Georgiadis MP, Johnson WO. Estimation of sensitivity and specificity of diagnostic tests and

disease prevalence when the true disease state is unknown. Prev Vet Med. 2000;45(1–2):61–81.

24.   Van der Kwast TH. The trade-off between sensitivity and specificity of clinical protocols for

identification of insignificant prostate cancer. Vol. 62, European urology. Citeseer; 2012. p. 469–71.

25.   El-Badawy IM, Omar Z, Singh OP. Cooperative classification of clean and deformed capnogram

segments using a voting approach: A trade-off between specificity and sensitivity. In: 2022 44th Annual

International Conference of the IEEE Engineering in Medicine & Biology Society (EMBC). IEEE; 2022.

p. 389–93.

26.   Bassuk SS, Rifai N, Ridker PM. High-sensitivity C-reactive protein: clinical importance. Curr Probl

Cardiol. 2004;29(8):439–93.

27.   Onde ME, Ozge A, Senol MG, Togrol E, Ozdag F, Saracoglu M, et al. The sensitivity of clinical

diagnostic methods in the diagnosis of diabetic neuropathy. Journal of International Medical Research.

2008;36(1):63–70.

28.   Joly P, Richard C, Gilbert D, Courville P, Chosidow O, Roujeau JC, et al. Sensitivity and specificity of

clinical, histologic, and immunologic features in the diagnosis of paraneoplastic pemphigus. J Am Acad

Dermatol. 2000;43(4):619–26.