

1 **Management and outcomes of fractures over cranial venous sinuses:**
2 **A scoping review protocol**

3
4 **Authors:** Berjo D. Takoutsing^{1,2}, Urbaine Wunde Njineck^{1,3}, Praise Senju Wah^{1,2},
5 Lorraine Sebopelo^{2,4}, Josiane Ngouanfo Tchoffo¹, Ella Shinyuy Bah^{1,5}, Ignatius
6 Esene^{1,3}

7
8 Berjo D. Takoutsing; takoutsingberjo@gmail.com; 0000-0002-8993-2060

9 Urbaine Wunde Njineck; ubrainewunde@gmail.com; 0000-0001-9662-6806

10 Praise Senju Wah; Wsenyuy2@gmail.com; 0000-0003-3617-6190

11 Lorraine Sebopelo; sebopelolorraine@gmail.com; 0000-0003-3905-4834

12 Josiane Ngouanfo Tchoffo; josianengouanfo@gmail.com; 0009-0006-7081-2807

13 Ella Shinyuy Bah; bahellashiynyuy@gmail.com;

14 Ignatius Esene; ignatiusesene@yahoo.co.uk; 0000-0001-8785-8875

15

16 **Affiliations:**

17 ¹Research Division, Winners Foundation, Yaounde, Cameroon

18 ²Research Department, Association of Future African Neurosurgeons, Yaounde,
19 Cameroon

20 ³Faculty of Health Sciences, University of Bamenda, Bambili, Cameroon

21 ⁴Princess Marina Hospital, Gaborone, Botswana

22 ⁵Faculty of Medicine and Biomedical Sciences, University of Garoua, Garoua,
23 Cameroon

24

25

26

27 **Correspondence to:**

28 Berjo D. Takoutsing, MD

29 Research Division, Winners Foundation, Yaounde, Cameroon

30 takoutsingberjo@gmail.com

31

32

33

34

35

36

37

38

39

40 **ABSTRACT**

41

42 **Introduction**

43 Fractures over venous sinuses (FOVS) are associated with difficulties in diagnosis, and
44 treatment resulting in a high level of morbidity, and mortality. Despite its importance,
45 there is limited aggregate data to guide the management of these fractures ultimately
46 inflicting a major challenge to neurosurgeons. This protocol describes the methodology
47 of a scoping review that aims to synthesise contemporary evidence on the management
48 and outcomes of FOVS.

49 **Methods and analysis**

50 The proposed study will be conducted in accordance with the Arksey and O'Malley's
51 framework for scoping reviews. The research question, eligibility criteria and search
52 strategy were developed based on the Population, Intervention, Comparator, Outcome
53 strategy. The following electronic bibliographic databases will be searched without
54 restrictions on language, and date of publication: PubMed, WHO Global Index Medicus,
55 African Journals Online, SCOPUS, Embase. All peer-reviewed studies of primary data

56 reporting on the management and outcomes of FOVS will be included. The data
57 extracted from included articles will be presented through descriptive statistics, pooled
58 statistics, and a narrative description.

59 **Ethics and dissemination:**

60 Because this study did not directly involve human individuals, ethical approval was not
61 necessary. Dissemination strategies will include publication in a peer-reviewed journal,
62 oral and poster presentations at local, regional, national and international conferences
63 and promotion over social media.

64

65 **Word count:** 208

66 **ARTICLE SUMMARY**

67 **Strengths and limitations of this study**

- 68 • To our knowledge this is the first scoping review focusing on the management
69 and outcomes of fractures over the venous sinuses
- 70 • This protocol will ensure transparency in the methodology of the proposed
71 review, thus will reduce the likelihood of reviewing bias.
- 72 • The search strategy will be conducted in five electronic databases commonly
73 used across both high- and low-income countries.
- 74 • There will be no restrictions on language, location, or publication date during the
75 screening process.
- 76 • Unpublished studies will not be sought.
- 77 • There will be no formal assessment on the quality of the included studies.

78

79

80 **Keywords:** Craniocerebral Trauma, Cranial Venous Sinuses, Evidence Gaps, Scoping
81 Review, Skull Fractures

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102 INTRODUCTION

103

104 Skull fractures following trauma continues to be a major public health problem
105 disproportionately affecting low- and middle-income countries (LMICs).¹ It is commonly
106 associated with traumatic brain injury (TBI) occurring in 28-37% of affected patients.^{1,2}
107 Several types of skull fractures exist, with depressed skull fractures and skull base
108 fractures usually associated with injury to the underlying dural venous sinuses.³ Despite
109 its importance, fractures over venous sinuses (FOVS) can be associated with a missed
110 or delayed diagnosis, and difficult management.⁴ These results in a high morbidity, and
111 mortality in the pre-operative, operative and perioperative periods.⁵

112 About 10% of skull fractures occur over the cranial venous sinuses, with the majority
113 occurring in the context of direct trauma, followed by road traffic accidents especially in
114 LMICs owing to safety measures not fully implemented.^{6,7} FOVS are associated with
115 high in-patient, mixed clinical presentation and are prone to high mortality.^{5,8} The
116 management can be conservative or surgical both of which have different indications
117 and risks.⁶ Due to the limited guidelines or consensus on the management of these
118 fractures coupled with the unavailability of imaging modalities such as CT scans, MRI in
119 resource limited settings, neurosurgeons are thus faced with a therapeutic dilemma
120 usually relying on good judgment and experience.⁹

121 There are ongoing investigations on novel therapeutic interventions to reduce mortality
122 associated with FOVS.^{8,10} However, they may be inaccessible in resource-limited
123 settings and emergency cases. There is therefore a need for a comprehensive overview
124 of the available literature on the management and outcomes of these fractures to
125 identify the various therapeutic modalities, assess the availability of various diagnostic
126 methods, and evaluate patients' outcomes. This would fill in key research gaps and
127 priorities, and inform future interventions.

128

129

130 STUDY AIMS

131 This protocol describes the methodology of a scoping review of the literature on the
132 management and outcomes of cranial venous sinus fractures. **Box 1** describes the
133 primary and secondary goals of the proposed review. The suggested review will provide
134 information to health care practitioners, policymakers, stakeholders, and healthcare
135 organisations involved in providing care to patients who have suffered venous sinus
136 fractures. The knowledge gained will aid in the identification of key priorities and the
137 establishment of larger research initiatives with a greater level of evidence.

138

139 **Box 1. Primary and secondary aims of the scoping review.**

Primary aim:

1. Describe the therapeutic modalities for FOVS.

Secondary aim:

2. Assess the availability of diagnostic methods including neuroimaging (MRI, CT scan), and fundoscopy.
3. Evaluate patient's clinical outcomes primarily defined as mortality and complication rates, Glasgow Outcome Score, Length of hospital stay, and clinical state on discharge.

140

141

142

143 **METHODS**

144 The planned scoping review will be carried out in accordance with the five stages of
145 Arksey and O'Malley's framework.¹¹ This review's protocol is formatted in accordance
146 with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses
147 (PRISMA) of Protocols.¹²

148

149 **Stage 1: identifying the research question**

150 The Patient, Intervention, Comparison, Outcome (PICO) framework was utilised to
151 determine the title, objectives, research questions, and eligibility criteria for the scoping
152 review. Our key research question is: How are venous sinus fractures managed
153 globally?

154

155 **Stage 2: identifying relevant studies**

156 The authors defined the keywords that will be employed in the search strategy as well
157 as the study eligibility criteria using the PICO approach.

158

159 **Databases and search strategy**

160 BDT developed the search strategy to identify research papers on fractures over the
161 venous sinus using multiple combinations of keywords, MeSH terms, and different
162 electronic bibliographic databases. Following discussion with the review team, the final
163 search strategy broadly consisted of variants of 'Skull fracture' and 'Venous sinus'
164 (**Supplementary Table 1**). The literature search was conducted from inception until 29
165 October 2023 without restrictions on language or date of publication in the following
166 databases: PubMed, WHO Global Index Medicus, African Journals Online, Scopus, and
167 Embase. A reviewer will hand search additional resources such as Google scholar, and
168 references from included studies to verify that all relevant material in the literature is
169 included.

170

171 **Eligibility criteria**

172 An eligibility criteria was developed following discussion with the reviewers to guide the
173 screening phase. The inclusion criteria adhere to the PICO strategy and are listed
174 below:

175

176 ***Population***

177 All studies of patients of all ages with a diagnosis of venous sinus fracture will be
178 included. Patients with the diagnosis of head injury or a pathology of the venous sinus
179 without any associated fracture will be excluded. Also, studies whose data cannot be
180 disaggregated to one particular country will be excluded.

181

182 ***Intervention***

183 Studies reporting on the diagnosis, and/or treatment (conservative or surgical treatment)
184 for fractures over the venous sinuses will be included. Patients diagnosed with a venous
185 sinus fracture whose method of diagnosis or treatment is unclear will be excluded.

186 ***Comparison***

187 We shall compare different modalities of treatment across various age groups (adults vs
188 paediatric patients), and income levels according to the World Bank Country and
189 Lending group.¹³

190 ***Outcome***

191 Studies reporting on (1) mortality rate (2) complication rate (3) Glasgow Outcome Score
192 (4) length of hospital admission (5) symptoms resolved/reduced/ unchanged/worsened
193 will be included.

194 ***Types of studies***

195 Will be included peer-reviewed publications such as: Cross-sectional studies, case
196 reports, case series, cohort studies, trials and audits. There will be no language
197 restriction and where needed, reports will be translated using online translation
198 platforms such as DeepL Translator ([https://www.deepl.com/en/ translator](https://www.deepl.com/en/translator)). The
199 following articles will be excluded if they do not have disaggregated data on fractures
200 over the venous sinus. Perspectives, letters, opinion pieces, editorials, comments,
201 guidelines, reviews, meta-analyses and articles published in non-peer-reviewed journals
202 will be excluded.

203

204

205 **Stage 3: Study selection**

206 The search results will be uploaded to Rayyan (<https://www.rayyan.ai>) to facilitate de-
207 duplication and independent, blinded screening.¹⁴ Initially, titles and abstracts will be
208 screened by two independent reviewers against the eligibility criteria. This will be
209 followed by full-text screening where the same reviewers will screen full-text studies
210 using the eligibility criteria mentioned above. Prior to these two steps, a calibration
211 exercise will be carried by the screeners to ensure an adequate understanding of the

212 eligibility criteria. During the full-text screening phase, corresponding authors of articles
213 will be contacted if the full-text cannot be accessed. A reminder will be sent after 2
214 weeks If no response is received. If still no avail, the article will be excluded and tagged
215 'full-text unavailable'. Unless otherwise stated, disagreements will be discussed and
216 resolved amongst the screeners following unblinding of their decisions on Rayyan.
217 Where consensus cannot be achieved, the lead or senior author (BT or IE) will arbitrate.
218 A PRISMA flow diagram (**figure 1**) will be presented reflecting the search process.

219

220 **Stage 4: Charting the data**

221 Data from studies who met the inclusion criteria will be independently extracted and
222 checked by two members of the review team using a standardised extraction proforma.
223 Any unresolved disagreements during this stage will be settled by a third reviewer (BT
224 or IE). This stage will be preceded by a pilot phase. In the pilot phase, all extractors will
225 extract data from the same ten randomly selected included articles. This will help
226 ensure the reliability of the data extraction sheet and that all authors accurately, and
227 homogeneously extract the data. Following this phase, necessary changes will be made
228 to the proforma to ensure that it reflects the included studies before initiation of the
229 extraction proper stage. Data items will include information on study and sample
230 characteristics; type, site and aetiology of venous sinus fracture; clinical presentation
231 and method of diagnosis; treatment mode, timing, and follow-up; and primary and
232 secondary outcomes. If data is insufficiently reported, we will contact the corresponding
233 author as described in '**Stage 3**'. All extracted data will be recorded and stored in a
234 spreadsheet. The expected data to be extracted are outlined in **Box 2**.

235 **Box 2. Key data to be extracted on the spreadsheet.**

Study characteristic

- First author
- Year of publication
- Country where the study was conducted

- Language of publication
- Year(s) of data collection
- Income level of study setting
- Region of study setting
- Study design
- Number of centers involved

- Number of with fracture over dural venous sinus

- Age (Mean, median, standard deviation and/or range)

- Sex (Female, male)

Characteristics of FOVS

- Mechanism of injury
- Type of fracture
- Fracture site
- Associated cranio-cerebral lesions

Management of FOVS

- Clinical presentation
- Method of diagnosis
- Treatment mode (Surgical vs non-surgical)
- Timing of treatment

Outcome of FOVS

- Mortality rate
- Complication rates
- Glasgow outcome score
- Functional outcome
- Length of hospital admission
- Symptoms resolved/ reduced/ unchanged/ worsened

236

237

238 **Stage 5: collating, summarising and reporting the results**

239 Following extraction, data will be transferred to SPSS V.26 for analysis. Results will be
240 primarily reported in a tabular form showing descriptive statistics. When appropriate, a
241 narrative description of the results will be presented by grouping the data into
242 meaningful summaries to allow comprehensive reporting of the findings. Pooled
243 statistics will be calculated and presented using measures of central tendency and
244 spread. Unless otherwise stated, the statistical significance will be set at $p < 0.05$. The
245 tables and figures may display information such as: The publication year of the studies,
246 the geographical area where the studies were conducted, study design and
247 demographic characteristics of included patients; aetiology, site, and type of venous
248 sinus fracture; clinical presentation, mode of diagnosis, and mode of treatment; the
249 length of hospital stay, timing of follow-up and outcomes.

250

251 **Risk of bias assessment**

252 Given the limited and varied evidence on the topic, the proposed scoping review seeks
253 to offer a comprehensive overview of the management and outcomes of venous sinus
254 fractures. As a result of the inherent standard biases associated with new areas of
255 clinical research, a formal bias risk assessment was judged unnecessary.

256

257 **Ethics and dissemination**

258 Because this study did not directly involve human individuals, ethical approval was not
259 necessary. Dissemination strategies will include publication in a peer-reviewed journal,
260 oral and poster presentations at local, regional, national and international conferences
261 and promotion over social media. This work will provide an overview of the available
262 evidence on the management and outcomes of venous sinus fractures globally. This will
263 help in the identification of key research priorities and uncovering any disparities among
264 countries. The proposed review may inform the development of new research questions
265 as well as interventions to systematically improve the management and outcomes of
266 patients sustaining a venous sinus fracture.

267

268 **Patient and public involvement**

269 None

270

271 **ACKNOWLEDGEMENTS**

272 None

273

274

275

276 **REFERENCES**

- 277 1. Tsai Y-C, Rau C-S, Huang J-F, et al. The association between skull bone
278 fractures and the mortality outcomes of patients with traumatic brain injury.
279 Emerg Med Int. 2022;2022:1296590.
280
- 281 2. UpToDate. <https://www.uptodate.com/contents/skull-fractures-in-adults>
282 (accessed 17 November 2023)
283
- 284 3. Skull Fracture - an overview | ScienceDirect Topics.
285 <https://www.sciencedirect.com/topics/psychology/skull-fracture> (accessed 17
286 November 2023)
287
- 288 4. Wang W, Lin J, Luo F, et al. Early Diagnosis and Management of Cerebral
289 Venous Flow Obstruction Secondary to Transsinus Fracture after Traumatic
290 Brain Injury. J Clin Neurol. 2013;9:259–68.
291
- 292 5. Kim Y-S, Jung S-H, Lim D-H, et al. Traumatic Dural Venous Sinus Injury. Korean
293 J Neurotrauma. 2015;11:118–23.
294
- 295 6. Aziz MM, Molla S, Abdelrahim HA, et al. Depressed Skull Fractures Overlying
296 Dural Venous Sinuses: Management Modalities and Review of Literature. Turk
297 Neurosurg. 2019;29:856–63.
298
- 299 7. Abdelaal MA, Saro ASEM, Fadl KN, et al. Management of Compound Depressed
300 Fractures Over Major Cranial Venous Sinuses. The Egyptian Journal of Hospital
301 Medicine. 2021;83:1177–82.
302
- 303 8. Qureshi AI, Sahito S, Liaqat J, et al. Traumatic Injury of Major Cerebral Venous
304 Sinuses Associated with Traumatic Brain Injury or Head and Neck Trauma:
305 Analysis of National Trauma Data Bank. J Vasc Interv Neurol. 2020;11:27–33.
306
- 307 9. Mohamed a. Thabit MD. Venous Sinus Thrombosis Following Repair of
308 Compound Depressed Fractures Overlying Sinuses. The Medical Journal of
309 Cairo University. 2019;87:3277–82.
310
- 311 10. Stein SC. The Evolution of Modern Treatment for Depressed Skull Fractures.
312 World Neurosurg. 2019;121:186–92.
313
- 314 11. Arksey H, O'Malley L. Scoping studies: towards a methodological framework.
315 International Journal of Social Research Methodology. 2005;8:19–32.

316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348

12. Shamseer L, Moher D, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ*. 2015;349:g7647.

13. World Bank Country and Lending Groups – World Bank Data Help Desk. <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups> (accessed 17 November 2023)

14. Ouzzani M, Hammady H, Fedorowicz Z, et al. Rayyan—a web and mobile app for systematic reviews. *Syst Rev*. 2016;5:210.

TWITTER: @BerjoDongmo, @PraiseWah, @ngouanfojosiane, @UbraineWunde, @LASEbopelo, @ignatiusesene

AUTHOR CONTRIBUTIONS

BT, and IE were responsible for conceiving the article. IE is the guarantor. BT, UWN, PSW, LS, and JNT wrote the manuscript. IE, BT, UWN, PSW, LS, JNT, and ESB provided a critical appraisal of the manuscript. All authors critically revised and approved the final manuscript.

DATA AVAILABILITY STATEMENT


FUNDING

This work did not receive any funding.

COMPETING INTERESTS

None declared.

Inception – Day search will be conducted



Identification

Screening

Eligibility

Included