Three components of glucose dynamics – mean, variance, and autocorrelation – are independently associated with coronary plaque vulnerability

Authors

Hikaru Sugimoto,[1] Ken-ichi Hironaka,[2] Tomoko Yamada,[3] Natsu Otowa-Suematsu,[3] Yushi Hirota,[3] Hiromasa Otake,[4] Ken-Ichi Hirata,[3] Kazuhiko Sakaguchi,[3] Wataru Ogawa,[3]* and Shinya Kuroda[1,2,5]*

Affiliations

[1]Department of Biochemistry and Molecular Biology, Graduate School of Medicine, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

[2]Department of Biological Sciences, Graduate School of Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

[3]Division of Diabetes and Endocrinology, Department of Internal Medicine, Kobe University Graduate School of Medicine, Kusunoki-cho 7-5-1, Chuo-ku, Kobe, Hyogo 650-0017, Japan

[4]Division of Cardiovascular Medicine, Department of Internal Medicine, Kobe University Graduate School of Medicine, Kusunoki-cho 7-5-1, Chuo-ku, Kobe, Hyogo 650-0017, Japan

[5]Lead contact

*Corresponding authors: Wataru Ogawa, ogawa@med.kobe-u.ac.jp; Shinya Kuroda, skuroda@bs.s.u-tokyo.ac.jp

ABSTRACT

Impaired glucose homeostasis leads to numerous complications, with coronary artery disease (CAD) being a major contributor to healthcare costs worldwide. Given the limited efficacy of current CAD screening methods, we investigated the association between glucose dynamics and a predictor of coronary events measured by virtual histology-intravascular ultrasound (%NC), with the aim of predicting CAD using easy-to-measure indices. We found that continuous glucose monitoring (CGM)-derived indices, particularly average daily risk ratio (ADRR) and AC_Var, exhibited stronger predictive capabilities for %NC compared to commonly used indices such as fasting blood glucose (FBG), hemoglobin A1c (HbA1c), and plasma glucose level at 120 min during oral glucose tolerance tests (PG120). Factor analysis identified three distinct components underlying glucose dynamics – mean, variance, and autocorrelation – each independently associated with %NC. ADRR was influenced by the first two components and AC_Var by the third. FBG, HbA1c, and PG120 were influenced only by the mean component, making them insufficient for %NC prediction. Our results were validated using data sets from Japan (n=64), America (n=53), and China (n=100). CGM-derived indices reflecting the three components of glucose dynamics can serve as more effective screening tools for CAD risk assessment, complementing or possibly replacing traditional diabetes diagnostic methods.

INTRODUCTION

Diabetes mellitus (DM) affects more than 400 million people globally, and coronary artery disease (CAD) is a major driver of morbidity, mortality and healthcare costs among individuals with type 2 DM (T2DM) (Bax et al., 2007). Various prognostic models (Fiarni et al., 2019; Ravaut et al., 2021) and diagnostic markers (Bax et al., 2007) for CAD have been developed; however, population-level screening remains imperfect, often being ineffective, costly, or labor-intensive (Bax et al., 2007; Young et al., 2009). There is a need for scalable approaches that can identify individuals at the highest risk using easily accessible clinical information.

Blood glucose levels are among the readily obtained predictors of the complications (Psoma et al., 2022). The disrupted conditions of glucose dynamics seen in impaired glucose tolerance (IGT) and T2DM are partly characterized by high concentrations of blood glucose levels (Monnier et al., 2008). High concentrations of blood glucose levels have been defined as having high hemoglobin A1c (HbA1c) levels, fasting blood glucose (FBG) levels, and plasma glucose concentration at 120 min during the oral glucose tolerance test (OGTT) (PG120) (Monnier et al., 2008). These indices, especially HbA1c, are associated with complications of T2DM (Selvin et al., 2010).

Recent studies have shown that glucose variability, in addition to absolute glucose concentration, significantly contributes to the prognosis of complications (Gerbaud et al., 2019; Gorst et al., 2015; Monnier et al., 2008; Psoma et al., 2022; Su et al., 2011; Zhou et al., 2018) and all-cause mortality (Cai et al., 2022). Continuous glucose monitoring (CGM) can estimate short-term glycemic variability (Service, 2013), and has been reported to predict T2DM complications (Tang et al., 2016). Standard deviation (Std) of glucose levels (CGM_Std), mean amplitude of glycemic excursion (MAGE), mean of daily difference (MODD), and continuous overlapping net glycemic action (CONGA) are established indices of glycemic variability, of which CGM_Std and MAGE are more highly correlated with coronary plaque properties (Otowa-Suematsu et al., 2018). Among glucose level-related indices, including HbA1c and FBG, MAGE is an independent determinant of coronary plaque instability (Okada et al., 2015).

Other CGM-derived indices such as average daily risk ratio (ADRR), lability index (LI), J-index, mean absolute glucose (MAG), and glycemic risk assessment in diabetes equation (GRADE) have also been developed (Hill et al., 2011). We recently showed that AC_Mean and AC_Var, which are calculated from the autocorrelation function of glucose levels measured by CGM, can detect decreased abilities in glucose regulation that cannot be captured by FBG, HbA1c, or the other conventional CGM-derived indices (Sugimoto et al.,

2025). The characteristics of glucose dynamics can also be estimated from insulin concentrations. The disposition index (DI), which is the product of insulin sensitivity and insulin secretion, reflects and predicts glycemic disability beyond FBG (Utzschneider et al., 2009). Several other glucose-related indices and the relationship between the indices have also been reported (Fabris et al., 2015, 2014; Keshet et al., 2023). Despite these advances, a comprehensive understanding of how these various indices can be optimally combined to predict T2DM complications, particularly CAD, remains elusive. Furthermore, the underlying factors that these indices represent and their individual associations with CAD remain to be fully elucidated.

This study aims to address these knowledge gaps through three objectives: (i) to determine which clinical parameters are effective predictors of coronary plaque vulnerability; (ii) to identify the factors underlying these indices; and (iii) to elucidate how these factors are associated with coronary plaque vulnerability. We investigated the characteristics of 14 CGM-derived indices: 12 relatively well-known CGM-derived indices (Hill et al., 2011) and 2 indices (AC_Mean and AC_Var) as well as OGTT-derived indices, and investigated the relationship between these parameters and coronary plaque vulnerability assessed by virtual histology-intravascular ultrasound (VH-IVUS). We showed that three factors, namely, mean, variance, and autocorrelation, underlie blood glucose level-related indices, and that the three are independently associated with coronary plaque vulnerability.

RESULTS

Mean, standard deviation, and autocorrelation function of glucose levels independently contribute to the prediction of coronary plaque vulnerability

We previously showed that AC_Var, a measure derived from the autocorrelation function of CGM time series, detects impaired glucose handling that is not reflected by the CGM-derived mean (CGM_Mean) or standard deviation (CGM_Std) (Sugimoto et al., 2025). Based on the study, we hypothesized that AC_Var could identify individuals with high %NC, a widely used parameter of plaque vulnerability, independently from CGM_Mean and CGM_Std. To test this hypothesis, we conducted a multiple regression analysis with CGM_Mean, CGM_Std, and AC_Var as input variables and %NC as the objective variable using a previously described dataset consisting of 8 individuals with normal glucose tolerance (NGT), 16 with IGT, and 29 with T2DM (Otowa-Suematsu et al., 2018).

The variance inflation factors (VIFs) for CGM_Mean, CGM_Std, and AC_Var were 1.1, 1.1, and 1.0, respectively, indicating low multicollinearity among these variables. The regression model including CGM_Mean, CGM_Std and AC_Var to predict %NC achieved an R^2 of 0.36 and an Akaike Information Criterion (AIC) of 321. Each of these indices showed a statistically significant independent positive correlation with %NC (**Fig. 1A**). In contrast, the model using conventional glycemic markers (FBG, HbA1c, and PG120) yielded an R^2 of only 0.05 and an AIC of 340 (**Fig. 1B**). Similarly, the model using the Framingham Risk Score for Hard Coronary Heart Disease (Wilson et al., 1998) showed limited predictive value, with an R^2 of 0.04 and an AIC of 330 (**Fig. 1C**). The AC_Var computed from 15-minute CGM sampling was nearly identical to that computed from 5-minute sampling (R = 0.99, 95% CI: 0.97-1.00) (**Fig. S1A**), and the regression using the 15-min features yielded almost the same performance (R^2 = 0.36; AIC = 321; **Fig. S1B**). Collectively, these results indicate that CGM_Mean, CGM_Std and AC_Var provide complementary and independent information for predicting coronary plaque vulnerability.

Association among clinical variables

To further characterize CGM-derived indices in estimating %NC, we examined Spearman's correlation coefficients (*r*) between CGM-derived indices and %NC (**Fig. 1D, E**). For comparison, we also investigated FBG, HbA1c, OGTT-derived indices, body mass index (BMI), triglycerides (TGs), low-density lipoprotein cholesterol (LDL-C), high-density lipoprotein cholesterol (HDL-C), systolic blood pressure (SBP), and diastolic BP (DBP). Given that this study enrolled individuals with well-controlled serum cholesterol and BP levels (**Fig. S1C**), weak correlations between these indices and %NC were expected.

A correlation network connecting relationships with Q < 0.05 showed that AC_Var was
statistically significantly correlated with %NC ($r = 0.35$; 95%CI, 0.09–0.57) (**Fig. 1D, E**). By
contrast, AC_Var showed relatively weak correlations with other indices, including
CGM_Mean ($r = –0.02$; 95% CI, –0.30–0.24) and CGM_Std ($r = 0.15$; 95% CI, –0.14–0.43)
(**Fig. 1F**). Twelve CGM-derived indices, ADRR, MAGE, JINDEX, CGM_Std, CGM_Mean,
GRADE, MVALUE, AC_Var, LI, HBGI, CONGA, and MODD, showed significant
correlations with %NC (**Fig. 1E, S1D**). By contrast, with the exception of the insulinogenic
index (I.I.), OGTT-derived indices, FBG and HbA1c showed relatively weak correlations
with %NC.
147
To assess the multicollinearity among the input variables, we examined the VIF for each
variable (**Fig. 1G, H**). We removed variables with the highest VIF one by one until all
variables had VIF values less than 10. This procedure retained 19 variables when considering
the full set of clinical and CGM measures (**Fig. 1G**) and eight variables when considering
only CGM-derived measures (**Fig. 1H**). AC_Var was retained in both sets, indicating
relatively low multicollinearity with the other indices.
154
To examine the reproducibility of the relationship among the input variables, we compared
VIF values calculated from this dataset (VIFt) with those from previously reported datasets
(VIFp1 (Sugimoto et al., 2025) and VIFp2 (Hall et al., 2018)). VIFt significantly correlated
with both VIFp1 and VIFp2 (**Fig. 1I, J**). VIFs for AC_Mean and AC_Var were consistently
low across the datasets, suggesting the reproducibility of our observation that AC_Mean and
AC_Var show relatively low multicollinearity with other CGM-derived indices.
161

CGM-derived indices, particularly ADRR, AC_Var, MAGE, and LI, contribute to the
prediction of coronary plaque vulnerability
164   To investigate which variables are particularly useful in estimating %NC, we used two
165   statistical techniques: Least Absolute Shrinkage and Selection Operator (LASSO) regression
166   and Partial Least Squares (PLS) regression (**Fig. 2**) (Tibshirani, 1996; Wold et al., 2001).
167   These regression models have been used for studies where the number of input variables is
168   large relative to the sample size (Pei et al., 2023; Wang et al., 2005).
169
LASSO uses L1 regularization to produce models with fewer parameters and has been widely
used for feature selection in predictive modeling (Wei et al., 2022). We included BMI, FBG,
HbA1c, OGTT-derived indices, and CGM-derived indices as the input variables. The leave-
one-out cross-validation identified the optimal regularization coefficient, lambda, as 0.849
(**Fig. 2A**). At the lambda, the coefficients of ADRR, AC_Var, MAGE, and LI were estimated
to be non-zero coefficients (**Fig. 2B, C**), suggesting that CGM-derived indices, particularly

ADRR, AC_Var, MAGE, and LI, contribute to the prediction of %NC. Even with the
inclusion of SBP, DBP, TG, LDL-C, and HDL-C as additional input variables, the results
remained consistent, with the coefficients of ADRR, AC_Var, MAGE, and LI still estimated
as non-zero coefficients (**Fig. S2A-C**).
180
To further validate the LASSO results and address potential instability, we performed PLS
regression and examined the Variable Importance in Projection (VIP) scores (**Fig. 2D**). PLS
regression is particularly useful when dealing with many input variables that may be highly
collinear (Wold et al., 2001). The VIP scores of ADRR, AC_Var, MAGE, and LI, which
were estimated to be non-zero coefficients by LASSO, were higher than 1, indicating that
these four variables especially contribute to the prediction of %NC. Conventional indices,
including FBG, HbA1c, PG120, I.I., Composite index, and Oral DI, did not contribute
significantly to the prediction compared to these CGM-derived indices. Even when SBP,
DBP, TG, LDL-C, and HDL-C were included as additional input variables, the results
remained consistent, and the VIP scores for ADRR, AC_Var, MAGE, and LI remained
greater than 1 (**Fig. S2D**).
192

Three components of glucose dynamics –mean, variance, and autocorrelation – are
associated with coronary plaque vulnerability
To elucidate the underlying factors of clinical parameters and their association with %NC, we
performed an exploratory factor analysis. Factor analysis reduces interrelated indices into a
smaller set of underlying common factors, and has been employed to examine the
interdependencies among various clinical parameters (Augstein et al., 2015; Cappelleri et al.,
2000; Oh et al., 2004) and DM complications (Guo et al., 2020).
200

The optimal number of underlying factors was determined using Bayesian information
criterion (BIC) and minimum average partial (MAP) methods, which indicated that five and
six factors were appropriate, respectively. We first set the number of underlying factors as
five. **Figure 3A** shows that FBG, HbA1c, PG120, I.I., oral DI, CGM_Mean, CONGA, HBGI,
MVALUE, GRADE, JINDEX, and ADRR were included in the first factor. Given that most
of these indices are related to the value of blood glucose concentration, factor 1 was labeled
"mean." CGM_Std, MAGE, LI, MAG, MODD, JINDEX, and ADRR were included in the
second factor. Given that these indices are related to glucose variability, factor 2 was labeled
"variance." Given that the definition of JINDEX is based on the sum of CGM_Mean and
CGM_Std, and that of ADRR is based on both high and low values of glucose, the result that
JINDEX and ADRR clustered in both factors 1 and 2 is plausible. Given that autocorrelation-
derived indices, AC_Mean and AC_Var, were included in the third factor, factor 3 was
labeled "autocorrelation." BMI, PG120, composite index, and oral DI were included in the

fourth factor. Factor 4 did not include any CGM-derived indices. Given that this combination of indices indicates a decrease in oral DI and associated increase in blood glucose due to decreased insulin sensitivity, factor 4 was labeled "sensitivity (without CGM)". PG120, I.I., oral DI, and MAG were included in the fifth factor. Factor 5 did not have positive loadings of any CGM-derived indices. Given that this combination of the indices indicates a decrease in oral DI and associated increase in blood glucose due to decreased insulin secretion (I.I.), factor 5 was labeled "secretion (without CGM)." Each factor explained 39%, 21%, 10%, 5%, and 5% of the total variance of the factors, respectively.

The validity of the factor analysis was assessed according to previous studies (Cappelleri et al., 2000; Guo et al., 2020). To evaluate the applicability of the factor analysis, the Kaiser-Meyer-Olkin (KMO) and Bartlett's spherical test were performed. The KMO test indicated that the value of the measure of sampling adequacy for this data was 0.64, and Bartlett's spherical test indicated that the variables were statistically significantly intercorrelated ($P < 0.01$), suggesting that this dataset was applicable for the factor analysis. To evaluate the internal consistency, Cronbach's α (**Fig. 3B**) and item–total correlations were calculated for each factor. Cronbach's α was 0.97 for factor 1, 0.93 for factor 2, 0.90 for factor 3, 0.72 for factor 4, and 0.66 for factor 5; these values were larger than 0.65, suggesting that the internal consistency was satisfactory. While Cronbach's α of factor 5 was relatively low, exclusion of MAG increased the Cronbach's α to 0.84, indicating that the association between factor 5 and decrease in oral DI and associated increase in blood glucose due to decreased insulin secretion could be considered reliable. Item–total correlations ranged from 0.63 to 0.97 for factor 1, 0.72 to 0.94 for factor 2, 0.82 for factor 3, 0.54 to 0.76 for factor 4, and 0.37 to 0.86 for factor 5. With the exception of MAG, item–total correlations ranged from 0.84 to 0.91 for factor 5. The correlation coefficient of MAG was 0.37, which can be considered a modest correlation (Cappelleri et al., 2000), and the item–total correlations were generally reasonably strong in demonstrating reliability.

We also investigated a 6-factor solution (**Fig. S3A**). Factors 1, 2, and 3 could be interpreted as mean, variance, and autocorrelation, respectively, similar to the 5-factor solution. Given that factor 6 had no factor loadings ≥ 0.3, we applied the 5-factor solution in the subsequent analysis. Furthermore, the inclusion of SBP, DBP, TG, LDL-C, and HDL-C into the input variables did not change the presence of the three components (mean, variance and autocorrelation) in glucose dynamics (**Fig. S3B**). Since we included only individuals with well-controlled serum cholesterol and BP levels in this study, we applied the 5-factor solution without these indices (**Fig. 3A**) to the following analysis.

To further examine the stability of the results of the factor analysis, we also conducted hierarchical clustering analysis (**Fig. S4**). The optimal number of clusters was determined based on silhouette analysis. A large positive silhouette coefficient indicates that each cluster is compact and distinct from the others. The analysis indicated that the four clusters were appropriate (**Fig. S4A**). Clusters 1, 2, and 3 can be interpreted as mean, variance, and autocorrelation, respectively (**Fig. S4B**).

To investigate the association between these underlying factors and %NC, we investigated the correlation between the factor scores and %NC (**Fig. 3C**). The factor mean, variance showed significant positive correlations with %NC, whereas autocorrelation showed a significant negative correlation. Factors 4 and 5, which were less related to the CGM-derived indices, showed weaker correlations with %NC. Collectively, we conclude that glucose dynamics have three components – mean, variance, and autocorrelation – and that these three components are associated with %NC.

To assess the robustness and generalizability of our factor analysis results, we performed similar analyses using previously published datasets from diverse populations (**Figs. S5, 6**). Factors that could be interpreted as representing the mean, variance, and autocorrelation of glucose dynamics were consistently observed across diverse populations, including Japanese (Sugimoto et al., 2025) (**Fig. S5A**), American (Hall et al., 2018) (**Fig. S5B**), and Chinese (Zhao et al., 2023) (**Fig. S6A**) datasets. In the Chinese dataset, all three factors were significantly different between individuals with and without diabetic macrovascular complications (**Fig. S6B**). Taken together, these results support the reproducibility and cross-cultural validity of our three-component model of glucose dynamics.

Overview of the three components of glucose dynamics – mean, variance, and autocorrelation –

We have shown that three components of glucose dynamics – mean, variance, and autocorrelation – are associated with %NC. To overview the characteristics of glucose dynamics with different values of the components, we simulated glucose fluctuations using a previously reported mathematical model (De Gaetano and Arino, 2000) (**Figs. 4, S7**). We could generate glucose fluctuations with almost the same standard deviation (Std) and AC_Var but with a different mean (**Fig. 4A**). Similarly, we could also simulate glucose fluctuations with almost the same mean and AC_Var but different Std, and with almost the same mean and Std but different AC_Var. These three components were independently adjustable by changing parameters within the range of values for NGT individuals (**Fig. S7B**). Individuals with higher AC_Var tended to have higher %NC (**Fig. 1**); however,

comparing the glucose dynamics with higher and lower AC_Var, the maximum value of

blood glucose was lower in individuals with higher AC_Var (**Fig. 4A**).

We then investigated the relationship between the three components and shapes of the

glucose response curve after OGTT. Patterns of the glucose response curve after OGTT were

heterogeneous, and four distinct patterns, denoted classes 1–4, were previously identified by

analyzing the glucose dynamics of 5861 individuals in Denmark (**Fig. 4B**) (Hulman et al.,

2018). The classes with a high mean did not necessarily have high Std or AC_Var (**Fig. 4C**).

The classes with high Std did not necessarily have high mean and AC_Var (**Fig. 4C**),

consistent with the result that mean, Std, and AC_Var had low multicollinearity with each

other. Compared to class 2, only class 1 was lower in mean and Std, and was higher in

AC_Var. Compared to class 2, only class 3 was higher in mean, Std, and AC_Var. Compared

to class 2, only class 4 was higher in mean and Std, and lower in AC_Var. Collectively, the

three components could characterize the previously reported four distinct patterns during

OGTT. Class 3 was characterized by normal FBG and PG120 values, but has been associated

with increased risk of diabetes and higher all-cause mortality rate, suggesting that subgroups

at high risk may not be identified by investigating only FBG and PG120 (Hulman et al.,

2018). Std and AC_Var were high in class 3 (**Fig. 4C**), suggesting that high Std and high

AC_Var indicate glycemic disability independent of FBG and PG120.

DISCUSSION

Our results show that glucose dynamics consist of three distinct components - mean, variance and autocorrelation - each of which is independently associated with coronary plaque vulnerability. We further show that traditional diabetes diagnostics rely on indices such as FBG, PG120, and HbA1c that primarily reflect only the mean component, and that incorporating all three components significantly improves the prediction of coronary plaque vulnerability. This result is partially consistent with a previous notion that glucose dynamics include two components: amplitude and timing (Cobelli and Facchinetti, 2018). Given our previous study demonstrating that AC_Var, an index reflecting autocorrelation, identifies impaired glucose regulation independently of CGM-derived indices representing mean and variance (Sugimoto et al., 2025), CGM-derived indices incorporating all three components may outperform conventional diabetes diagnostics in predicting both glycemic control capacity and coronary plaque vulnerability.

We also showed that CGM-derived indices, especially ADRR and AC_Var, contribute to the prediction of %NC by using LASSO and PLS. Given that the definition equation for ADRR is affected by both high and low concentrations of blood glucose (Hill et al., 2011), it is likely affected by both the mean and variance components. Factor analysis confirmed that ADRR was included in both factor 1 (mean) and factor 2 (variance). Since three factors, mean, variance, and autocorrelation, contribute independently to the prediction of the complication, it would be useful to examine ADRR, which is influenced by both mean and variance, and AC_Var, which is influenced by autocorrelation, in predicting %NC with a minimal number of variables.

This study also provided evidence that autocorrelation can vary independently from the mean and variance components using simulated data. In addition, simulated glucose dynamics indicated that even individuals with high AC_Var did not necessarily have high maximum and minimum blood glucose levels. This study also indicated that these three components qualitatively corresponded to the four distinct glucose patterns observed after glucose administration, which were identified in a previous study (Hulman et al., 2018). Thus, the inclusion of autocorrelation in addition to mean and variance may improve the characterization of inter-individual differences in glucose regulation and improve the predictive accuracy of various clinical outcomes.

Despite increasing evidence linking glycemic variability to oxidative stress and endothelial dysfunction in T2DM complications (Ceriello et al., 2008; Monnier et al., 2008), the biological mechanisms underlying the independent predictive value of autocorrelation remain to be elucidated. Our previous work has shown that glucose autocorrelation is influenced by

insulin clearance (Sugimoto et al., 2025), a process known to be associated with

cardiovascular disease risk (Randrianarisoa et al., 2020). Therefore, the molecular pathways

linking glucose autocorrelation to cardiovascular disease may share common mechanisms

with those linking insulin clearance to cardiovascular disease. Although previous studies have

primarily focused on investigating the molecular mechanisms associated with mean glucose

levels and glycemic variability, our findings open new avenues for exploring the molecular

basis of glucose autocorrelation, potentially revealing novel therapeutic targets for preventing

diabetic complications.

The current study had several limitations. Although our LASSO and factor analysis indicated

that CGM-derived measures were strong predictors of %NC, this does not mean that other

clinical parameters, such as lipids and blood pressure, are irrelevant in T2DM complications.

Our study specifically focused on characterizing glucose dynamics, and we analyzed

individuals with well-controlled serum cholesterol and blood pressure to reduce confounding

effects. While we anticipate that inclusion of a more diverse population would not alter our

primary findings regarding glucose dynamics, it is likely that a broader data set would reveal

additional predictive contributions from lipid and blood pressure parameters. In our analysis

of demographic factors, we found that age and gender had minimal influence on glucose

dynamics characteristics (**Fig. S8A, B**), suggesting that our findings regarding the

relationship between glucose dynamics and coronary risk are robust across different

demographic groups within our dataset. Future studies involving larger and more diverse

populations would be valuable to comprehensively elucidate the potential influence of age,

gender, and other demographic factors on glucose dynamics characteristics and their

relationship to cardiovascular risk.

A previous study identified components of interday variability and hypoglycemia in CGM-

derived indices (Augstein et al., 2015) that were not observed in our analysis. This

discrepancy may be due to the relatively small number of T2DM individuals in our study. We

acknowledge that factor analyses of data from longer measurement periods, including more

patients with T1DM and T2DM, could potentially yield different results. However, it is

noteworthy that our analysis of longer-term CGM data sets from Japanese and American

populations confirmed the existence of the same three factors - mean, variance, and

autocorrelation - in glucose dynamics. Moreover, even with the short measurement period,

CGM-derived indices reflecting these three factors demonstrated superior predictive accuracy

for %NC compared to traditional indices such as FBG, HbA1c, and PG120, underscoring the

potential utility of CGM. Although time in range (TIR) was not included in the main analyses

due to the relatively small number of T2DM patients and the predominance of participants

with TIR >70%, our results demonstrate that CGM-derived indices outperformed

conventional markers such as FBG, HbA1c, and PG120 in predicting %NC. Furthermore,

multiple regression analysis between factor scores and TIR revealed that only factor 1 (mean)

and factor 2 (variance) were significantly associated with TIR (**Fig. S8C, D**). This finding

confirms the presence of three distinct components in glucose dynamics and highlights the

added value of examining AC_Var as an independent glycemic feature beyond conventional

CGM-derived measures. Some indices, such as HBGI, showed variation in classification

across datasets, with some populations showing higher factor loadings in the "mean"

component and others in the "variance" component. This variation occurs because HBGI

calculations depend on the number of glucose readings above a threshold. In populations

where mean glucose levels are predominantly below this threshold, the HBGI is more

sensitive to glucose variability (**Fig. S5A**). Conversely, in populations with a wider range of

mean glucose levels, the HBGI correlates more strongly with mean glucose levels (**Fig. 3A**).

Despite these differences, our validation analyses confirm that CGM-derived indices

consistently cluster into three components: mean, variance, and autocorrelation.

We acknowledge the potential concern of multiple testing in our study. However, even after

adjusting for multiple comparisons, the CGM-derived indices retained significant correlations

with %NC (**Fig. 1E**). The consistency of our findings across different analytical approaches

(Lasso, PLS, and factor analysis) and different data sets further supports the robustness of our

conclusions regarding the characteristics of glucose dynamics. While we used methods that

assume linearity, such as LASSO, we also examined nonlinear relationships using

Spearman's correlation for index relationships and factor loadings with %NC and found

significant associations. We acknowledge that some significant correlations appear to be

relatively small. However, these findings, combined with our predictive models showing

improved accuracy using CGM compared to traditional diabetes diagnostic indices, and the

theoretical framework showing that conventional markers only consider the "mean"

component of glucose dynamics, can demonstrate the clinical significance. Although our

analysis included four datasets with a total of 270 individuals, and our sample size of 53 met

the required threshold based on power calculations with a type I error of 0.05, a power of 0.8,

and an expected correlation coefficient of 0.4, we acknowledge that the sample size may still

be considered relatively small for a comprehensive assessment of these relationships. To

further validate these findings, larger prospective studies with diverse populations are needed.

While CGM-derived indices such as AC_Var and ADRR hold promise for CAD risk

assessment, their complexity may present challenges for routine clinical implementation. To

improve usability, we have developed a web-based calculator that automates these

calculations. However, defining clinically relevant thresholds and reference ranges requires

further validation in larger cohorts.

In conclusion, glucose dynamics have three components: mean, variance, and autocorrelation.

These three components are associated with coronary plaque vulnerability. CGM-derived

indices reflecting these three components can be valuable predictive tools for T2DM

complications, compared to conventional diabetes diagnostic markers reflecting only the

mean component. This new predictive model has the potential to improve the diagnosis and

management of diabetes worldwide. To facilitate this CGM-derived prediction, we created a

web application that performs a multiple regression model with these three components as

input variables (https://cgmregressionapp2.streamlit.app/).

METHODS

Key Resources Table				
Reagent type	Designation	Source or reference	Identifiers	Additional information
software, algorithm	SciPy v1.10.1	(Virtanen et al., 2020)		
software, algorithm	scikit-learn v1.0.2	https://scikit-learn.org/stable/		

Study design and population

This retrospective observational study was approved by the ethics committee of Kobe University Graduate School of Medicine (UMIN000018326; Kobe, Japan), as described previously (Otowa-Suematsu et al., 2018). The study included 53 participants who underwent a 75-g oral glucose tolerance test (OGTT), continuous glucose monitoring (CGM) with the use of an iPro2 CGM system (Medtronic, Northridge, CA, USA), and percutaneous coronary intervention (PCI). During PCI, VH-IVUS was carried out to assess the plaque components. All individuals underwent CGM for at least three consecutive days within the seven-day period prior to the PCI procedure.

Among the 53 participants, eight, 16, and 29 individuals were categorized as having normal glucose tolerance (NGT), impaired glucose tolerance (IGT) and T2DM, respectively. Of note, with a type I error of 0.05, a power of 0.8, and an expected correlation coefficient of 0.4, a sample size of 47 was required to detect a significant difference from zero in the correlation coefficient. T2DM was defined as HbA1c $\geq 6.5\%$, fasting plasma glucose (FPG) ≥ 126 mg/dL or 2-h plasma glucose during a 75-g OGTT (PG120) ≥ 200 mg/dL. IGT was defined as HbA1c 6.0–6.4%, FPG 110–125 mg/dL or PG120 140–199 mg/dL. NGT was defined as values below all prediabetes thresholds (HbA1c $< 6.0\%$, FPG < 110 mg/dL and PG120 < 140 mg/dL).

Detailed participant characteristics have been reported in the previous study (Otowa-Suematsu et al., 2018). Briefly, participants aged 20–80 years with LDL-C levels < 120 mg/dL under statin administration or < 100 mg/dL under other treatments for dyslipidemia, including lifestyle intervention, were included in this study. Participants with acute coronary syndrome, unsuitable anatomy for VH-IVUS, poor imaging by VH-IVUS, hemodialysis, inflammatory disease, shock, low cardiac output, or concurrent malignant disease were excluded from this study.

To validate the findings on glucose dynamics, a previously reported data set from Japan (Sugimoto et al., 2025) was analyzed. The study was conducted in accordance with the Declaration of Helsinki and approved by the Kobe University Hospital Ethics Committee (Approval No. 1834). Briefly, individuals aged ≥ 20 years with no previous diagnosis of diabetes were recruited from Kobe University Hospital between January 2016 and March 2018. Exclusion criteria included the use of medications that affect glucose metabolism (e.g., steroids, β-blockers), psychiatric disorders, pregnancy or lactation, and ineligibility as determined by treating physicians. Participants wore a CGM device (iPro; Medtronic, USA). The study included 52 individuals with NGT, nine with IGT, and three with T2DM.

Further validation of the glucose dynamics findings was performed using a previously reported CGM dataset (Dexcom G4 CGM System; Dexcom, Fort Lauderdale, FL, USA) obtained in the United States (Hall et al., 2018). The study included 53 individuals aged 30–70 years with a BMI of 23–40 kg/m^2. Exclusion criteria included prior diagnosis of diabetes, major organ diseases, uncontrolled hypertension, malignancy, chronic inflammatory conditions, use of any medications known to alter blood glucose or insulin sensitivity, haematocrit < 30, creatinine above the normal range, and ALT levels more than twice the upper limit of normal.

In addition, the present study analyzed a previously reported CGM (FreeStyle Libre H, Abbott Diabetes Care, Witney, UK) dataset from China (Zhao et al., 2023). Participants were recruited from the DiaDRIL registry at Shanghai East Hospital (September 2019 to March 2021) and Shanghai Fourth People's Hospital (June 2021 to November 2021). Inclusion criteria for this dataset were: diagnosis of diabetes according to the 1999 World Health Organization (WHO) criteria, age 18 years or older, willingness to provide informed consent, and CGM recording for at least three days. Exclusion criteria included reported alcohol or drug abuse, inability to comply with study protocols, or inability to participate as determined by the investigators. We extracted and analyzed glucose profile characteristics from the first three days of CGM data for each participant. 100 individuals with T2DM were analyzed in this study. Our primary objective was to explore the relationship between the CGM-derived indices and the presence of diabetic macrovascular complications. Of note, as the original reports document, the validation datasets did not specify explicit cutoffs for blood pressure or cholesterol. Consequently, they included participants with suboptimal control of these parameters.

The data collection protocol for the Chinese dataset was previously documented (Zhao et al., 2023). Briefly, trained research staff used standardized questionnaires to collect demographic

and clinical information, including diabetes diagnosis, treatment history, comorbidities, and medication use. CGM records interstitial glucose levels at 15-minute intervals for up to 14 days. Laboratory measurements, including metabolic panels, lipid profiles, and renal function tests, were obtained within six months of CGM placement. While previous studies have linked necrotic core to macrovascular events, we acknowledge the limitations of the cardiovascular outcomes in the Chinese data set. These outcomes were extracted from medical records rather than from standardized diagnostic procedures or imaging studies.

Calculation of clinical indices

CGM-derived indices:

The CGM data collected in this study were obtained at different sampling frequencies in the different data sets. For the two Japanese datasets and the American dataset, glucose measurements were collected at 5-minute intervals. In contrast, the Chinese dataset collected glucose measurements at 15-minute intervals.

Fourteen CGM-derived indices were evaluated in this study: twelve well-established CGM-derived indices (Hill et al., 2011) and two indices (AC_Mean and AC_Var) that have been reported to capture glucose handling capacity independently of the aforementioned twelve indices (Sugimoto et al., 2025). For the datasets with 5-minute sampling intervals (the two Japanese datasets and the American dataset), AC_Mean and AC_Var were calculated as the mean and variance of the autocorrelation functions at lags 1-30 of the glucose levels, respectively. Of note, autocorrelation refers to the relationship between a variable and its past values over time. In the context of glucose dynamics, it reflects how current glucose levels are influenced by past levels, capturing patterns such as sustained hyperglycemia or recurrent fluctuations. For example, if an individual experiences sustained high glucose levels after a meal, the strong correlation between successive glucose readings indicates high autocorrelation. For the data set with 15-minute sampling intervals (the Chinese data set), AC_Mean and AC_Var were calculated as the mean and variance of the autocorrelation functions at lags 1-10 of the glucose levels, respectively. These calculations correspond to a time window of 150 minutes. CGM_Mean and CGM_Std indicate the mean value and standard deviation of glucose levels measured by CGM, respectively. CONGA, LI, JINDEX, HBGI, GRADE, MODD, MAGE, ADRR, MVALUE, and MAG were calculated using EasyGV software (Hill et al., 2011). The calculating formulae of these indices are shown in **Table S1**.

In the two Japanese datasets and the American dataset, there was a relatively small proportion of T2DM patients and a substantial number of individuals with time in range (TIR) (Battelino et al., 2023; Larkin et al., 2019) values (glucose levels between 70 and 180 mg/dL) greater

536 than 70% (Hall et al., 2018; Otowa-Suematsu et al., 2018; Sugimoto et al., 2025). Given this
537 distribution, we decided not to include TIR in our primary analyses. However, given the
538 potential interest of this metric, we provided correlations between TIR and other indices as
539 supplementary information.
540
OGTT-derived indices:
Three OGTT-derived indices were calculated as previously described (Matsuda and
DeFronzo, 1999; Otowa-Suematsu et al., 2018). The insulinogenic index (I.I.) indicates
insulin secretion and is calculated from the ratio of the increment of immunoreactive insulin
(IRI) to that of plasma glucose at 30 min after onset of the OGTT. The composite index
indicates insulin sensitivity, which can be calculated from fasting plasma glucose, fasting IRI,
mean blood glucose levels, and mean serum IRI concentrations during the OGTT. The oral
disposition index (Oral DI) was calculated from the product of the composite index and the
ratio of the area under the insulin concentration curve from 0 to 120 minutes to that for
plasma glucose from 0 to 120 minutes, without using the data measured at 90 min, in the
OGTT.
552
VH-IVUS-derived index:
VH-IVUS was carried out using the Eagle Eye Platinum 3.5-Fr 20-MHz catheter (Volcano,
Rancho Cordova, CA, USA), as previously described (Otowa-Suematsu et al., 2018). The
intraclass correlation coefficients for interobserver and intraobserver reliability of external
elastic membrane volume were 0.95 and 0.97, respectively (Otowa-Suematsu et al., 2018),
indicating high reproducibility. The VH-IVUS categorized plaque into four components:
fibrous, fibrofatty, necrotic core, and dense calcium. Following the previous study, our
investigation focused specifically on the ratio of necrotic core to total plaque volume (%NC),
a widely used parameter of plaque vulnerability. For patients with multiple plaques, the
mean %NC was calculated.
563
Prediction models and statistical analyses
We modelled %NC using multiple linear regression, LASSO regression, and PLS regression.
The input variables in these models included the following 26 variables: BMI, SBP, DBP,
TGs, LDL-C, HDL-C, FBG, HbA1c, PG120, I.I., composite index, oral DI, CGM_Mean,
CGM_Std, CONGA, LI, JINDEX, HBGI, GRADE, MODD, MAGE, ADRR, MVALUE,
MAG, AC_Mean, and AC_Var. In conducting these models, z-score normalization on each
input variable was performed.
571
572 The predictive performance of multiple linear regression was evaluated by the coefficient of
573 determination (R^2), the adjusted coefficient of determination (Adj R^2), or AIC. The

multicollinearity of the input variables was estimated by VIF. LASSO regression is a kind of linear regression with L1 regularization (Tibshirani, 1996). The optimal regularization coefficient, lambda, was based on leave-one-out cross-validation and mean-squared error (MSE). The importance of the input variables in predicting %NC was evaluated by the VIP scores (Wold et al., 2001) that were generated from PLS regression. LASSO regression was chosen for its ability to perform feature selection by identifying the most relevant predictors. Unlike Ridge regression, which simply shrinks coefficients toward zero without reaching exactly zero, LASSO produces sparse models, which is consistent with our goal of identifying the most critical features of glucose dynamics associated with coronary plaque vulnerability. In addition, we implemented PLS regression as a complementary approach due to its effectiveness in dealing with multicollinearity, which was particularly relevant given the high correlation among several CGM-derived measures. These models were conducted using scikit-learn v1.0.2, a Python-based toolkit (https://scikit-learn.org/stable/).

Relationships among indices were also evaluated using Spearman's correlation coefficients (r), and the correlation coefficients were reported with 95% CIs through bootstrap resampling. The number of resamples performed to form the distribution was set at 10000. Benjamini–Hochberg's multiple comparison test was also performed with a significance threshold of $Q < 0.05$.

Factor analysis and hierarchical clustering analysis
The intercorrelations of the clinical parameters and their associations with %NC were assessed using exploratory factor analyses and hierarchical clustering analyses. We followed the previously reported approach (Cappelleri et al., 2000; Guo et al., 2020; Lakka et al., 2002; Oh et al., 2004) with some modifications in conducting our exploratory factor analyses. BIC and MAP methods were used to determine the number of underlying factors. Variables with absolute factor loadings of ≥ 0.30 were used in interpretation. To improve the interpretation, orthogonal (varimax) rotation was used. To evaluate the applicability of the factor analysis, KMO and Bartlett's spherical test were performed. To evaluate the internal consistency of each factor, Cronbach's α and item–total correlations were calculated. The association of the factor scores with %NC was assessed using Spearman's correlation.

Hierarchical clustering analysis was conducted using a method that combines a Euclidean distance measure and Ward linkage. I.I., composite index, oral DI, and AC_Mean were inverted negatively so that the value of indices increased in individuals with abnormalities. The quality of the hierarchical clustering analysis was evaluated based on silhouette analysis (Rousseeuw, 1987). These analyses were performed after Z score normalization using scikit-learn v1.0.2, a Python-based toolkit (https://scikit-learn.org/stable/).

Mathematical models used for simulating the characteristics of glucose dynamics

In simulating the characteristics of glucose dynamics, we used a simple and stable model (De Gaetano and Arino, 2000), which can be written as follows:

$$\frac{dG}{dt} = -k_{\text{glu}}G - k_{\text{sen}}IG + k_{\text{pro}} + f$$

$$\frac{dI}{dt} = \frac{k_{\text{sec}}}{k_{\text{tim}}} \int_{t-k_{\text{tim}}}^{t} G\,ds - k_{\text{cle}}I$$

where the variables G and I denote blood glucose and insulin concentrations, respectively. We simulated 240-minute profiles of G, and calculated the mean, Std, and AC_Var of G. The parameters were changed within the range participants could take (De Gaetano and Arino, 2000). Five mg/dL/min glucose was applied for 10 minutes at 30 minutes as the external input of glucose f. The simulations were conducted using SciPy v1.10.1 (Virtanen et al., 2020).

Characterization of glucose patterns during the OGTT

The relation between the three components of glucose dynamics and the characteristics of previously reported glucose patterns during the OGTT (Hulman et al., 2018) was investigated. In the study, 5861 individuals without diabetes in Denmark underwent the OGTT with measurements of glucose levels at three time points (0, 30, and 120 min), and four distinct glucose patterns associated with long-term outcomes including diabetes onset, cardiovascular disease, and all-cause mortality rate were identified. For the calculation of mean, Std, and AC_Var of glucose levels, each time point was linearly interpolated. AC_Var was calculated using the autocorrelation function at lags 1–20 because glucose data were available only for 2 hours after the OGTT.

Acknowledgments

Author Contributions. H.S. analyzed the data. H.S., K.H., T.Y., N.O.S., Y.H., H.O., K.H., K.S., W.O., and S.K. wrote the manuscript. W.O. and S.K. supervised the study.

Conflict of Interest. The authors have no conflicts of interest to declare.

Funding and Assistance. This study was supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI (JP21H04759), CREST, the Japan Science and Technology Agency (JST) (JPMJCR2123), The Uehara Memorial Foundation, and The Takeda Science Foundation.

Data and code availability

We used only previously published data sets. The code that calculates AC_Mean and AC_Var and that performs regression analysis with CGM-derived indices as input variables

647   are available from the repository (https://github.com/HikaruSugimoto/CGM_regression_app)

648   and the web application (https://cgmregressionapp2.streamlit.app/).

     Figure Legends

Figure 1. Relations among clinical measures.

(A) Multiple regression analysis between %NC and CGM_Mean, CGM_Std, and AC_Var.

Scatter plots for predicted %NC versus measured %NC (the left). Each point corresponds to

the values for a single individual. Bars represent the 95% confidence intervals (CIs) of the

regression coefficients (the right).

(B) Multiple regression analysis between %NC and FBG, HbA1c, and PG120.

(C) Linear regression analysis between %NC and the Framingham Risk Score for Hard

Coronary Heart Disease.

(D) A spring layout of the correlation network involving %NC (black), 14 CGM-derived

indices (red), 3 blood glucose level-related indices (magenta), 3 insulin sensitivity or

secretion-related indices (blue), and 6 other clinical indices (green) obtained from a single

blood test or physical measurement. Connections denote relationships with $Q < 0.05$. The

width of the edges is proportional to the corresponding correlation coefficient.

(E) The absolute values of Spearman's correlation coefficients between clinical parameters

and %NC. Bars represent the 95% CIs.

(F) Scatter plots for AC_Var versus CGM_Mean (the left), and AC_Var versus CGM_Std
(the right). Each point corresponds to a single individual's values. Individuals are colored
based on their %NC value. R is Spearman's correlation coefficient, and the value in
parentheses is 95% CI.
(G, H) VIF of each variable remaining after removing the variables with the highest VIF one
by one until the VIFs of all variables are less than 10. The input variables include the
following 26 variables: BMI, SBP, DBP, TG, LDL-C, HDL-C, FBG, HbA1c, PG120, I.I.,
composite index, Oral DI, CGM_Mean, CGM_Std, CONGA, LI, JINDEX, HBGI, GRADE,
MODD, MAGE, ADRR, MVALUE, MAG, AC_Mean, and AC_Var. The input variables of
(H) included only the CGM-derived measures from the above 26 indices.
(I) Scatter plot of the VIF of the indices measured in this study (VIFt) versus that of indices
measured in a previous study (VIFp1) (Sugimoto et al., 2025). R is Spearman's correlation
coefficient, and the value in parentheses is the 95% CI.
(J) Scatter plot of the VIF of the indices measured in this study (VIFt) versus that of the
indices measured in a previous study (Hall et al., 2018) (VIFp2).
680

Figure 2. LASSO and PLS regression analyses for predicting %NC.

(A) Relationship between regularization coefficients (lambda) and the MSE based on the

leave-one-out cross-validation in predicting %NC. Dotted vertical line indicates the optimal

lambda, which provides the lowest MSE. The optimal lambda was 0.849.

(B) LASSO regularization paths along the lambda in predicting %NC. Cyan, magenta, and

gray lines indicate the estimated coefficients of AC_Mean, AC_Var, and the other input

variables, respectively. Dotted vertical line indicates the optimal lambda.

(C) Estimated coefficients with the optimal lambda. Only variables with non-zero coefficients

are shown. Input variables include the following 21 variables: BMI, FBG, HbA1c, PG120,

I.I., Composite index, Oral DI, CGM_Mean, CGM_Std, CONGA, LI, JINDEX, HBGI,

GRADE, MODD, MAGE, ADRR, MVALUE, MAG, AC_Mean, and AC_Var.

(D) VIP generated from the PLS regression predicting %NC. Variables with a VIP $\geq$ 1 (the

dotted line) were considered to significantly contribute to the prediction.

Figure 3. Factor analysis of the clinical parameters.

(A) Factor analysis after orthogonal rotation. The values and colors are based on the factor loadings. The columns represent each factor. The rows represent input indices.

(B) Cronbach's α for each factor. Bars represent the 95% CI.

(C) Scatter plots and fitted linear regression lines for factor scores versus %NC. Each point corresponds to the values for a single individual. R is Spearman's correlation coefficient, and the value in parentheses is the 95% CI.

Figure 4. Overview of the three components of glucose dynamics.

(A) 240 min simulated glucose concentration. The colors of the lines are based on the mean

(Mean), Std, and AC_Var of the simulated blood glucose. Red and gray dotted horizontal

lines indicate the minimum and maximum blood glucose values, respectively.

(B) Previously reported patterns of blood glucose during the OGTT (Hulman et al., 2018).

Green, class 1; light blue, class 2; dark blue, class 3; red, class 4.

(C) Mean, Std, and AC_Var of the glucose during the OGTT. Colors are based on the classes

shown in Figure 4B.

References

Augstein P, Heinke P, Vogt L, Vogt R, Rackow C, Kohnert K-D, Salzsieder E. 2015. Q-Score: development of a new metric for continuous glucose monitoring that enables stratification of antihyperglycaemic therapies. *BMC Endocr Disord* **15**:22.

Battelino T, Alexander CM, Amiel SA, Arreaza-Rubin G, Beck RW, Bergenstal RM, Buckingham BA, Carroll J, Ceriello A, Chow E, Choudhary P, Close K, Danne T, Dutta S, Gabbay R, Garg S, Heverly J, Hirsch IB, Kader T, Kenney J, Kovatchev B, Laffel L, Maahs D, Mathieu C, Mauricio D, Nimri R, Nishimura R, Scharf M, Del Prato S, Renard E, Rosenstock J, Saboo B, Ueki K, Umpierrez GE, Weinzimer SA, Phillip M. 2023. Continuous glucose monitoring and metrics for clinical trials: an international consensus statement. *Lancet Diabetes Endocrinol* **11**:42–57.

Bax JJ, Young LH, Frye RL, Bonow RO, Steinberg HO, Barrett EJ, ADA. 2007. Screening for coronary artery disease in patients with diabetes. *Diabetes Care* **30**:2729–2736.

Cai J, Yang Q, Lu J, Shen Y, Wang C, Chen L, Zhang L, Lu W, Zhu W, Xia T, Zhou J. 2022. Impact of the complexity of glucose time series on all-cause mortality in patients with type 2 diabetes. *J Clin Endocrinol Metab*. doi:10.1210/clinem/dgac692

Cappelleri JC, Gerber RA, Kourides IA, Gelfand RA. 2000. Development and factor analysis of a questionnaire to measure patient satisfaction with injected and inhaled insulin for type 1 diabetes. *Diabetes Care* **23**:1799–1803.

Ceriello A, Esposito K, Piconi L, Ihnat MA, Thorpe JE, Testa R, Boemi M, Giugliano D. 2008. Oscillating glucose is more deleterious to endothelial function and oxidative stress than mean glucose in normal and type 2 diabetic patients. *Diabetes* **57**:1349–1354.

Cobelli C, Facchinetti A. 2018. Yet Another Glucose Variability Index: Time for a Paradigm Change? *Diabetes Technol Ther* **20**:1–3.

Dalla Man C, Rizza RA, Cobelli C. 2007. Meal simulation model of the glucose-insulin system. *IEEE Trans Biomed Eng* **54**:1740–1749.

De Gaetano A, Arino O. 2000. Mathematical modelling of the intravenous glucose tolerance test. *J Math Biol* **40**:136–168.

Fabris C, Facchinetti A, Fico G, Sambo F, Arredondo MT, Cobelli C, MOSAIC EU Project Consortium. 2015. Parsimonious Description of Glucose Variability in Type 2 Diabetes by Sparse Principal Component Analysis. *J Diabetes Sci Technol* **10**:119–124.

Fabris C, Facchinetti A, Sparacino G, Zanon M, Guerra S, Maran A, Cobelli C. 2014. Glucose variability indices in type 1 diabetes: parsimonious set of indices revealed by sparse principal component analysis. *Diabetes Technol Ther* **16**:644–652.

Fiarni C, Sipayung EM, Maemunah S. 2019. Analysis and Prediction of Diabetes Complication Disease using Data Mining Algorithm. *Procedia Comput Sci* **161**:449–457.

Gerbaud E, Darier R, Montaudon M, Beauvieux M-C, Coffin-Boutreux C, Coste P, Douard H, Ouattara A, Catargi B. 2019. Glycemic Variability Is a Powerful Independent Predictive Factor of Midterm Major Adverse Cardiac Events in Patients With Diabetes With Acute Coronary Syndrome. *Diabetes Care* **42**:674–681.

Gorst C, Kwok CS, Aslam S, Buchan I, Kontopantelis E, Myint PK, Heatlie G, Loke Y, Rutter MK, Mamas MA. 2015. Long-term Glycemic Variability and Risk of Adverse Outcomes: A Systematic Review and Meta-analysis. *Diabetes Care* **38**:2354–2369.

Guo W, Zhou Q, Jia Y, Xu J. 2020. Cluster and Factor Analysis of Elements in Serum and Urine of Diabetic Patients with Peripheral Neuropathy and Healthy People. *Biol Trace Elem Res* **194**:48–57.

Hall H, Perelman D, Breschi A, Limcaoco P, Kellogg R, McLaughlin T, Snyder M. 2018. Glucotypes reveal new patterns of glucose dysregulation. *PLoS Biol* **16**:e2005143.

Hill NR, Oliver NS, Choudhary P, Levy JC, Hindmarsh P, Matthews DR. 2011. Normal reference range for mean tissue glucose and glycemic variability derived from continuous glucose monitoring for subjects without diabetes in different ethnic groups. *Diabetes Technol Ther* **13**:921–928.

Hulman A, Vistisen D, Glümer C, Bergman M, Witte DR, Færch K. 2018. Glucose patterns during an oral glucose tolerance test and associations with future diabetes, cardiovascular disease and all-cause mortality rate. *Diabetologia* **61**:101–107.

Keshet A, Shilo S, Godneva A, Talmor-Barkan Y, Aviv Y, Segal E, Rossman H. 2023. CGMap: Characterizing continuous glucose monitor data in thousands of non-diabetic individuals. *Cell Metab* **35**:758-769.e3.

Lakka H-M, Laaksonen DE, Lakka TA, Niskanen LK, Kumpusalo E, Tuomilehto J, Salonen JT. 2002. The metabolic syndrome and total and cardiovascular disease mortality in middle-aged men. *JAMA* **288**:2709–2716.

Larkin ME, Nathan DM, Bebu I, Krause-Steinrauf H, Herman WH, Higgins JM, Tiktin M, Cohen RM, Lund C, Bergenstal RM, Johnson ML, Arends V, GRADE Research Group. 2019. Rationale and Design for a GRADE Substudy of Continuous Glucose Monitoring. *Diabetes Technol Ther* **21**:682–690.

Matsuda M, DeFronzo RA. 1999. Insulin sensitivity indices obtained from oral glucose tolerance testing: comparison with the euglycemic insulin clamp. *Diabetes Care* **22**:1462–1470.

Monnier L, Colette C, Owens DR. 2008. Glycemic variability: the third component of the dysglycemia in diabetes. Is it important? How to measure it? *J Diabetes Sci Technol* **2**:1094–1100.

Oh J-Y, Hong YS, Sung Y-A, Barrett-Connor E. 2004. Prevalence and factor analysis of
metabolic syndrome in an urban Korean population. *Diabetes Care* **27**:2027–2032.
Okada K, Hibi K, Gohbara M, Kataoka S, Takano K, Akiyama E, Matsuzawa Y, Saka K,
Maejima N, Endo M, Iwahashi N, Tsukahara K, Kosuge M, Ebina T, Fitzgerald PJ,
Honda Y, Umemura S, Kimura K. 2015. Association between blood glucose
variability and coronary plaque instability in patients with acute coronary syndromes.
Cardiovasc Diabetol **14**:111.
Otowa-Suematsu N, Sakaguchi K, Komada H, Nakamura T, Sou A, Hirota Y, Kuroda M,
Shinke T, Hirata K-I, Ogawa W. 2018. Comparison of the relationship between
multiple parameters of glycemic variability and coronary plaque vulnerability
assessed by virtual histology-intravascular ultrasound. *J Diabetes Investig* **9**:610–615.
Pei X, Qi D, Liu J, Si H, Huang S, Zou S, Lu D, Li Z. 2023. Screening marker genes of type
2 diabetes mellitus in mouse lacrimal gland by LASSO regression. *Sci Rep* **13**:6862.
Psoma O, Makris M, Tselepis A, Tsimihodimos V. 2022. Short-term Glycemic Variability
and Its Association With Macrovascular and Microvascular Complications in Patients
With Diabetes. *J Diabetes Sci Technol* 19322968221146810.
Randrianarisoa E, Lehn-Stefan A, Hieronimus A, Wagner R, Maucher J, Rittig K,
Balletshofer B, Birkenfeld AL, Peter A, Stefan N, Häring H-U, Fritsche A, Heni M.
2020. Reduced insulin clearance is linked to subclinical atherosclerosis in individuals
at risk for type 2 diabetes mellitus. *Sci Rep* **10**:22453.
Ravaut M, Sadeghi H, Leung KK, Volkovs M, Kornas K, Harish V, Watson T, Lewis GF,
Weisman A, Poutanen T, Rosella L. 2021. Predicting adverse outcomes due to
diabetes complications with machine learning using administrative health data. *NPJ*
Digit Med **4**:24.
Rousseeuw PJ. 1987. Silhouettes: A graphical aid to the interpretation and validation of
cluster analysis. *J Comput Appl Math* **20**:53–65.
Selvin E, Steffes MW, Zhu H, Matsushita K, Wagenknecht L, Pankow J, Coresh J, Brancati
FL. 2010. Glycated hemoglobin, diabetes, and cardiovascular risk in nondiabetic
adults. *N Engl J Med* **362**:800–811.
Service FJ. 2013. Glucose variability. *Diabetes* **62**:1398–1404.
Su G, Mi S, Tao H, Li Z, Yang H, Zheng H, Zhou Y, Ma C. 2011. Association of glycemic
variability and the presence and severity of coronary artery disease in patients with
type 2 diabetes. *Cardiovasc Diabetol* **10**:19.
Sugimoto H, Hironaka K-I, Nakamura T, Yamada T, Miura H, Otowa-Suematsu N, Fujii M,
Hirota Y, Sakaguchi K, Ogawa W, Kuroda S. 2025. Improved detection of decreased
glucose handling capacities via continuous glucose monitoring-derived indices.
Commun Med (Lond) **5**:103.

Tang X, Li S, Wang Y, Wang M, Yin Q, Mu P, Lin S, Qian X, Ye X, Chen Y. 2016. Glycemic variability evaluated by continuous glucose monitoring system is associated with the 10-y cardiovascular risk of diabetic patients with well-controlled HbA1c. *Clin Chim Acta* **461**:146–150.

Tibshirani R. 1996. Regression shrinkage and selection via the lasso. *J R Stat Soc* **58**:267–288.

Utzschneider KM, Prigeon RL, Faulenbach MV, Tong J, Carr DB, Boyko EJ, Leonetti DL, McNeely MJ, Fujimoto WY, Kahn SE. 2009. Oral disposition index predicts the development of future diabetes above and beyond fasting and 2-h glucose levels. *Diabetes Care* **32**:335–341.

Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, Burovski E, Peterson P, Weckesser W, Bright J, van der Walt SJ, Brett M, Wilson J, Millman KJ, Mayorov N, Nelson ARJ, Jones E, Kern R, Larson E, Carey CJ, Polat İ, Feng Y, Moore EW, VanderPlas J, Laxalde D, Perktold J, Cimrman R, Henriksen I, Quintero EA, Harris CR, Archibald AM, Ribeiro AH, Pedregosa F, van Mulbregt P, SciPy 1.0 Contributors. 2020. SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat Methods* **17**:261–272.

Wang C, Kong H, Guan Y, Yang J, Gu J, Yang S, Xu G. 2005. Plasma phospholipid metabolic profiling and biomarkers of type 2 diabetes mellitus based on high-performance liquid chromatography/electrospray mass spectrometry and multivariate statistical analysis. *Anal Chem* **77**:4108–4116.

Wei H, Sun J, Shan W, Xiao W, Wang B, Ma X, Hu W, Wang X, Xia Y. 2022. Environmental chemical exposure dynamics and machine learning-based prediction of diabetes mellitus. *Sci Total Environ* **806**:150674.

Wilson PW, D'Agostino RB, Levy D, Belanger AM, Silbershatz H, Kannel WB. 1998. Prediction of coronary heart disease using risk factor categories. *Circulation* **97**:1837–1847.

Wold S, Sjöström M, Eriksson L. 2001. PLS-regression: a basic tool of chemometrics. *Chemometrics Intellig Lab Syst* **58**:109–130.

Young LH, Wackers FJT, Chyun DA, Davey JA, Barrett EJ, Taillefer R, Heller GV, Iskandrian AE, Wittlin SD, Filipchuk N, Ratner RE, Inzucchi SE, DIAD Investigators. 2009. Cardiac outcomes after screening for asymptomatic coronary artery disease in patients with type 2 diabetes: the DIAD study: a randomized controlled trial. *JAMA* **301**:1547–1555.

Zhao Q, Zhu J, Shen X, Lin C, Zhang Y, Liang Y, Cao B, Li J, Liu X, Rao W, Wang C. 2023. Chinese diabetes datasets for data-driven machine learning. *Sci Data* **10**:35.

859   Zhou JJ, Schwenke DC, Bahn G, Reaven P, VADT Investigators. 2018. Glycemic Variation

860   and Cardiovascular Risk in the Veterans Affairs Diabetes Trial. *Diabetes Care*

861   **41**:2187–2194.