1  **Title:** Exposure to heavy metals *in utero* and autism spectrum disorder at age 3: A meta-
2  analysis of two cohorts with enriched likelihood of autism
3

4  **Authors:** John F. Dou[1], Rebecca J. Schmidt[2], Heather E. Volk[3], Manon M. Nitta[1], Jason I.
5  Feinberg[3], Craig J. Newschaffer[4], Lisa A. Croen[5], Irva Hertz-Picciotto[2], M. Daniele Fallin[6], Kelly
6  M. Bakulski[1]
7
**Affiliations:**
[1]University of Michigan, Ann Arbor, Michigan, USA
[2]University of California Davis, Davis, California, USA
[3]Johns Hopkins University, Baltimore, Maryland, USA
[4]Penn State University, State College, PA, USA
[5]Division of Research, Kaiser Permanente Northern California, Oakland, CA, USA
[6]Rollins School of Public Health, Emory University, Atlanta, GA, USA
15
16
17  **Corresponding author:**
18  Kelly M. Bakulski, email: bakulski@umich.edu, phone: (734) 615 5899, address: M5511 SPH2,
19  1415 Washington Heights Ann Arbor, MI 48109

21  **Conflicts of Interest**: The authors declare they have no conflicts of interest related to this work
22  to disclose.
23
**Funding Acknowledgements:**
Funding for the EARLI study was provided by the National Institutes of Health (R01ES016443,
R24ES030893) and Autism Speaks (003953)**.** Funding for the MARBLES study was provided by
the National Institutes of Health (R01ES020392, R01ES028089, R/U24ES028533, and
P01ES011269) and the United States Environmental Protection Agency Science to Achieve
Results program (#RD-83329201). Funding for metals measures and this work was supported
by the National Institutes of Health **(**R01ES025531). The content is solely the responsibility of
the authors and does not necessarily represent the official views of the National Institutes of
Health.

**Abstract**

**Background:** Autism spectrum disorder (ASD) is a prevalent and heterogeneous
neurodevelopmental disorder. Risk is attributed to genetic and prenatal environmental factors,
though the environmental agents are incompletely characterized.
37
**Methods:** In Early Autism Risk Longitudinal Investigation (EARLI) and Markers of Autism Risk
in Babies Learning Early Signs (MARBLES), two pregnancy cohorts with high likelihood of ASD,
maternal urinary metals concentrations at two time points during pregnancy were measured
using inductively coupled plasma mass spectrometry. At age three, clinicians assessed ASD
with DSM-5 criteria. Using multivariable log binomial regression, we examined each metal for
association with ASD status, adjusting for gestational age at urine sampling, child sex, maternal
age, and maternal education, and meta-analyzed across the two cohorts.
45
**Results:** In EARLI (n=170) 17.6% of children were diagnosed with ASD, and an additional
43.5% were classified as having other non-neurotypical development (Non-TD). In MARBLES
(n=156), 22.7% were diagnosed with ASD, while an additional 11.5% had Non-TD. In earlier
pregnancy metals measures, having cadmium concentration over the level of detection was
associated with 1.78 (1.19, 2.67) times higher risk of ASD, and 1.43 (1.06, 1.92) times higher
risk of Non-TD. A doubling of early pregnancy cesium concentration was marginally associated
with 1.81 (0.95, 3.42) times higher risk of ASD, and 1.58 (0.95, 2.63) times higher risk of Non-
TD.
54
55   **Conclusion:** Exposure *in utero* to elevated levels of cadmium and cesium, as measured in
56   maternal urine collected during pregnancy, was associated with increased risk of developing
57   ASD.
58

59   **Keywords:** Metals exposure, autism spectrum disorder, pregnancy cohort, epidemiology,
60   cadmium

**Introduction**

Autism Spectrum Disorder (ASD) presents a major public health concern. ASD is a neurodevelopmental disorder characterized by impairments in social communication, social interaction, and restrictive and repetitive behavioral patterns and interests (Diagnostic and statistical manual of mental disorders : DSM-5. 2013). In the United States, 1 in 36 children are affected by ASD, with the prevalence among boys 3.8 times greater than among girls (Maenner et al. 2023). Individuals with ASD and their families face significant social and financial burdens, with higher costs for individuals with more severe ASD (Rogge and Janssen 2019). The social cost of ASD was greater than $7 trillion between the years 1990 – 2019 and is projected to be an additional $4 to $15 trillion by 2029 (Cakir et al. 2020). Increases in ASD prevalence have been attributed to increasing social awareness (Keyes et al. 2012), changes to diagnostic criteria (Hansen et al. 2015), and participation in early intervention services (Worley et al. 2011). However, the full source of this increase is largely unknown, suggesting incidence may be rising. Environmental exposures may play a role in this increase. Understanding modifiable risk factors for ASD could play a major role in guiding public health interventions.

Metals are a potential modifiable risk factor in ASD. In the United States, women of childbearing age experience widespread environmental exposure to metals, and higher concentrations have been observed in pregnant woman compared to non-pregnant women (Martin and Fry 2018; Watson et al. 2020). Important neurodevelopmental processes occur during pregnancy (Estes and McAllister 2016), and exposure to environmental factors such as metals are suggested to have a role in ASD etiology (Bölte et al. 2019; Heyer and Meredith 2017; Lyall et al. 2014). Among children diagnosed with ASD relative to controls, higher blood levels of arsenic (Ding et al. 2023), mercury (Ding et al. 2023; Jafari et al. 2017; Zhang et al. 2021), lead (Rashaid et al. 2021; Saghazadeh and Rezaei 2017; Zhang et al. 2021), and cadmium (Baj et al. 2021) have been observed. Although these findings are suggestive, exposure to metals was measured after ASD diagnosis. In studies where child levels of metals

were measured after ASD was diagnosed, it is not known if elevated exposure levels preceded

ASD. Some studies have examined exposure during pregnancy. Poorer performance on social

and behavioral tests among children at age 3 was associated with elevated manganese levels

in infant toenails and arsenic in maternal toenails (Doherty et al. 2020), and prenatal maternal

blood lead levels (Fruh et al. 2019). In contrast, elevated copper levels in maternal urine or

blood during pregnancy was associated with decreased behavior problems assessed in children

aged 3-7 years (Jedynak et al. 2021). Studies examining ASD diagnosis and metals exposure

during pregnancy are less common. One nested case-control study in the Norwegian Mother,

Father, and Child Cohort Study linked with the Norwegian Patient Registry examined maternal

blood metals concentrations during pregnancy, finding elevated arsenic, cadmium, and

manganese were associated with ASD, and lower levels of cesium, copper, mercury, and zinc

were associated with ASD (Skogheim et al. 2021). There is limited study on prenatal metals

exposure and ASD, and more prospective cohorts with exposure measures of multiple metals

are needed.

This study was conducted in two pregnancy cohorts with increased likelihood of ASD,

the Early Autism Risk Longitudinal Investigation (EARLI) and the Markers of Autism Risk in

Babies - Learning Early Signs (MARBLES) study. The goal of this study was to examine the

associations between concentrations of a panel of twenty-two metals measured during

pregnancy with ASD diagnosis in children at age 3 years.

**Methods**

*Study sample*

The Early Autism Risk Longitudinal Investigation (EARLI) is a prospective pregnancy

cohort to study autism etiology (Newschaffer et al. 2012). The EARLI study was reviewed and

approved by Human Subjects Institutional Review Boards (IRBs) from each of the four study

sites (Johns Hopkins University, Drexel University, University of California Davis, and Kaiser

Permanente Northern California). Markers of Autism Risk Learning Early Signs (MARBLES) is

also an enriched-familial risk prospective pregnancy cohort to study autism etiology (Hertz-

Picciotto et al.). The MARBLES protocol was reviewed and approved by the Human Subjects

IRB from University of California Davis. Secondary data analysis for this manuscript was

approved by the Human Subjects IRB for the University of Michigan. These studies recruited

mothers of children with clinically confirmed ASD (probands) who were early in a subsequent

pregnancy or were trying to become pregnant. In EARLI there were 232 mothers with a

subsequent child (sibling) born during the study between November 2009 and March 2012. In

MARBLES there were 389 enrolled mothers that gave birth to 425 subsequent children (sibling)

between December 1, 2006 and July 1, 2016.

*Covariate and outcome assessment*

Demographics, pregnancy behaviors, and medical history were all collected via maternal

questionnaire at enrollment. For the children born during the study (siblings), at age three years

clinicians assessed them with DSM-5 criteria. Children were categorized into three groups:

typically developing, ASD, or other non-typical development. Outcome categorization, based on

a previously published algorithm using the Autism Diagnostic Observation Schedule (ADOS)

and the Mullen Scales of Early Learning (MSEL) (Ozonoff et al. 2014), has been described in

these cohorts previously (Mordaunt et al. 2019; Philippat et al. 2018). In brief, the typical

development group did not meet diagnostic criteria for ASD. Those that did not meet diagnostic

criteria, but had ADOS scores within three points of the cutoff or MSEL scores 1.5 to 2 standard

deviations below average, were categorized in the non-typical development group. Finally,

those who met diagnostic DSM-5 criteria and ADOS scores over the cutoff were categorized in

the ASD group.

*Exposure assessment*

Maternal urine samples collected at two time points during pregnancy (earliest timepoint

approximately mean 19 weeks of pregnancy, latest timepoint approximately mean 32 weeks

pregnancy) had urinary concentrations of a panel of metals measured using inductively coupled

plasma mass spectrometry by NSF International (Centers for Disease Control and Prevention

method 3018.3, with modifications for the expanded metals panel and the Thermo Scientific

iCAP RQ instrument). Metals measured include antimony, arsenic, barium, beryllium, cadmium,

cesium, chromium, cobalt, copper, lead, manganese, mercury, molybdenum, nickel, platinum,

selenium, thallium, tin, tungsten, uranium, vanadium, and zinc. Samples for both cohorts were

randomized together into two laboratory runs and runs had variable limits of detection (LOD)

(**Supplemental Table 1**). For example, for lead the LOD was either 0.1 ppb (83 samples in

EARLI, 36 samples in MARBLES) or 0.2 ppb (262 samples in EARLI, 369 samples in

MARBLES), depending on batch. To assess urinary dilution, specific gravity was measured by

NSF International using an ATAGO handheld digital refractometer model PAL-10S. After

excluding samples involved in a multiple birth (n=23), related siblings from non-multiple births

(selecting one randomly to keep, n=18 samples dropped), or missing gestational age at

collection (n=3), EARLI had 165 mothers with metals measures from two timepoints, and 15

mothers with a measure from one timepoint. MARBLES had 154 mothers with metals measures

from two timepoints, and 97 mothers with a measure from one timepoint. Distribution of

gestational age at urine sample collection are shown in **Supplemental Figure 1**.

As a sensitivity analysis, maternal blood concentrations during pregnancy of cadmium,

manganese, lead, selenium, and total mercury were also measured in EARLI. Maternal venous

blood samples were collected in trace metal free EDTA tubes. Blood samples from the first

study visit (n=215) were used. Metal concentrations in maternal blood samples were measured

by inductively coupled dynamic reaction cell plasma mass spectrometry by the US Centers for

Disease Control and Prevention (ELAN DRC II, PerkinElmer Norwalk, CT) (method DLS 3016.8,

Centers for Disease Control and Prevention). Micro-clotting of the archived blood prevented

measures in half of samples, leaving n=104 with measured concentrations.

Data used in this manuscript is publicly available through the National Institute of Mental

Health Data Archive (EARLI cohort repository: 1600, MARBLES cohort repository: 1946,

EARLI/MARBLES metals repository: 2462) and through data requests to the Principal

Investigators of cohorts (EARLI: MDF, MARBLES: RJS).

*Statistical analyses*

We used R statistical software (version 4.0.2) for statistical analysis. Code to produce

analyses is available (https://github.com/bakulskilab). Metals with less than approximately 10%

of samples above the LOD were dropped from analysis (beryllium, platinum, tungsten, uranium,

vanadium). Metals with less than 75% of samples above the LOD (antimony, cadmium,

chromium, lead) were treated as binary variables, based on whether a sample was above or

below the LOD. For the rest of the metals, concentrations were used as continuous variables.

We substituted all urinary metals measures quantitated with values below the LOD with the

value of the LOD/square root of two (Hornung and Reed 1990). Metal concentrations were

adjusted for specific gravity by multiplying concentrations by the ratio of [the median specific

gravity – 1] and [sample specific gravity – 1] (Middleton et al. 2019). We then $\log_2$ transformed

the continuous concentrations. Outlier metals concentrations >5 standard deviations from the

mean were dropped from analyses. The number of samples dropped per metal are listed in

**Supplemental Table 2**.

We separated urinary measures into earlier and later pregnancy timepoints. For those

with two measures, the sample with lowest gestational age at collection was put into the earlier

timepoint sample. For those with only one sample, gestational age at collection >= 28 weeks

(third trimester) was considered late pregnancy, and otherwise samples were considered earlier

pregnancy. We compared exposure levels in early and late pregnancy with Spearman

correlation tests.

We calculated univariate descriptive statistics on each cohort using mean and standard

deviation for continuous variables and count and frequency for categorical variables. The

distributions of metal concentrations were described using mean, median, standard deviation,

interquartile range, and the number and percent above the limit of detection. We calculated

Spearman correlation of metals concentrations within each cohort. Separately for each cohort,

we compared the bivariate sample characteristics by neurodevelopmental outcome (ASD, non-

typically developing, typically developing) using ANOVA tests for continuous variables and chi-

square tests for categorical variables.

To estimate the adjusted associations between urinary metals concentration in

pregnancy and neurodevelopmental status, we used log binomial models to get estimates of

relative risk. Due to convergence issues, we used the delta-method normal approximations for

fitting models using the epitools package (Muller and MacLehose 2014). We estimated the

association of each metal with ASD status relative to the typically developing group using the

$\log_2$ transformed concentrations, adjusting for gestational age at urine sampling, child sex,

maternal age, and maternal education. We also tested metals associations with non-typically

developing status (typically developing as reference) in separate log binomial models.

Models were fit separately for each cohort, then meta analyzed together using the

inverse variance method in the R meta package. We reported risk ratios (RR) and 95%

confidence intervals (95% CI) for each association and visualized the results using forest plots.

For metals that were modeled continuously, since concentrations were log-transformed, the

reported associations are for a doubling in concentration. For metals that were modeled as

binary, we reported the risk ratio for above versus below the limit of detection. To account for

multiple comparisons, we also reported false discovery rate adjusted p-values.

We performed several sensitivity analyses to assess the robustness of our findings.

Since runs for metals measures had variable limits of detection, which impacts binary

categorization and imputation for values below limit of detection, we ran models adjusted for

batch. We also performed multivariable logistic regression for each of our models to generate

adjusted odds ratios (OR) that may be compared to the risk ratios and to prior findings in the

literature. Lastly, we performed analyses on the subset of EARLI samples with maternal blood

metals measures available and compared the findings to the findings in urinary metals.

**Results**

*Sample descriptive statistics*

At the earliest pregnancy timepoint, urinary metal concentrations were above the limit of

detection in greater than 75% of the samples for 13 metals in each cohort (arsenic, barium,

cesium, cobalt, copper, manganese, mercury, molybdenum, nickel, selenium, thallium, tin, and

zinc) (**Supplemental Table 3**). In both EARLI and MARBLES, cobalt (Co) and nickel (Ni)

concentrations had the strongest correlation (Spearman r=0.57 in EARLI, r=0.59 in MARBLES)

(**Figure 1**).

In the earlier pregnancy timepoint there were metals concentrations available from 170

urine samples in EARLI (66 typically developing, 74 non-typically developing, 30 ASD) and 158

in MARBLES (104 typically developing, 18 non-typically developing, 36 ASD) (**Table 1**). In

EARLI, maternal education and child sex assigned at birth were associated with child

neurodevelopmental status. The typically developing group had higher levels of maternal

education (71.2% with college degree), compared to the non-typically developing (54.1%) and

ASD groups (43.3%). The typically developing and non-typically developing groups had a

similar proportion of males (42.4% and 47.3%) but lower proportion than the ASD group

(76.7%). In MARBLES, compared to the typically developing group (51.9% male), both the non-

typically developing group (66.7% male) and ASD (75.0% male) had higher proportion of males

(**Table 1**).

At the later pregnancy timepoint, urinary metal concentrations were above the limit of

detection in slightly less than 75% of the sample for manganese, mercury, and tin

(**Supplemental Table 4**), however they were modelled as continuous for comparability with the

early timepoint results. In late pregnancy, cobalt and nickel remained the strongest correlated

metals in MARBLES (Spearman r=0.74), but not in EARLI. In both cohorts, lead and copper

(r=0.43 in EARLI, r=0.49in MARBLES) as well as lead and manganese (r=0.45 in both) were

correlated (**Supplemental Figure 2**).

At the later pregnancy timepoint, there were 171 samples with urinary metal

concentrations available in EARLI (65 typically developing, 75 non-typically developing, 31

ASD) and 231 in MARBLES (146 typically developing, 34 non-typically developing, 51 ASD)

(**Supplemental Table 5**). For mothers with two timepoints, correlation between the two were

strongest for measured cadmium (r=0.48 in EARLI, r=0.42 in MARBLES), cesium (r=0.42 in

both cohorts), mercury (r=0.48 in EARLI, r=0.63 in MARBLES), tin (r=0.57 in EARLI, r=0.59 in

MARBLES), and zinc (r=0.53 in EARLI, r=0.47 in MARBLES). Cross timepoint correlation was

weakest for antimony (r=0.29 in EARLI, r=0.15 in MARBLES) and manganese (r=0.21 in EARLI,

r=0.12 in MARBLES) (**Supplemental Table 6**).

*Urinary Metal Association with Autism Spectrum Disorder Status*

We examined associations between urinary metals in the earlier pregnancy timepoint

and ASD. In meta-analysis, comparing ASD to typical development, having urine cadmium

concentration above the limit of detection was associated with RR=1.78 (95% CI 1.19, 2.67)

times higher risk for ASD (EARLI RR=1.92, 95% CI 1.05, 3.51; MARBLES RR=1.68, 95% CI

0.97, 2.89). (**Figure 2, Table 2**). A doubling in arsenic was associated with lower ASD risk

(RR=0.85, 95% CI 0.76, 0.94), driven by the EARLI cohort (EARLI RR=0.82, 95% CI 0.73, 0.92;

MARBLES RR=1.01, 95% CI 0.78, 1.31). Marginal associations were observed with cesium,

where a doubling in urinary concentration was estimated to have RR=1.81 (95% CI 0.95, 3.42).

Thallium concentration doubling was marginally associated with RR=1.16 (95% CI 1.08, 1.26),

with stronger effect in MARBLES (RR=1.15, 95% CI 1.10, 1.32) than in EARLI (RR=1.00, 95%

CI 0.68, 1.48). The associations for cadmium (FDR=0.085) and arsenic (FDR=0.051) reached

FDR < 0.1 when adjusting for multiple comparisons. No associations were observed between

the remaining urinary metal concentrations and ASD status at this early pregnancy time point.

At the later pregnancy timepoint, we estimated the association between each metal

concentration and ASD. Comparing ASD to typically developing in meta-analyses, a doubling in

cesium was associated with RR=1.71 (95% CI 1.01, 2.9) in meta-analysis (EARLI RR=2.10,

95% CI 0.85, 5.22.; MARBLES RR=1.55, 95% CI 0.81, 2.95) (**Figure 2, Table 2**). A doubling in

thallium was associated with ASD with RR=1.16 (95% CI 1.10,1.23), though effects were

different between cohorts (EARLI RR=0.84, 95% CI 0.52, 1.37.; MARBLES RR=1.17, 95% CI

1.10, 1.24). A doubling in tin was associated with RR=1.12 (95% CI 1.01,1.24). A doubling in

mercury was marginally associated with RR=1.09 (95% CI 1.00, 1.20), with differences between

EARLI (RR=1.14, 95% CI 1.03, 1.26) and MARBLES (RR=0.91, 95% CI 0.73, 1.13). The

association with thallium reached FDR < 0.1. No associations were observed between the

remaining urinary metal concentrations and ASD status at this later pregnancy time point.

*Urinary Metal Association with Non-Typically Developing Status*

We repeated the adjusted regression analyses estimating the association of earlier

pregnancy urinary metals and non-typically developing status. Urine cadmium concentrations

above the LOD was associated with elevated risk of non-typical development, with RR=1.43

(95% CI 1.06, 1.92). (**Figure 3, Table 3**). A doubling of cesium urinary concentration was

marginally associated with RR=1.58 (95% CI 0.95, 2.63) times higher risk of non-typical

development. A marginal relationship with molybdenum was also observed, where a doubling in

concentration was related to RR=1.47 (95% CI 0.95, 2.28). A doubling of nickel was associated

with RR=1.40 (95% CI 1.01, 1.94), driven by the EARLI cohort with RR=1.55 (95% CI 1.08,

2.23). No associations were observed between the remaining urinary metal concentrations and

non-typically developing status at this early pregnancy time point.

We examined associations between non-typically developing and the later pregnancy

metals measures. Though not statistically significant, having urine cadmium concentration

above the limit of detection was associated with RR=1.24 (95% CI 0.95, 1.63) times higher risk

of non-typically developing status in meta-analysis (**Supplemental Figure 3, Table 3**). A

doubling of the essential metal selenium concentration was associated in meta-analysis with

RR=0.85 (95% CI 0.82, 0.88) times lower risk of non-typically developing status, driven by

precision of results in MARBLES and had opposite directions of effect by cohort (EARLI

RR=1.31, 95% CI 0.64, 2.69; MARBLES RR=0.85, 95% CI 0.82, 0.88). A doubling of the

essential metal zinc concentration was associated with RR=0.97 (95% CI 0.95, 0.98) lower risk

of non-typically developing status. The zinc and selenium associations reached FDR < 0.1. No

associations were observed between the remaining urinary metal concentrations and non-

typically developing status at this late pregnancy time point.

*Maternal Blood Metal Association with Neurodevelopmental Status*

In EARLI, 92 maternal blood samples collected during pregnancy had available

covariate and blood metals measures (41 typically developing, 32 non- typically developing, 19

ASD) (**Supplemental Table 7**). A doubling in maternal blood cadmium was marginally

associated with RR=1.11 (95% CI 0.96, 1.29) higher risk of ASD, and a doubling in maternal

blood lead was associated with RR=1.23 (95% CI 1.01, 1.54) higher risk of ASD (**Supplemental**

**Figure 3**). A doubling in cadmium was also associated with RR=1.10 (95% CI 1.02, 1.19) higher

risk of non-typical development. A doubling in maternal blood lead was associated with

RR=1.16 (95% CI 1.00, 1.35) higher risk of non-typical development (**Supplemental Figure 2**).

No associations were observed between the remaining blood metal concentrations (mercury,

selenium, manganese) and neurodevelopmental status.

*Sensitivity Analysis*

With batch as a covariate (**Supplemental Tables 8 and 9)**, the cadmium association in

early pregnancy with ASD remained consistent where being over the limit of detection was

associated with RR=1.78 (95% CI 1.19, 2.65) times higher risk of ASD. The cadmium

associations with non-typical development also remained consistent. With batch adjustment,

antimony in earlier pregnancy was associated with ASD, with RR=1.58 (95% CI 1.01, 2.48). The

associations between cesium and ASD as well as non-typical development remained consistent

with slight attenuation, as did associations between earlier pregnancy molybdenum and nickel

with non-typical development. On the other hand, the relationships with thallium and ASD were

attenuated.

Using logistic regression models, consistency to the previous log binomial findings was

observed for cadmium and cesium. In general, estimates on the odds ratio scale were higher in

magnitude and significance for cadmium and cesium. Strength of relationships between arsenic,

mercury, thallium, and tin with ASD were attenuated with larger confidence intervals when using

logistic regression (**Supplemental Table 10**)**.** Earlier pregnancy molybdenum and nickel

associations with non-typical development remained consistent in logistic regression, while

selenium and zinc associations with non-typical development were attenuated (**Supplemental**

**Table 11**). In logistic regression, earlier pregnancy lead had stronger association with non-

typical development, where lead over the LOD was marginally associated with OR=1.68 (95%

CI 0.92, 3.07).

**Discussion**

In these two enriched-likelihood prospective birth cohorts, we measured maternal urinary

metals levels during two timepoints in pregnancy and examined relationships to ASD or non-

typical development status at age 3 in offspring. To our knowledge, this is the first study to

report associations between maternal prenatal urine heavy metal concentrations and ASD

diagnosis in children at 3 years of age in enriched-likelihood prospective cohorts. Our most

consistent finding was heightened risk of atypical neurodevelopment related to cadmium

exposure. Although the relationships were not significant in late pregnancy, the directions of

effect were consistent across time periods. Furthermore, similar findings were observed in the

maternal blood subsample. Cesium related to atypical neurodevelopment was also notable, with

consistency across ASD and non-typical development outcomes and timepoints, with exception

of later pregnancy cesium and non-typical development. Cadmium and cesium associations

were also the most robust to different modelling strategies. This study suggests metals

exposure during pregnancy may be related to risk of ASD or non-typical development status at

age 3.

Existing studies have examined the relationship between heavy metals exposure and

ASD with considerable heterogeneity in exposure timing and matrices measured (Campbell et

al. 2021). A systematic review and meta-analysis of lead concentrations in children with ASD

from cross-sectional and case-control studies showed significant difference in blood lead levels

compared to controls, but not in urinary lead levels (Nakhaee et al. 2023)**.** This mirrors our

results in maternal measures during pregnancy, where we found maternal blood lead levels

were associated with risk of ASD or non-typical development in offspring, but not maternal

urinary lead levels. The study in the Norwegian Mother, Father, and Child Cohort Study found

higher odds of ASD for children in the highest quartile of cadmium exposure measured in

maternal blood during pregnancy (Skogheim et al. 2021), matching results from the present

study. The same study found the highest quartile of maternal blood cesium levels had lower

odds of ASD compared to the lowest quartile, while in contrast our study suggests higher risk of

ASD with higher maternal urinary cesium. Our results for selenium were mixed, doubling of late

pregnancy selenium concentration was associated with lower risk of non-typical development,

however there were opposite effect estimates between cohorts. Selenium supplementation in an

animal model attenuated autism phenotype (Wu et al. 2022a), and studies measuring selenium

cross-sectionally in children in Saudi Arabia (El-Ansary et al. 2017) and China (Wu et al. 2022b)

found lower selenium levels in those with ASD. On the other hand, two-sample Mendelian

randomization analysis using genetic instruments of blood and blood-toenail selenium suggest

selenium levels are associated with increased risk of ASD (Guo et al. 2023), and in the Boston

Birth Cohort maternal red blood cell selenium levels measured at near delivery were associated

with increased odds of ASD in children (Lee et al. 2021). Considerable heterogeneity in

direction of association between selenium and ASD exist in our study and in the literature, along

with heterogeneity in timing and tissue of measurements. Our findings add to a growing body of

evidence of the neurodevelopmental impacts of metals exposure during pregnancy.

This study has several strengths. We were able to assay a wide array of metals with

high detection rates in two different birth cohorts, at two different timepoints. In one cohort, we

were also able to evaluate five metals in a different exposure matrix: maternal blood during

pregnancy. The longitudinal design allowed examination of exposure measures during

pregnancy that preceded subsequent ASD outcome 36 months after birth. The enriched risk

cohort design ensured all participants were clinically assessed using gold standards for ASD

diagnosis.

This study modeled metals as linear or dichotomous, but some metals may have non-

linear relationships. While the enriched likelihood cohort design allowed for an extensively

phenotyped sample, our findings may not be generalizable to populations without increased

propensity for ASD, thus it would be important to also compare to results found in population-

based samples. Future studies should also consider other exposure matrices or timepoints. The

choice of exposure matrix is also important for exposure timing. For example, blood cadmium

levels reflect recent exposure, while urinary cadmium reflects a longer, cumulative exposure

(Agency for Toxic Substances and Disease Registry (ATSDR) 2012). Certain exposure matrices

may be more reliable for some metals. For example, blood lead is a more reliable measure of

recent exposure compared to urinary or hair lead levels (Agency for Toxic Substances and

Disease Registry (ATSDR) 2020), which may explain our findings of stronger blood lead ASD

associations than those seen with urinary lead.

This study suggests that prenatal exposure to metals, such as cadmium, impacts risk of

ASD or non-typical development in offspring. Public health measures to reduce exposure to

heavy metals during pregnancy may be an important preventative strategy for

neurodevelopmental disorders, though larger longitudinal studies are needed.

**References**

Agency for Toxic Substances and Disease Registry (ATSDR). 2012. Toxicological profile for Cadmium.

Agency for Toxic Substances and Disease Registry (ATSDR). 2020. Toxicological profile for Lead.

Baj J, Flieger W, Flieger M, Forma A, Sitarz E, Skórzyńska-Dziduszko K, et al. 2021. Autism spectrum disorder: Trace elements imbalances and the pathogenesis and severity of autistic symptoms. Neuroscience and Biobehavioral Reviews 129:117–132; doi:10.1016/j.neubiorev.2021.07.029.

Bölte S, Girdler S, Marschik PB. 2019. The contribution of environmental exposure to the etiology of autism spectrum disorder. Cellular and Molecular Life Sciences 76:1275–1297; doi:10.1007/s00018-018-2988-4.

Cakir J, Frye RE, Walker SJ. 2020. The lifetime social cost of autism: 1990–2029. Research in Autism Spectrum Disorders 72:1–18; doi:10.1016/j.rasd.2019.101502.

Campbell KA, Hickman R, Fallin MD, Bakulski KM. 2021. Prenatal exposure to metals and autism spectrum disorder: Current status and future directions. Current Opinion in Toxicology 26:39–48; doi:10.1016/j.cotox.2021.04.001.

Diagnostic and statistical manual of mental disorders□: DSM-5. 2013.5th ed. American Psychiatric Association:Arlington, VA.

Ding M, Shi S, Qie S, Li J, Xi X. 2023. Association between heavy metals exposure (cadmium, lead, arsenic, mercury) and child autistic disorder: a systematic review and meta-analysis. Front Pediatr 11:1169733; doi:10.3389/fped.2023.1169733.

Doherty BT, Romano ME, Gui J, Punshon T, Jackson BP, Karagas MR, et al. 2020. Periconceptional and prenatal exposure to metal mixtures in relation to behavioral development at 3 years of age. Environmental Epidemiology 4; doi:10.1097/EE9.0000000000000106.

El-Ansary A, Bjørklund G, Tinkov AA, Skalny AV, Al Dera H. 2017. Relationship between selenium, lead, and mercury in red blood cells of Saudi autistic children. Metab Brain Dis 32:1073–1080; doi:10.1007/s11011-017-9996-1.

Estes ML, McAllister AK. 2016. Maternal immune activation: Implications for neuropsychiatric disorders. Science 353:772–777; doi:10.1126/science.aag3194.

Fruh V, Rifas-Shiman SL, Amarasiriwardena C, Cardenas A, Bellinger DC, Wise LA, et al. 2019. Prenatal lead exposure and childhood executive function and behavioral difficulties in project viva. NeuroToxicology 75:105–115; doi:10.1016/j.neuro.2019.09.006.

Guo X, Tang P, Hou C, Li R. 2023. Mendelian randomization investigation highlights different roles of selenium status in mental disorders. Prog Neuropsychopharmacol Biol Psychiatry 122:110694; doi:10.1016/j.pnpbp.2022.110694.

Hansen SN, Schendel DE, Parner ET. 2015. Explaining the increase in the prevalence of autism
spectrum disorders: The proportion attributable to changes in reporting practices. JAMA
Pediatrics 169:56–62; doi:10.1001/jamapediatrics.2014.1893.

Hertz -Picciotto Irva, Schmidt RJ, Walker CK, Bennett DH, Oliver M, Shedd -Wise Kristine M., et
al. A Prospective Study of Environmental Exposures and Early Biomarkers in Autism
Spectrum Disorder: Design, Protocols, and Preliminary Data from the MARBLES Study.
Environmental Health Perspectives 126:117004; doi:10.1289/EHP535.

Heyer DB, Meredith RM. 2017. Environmental toxicology: Sensitive periods of development and
neurodevelopmental disorders. NeuroToxicology 58:23–41;
doi:10.1016/j.neuro.2016.10.017.

Hornung RW, Reed LD. 1990. Estimation of Average Concentration in the Presence of
Nondetectable Values. Applied Occupational and Environmental Hygiene 5:46–51;
doi:10.1080/1047322X.1990.10389587.

Jafari T, Rostampour N, Fallah AA, Hesami A. 2017. The association between mercury levels
and autism spectrum disorders: A systematic review and meta-analysis. Journal of Trace
Elements in Medicine and Biology 44:289–297; doi:10.1016/j.jtemb.2017.09.002.

Jedynak P, Maitre L, Guxens M, Gützkow KB, Julvez J, López-Vicente M, et al. 2021. Prenatal
exposure to a wide range of environmental chemicals and child behaviour between 3
and 7 years of age – An exposome-based approach in 5 European cohorts. Science of
the Total Environment 763; doi:10.1016/j.scitotenv.2020.144115.

Keyes KM, Susser E, Cheslack-postava K, Fountain C, Liu K, Bearman PS. 2012. Cohort
effects explain the increase in autism diagnosis among children born from 1992 to 2003
in california. International Journal of Epidemiology 41:495–503; doi:10.1093/ije/dyr193.

Lee ASE, Ji Y, Raghavan R, Wang G, Hong X, Pearson C, et al. 2021. Maternal prenatal
selenium levels and child risk of neurodevelopmental disorders: A prospective birth
cohort study. Autism Res 14:2533–2543; doi:10.1002/aur.2617.

Lyall K, Schmidt RJ, Hertz-Picciotto I. 2014. Maternal lifestyle and environmental risk factors for
autism spectrum disorders. International Journal of Epidemiology 43:443–464;
doi:10.1093/ije/dyt282.

Maenner MJ, Warren Z, Williams AR, Amoakohene E, Bakian AV, Bilder DA, et al. 2023.
Prevalence and Characteristics of Autism Spectrum Disorder Among Children Aged 8
Years - Autism and Developmental Disabilities Monitoring Network, 11 Sites, United
States, 2020. MMWR Surveill Summ 72:1–14; doi:10.15585/mmwr.ss7202a1.

Martin EM, Fry RC. 2018. Environmental Influences on the Epigenome: Exposure- Associated
DNA Methylation in Human Populations. Annual Review of Public Health 39:309–333;
doi:10.1146/annurev-publhealth-040617-014629.

Middleton DRS, Watts MJ, Polya DA. 2019. A comparative assessment of dilution correction
methods for spot urinary analyte concentrations in a UK population exposed to arsenic in
drinking water. Environment International 130:104721; doi:10.1016/j.envint.2019.03.069.

Mordaunt CE, Park BY, Bakulski KM, Feinberg JI, Croen LA, Ladd-Acosta C, et al. 2019. A
meta-analysis of two high-risk prospective cohort studies reveals autism-specific
transcriptional changes to chromatin, autoimmune, and environmental response genes
in umbilical cord blood. Mol Autism 10:36; doi:10.1186/s13229-019-0287-z.

Muller CJ, MacLehose RF. 2014. Estimating predicted probabilities from logistic regression:
different methods correspond to different target populations. International Journal of
Epidemiology 43:962–970; doi:10.1093/ije/dyu029.

Nakhaee S, Amirabadizadeh A, Farnia V, Ali Azadi N, Mansouri B, Radmehr F. 2023.
Association Between Biological Lead Concentrations and Autism Spectrum Disorder
(ASD) in Children: a Systematic Review and Meta-Analysis. Biol Trace Elem Res
201:1567–1581; doi:10.1007/s12011-022-03265-9.

Newschaffer CJ, Croen LA, Fallin MD, Hertz-Picciotto I, Nguyen DV, Lee NL, et al. 2012. Infant
siblings and the investigation of autism risk factors. Journal of Neurodevelopmental
Disorders 4:7; doi:10.1186/1866-1955-4-7.

Ozonoff S, Young GS, Belding A, Hill M, Hill A, Hutman T, et al. 2014. The Broader Autism
Phenotype in Infancy: When Does It Emerge? J Am Acad Child Adolesc Psychiatry
53:398-407.e2; doi:10.1016/j.jaac.2013.12.020.

Philippat C, Barkoski J, Tancredi DJ, Elms B, Barr D, Ozonoff S, et al. 2018. Prenatal exposure
to organophosphate pesticides and risk of autism spectrum disorders and other non-
typical development at 3 years in a high-risk cohort. Int J Hyg Environ Health 221:548–
555; doi:10.1016/j.ijheh.2018.02.004.

Rashaid AHB, Nusair SD, Alqhazo MT, Adams JB, Abu-Dalo MA, Bashtawi MA. 2021. Heavy
metals and trace elements in scalp hair samples of children with severe autism spectrum
disorder: A case-control study on Jordanian children. Journal of Trace Elements in
Medicine and Biology 67; doi:10.1016/j.jtemb.2021.126790.

Rogge N, Janssen J. 2019. The Economic Costs of Autism Spectrum Disorder: A Literature
Review. Journal of Autism and Developmental Disorders 49:2873–2900;
doi:10.1007/s10803-019-04014-z.

Saghazadeh A, Rezaei N. 2017. Systematic review and meta-analysis links autism and toxic
metals and highlights the impact of country development status: Higher blood and
erythrocyte levels for mercury and lead, and higher hair antimony, cadmium, lead, and
mercury. Progress in Neuro-Psychopharmacology and Biological Psychiatry 79:340–
368; doi:10.1016/j.pnpbp.2017.07.011.

Skogheim TS, Weyde KVF, Engel SM, Aase H, Surén P, Øie MG, et al. 2021. Metal and
essential element concentrations during pregnancy and associations with autism
spectrum disorder and attention-deficit/hyperactivity disorder in children. Environment
International 152; doi:10.1016/j.envint.2021.106468.

Watson CV, Lewin M, Ragin-Wilson A, Jones R, Jarrett JM, Wallon K, et al. 2020.
Characterization of trace elements exposure in pregnant women in the United States,
NHANES 1999–2016. Environmental Research 183; doi:10.1016/j.envres.2020.109208.

Worley JA, Matson JL, Sipes M, Kozlowski AM. 2011. Prevalence of autism spectrum disorders
in toddlers receiving early intervention services. Research in Autism Spectrum Disorders
5:920–925; doi:10.1016/j.rasd.2010.10.007.

Wu H, Zhao G, Liu S, Zhang Q, Wang P, Cao Y, et al. 2022a. Supplementation with selenium
attenuates autism-like behaviors and improves oxidative stress, inflammation and
related gene expression in an autism disease model. J Nutr Biochem 107:109034;
doi:10.1016/j.jnutbio.2022.109034.

Wu J, Wang D, Yan L, Jia M, Zhang J, Han S, et al. 2022b. Associations of essential element
serum concentrations with autism spectrum disorder. Environ Sci Pollut Res Int
29:88962–88971; doi:10.1007/s11356-022-21978-1.

Zhang J, Li X, Shen L, Ullah N, Zhang X, Chen L, et al. 2021. Journal of Trace Elements in
Medicine and Biology Trace elements in children with autism spectrum disorder□: A
meta-analysis based on case-control studies. 67.

535
536

**Table 1.** Maternal and child characteristics of participants in the primary analytic sample with measures of metal exposure in urine from early pregnancy. Data are split by cohort and comparted by neurodevelopmental status of the sibling. Distributions of categorical variables are compared with a chi-square test and continuous variables are compared with ANOVA test.

| EARLI cohort | Typically developing *N=66* | Non-typically developing *N=74* | Autism spectrum disorder *N=30* | P-value |
|---|---|---|---|---|
| Maternal Education | | | | 0.020 |
|   College Degree | 47 (71.2%) | 40 (54.1%) | 13 (43.3%) | |
|   No Degree | 19 (28.8%) | 34 (45.9%) | 17 (56.7%) | |
| Maternal Age | 35.1 (4.68) | 33.3 (4.52) | 33.8 (3.87) | 0.057 |
| Infant Sex | | | | 0.006 |
|   F | 38 (57.6%) | 39 (52.7%) | 7 (23.3%) | |
|   M | 28 (42.4%) | 35 (47.3%) | 23 (76.7%) | |
| Weeks of Gestation at Sample Collection | 19.1 (5.14) | 18.3 (6.44) | 19.4 (7.76) | 0.607 |
| Infant Gestational Age at Birth | 39.5 (1.40) | 39.5 (1.36) | 38.9 (2.00) | 0.094 |
| **MARBLES cohort** | **Typically developing** *N=104* | **Non-typically developing** *N=18* | **Autism spectrum disorder** *N=36* | **P-value** |
| Maternal Education | | | | 0.259 |
|   College Degree | 62 (59.6%) | 9 (50.0%) | 16 (44.4%) | |
|   No Degree | 42 (40.4%) | 9 (50.0%) | 20 (55.6%) | |
| Maternal Age | 34.6 (4.73) | 33.8 (4.71) | 34.6 (5.01) | 0.822 |
| Infant Sex | | | | 0.041 |
|   Female | 50 (48.1%) | 6 (33.3%) | 9 (25.0%) | |
|   Male | 54 (51.9%) | 12 (66.7%) | 27 (75.0%) | |
| Weeks of Gestation at Sample Collection | 19.3 (3.99) | 18.7 (3.99) | 19.0 (3.98) | 0.803 |
| Infant Gestational Age at Birth | 38.9 (1.39) | 38.9 (1.54) | 39.3 (1.04) | 0.296 |

*Acronyms: Early Autism Risk Longitudinal Investigation (EARLI), Markers of Autism Risk in Babies-Learning Early Signs (MARBLES)*

**Table 2.** Adjusted risk ratios for the associations between maternal urinary metal concentrations measured during pregnancy and risk of autism spectrum disorder, relative to typically developing. Log binomial models were adjusted for gestational age, child sex, maternal age, and maternal education. Four metals were modeled categorically (above versus below the limit of detection) and the remaining metals were $\log_2$ transformed and modeled continuously.

| Metal | Time | EARLI RR | EARLI CI | EARLI P | MARBLES RR | MARBLES CI | MARBLES P | Meta-Analysis RR | Meta-Analysis CI | Meta-Analysis P | Meta-Analysis FDR |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | | RR | CI | P | RR | CI | P | RR | CI | P | FDR |
| **Modeled categorically** | | | | | | | | | | | |
| Antimony | Early | 1.10 | (0.59,2.07) | 0.76 | 1.25 | (0.63,2.46) | 0.52 | 1.17 | (0.74,1.85) | 0.51 | 0.67 |
| | Late | 1.08 | (0.59,1.97) | 0.81 | 1.21 | (0.66,2.20) | 0.54 | 1.14 | (0.75,1.75) | 0.54 | 0.88 |
| Cadmium | Early** | 1.92 | (1.05,3.51) | 0.034 | 1.68 | (0.97,2.89) | 0.062 | 1.78 | (1.19,2.67) | 0.005 | 0.085 |
| | Late | 1.55 | (0.91,2.64) | 0.11 | 0.97 | (0.56,1.68) | 0.90 | 1.23 | (0.84,1.81) | 0.28 | 0.79 |
| Chromium | Early | 1.34 | (0.76,2.39) | 0.31 | 1.35 | (0.71,2.56) | 0.37 | 1.34 | (0.88,2.06) | 0.18 | 0.45 |
| | Late | 1.43 | (0.81,2.51) | 0.22 | 0.92 | (0.42,2.03) | 0.84 | 1.23 | (0.78,1.95) | 0.38 | 0.88 |
| Lead | Early | 1.73 | (0.90,3.33) | 0.10 | 1.09 | (0.61,1.93) | 0.78 | 1.33 | (0.86,2.05) | 0.20 | 0.45 |
| | Late | 1.64 | (0.87,3.08) | 0.13 | 1.08 | (0.67,1.73) | 0.75 | 1.25 | (0.86,1.83) | 0.24 | 0.74 |
| **Modeled continuously** | | | | | | | | | | | |
| Arsenic | Early** | 0.82 | (0.73,0.92) | <0.001 | 1.01 | (0.78,1.31) | 0.93 | 0.85 | (0.76,0.94) | 0.0015 | 0.051 |
| | Late | 1.15 | (0.91,1.46) | 0.24 | 1.10 | (0.89,1.36) | 0.39 | 1.12 | (0.96,1.32) | 0.15 | 0.60 |
| Barium | Early | 1.19 | (0.91,1.56) | 0.20 | 1.07 | (0.84,1.36) | 0.58 | 1.12 | (0.94,1.34) | 0.20 | 0.45 |
| | Late | 0.98 | (0.81,1.19) | 0.85 | 0.99 | (0.83,1.16) | 0.86 | 0.98 | (0.87,1.11) | 0.80 | 0.92 |
| Cesium | Early* | 2.37 | (0.80,7.04) | 0.12 | 1.56 | (0.71,3.44) | 0.27 | 1.81 | (0.95,3.42) | 0.069 | 0.35 |
| | Late** | 2.10 | (0.85,5.22) | 0.11 | 1.55 | (0.81,2.95) | 0.19 | 1.71 | (1.01,2.9) | 0.045 | 0.31 |
| Cobalt | Early | 0.83 | (0.51,1.34) | 0.43 | 0.74 | (0.42,1.29) | 0.28 | 0.79 | (0.55,1.13) | 0.20 | 0.45 |
| | Late | 0.96 | (0.67,1.36) | 0.81 | 1.00 | (0.73,1.36) | 0.99 | 0.98 | (0.78,1.24) | 0.87 | 0.92 |
| Copper | Early | 1.37 | (0.61,3.08) | 0.45 | 1.09 | (0.64,1.85) | 0.76 | 1.17 | (0.75,1.82) | 0.50 | 0.67 |
| | Late | 3.55 | (1.23,10.3) | 0.02 | 0.93 | (0.72,1.19) | 0.55 | 0.99 | (0.78,1.27) | 0.96 | 0.96 |
| Manganese | Early | 0.95 | (0.67,1.35) | 0.79 | 1.05 | (0.80,1.38) | 0.71 | 1.01 | (0.82,1.26) | 0.90 | 0.96 |
| | Late | 0.94 | (0.70,1.25) | 0.66 | 1.00 | (0.79,1.28) | 0.97 | 0.98 | (0.81,1.17) | 0.79 | 0.92 |
| Mercury | Early | 1.14 | (1.01,1.29) | 0.039 | 0.82 | (0.60,1.11) | 0.19 | 1.09 | (0.97,1.22) | 0.16 | 0.45 |
| | Late* | 1.14 | (1.03,1.26) | 0.013 | 0.91 | (0.73,1.13) | 0.40 | 1.09 | (1.00,1.20) | 0.061 | 0.35 |
| Molybdenum | Early | 1.46 | (0.73,2.91) | 0.29 | 0.97 | (0.68,1.38) | 0.88 | 1.06 | (0.77,1.45) | 0.73 | 0.85 |
| | Late | 0.93 | (0.79,1.10) | 0.43 | 1.08 | (0.78,1.49) | 0.64 | 0.96 | (0.83,1.12) | 0.62 | 0.88 |
| Nickel | Early | 1.05 | (0.72,1.53) | 0.8 | 1.04 | (0.57,1.88) | 0.90 | 1.05 | (0.76,1.44) | 0.78 | 0.86 |
| | Late | 1.07 | (0.72,1.60) | 0.73 | 1.01 | (0.70,1.44) | 0.96 | 1.04 | (0.79,1.35) | 0.79 | 0.92 |
| Selenium | Early | 0.95 | (0.65,1.40) | 0.8 | 0.91 | (0.75,1.10) | 0.33 | 0.92 | (0.77,1.09) | 0.32 | 0.55 |
| | Late | 2.13 | (0.69,6.55) | 0.19 | 0.92 | (0.72,1.18) | 0.52 | 0.96 | (0.75,1.22) | 0.73 | 0.92 |
| Thallium | Early* | 1 | (0.68,1.48) | 0.98 | 1.15 | (1.01,1.32) | 0.038 | 1.14 | (1.00,1.29) | 0.05 | 0.34 |
| | Late** | 0.84 | (0.52,1.37) | 0.5 | 1.17 | (1.10,1.24) | <0.001 | 1.16 | (1.10,1.23) | <0.001 | <0.001 |
| Tin | Early | 1.17 | (1,1.36) | 0.043 | 0.98 | (0.77,1.24) | 0.85 | 1.11 | (0.98,1.26) | 0.11 | 0.42 |

| Metal | Time | EARLI | | | MARBLES | | | Meta-Analysis | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | | RR | CI | P | RR | CI | P | RR | CI | P | FDR |
| | Late** | 1.18 | (1.04,1.33) | 0.01 | 1.01 | (0.84,1.22) | 0.89 | 1.12 | (1.01,1.24) | 0.027 | 0.23 |
| Zinc | Early | 1.93 | (1.01,3.70) | 0.048 | 0.95 | (0.82,1.09) | 0.45 | 0.98 | (0.85,1.13) | 0.75 | 0.85 |
| | Late | 1.51 | (0.75,3.01) | 0.25 | 1.09 | (0.81,1.48) | 0.56 | 1.15 | (0.87,1.52) | 0.32 | 0.84 |

*Symbols: ** meta-analysis p-value < 0.05. * meta-analysis p-value < 0.1.*

**Table 3.** Adjusted risk ratios for the associations between maternal urinary metal concentrations measured during pregnancy and risk of non-typically developing, relative to typically developing. Log binomial models were adjusted for gestational age, child sex, maternal age, and maternal education. Four metals were modeled categorically (above versus below the limit of detection) and the remaining metals were $\log_2$ transformed and modeled continuously.

| Metal | Time | EARLI | | | MARBLES | | | Meta-Analysis | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | | RR | CI | P | RR | CI | P | RR | CI | P | FDR |
| | | | | | Modeled categorically | | | | | | |
| Antimony | Early | 1.18 | (0.85,1.64) | 0.32 | 0.70 | (0.18,2.81) | 0.62 | 1.15 | (0.84,1.58) | 0.39 | 0.55 |
| | Late | 0.96 | (0.67,1.38) | 0.84 | 1.68 | (0.85,3.35) | 0.14 | 1.09 | (0.79,1.50) | 0.61 | 0.88 |
| Cadmium | Early** | 1.28 | (0.94,1.76) | 0.12 | 3.12 | (1.34,7.3) | 0.008 | 1.43 | (1.06,1.92) | 0.018 | 0.20 |
| | Late | 1.10 | (0.81,1.49) | 0.55 | 2.02 | (1.1,3.71) | 0.023 | 1.24 | (0.95,1.63) | 0.12 | 0.58 |
| Chromium | Early | 0.97 | (0.68,1.37) | 0.86 | 1.47 | (0.54,3.99) | 0.45 | 1.01 | (0.73,1.41) | 0.93 | 0.96 |
| | Late | 1.01 | (0.67,1.51) | 0.97 | 1.50 | (0.70,3.21) | 0.29 | 1.1 | (0.77,1.58) | 0.60 | 0.88 |
| Lead | Early | 1.11 | (0.80,1.55) | 0.53 | 2.50 | (1.04,6.00) | 0.041 | 1.23 | (0.9,1.67) | 0.19 | 0.45 |
| | Late | 0.87 | (0.64,1.18) | 0.36 | 1.09 | (0.59,2.00) | 0.79 | 0.91 | (0.69,1.19) | 0.49 | 0.88 |
| | | | | | Modeled continuously | | | | | | |
| Arsenic | Early | 1.10 | (0.95,1.28) | 0.19 | 0.87 | (0.60,1.27) | 0.48 | 1.07 | (0.93,1.23) | 0.33 | 0.55 |
| | Late | 1.08 | (0.93,1.25) | 0.34 | 0.81 | (0.68,0.97) | 0.025 | 0.96 | (0.86,1.08) | 0.49 | 0.88 |
| Barium | Early | 1.00 | (0.88,1.13) | 0.99 | 1.04 | (0.7,1.53) | 0.85 | 1.00 | (0.89,1.13) | 0.96 | 0.96 |
| | Late | 0.97 | (0.87,1.07) | 0.53 | 0.96 | (0.77,1.2) | 0.74 | 0.97 | (0.88,1.06) | 0.48 | 0.88 |
| Cesium | Early* | 1.63 | (0.90,2.96) | 0.11 | 1.44 | (0.53,3.87) | 0.47 | 1.58 | (0.95,2.63) | 0.079 | 0.35 |
| | Late | 1.07 | (0.82,1.39) | 0.63 | 1.06 | (0.55,2.03) | 0.86 | 1.07 | (0.84,1.36) | 0.61 | 0.88 |
| Cobalt | Early | 1.09 | (0.91,1.30) | 0.34 | 1.06 | (0.51,2.19) | 0.88 | 1.09 | (0.92,1.29) | 0.33 | 0.55 |
| | Late | 0.9 | (0.72,1.12) | 0.33 | 0.77 | (0.50,1.18) | 0.23 | 0.87 | (0.71,1.06) | 0.16 | 0.60 |
| Copper | Early | 1.19 | (0.74,1.91) | 0.47 | 1.33 | (0.54,3.28) | 0.53 | 1.22 | (0.80,1.85) | 0.35 | 0.55 |
| | Late | 1.01 | (0.78,1.29) | 0.97 | 0.87 | (0.65,1.17) | 0.35 | 0.95 | (0.78,1.14) | 0.57 | 0.88 |
| Manganese | Early | 0.88 | (0.69,1.13) | 0.32 | 1.01 | (0.64,1.58) | 0.98 | 0.91 | (0.73,1.13) | 0.39 | 0.55 |
| | Late | 0.92 | (0.77,1.11) | 0.39 | 1.05 | (0.79,1.40) | 0.74 | 0.96 | (0.82,1.11) | 0.58 | 0.88 |
| Mercury | Early | 1.04 | (0.95,1.14) | 0.40 | 0.76 | (0.49,1.19) | 0.23 | 1.03 | (0.94,1.12) | 0.57 | 0.72 |
| | Late | 1.03 | (0.93,1.14) | 0.54 | 0.83 | (0.61,1.12) | 0.22 | 1.01 | (0.92,1.11) | 0.84 | 0.92 |
| Molybdenum | Early* | 1.54 | (0.91,2.59) | 0.10 | 1.32 | (0.59,2.95) | 0.49 | 1.47 | (0.95,2.28) | 0.083 | 0.35 |
| | Late | 1.04 | (0.82,1.33) | 0.72 | 0.93 | (0.71,1.22) | 0.62 | 0.99 | (0.83,1.19) | 0.95 | 0.96 |
| Nickel | Early** | 1.55 | (1.08,2.23) | 0.018 | 0.92 | (0.44,1.93) | 0.82 | 1.40 | (1.01,1.94) | 0.043 | 0.34 |
| | Late | 1.23 | (0.91,1.66) | 0.19 | 0.74 | (0.59,0.94) | 0.012 | 0.89 | (0.74,1.07) | 0.23 | 0.74 |
| Selenium | Early | 1.38 | (0.76,2.51) | 0.29 | 1.13 | (0.53,2.43) | 0.75 | 1.28 | (0.80,2.05) | 0.30 | 0.55 |
| | Late** | 1.31 | (0.64,2.69) | 0.46 | 0.85 | (0.82,0.88) | <0.001 | 0.85 | (0.82,0.88) | <0.001 | <0.001 |
| Thallium | Early | 0.71 | (0.41,1.23) | 0.22 | 0.96 | (0.56,1.66) | 0.89 | 0.83 | (0.56,1.22) | 0.34 | 0.55 |
| | Late | 1.01 | (0.87,1.19) | 0.87 | 1.02 | (0.71,1.47) | 0.92 | 1.01 | (0.88,1.17) | 0.85 | 0.92 |
| Tin | Early | 1.05 | (0.96,1.16) | 0.27 | 0.83 | (0.54,1.3) | 0.42 | 1.04 | (0.95,1.14) | 0.36 | 0.55 |

| Metal | Time | EARLI | | | MARBLES | | | Meta-Analysis | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | | RR | CI | P | RR | CI | P | RR | CI | P | FDR |
| | Late | 1.03 | (0.94,1.13) | 0.52 | 0.84 | (0.63,1.13) | 0.26 | 1.01 | (0.93,1.11) | 0.78 | 0.92 |
| Zinc | Early | 0.98 | (0.88,1.09) | 0.73 | 0.96 | (0.67,1.36) | 0.80 | 0.98 | (0.88,1.08) | 0.69 | 0.84 |
| | Late** | 0.97 | (0.95,0.98) | <0.001 | 0.92 | (0.82,1.03) | 0.14 | 0.97 | (0.95,0.98) | <0.001 | <0.001 |

*Symbols: ** meta-analysis p-value < 0.05. * meta-analysis p-value < 0.1.*

**Figure 1.** Spearman correlations of urinary metals concentrations, measured during early pregnancy, stratified by cohort. The upper right triangle shows the EARLI cohort. The lower left triangle shows the MARBLES cohort. Metals are represented by their chemical symbol along the diagonal.

**Figure 2**. Adjusted risk ratios for the associations between maternal urinary metals concentrations measuring during pregnancy and risk of autism spectrum disorder, relative to typically developing. Antimony, cadmium, chromium, and lead compare over limit of detection vs under the limit of detection for that metal. Remaining metals show the risk ratio for a doubling in metal concentration. Analyses were performed stratified by cohort (EARLI and MARBLES) and then meta-analyzed across cohorts. Red denotes a nominal meta-analysis p-value < 0.05, and orange a nominal meta-analysis p-value < 0.10.

**Figure 3**. Adjusted risk ratios for the associations between maternal urinary metals concentrations measuring during pregnancy and risk of non-typically developing, relative to typically developing. Antimony, cadmium, chromium, and lead compare over limit of detection vs under the limit of detection for that metal. Remaining metals show risk ratio for a doubling in metal concentration. Analyses were performed stratified by cohort (EARLI and MARBLES) and then meta-analyzed across cohorts. Red denotes a nominal meta-analysis p-value < 0.05, and orange a nominal meta-analysis p-value < 0.10.