Full title:

Development of the Advised Protocol for OCT Study Terminology and Elements Anterior

Segment OCT extension reporting guidelines (APOSTEL-AS): Study protocol

Short title

APOSTEL-AS extension

Authors: Ameenat L Solebo PhD[1,2], Marcus Ang PhD[3,4], Alice Bellchambers MBBS[1,5], Colin

J Chu PhD[6,7], Prof Alastair K Denniston FRCOphth[8,9], Prof Laura E Downie PhD[10],Thomas

Evans[11], Alexander S Fraser MD[12], Scott Hau MCOptom[6], Prof Alex S Huang MD[13], Prof

Pearse A. Keane FRCOphth[6,7], Xiaoxuan Liu PhD[8,9,14], Prof Jodhbir S Mehta PhD[3,15],

Giovanni Ometto PhD[5,6,16], Prof Axel Petzold MD PhD[6,17,18], Edmund Tsui MD[19], Tamara S

Fraser MD[20], Prof Benjamin Xu PhD[21]

1. Population, Policy and Practice Department, UCL GOS Institute of Child Health,

London, UK; Great Ormond Street Hospital, London, UK,

2. Great Ormond Street NIHR Biomedical Research Centre, London, UK

3. Cornea and External Eye Disease Service and Refractive Service, Singapore National

Eye Center, Singapore

4. DUKE-NUS Department of Ophthalmology and Visual Sciences, Singapore

5. University Hospitals Sussex NHS Foundation Trust, UK

6. Moorfields NIHR Biomedical Research Centre, London, UK

7. UCL Institute of Ophthalmology, London, UK

8. NIHR Birmingham Biomedical Research Centre, Birmingham, UK

9. College of Medicine and Health, University of Birmingham, Birmingham, UK

10. Department of Optometry and Vision Sciences, The University of Melbourne, Parkville, Victoria, Australia

11. Department of Ophthalmology, Sheffield Teaching Hospitals NHS Foundation Trust, Sheffield, UK

12. Trinidad Eye Hospital, San Fernando, Trinidad and Tobago

13. Hamilton Glaucoma Center, The Viterbi Family Department of Ophthalmology, Shiley Eye Institute, University of California, San Diego, CA, USA

14. Birmingham Health Partners Centre for Regulatory Science and Innovation

15. Singapore Eye Research Institute, Singapore.

16. City St George's, University of London, Department of Optometry & Visual Sciences, London, UK.

17. University College London, UCL Institute of Neurology, London UK

18. Departments of Ophthalmology and Neurology, Amsterdam UMC, NL.

19. Ocular Inflammatory Disease Center, UCLA Stein Eye Institute, David Geffen School of Medicine at UCLA, Los Angeles, California, USA

20. Mid South Essex NHS Foundation Trust, Essex, UK

21. USC Roski Eye Institute, Keck School of Medicine at USC, Los Angeles, California, USA

Corresponding author: Ameenat Lola Solebo, UCL GOS Institute of Child Health, 30 Guilford Street, London, WC1N 1EH, UK. Email a.solebo@ucl.ac.uk

Abstract

Background:

Anterior segment optical coherence tomography (AS-OCT) is emerging as a valuable

diagnostic, monitoring and predictive tool. Clinical utility has been suggested for ophthalmic

disorders such as glaucoma, corneal disease, cataract and uveitis, which taken together

comprise the majority of the blinding conditions affecting working age individuals globally.

The current absence of guidelines for reporting OCT studies that quantitatively assess the

anterior segment is an obstacle to reproducibility and interoperability. To provide this

guidance, we aim to extend the existing Advised Protocol for OCT Study Terminology and

Elements (APOSTEL) guidelines, to ensure applicability to AS-OCT.

Methods

In line with EQUATOR Network guidance for the development of reporting guidelines,

APOSTEL-AS will be developed through a staged consensus process involving literature

review and Delphi consensus across an international multi-disciplinary stakeholder group,

overseen by a multi-disciplinary multi-national Steering Committee. The systematic scoping

review will be used to generate candidate items, support the development of a consensus

nomenclature for AS-OCT representation of ocular structure, and to form Delphi group

membership. Delphi methodology, used to consider items for inclusion, rewording or

exclusion, will be undertaken in line with ACCORD (ACcurate COnsensus Reporting

Document) guidance, with at least two rounds of Delphi survey, inclusion consensus

threshold set at 80%, and steering committee reviews between rounds. The resultant

APOSTEL-AS guideline will undergo piloting before dissemination of the final version.

Discussion

The APOSTEL-AS checklist, with minimum and recommended items to be reported about

study methods, should provide timely support for researchers to ensure future standardisation,

74 interoperability and reproducibility of reported work, hastening implementation and the

75 translation of knowledge into clinically beneficial action.

Summary

Anterior segment optical coherence tomography (AS-OCT) is emerging as a valuable diagnostic, monitoring and predictive tool. To address the current absence of guidelines for reporting OCT studies which quantitatively assess the anterior segment, we aim to extend the existing Advised Protocol for OCT Study Terminology and Elements (APOSTEL) recommendations, to enable applicability to studies involving AS-OCT. Following steering committee formation, literature review will be used to generate items additional to those within the current APOSTEL guidance, to create a consensus nomenclature and the Delphi membership. A multi-round Delphi exercise, with the inclusion consensus threshold set at 80%, will be used to generate reporting guidance items, with process overview by the steering committee. The resultant guideline will be piloted prior to dissemination, providing a timely support tool for researchers to ensure future standardisation, interoperability and reproducibility of reported work, hastening the implementation and the translation of knowledge on the utility of AS-OCT into clinical benefit.

Introduction

Optical coherence tomography (OCT) has changed the face of ophthalmic clinical practice

and medical research. The research and clinical utility of ocular OCT biomarkers has been

demonstrated within ophthalmology,(1–3) and beyond the eye.(4) Quantitative data from

posterior segment ocular OCT, such as retinal nerve fibre layer or central macular thickness,

have been used as diagnostic, predictive, prognostic and monitoring biomarkers for

inflammatory, degenerative, vascular, and metabolic disorders.(4–7)

The validation and adoption of these biomarkers has been enabled by the standardisation and

harmonisation of the reporting of research findings within the underpinning evidence base.

The 2016 and 2021 Advised Protocol for OCT Study Terminology and Elements

(APOSTEL) and APOSTEL2.0 recommendations(8,9) for reporting quantitative retinal OCT

studies strengthened reproducibility and interoperability across OCT studies of the posterior

segment of the eye. However, some aspects of the current APOSTEL guidance are not

applicable to anterior segment OCT (AS-OCT) studies. AS-OCT is being used to derive

quantitative data from images of the normal physiology of the anterior ocular structure, data

which carry clinical utility for the diagnosis and management of diseases involving those

structures. These diseases include tear film disorders, corneal pathology, glaucoma, cataract

and anterior uveitis,(10–16) disorders which, collectively, constitute the most common causes

of working aged blindness globally.(17)

In order to improve the reproducibility and interoperability of AS-OCT studies, we aim to

develop a checklist of the core information that should be provided when reporting studies in

which the quantitatively assessment of AS OCT structures is undertaken. This checklist will

be developed using the framework of the current APOSTEL recommendations.(8)

Materials and methods

APOSTEL-AS has been registered as a reporting guideline under development on the

EQUATOR (Enhancing the QUality and Transparency Of health Research) library of

reporting guidelines and will be developed in accordance with the EQUATOR Network's

methodological framework. A steering committee with representation of expertise in tear

film, corneal, lens, glaucoma and uveitic disorders, expertise in multinational, multi-

disciplinary consensus exercises, and including investigators involved in developing the

APOSTEL guidelines has been formed to oversee the conduct and methodology of the study.

A multi-stage Delphi consensus approach will be used (figure 1).

Figure 1. APOSTEL-AS development process

Further Delphi rounds may be needed to reach consensus

Literature review and candidate item generation

An initial list of candidate items for the APOSTEL-AS checklist will be generated through

review of the published literature and consultation with the steering committee (SC). A

database search will be performed using the terms "anterior segment optical coherence

tomography" (see supplemental document, search approach) to identify eligible studies.

Eligibility criteria comprise publications reporting the analysis of quantitative data derived

from AS-OCT acquired from human eyes of living participants. Specific considerations for

Screening, selection AS-OCT will be identified from these studies and extracted as potential

reporting items using a specific data collection form *developed de novo* by the SC. and data

extraction to the data collection form (DCF) will be undertaken independently by at least two

investigators. Items will be reframed as candidate reporting items ahead of the consensus

process. The SC will also create a consensus nomenclature for ocular anterior segment

structures visible on axial cross sectional OCT imaging. This consensus nomenclature will be

developed by the APOSTEL-AS SC using the nominal group technique, with consensus

reached through discussion and majority voting.

Delphi consensus group membership:

Membership of the Delphi consensus group will be informed by the findings of the

systematic review. Corresponding authors of studies reporting quantitative anterior segment

OCT results will be contacted by email, requesting a response within six weeks. Reminder

emails will be sent three weeks later. Records will be kept of attempted email and confirmed

contact. Response rates will be reported using those who complete the Delphi as the

nominator, and those for whom receipt of the email can be confirmed as the denominator.

Other stakeholders to be identified and invited by the SC to join the Delphi consensus group

comprise eye healthcare professionals, patients, methodologists, statisticians, computer

scientists, industry representatives, health informaticists, and journal editors. Experts will be

allowed to support snowball recruitment (i.e., to suggest additional experts). Records will be

kept of all those successfully contacted, either directly by the SC or as corresponding authors,

or indirectly through snowball recruitment.

Online Questionnaire 1 (Delphi 1):

Participants will be given written information about the study and asked to state their

professional role, region of origin, and level of expertise (using years active). The candidate

reporting items generated by the systematic review will be presented to the Delphi consensus

group members, alongside the existing APOSTEL 2.0 items. Members will be asked to vote

on each item using a 3-point scale: not important, important but not critical, and important

and critical. They will also be asked about their agreement on the wording of each item and

given the opportunity to submit free text comments about suggested amendments to wording

of additional items, and invited to suggest additional items.

OQ2 (Expert Group virtual meeting):

We will then form a panel of experts comprising the SC and selected members of the Delphi

consensus group (ensuring representation of stakeholder groups), who will convene at a

virtual meeting. The aggregated results of the initial questionnaire, comments made by

participants) will be reviewed by the panel. Following open discussion of the Delphi results,

anonymous electronic voting will be used to include or exclude each reworded, amended, or

newly suggested item. An 80% threshold will be used to demonstrate majority consensus for

the inclusion of an item.

OQ3 (Delphi 2): The included items will be considered in a second Delphi round by the

original Delphi consensus group. In this round, participants will be asked to approve or reject

the list of APOSTEL-AS items. Items achieving consensus ≥80% will be included. Any item

which does not achieve consensus but is approved by the majority (51-79% inclusive) will be

reconsidered by the expert group for consideration following further amendment or

refinement, and representation in a subsequent round (Delphi 3).

OQ4 (Delphi 3): Those items that reached majority consensus but did not achieve the 80%

consensus threshold will be represented to the Delphi consensus group following amendment

by the SC for one final Delphi round. Participants will again be asked to approve or reject the

items. Again, items achieving consensus ≥80% will be included. Any item that does not

achieve consensus but is approved by the majority (51-79% inclusive) will be reconsidered

within subsequent Delphi rounds (OQ5+) by the expert group for consideration for further

amendment or refinement prior to inclusion in the penultimate list.

Guideline pilot

The penultimate guideline will be checked by the Operations team (ALS, SD, AB, AF, TT) to

ensure its clarity. It will then be presented to at least five stakeholders for piloting. The

stakeholders participating in this pilot will comprise Delphi consensus participants who were

not part of the expert group, or additional external experts identified following the start of the

Delphi process. The Operations team will use feedback from this pilot to refine the guideline

and produce the final version of the checklist.

Ethics approval for this work has been granted by the University College London

Institutional Review Board (UCL Research Ethics Committee Approval ID Number:

26253/001). Informed electronic consent will be obtained from consensus group members.

Discussion

The APOSTEL-AS recommendations will be an extension to the current APOSTEL

guidelines, adding specific parameters for the reporting of AS-OCT data. AS-OCT is

increasingly being used to derive quantitative data from images of normal physiology and for

diseases involving the anterior segment of the eye. There are several differences in the

acquisition and analysis of anterior versus posterior segment OCT images. Whilst posterior

segment OCT (PS-OCT) concerns images of tissues with similar refractive indices and

typically measuring no more than 1mm in depth (as measured from the inner limiting

membrane to the sclera), with the *'en face'* area of interest typically being 4mm^2, the ocular

structures imaged in AS-OCT are arranged over a greater depth (at least 10mm, as measured

from the corneal epithelium to the anterior hyaloid area) and with a much wider area of

interest depending on the clinical or research need.(18) Furthermore, whilst the posterior

segment tissues minimally scatter light, there are unique features related to light

transmissibility of the different anterior ocular structures, from the optically clear cornea to

the optically opaque sclera.(18) These considerations differentiate AS-OCT from PS-OCT,

but it is also likely that these challenges result in heterogeneity of study methods, including

acquisition processes, image quality assessment, data extraction and image analysis across

the AS-OCT literature. APOSTEL-AS will provide timely support for users of AS-OCT, to

ensure future standardisation, interoperability and reproducibility of reported research and

clinical findings.

A limitation of the approach described in this protocol is the restriction of the identification

of potential consensus group members to those researchers who have published eligible

studies within the preceding four years. High impact, novel, innovative imaging studies may

require lengthy recruitment periods, which may mean that our time-restricted approach does

not involve scientists whose work is not yet disseminated through peer-reviewed

publications. The use of 'snowball' extension (i.e. consensus group members identifying

other potential members) should help to address this limitation. Strengths of the approach

include the involvement of a wide range of stakeholder expertise and experience, across

multiple countries, with collection of information about their degree of experience and

expertise to inform the contextualisation of responses. Development of items will be

underpinned by a comprehensive systematic review, with collection of data from manuscripts

published over the last 20 years. It is acknowledged that this comprehensiveness may result

in the collection of historical data from imaging platforms that are no longer in use, due to

being superseded by more contemporary devices.

Several consensus methods exist for the capture and description of consensus across multi-

disciplinary groups. The nominal group technique (NGT) was selected for the initial activities

of the SC, in view of its structured sequence of events, principal of equal participation, and

promotion of diversity in opinion, which should support the work of this smaller group.(19)

NGT consensus exercises can be time consuming, particularly for large groups, but this

process will benefit from facilitation undertaken by subject matter experts with experience in

supporting consensus initiatives.(20,21) The selection of the Delphi method for the

development of the checklist enables this activity to be undertaken by a large, geographically

and scientifically diverse group with different degrees and spheres of experience and

expertise. A recently published EQUATOR (Enhancing the QUality and Transparency Of

health Research) guideline for reporting consensus-based methods in biomedical research and

clinical practice (the ACcurate COnsensus Reporting Document, or ACCORD) will be used

to report the study findings.(19)

The original APOSTEL checklist was developed following concerns around the absence of

consensus around quality control criteria for posterior segment OCT imaging.(9) Similar

concerns are emerging for AS-OCT imaging, with recognition of this issue being a particular

obstacle to the potentially transformative role of artificial intelligence (AI) approaches to

image analysis and the development of imaging-based biomarkers.(22,23) AI approaches

hold the promise of dramatically increasing the utility of clinical images, with deep learning

(DL) based image analysis in particular proving non-inferior to expert human

assessment.(12,24,25) Human analysis of the quantitative information that can be extracted

from anterior segment structures is subjective, open to inter- and intraobserver variability,

and time consuming.(26) DL image analysis methods would provide objective, repeatable,

reliable OCT biomarkers, but these approaches require large (or harmonised), diverse,

annotated datasets. The current absence of explicit and comprehensive reporting of AS-OCT

study methods is an obstacle to comparison or collation of existing annotated imaging

datasets. Through the wide dissemination and uptake of the APOTEL-AS guidance, enabled

by ensuring accessibility in open access formats and translation into other languages as

directed and supported by stakeholders, this obstacle should be addressed.

Authors' contributions

274 Conceptualisation and manuscript drafting: AL Solebo

275 All authors contributed to study design, manuscript editing and final approval for submission.

Acknowledgments

278 We thank the Enhancing the Quality and Transparency of health research Network

279 (EQUATOR Network) team for their support with study development. This work was

280 supported by an NIHR Clinician Scientist award (CS-2018-18-ST2-005) and Fight for Sight

281 (SGAGUK2207).

References

1. Hui VWK, Szeto SKH, Tang F, Yang D, Chen H, Lai TYY, et al. Optical Coherence

Tomography Classification Systems for Diabetic Macular Edema and Their

Associations With Visual Outcome and Treatment Responses - An Updated Review.

Asia Pac J Ophthalmol (Phila). 2022 Jun 1;11(3):247–57.

2. Hood DC, La Bruna S, Tsamis E, Thakoor KA, Rai A, Leshno A, et al. Detecting

glaucoma with only OCT: Implications for the clinic, research, screening, and AI

development. Prog Retin Eye Res. 2022 Sep;90:101052.

3. Virgili G, Menchini F, Casazza G, Hogg R, Das RR, Wang X, et al. Optical coherence

tomography (OCT) for detection of macular oedema in patients with diabetic

retinopathy. Cochrane Database Syst Rev. 2015 Jan 7;1:CD008081.

4. Wagner SK, Fu DJ, Faes L, Liu X, Huemer J, Khalid H, et al. Insights into Systemic

Disease through Retinal Imaging-Based Oculomics. Transl Vis Sci Technol. 2020 Feb

12;9(2):6.

5. Bostan M, Li C, Sim YC, Bujor I, Wong D, Tan B, et al. Combining retinal structural

and vascular measurements improves discriminative power for multiple sclerosis

patients. Ann N Y Acad Sci. 2023 Sep 1;

6. Chen S, Zhang D, Zheng H, Cao T, Xia K, Su M, et al. The association between retina

thinning and hippocampal atrophy in Alzheimer's disease and mild cognitive

impairment: a meta-analysis and systematic review. Front Aging Neurosci.

2023;15:1232941.

7. Wagner SK, Romero-Bascones D, Cortina-Borja M, Williamson DJ, Struyven RR,

Zhou Y, et al. Retinal Optical Coherence Tomography Features Associated With

Incident and Prevalent Parkinson Disease. Neurology. 2023 Aug 21;

8. Aytulun A, Cruz-Herranz A, Aktas O, Balcer LJ, Balk L, Barboni P, et al. APOSTEL

2.0 Recommendations for Reporting Quantitative Optical Coherence Tomography

Studies. Neurology. 2021 Jul 13;97(2):68–79.

9. Cruz-Herranz A, Balk LJ, Oberwahrenbrock T, Saidha S, Martinez-Lapiscina EH,

Lagreze WA, et al. The APOSTEL recommendations for reporting quantitative optical

coherence tomography studies. Neurology. 2016 Jun 14;86(24):2303–9.

10. Chen C, Lee Y, Gerber MJ, Cheng H, Yang Y, Govetto A, et al. Intraocular robotic

interventional surgical system (IRISS): Semi-automated OCT-guided cataract removal.

The International Journal of Medical Robotics and Computer Assisted Surgery. 2018

Dec 28;14(6).

11. Sharma R, Sharma A, Arora T, Sharma S, Sobti A, Jha B, et al. Application of anterior

segment optical coherence tomography in glaucoma. Surv Ophthalmol. 2014

May;59(3):311–27.

12. Santos VA dos, Schmetterer L, Stegmann H, Pfister M, Messner A, Schmidinger G, et

al. CorneaNet: fast segmentation of cornea OCT scans of healthy and keratoconic eyes

using deep learning. Biomed Opt Express. 2019 Feb 1;10(2):622.

13. Etherton K, Rahi JS, Petrushkin H, Dick AD, Akbarali S, Pattani R, et al. Quantitative

and qualitative assessment of anterior segment optical coherence tomography capture

of disease state in childhood anterior uveitis. Br J Ophthalmol. 2023 Jul;107(7):966–

72.

14. Tsui E, Chen JL, Jackson NJ, Leyva O, Rasheed H, Baghdasaryan E, et al.

Quantification of Anterior Chamber Cells in Children With Uveitis Using Anterior

Segment Optical Coherence Tomography. Am J Ophthalmol. 2022 Sep;241:254–61.

15. Bao YK, Xu BY, Friedman DS, Cho A, Foster PJ, Jiang Y, et al. Biometric Risk

Factors for Angle Closure Progression After Laser Peripheral Iridotomy. JAMA

Ophthalmol. 2023 Jun 1;141(6):516–24.

16. Xu BY, Friedman DS, Foster PJ, Jiang Y, Porporato N, Pardeshi AA, et al. Ocular

Biometric Risk Factors for Progression of Primary Angle Closure Disease: The

Zhongshan Angle Closure Prevention Trial. Ophthalmology. 2022 Mar;129(3):267–75.

17. Flaxman SR, Bourne RRA, Resnikoff S, Ackland P, Braithwaite T, Cicinelli M V, et

al. Global causes of blindness and distance vision impairment 1990–2020: a systematic

review and meta-analysis. Lancet Glob Health. 2017 Dec;5(12):e1221–34.

18. Xie X, Corradetti G, Song A, Pardeshi A, Sultan W, Lee JY, et al. Age- and refraction-

related changes in anterior segment anatomical structures measured by swept-source

anterior segment OCT. PLoS One. 2020 Oct 23;15(10):e0240110.

19. Gattrell WT, Hungin AP, Price A, Winchester CC, Tovey D, Hughes EL, et al.

ACCORD guideline for reporting consensus-based methods in biomedical research

and clinical practice: a study protocol. Res Integr Peer Rev. 2022 Dec 7;7(1):3.

20. Solebo AL, Rahi JS, Dick AD, Ramanan A V, Ashworth J, Edelsten C. Areas of

agreement in the management of childhood non-infectious chronic anterior uveitis in

the UK. British Journal of Ophthalmology. 2020 Jan;104(1):11–6.

21. Solebo AL, Kellett S, Rahi J, Pattani R, Edelsten C, Dick AD, et al. Development of a

Nationally Agreed Core Clinical Dataset for Childhood Onset Uveitis. Front Pediatr.

2022 Jun 21;10.

22. Tewarie P, Balk L, Costello F, Green A, Martin R, Schippling S, et al. The OSCAR-IB

Consensus Criteria for Retinal OCT Quality Assessment. PLoS One. 2012 Apr

19;7(4):e34823.

23. Petzold A, Albrecht P, Balcer L, Bekkers E, Brandt AU, Calabresi PA, et al. Artificial

intelligence extension of the OSCAR-IB criteria. Ann Clin Transl Neurol. 2021 Jul

19;8(7):1528–42.

24. Bolo K, Apolo Aroca G, Pardeshi AA, Chiang M, Burkemper B, Xie X, et al.

Automated expert-level scleral spur detection and quantitative biometric analysis on

the ANTERION anterior segment OCT system. Br J Ophthalmol. 2024 May

21;108(5):702–9.

25.   Xu BY, Chiang M, Pardeshi AA, Moghimi S, Varma R. Deep Neural Network for

362   Scleral Spur Detection in Anterior Segment OCT Images: The Chinese American Eye

363   Study. Transl Vis Sci Technol. 2020 Mar;9(2):18.

364   26.   Akbarali S, Rahi JS, Dick AD, Parkash K, Etherton K, Edelsten C, et al. Imaging-

365   Based Uveitis Surveillance in Juvenile Idiopathic Arthritis: Feasibility, Acceptability,

366   and Diagnostic Performance. Arthritis Rheumatol. 2021 Feb;73(2):330–5.

369 # **Supporting documents**

370 Supplemental document 1: Literature search strategy