A distinct T cell receptor signature associates with cardiac outcome in myocardial

infarction patients

Kenz Le Gouge*[1], Diyaa Ashour*[3,4], Margarete Heinrichs**[3,4], Paul Stys**[1], Pierre

Barennes**[1], Verena Stangl[5], Lavinia Rech[6], Gerald Hoefler[5], Karl Kashofer[5], Tobias

Gassenmaier[7], Valerie Boivin-Jahns[8], Roland Jahns[9], Ulrich Hofmann[3,4], Dominik Schmitt[3],

Anna Frey[3,4], Stefan Störk[3,4], Stefan Frantz[3,4], Peter P. Rainer,[6,10,11]§, Gustavo Campos

Ramos[3,4]§, Encarnita Mariotti-Ferrandiz[1,2]§

[1] Sorbonne Université, INSERM, UMRS959, Immunology-Immunopathology-Immunotherapy (i3),

Paris, France

[2] Institut Universitaire de France (IUF)

[3] Department of Internal Medicine I, University Hospital Würzburg, Würzburg, Germany

[4] Comprehensive Heart Failure Center, University Hospital Würzburg, Würzburg, Germany

[5] Diagnostic and Research Institute of Pathology, Medical University of Graz

[6] Division of Cardiology, Medical University of Graz, Graz, Austria

[7] Department of Diagnostic and interventional Radiology, University Hospital of Würzburg, Germany

[8] Institute of Pharmacology and Toxicology, University of Würzburg, Würzburg, Germany.

[9] Interdisciplinary Bank of Biomaterials and Data Würzburg (IBDW), University Hospital of Würzburg,

Würzburg, Germany.

[10] BioTechMed Graz, Graz, Austria

[11] St. Johann in Tirol General Hospital, St. Johann in Tirol, Austria

* Equally contributed

** Equally contributed

§ **Corresponding authors**

Peter P. Rainer
CardioScience Lab
Division of Cardiology
Medical University of Graz
A-8036 Graz, Austria
Phone: +43 316 385 12544
Fax: +43 316 385 13733
Email:
peter.rainer@medunigraz.at
35
36

Gustavo Ramos
Immunocardiology Lab
University Hospital Würzburg,
Department of Internal Medicine
I,
Am Schwarzenberg 15, Haus
A15
D-97078 Würzburg, Germany
Phone: +49 931 201 46477
Fax: +49 931 201 46485
Email: Ramos_G@ukw.de
48

Encarnita Mariotti-Ferrandiz
UMRS959, Immunology-
Immunopathology-
Immunotherapy lab
Sorbonne Université/INSERM
83 boulevard de l'Hôpital
75013 Paris
France
Phone: +33 1 42 17 74 68
Email:
encarnita.mariotti@sorbonne-
universite.fr
61

Keywords: Myocardial infarction, ejection fraction, T cell, TCR, lymphocytes.

Article type: Rapid communication

List of non-standard abbreviations:

MRI: cardiac magnetic resonance imaging, **ESV**: end-systolic volume, **EDV**: end-diastolic volume

FUP: follow-up visit, **HF:** Heart failure, **LVEF:** left-ventricular ejection fraction, **MI**: myocardial

infarction, **STEMI**: ST-elevation myocardial infarction, **TCR:** T cell receptor, **TRA**: T cell receptor,

alpha chain, **TRB**: T cell receptor, beta chain

Abstract

Myocardial infarction (MI) is associated with an inflammatory process mainly attributed to

innate immune components. Very recently, the role of T-cells in both inflammation and

healing has been suggested through various human and mouse studies. Previous studies

showed that CD4+ and CD8+ T cells affect post-MI repair but did not investigate how to

leverage T cell biology to predict post-MI outcomes in patients. For instance, the antigenic

trigger of T-cells is still unknown in human. Indeed, others and we identified T-cell specific

for myosin infiltrating the myocardium in mouse models of MI, recent studies identified

expanded clones in human myocardium, altogether suggesting a tissue-specific T-cell

activation. However, it is still unclear how acute post-MI immune responses shape long-term

cardiac functional outcomes in individual patients. In this study, we analyzed the role of T-

cell in predicting post-MI repair by analyzing the T-cell receptor (TCR) repertoire. Indeed, the

TCR repertoire is now considered as a marker of the clinical status of individuals. Previous

studies in infectious but also autoimmune contexts showed the potential of the TCR

repertoire to predict the disease. Therefore, assessing the dynamic changes in global TCR

repertoires may provide valuable information about the antigen-specific immune responses

underlying post-MI healing. In our study, we carefully selected patients that suffered from MI

on a prospective cohort. The TCR repertoire has been analyzed by next generation

sequencing at the index hospitalization with the aim to identify features predict of their

healing outcome assessed at 12 months post-MI. While no major variations have been found

in diversity of TCR gene usage, we identified unique TCR signatures predicting one-year

cardiac functional outcomes. Our result enables early immune-based risk stratification of MI

patients and calls for larger studies to develop novel predictive biomarkers and possibly new

therapeutics.

Main text

Inflammatory processes govern post-myocardial infarction (MI) healing and remodelling. Yet,

we cannot predict how acute post-MI immune responses shape long-term cardiac functional

outcomes in individual patients. Previous studies showed that CD4$^+$ and CD8$^+$ T cells are

key actors of the post-MI repair regulation(1, 2) but did not investigate how they could be

used as predictors of the cardiac outcome. Here, we sought to identify T cell receptor (TCR)

signatures predicting cardiac functional outcomes in a well-characterized MI patient
cohort(3).

TCRs are heterodimeric antigen-specific receptors expressed by T cells, generated through
somatic recombination of multiple gene segments in a process that generates a potential
repertoire of 10^{19} unique receptors(4). Thus, we hypothesised that analysing TCR
repertoires in patients after MI could provide valuable information on the individual
immunological status underlying post-MI healing outcomes.

First, we selected acute ST-elevation MI-patients from the ETiCS cohort (**E**tiology, **Ti**tre-
Course, and effect of autoimmunity on **S**urvival study(3), Würzburg arm) exhibiting reduced
left ventricular ejection fraction (LVEF) assessed by cardiac magnetic resonance (cMRI) on
day 4 post-MI (LVEF <50%, 54/150 patients). Amongst those, we further selected patients
with complete serial cMRI scans at baseline and 12 months of follow-up (FU) (38/54
patients) (**Figure 1A**). Next, we stratified these patients into "good" versus "poor" healers
based on a priori defined cMRI criteria(5)). In brief, patients showing a ΔLVEF <13%
between baseline and follow-up (FUP) were defined as "poor healers" (25/38), whereas
those showing greater improvement (ΔLVEF >13%) were considered "good healers" (13/38)
(**Figure 1B**). In addition to lower ΔLVEF values (P<0.0001), poor healers had significantly
greater Δ end-systolic volumes (ESV) (P<0.0001), whereas Δ end-diastolic volumes (EDV)
did not differ. Age, BMI, infarct size, and routine blood biomarkers were similar in poor and
good healers at baseline (**Figure 1B**). After defining groups of MI-patients with diverging
healing phenotypes, we extracted RNA of cryopreserved whole blood samples that were
collected at hospital admission from all 25 poor and 13 good healers (Nucleospin RNA blood
kit, Macherey-Nagel, Düren, Germany). Due to technical limitations inherent to the
processing of archived biomaterial, we obtained RNA with sufficient quality from 19 poor and
9 good healers, which were then used for RT-PCR, amplification of TCR alpha and beta
chains (TRA and TRB respectively), library preparation and sequencing(6).

To evaluate potential global changes in the TCR repertoires found in blood sampled at
baseline, we first measured the diversity of TCRs in each group by computing the Shannon
entropy and the gene usage frequency (**Figure 1C-D**), two canonical measures of immune
repertoire diversity(7). These basic analyses did not reveal any significant differences
between groups. We then hypothesised that distinct TCR repertoire composition may explain
healing outcomes. Thus, we devised a refined strategy to identify a set of TCRs that
effectively differentiates between cardiac functional outcomes (**Figure 1E**). Accordingly,
using statistical modelling from Pogorely *et al.*(8) (**Figure 1E,** panel 1), we computed the

probability of each TCR being present in the dataset (Pdata) and identified TCRs that were
statistically enriched compared to their expected generation (Pgen). Additionally, we
searched for TCRs that were differentially represented between good and bad healers(9)
(**Figure 1E,** panel 2). Finally, we selected TCRs that met both criteria: being more
represented in one group than the other and having a statistically higher probability of being
present in the dataset than expected (**Figure 1E,** panel 3). As a result, we discovered a
signature of 348 unique TCRs (**Figure 1F**). Hierarchical clustering based on the
presence/absence of this signature allowed a separation of good from poor healers (cluster
purity = 0.93). Interestingly, this signature comprising 237 unique TRA and 111 unique TRB
was primarily present among the good rather than the poor healers. We then assessed the
relevance of the signature by comparing its cumulative frequency in the ETiCS patients
versus a group of healthy subjects (n = 27). **Figure H** shows that the cumulative frequency
of the signature is significantly higher in good healers when compared with poor healers and
healthy volunteers ($p<0.001$ for both TRA and TRB chains). Moreover, the TCR signature
was more enriched in healthy subjects when compared with poor healers ($p<0.001$ for TRA
and $p<0.01$ for TRB), suggesting that the clones found in the signature are specifically
depleted in MI patients with poor cardiac functional recovery (**Figure 1G**).

In summary, we identified a distinct TCR signature associated with cMRI-assessed post-MI
cardiac healing outcomes. Specifically, we found that MI-patients showing a substantial post-
MI LVEF improvement were enriched for a set of 348 unique TCRs in the peripheral blood
already at baseline (hospitalization). These exploratory findings provide clinical evidence that
the individual TCR profile at early stages post-MI contributes to long-term healing outcomes
in humans. This supports the notion that modulation of T cell responses may eventually help
to improve post-MI remodelling(10) and further confirms that the circulating T cell
compartment is impacted by the ongoing cardiac repair. Moreover, the present findings
provide a compelling example on the value of immune-based diagnostics in cardiology. Still,
our study has some important limitations. First, it was a rather small - though well stratified
and characterized - retrospective patient collective. Moreover, since we analysed
cryopreserved whole blood samples, we were unable to phenotype T cell subsets that might
have accounted for the identified TCR signature. Furthermore, as it is currently not possible
to infer which antigens are recognised by the TCRs comprised in this distinct signature, the
mechanisms underlying their association with the cardiac functional outcomes remain
unknown. Still, our findings underline the potential predictive value of TCR repertoire
analyses in MI and might encourage further mechanistic investigations and confirmation in
larger patient cohorts.

Author contribution: RJ, VJ, DS, AA, SS recruited the patients for this study and coordinated the clinical study; TG analysed the cMRI images, MH, DA, LV, VS, PPR and GCR retrospectively stratified and selected samples; DA, KLG, PS, PB, and EMF processed the RNA samples, developed bioinformatic tools and analysed the TCR-seq data. DA, MH, KLG, GH, UH, SF, PPR, EMF, GCR made substantial contributions to the conception and design of the present work. KLG, DA, PPR, EMF, and GCR drafted the manuscript. PPR, EMF, and GCR supervised this study. All co-authors critically revised it, made significant intellectual contribution, and approved the manuscript in its final version to be published.

Acknowledgements:
This work benefited from equipment and services from the iGenSeq core facility at Institut du Cerveau (ICM, Paris, France) for all the TCR data production.

Data availability statement
The anonymised TCR-seq raw files can be made available upon reasonable request.

Conflict of interest statement
The authors declare no conflict of interest.

Funding statement
This work was supported by the European Research Area Network—Cardiovascular Diseases [ERANET-CVD JCT2018, AIR-MI Consortium to GCR (01KL1902), PPR (4168-B) and EMF (ANR-18-ECVD-0001)]. The ETiCS study was supported by the Bundesministerium für Bildung und Forschung (BMBF), grant "Molecular Diagnostics" FKZ 01ES0901 and FKZ 01ES0802 (to V.B.-J. and R.J.). KLG doctoral fellowship was supported by ERANET-CVD JCT2018 (ANR-18-ECVD-0001) and additional support from Sorbonne Université. EMF work was supported by the iReceptorPlus (H2020 Research and Innovation Programme 825821) and SirocCo (ANR-21-CO12-0005-01) grants and the Institut Universitaire de France. GCR is supported by the Interdisciplinary Centre for Clinical Research Würzburg [E-354]. GCR, UH, and SF received funding from the German Research foundation (through the Collaborative Research Centre "Cardio-Immune interfaces" SFB1525, grant number 453989101).

References

1. Forte E, et al. Cross-Priming Dendritic Cells Exacerbate Immunopathology After Ischemic

Tissue Damage in the Heart. *Circulation*. 2021;143(8):821–836.

2. Rieckmann M, et al. Myocardial infarction triggers cardioprotective antigen-specific T

helper cell responses. *Journal of Clinical Investigation*. 2019;129(11):4922–4936.

3. Deubner N, et al. Cardiac beta1-adrenoceptor autoantibodies in human heart disease:

rationale and design of the Etiology, Titre-Course, and Survival (ETiCS) Study. *Eur J Heart*

Fail. 2010;12(7):753–762.

4. Bradley P, Thomas PG. Using T Cell Receptor Repertoires to Understand the Principles

of Adaptive Immune Recognition. *Annu Rev Immunol*. 2019;37(1):547–570.

5. Bulluck H, et al. Defining left ventricular remodeling following acute ST-segment elevation

myocardial infarction using cardiovascular magnetic resonance. *Journal of Cardiovascular*

Magnetic Resonance. 2017;19(1):26.

6. Barennes P, et al. Benchmarking of T cell receptor repertoire profiling methods reveals

large systematic biases. *Nat Biotechnol*. 2021;39(2):236–245.

7. Six A, et al. The Past, Present, and Future of Immune Repertoire Biology – The Rise of

Next-Generation Repertoire Analysis. *Front Immunol*. 2013;4.

https://doi.org/10.3389/fimmu.2013.00413.

8. Pogorelyy MV, et al. Method for identification of condition-associated public antigen

receptor sequences. *eLife*. 2018;7:e33050.

9. Emerson RO, et al. Immunosequencing identifies signatures of cytomegalovirus exposure

history and HLA-mediated effects on the T cell repertoire. *Nat Genet*. 2017;49(5):659–665.

10. Zhao TX, et al. Regulatory T-Cell Response to Low-Dose Interleukin-2 in Ischemic Heart

Disease. *NEJM Evidence*. 2022;1(1):EVIDoa2100009.

Figure 1: A distinct T cell receptor signature associates with cardiac functional outcomes in myocardial infarction patients. (A) Patient selection and stratification into good vs poor healers based on cMRI findings. The number of patients is indicated within the boxes (**B**) cMRI findings, including LVEF at baseline and FUP 12 months in good vs poor healers, ΔLVEF (%), ΔESV and ΔEDV (mL), and distribution of confounding factors between groups (BMI: body-mass index; LDH: lactate dehydrogenase; Time: time from pain to intervention; LGE: infarct size assessed by late gadolinium enhancement, CRP: C-reactive protein, WBC: white blood cell count; RBC: red blood cell count, Hgb: hemoglobin; Plat: platelets count; NTproBNP; N-terminal pro-brain natriuretic peptide; Chol: cholesterol; Creat: creatinine; SBP: systolic blood pressure; Alb: albumin. Values are expressed as fold change from the average levels found in the "good healer" group). (**C**) Shannon index based on clone counts from aligned TCR samples for the alpha and beta chain. P-value was computed using Wilcoxon U-test. (**D**) Non-metric dimensional scaling of the TRB variable (TRBV) and joining (TRBJ) gene usage frequencies. (**E**) Strategy used to identify Complementarity-determining region 3 (CDR3, antigen-binding domain) signature. (**F**) Heatmap of the 348 CDR3 signature found in patients. CDR3, by column, were clustered independently by chain. (**G**) Sample's cumulative frequencies (in %) of the CDR3 from the signature. Healthy volunteers are depicted in yellow.