Associations between Social Contact, Sleep and Dietary Patterns Among
Children: A Cross-Sectional Study

Christophe Mühlematter[1], Matthieu Beaugrand[1], Andjela Markovic[1], Salome Kurth[1,2]

[1] Department of Psychology, University of Fribourg, Fribourg, Switzerland

[2] Department of Pulmonology, University Hospital Zurich, Zurich, Switzerland

Corresponding author:

Salome Kurth, PhD

Department of Psychology, University of Fribourg

Rue de Faucigny 2, 1700 Fribourg

Switzerland

salome.kurth@unifr.ch

Number of words: 4090

Number of references: 25

Running head: Social Contact, Sleep, and Diet Patterns in Childhood

Abstract

Social isolation in adults can be associated with poor sleep and altered eating behavior. This
study aimed to investigate the interactions between the extent of social contact, eating
behavior and sleep in infants and preschool children. In an observational study, caregivers of
726 children aged 0-6 years provided information on sleep (i.e., duration, latency, bedtimes
and nighttime awakenings), eating behaviors (i.e., meal size, consumption of sweet snacks,
salty snacks, fruits and vegetables) and social contact (i.e., quarantine status, household
size, social activities) during the Covid-19 pandemic (April 2020). In infants (0-3 years), the
change in meal size and consumption of snacks, fruits, and vegetables did not significantly
relate to the extent of social contact. For preschool children (3-6 years), a trend was
observed, suggesting that quarantine status was associated with increased meal size.
Changes in sleep duration, sleep latency, bedtimes and nighttime awakenings from before to
during the pandemic were not significantly linked to the 3 variables quantifying social contact
in both age groups. The study highlights that, contrary to expectations, the extent of social
contact has negligible associations with infants' and preschool children's sleep and eating
behaviors. These findings indicate that other factors beyond social isolation play a larger role
in shaping young children's eating habits and sleep patterns.
38
39
40 **Keywords:** Poor sleep, eating habits, social activities, early development, early childhood,
41 pandemic effects

Introduction

Social relationships are fundamental to well-being, mental and physiological health in adults (Cacioppo et al., 2002; Holt-Lunstad et al., 2015) and adolescents (Qualter et al., 2013). Reported associations between social relationships and physical health mainly focus on two constructs: social support and social integration. Social support represents the psychological and material resources provided by the social network to help cope with stress, while social integration represents the participation in social relationships (Cohen, 2004). More generally, frequency and duration of social interactions have been linked with physical and mental health through changes in health-related behaviors, such as diet, exercise, or smoking (Berkman et al., 2000).

Social isolation in adults generates higher risk for depressive symptomology (Cacioppo et al., 2010), mortality (Holt-Lunstad et al., 2015) and reduced sleep quality (Azizi-Zeinalhajlou et al., 2022; Cacioppo et al., 2002). One relevant factor intertwined with these observations may lie in altered eating behavior, based on indications that social isolation can be associated with eating disorders and binge eating (Mason et al., 2016; Wright & Pritchard, 2009). The effect of social isolation on sleep and eating behavior has been experimentally examined in Drosophila, demonstrating that separating adult flies from the group led to increased food consumption and shorter sleep (Li et al., 2021).

The COVID-19 pandemic created a unique quasi-experimental framework to investigate the effects of enforced confinement on a global scale, downsizing direct social contact within populations. In this framework, social isolation significantly impacted mental health of children and parents (Ehrler et al., 2022). In addition, children and adolescents consumed more snacks and processed foods during the pandemic, leading to weight gain (González-Monroy et al., 2021; Pujia et al., 2021). Children and adolescents also experienced shortening of sleep (Markovic et al., 2021; Teixeira et al., 2021). Yet controversial results remain, for example as part of the COVID-19 pandemic, an increase in fruit and vegetable consumption was reported in Italian children ages 5 to 14 years (Pujia et al., 2021). The pandemic-induced reduction of social contacts could explain the changes in eating and sleeping behavior towards more "unhealthy" patterns, which aligns with findings from observational studies on social isolation. Indeed, the contrasting findings might be explained through differences in effective social isolation determined by the household structure, family size, extent of isolation orders by country, or geography (urban vs. rural regions). Thus, quantifying individual levels of social contact might contribute to a more precise understanding of the interactions among social contact, eating and sleep behaviors in children. Moreover, these interactions have not yet been examined in preschool-age children, who might respond differently to changes in social contacts due to their reliance on interactions with caregivers rather school-age children.

In this investigation, we thus tested the hypothesis that infants and preschool children who experienced increased social isolation (i.e., being in quarantine, engaging in activities alone rather than together with other family members and small household size) have worsened sleep quality (shorter sleep duration, increased night awakenings, longer sleep latency), and more "unhealthy" eating patterns (increased meal size, decreased consumption of fruits and vegetables, and increased consumption of sweet and salty snacks).

Methods

Study Design

During the start of the COVID-19 pandemic from April to July 2020, caregivers of children
below 6 years were invited to participate in a study about children's sleeping behavior.
Recruitment was realized through large-scale distribution of digital flyers and a video
invitation, distributed on social media platforms, childcare institutions, medical practices, and
personal channels. 726 primary caregivers (687 females, 94.62%, age 38.14 ± 4.83 years,
mean ± SD) completed an online survey (SoSci Survey; Leiner, 2021) with questions about
their children's demographic variables, sleeping behavior, eating behavior and social
isolation. The survey was compiled in English, and then translated to Italian, Spanish,
French, and German by the authors, which was then checked by at least two native
speakers. For families with multiple children, questions were looped to assess data for each
child individually. We collected a total dataset of 522 (232 females, 49.89%) infants and
toddlers (between 0 and 35 months old), and 401 (155 females, 50.16%) preschool children
(age 36 to 72 months). The institutional ethics board of the University of Fribourg,
Switzerland, approved the study, and parents gave informed consent before completing the
survey.

Sleep Behavior change

According to age, children's sleep was either assessed with the Brief Infant Sleep
Questionnaire (BISQ; 0-35 mo) (Sadeh, 2004) or with the Children's Sleep Habits
Questionnaire (CSHQ; 36-72 mo) (Owens et al., 2000). Caregivers reported on the child's
sleep behavior before (retrospectively) and during the confinement. The change in four
aspects of sleep behavior was analyzed by means of a difference during-before the
lockdown: sleep duration, sleep latency, bedtimes and nighttime awakenings (as in Markovic
et al., 2021). For the infant group (0-36 months), parents reported infants' sleep duration for
nighttime and daytime sleep, which were summed for the total sleep time (in minutes),
number of nighttime awakenings, sleep latency (in minutes), and bedtimes (in minutes). In
the preschool group (36-72 months), sleep duration was quantified from the item "*My child
sleeps about the same duration each 24-h-day (nighttime sleep and naps combined)*", sleep
latency from the item "*My child falls asleep within 20 min after going to bed*", nighttime
awakenings from "*How often does your child wake up during the night*" and bedtime from
"*My child goes to bed at the same time at night*". Each item for preschool children was
captured with ratings from 1 to 5, such that 1 referred to never (being the case, *i.e.*, at 0
days/week), 2 to rarely (1 day/week), 3 to sometimes (2-4 days/week), 4 to usually (5-6
days/week) and 5 to always (7 days/week) being the case.

Social contact

With three variables we captured the individual extent of social contact during the
confinement: First, *quarantine status* indicating whether the caregiver was in quarantine at
the time of survey completion. Second, *household size* indicating the number of people living
in the household. And third, *social activities* where parents reported the duration their child
engaged in several activities (e.g., watching television or playing), and whether the child was
doing these activities alone or with someone else (parents, siblings or friends, for details see

Supplementary Material). The percentage of activity spent engaged with the parents
compared to alone was computed. For example, a child who spent half of the time engaging
in activities with someone else would be attributed a 50% participation rate in social
activities.

Eating behavior change

We assessed the change in meal size from before the confinement to during the
confinement with a 5-point Likert scale capturing whether the child's overall meal size was
"much smaller", "a bit smaller", "same as before", "a bit larger" or "much larger". Moreover,
parents rated the child's change in the consumption of salty and sweet snacks (between
main meals), and the consumption of fruits and vegetables by means of "much less", "a bit
less", "the same amount", "a bit more" or "much more".

Statistical Analysis

Statistical analyses were performed in R version 4.0.5 with packages dplyr (Wickham et al.,
2022), and figures were created by using ggplot2 (Hadley Wickham, 2016). Due to a small
number of responses for meal-sizes "much smaller" (n = 8 for the infant group, n = 5 for
preschool children) and "much larger" (n = 5 for infants, n = 4 for preschool children), the
scales were consolidated into three response groups "decreased", "no change", "increased".
Accordingly, the consumption of salty and sweet snacks, fruits and vegetables were
composed into a 3-point scale ("decreased", "no change", "increased").

To examine the relationship between the extent of social contact and sleep behavior, we
separately computed generalized linear models for the two age groups, with sleep duration,
sleep latency, bedtimes and nighttime awakenings as the outcome and the extent of social
contact (quarantine status, household size, social activities) as the predictor. For both age
groups, the association between the extent of social contact and eating was tested with
change in meal size, snacks, fruits and vegetables as outcome and social contact as
predictor. Age and sex were included as control factors, and parental stress was included as
a covariate in alignment with previous work demonstrating an association with children's
sleep (Markovic et al., 2021). Change in parental stress was captured with ratings from 1 to
5 with the question "*Did your behavior change across the time that passed since the
lockdown, such that your level of stress: 1 decreased a lot, 2 decreased a little, 3 did not
change, 4 increased a little, or 5 increased a lot.*" Alpha level was set to $p < 0.05$ and p-
values were corrected for multiple testing using the false discovery rate method (Benjamini &
Hochberg, 1995).

Results

Study population

Demographic variables revealed that household size and percentage of quarantined families
were similar in the age groups (Table 1). Percentage of shared activities was by 10% larger
in the infant group compared to the preschool children. The control variable change in
parental stress was comparable between groups.

Demographic and control variables	Infants	Preschoolers
Number of children	522	401
Number of girls	232 (49.89%)	155 (50.16%)
Age (years)	1.50 ± 0.82	4.48 ± 0.81
Change in parental stress	0.56 ± 1.08	0.48 ± 1.17

Social contact	Infants	Preschoolers
Household size (persons)	3.7 ± 0.82	3.7 ± 0.81
Quarantined families	187 (37.63%)	136 (35.51%)
Shared activities (percentage)	75.78 ± 19.23%	65.52 ± 19.16%

169
Table 1. Demographic and control variables included in the analysis, indicated for the infant
(n=522) and preschool-age group (n=401). Data are reported as mean and standard
deviation.

Sleep behavior
Before the lockdown, sleep duration ranged from 540 to 1020 minutes for infants, during the
lockdown it ranged between 510 and 990. Sleep latency before the lockdown was between 0
and 60 minutes, during the lockdown it was between 0 and 90 minutes. Before lockdown,
bedtime was between 6 and 11pm, it ranged from 6:30pm to 11:30pm during the lockdown.
Infants used to be awakened between 0 and 75 minutes during the night before the
lockdown, which was then comprised between 0 and 60 minutes during the lockdown.
Regular sleeping times on 5-6 days a week were reported for 59.93% of preschool children
before the lockdown and 53.19% of preschool children after the lockdown. The number of
times they fell asleep in less than 20 minutes 5-6 days a week was 46.80% before lockdown
and went down to 38.72%. Before the lockdown, for most preschoolers, bedtimes were
regular 5-6 days a week 63.29%, which decreased to 49.83% during the lockdown. For
40.74% they woke up during the night once a week before the lockdown and 37.37% during
the lockdown.

Next, the change in sleep from before to during the pandemic was computed. This revealed
that more infants experienced a decrease in sleep duration (21.05% increase, 30.89%
decrease), a prolongation of sleep latency (34.79% longer, 11.52% shorter), later bedtimes
(51.14% later, 13.01% earlier), and more night awakenings (19.86% increase, 13.47%
decrease). For preschool children, the shift was in the same direction with less regular sleep
duration (6.73% more regular, 17.17% less regular), a more frequent long sleep latency
(27.95% more frequent, 8.08% less frequent), more irregular bedtimes (29.29% less regular,
6.73% more regular), and more frequent night awakenings (16.16% more frequent, 8.75%
less frequent, Table 2).

Sleep Variables	Before		During		Change	
	Infants	Preschoolers	Infants	Preschoolers	Infants	Preschoolers
Sleep duration	770.25 ± 130.63	4.26 ± 0.62	752.08 ± 119.17	4.1 ± 0.77	-17.96 ± 95.32	-0.16 ± 0.69
Sleep latency	49.8 ± 125.89	3.93 ± 1.00	63.32 ± 165.41	3.63 ± 1.11	15.37 ± 103.19	-0.29 ± 0.85
Bedtime	1147.19 ± 263.58	2.29 ± 1.02	1167.85 ± 270.32	2.6 ± 1.17	20.31 ± 259.3	0.31 ± 0.86
Night awakenings	30.41 ± 79.46	2.17 ± 1.24	33.59 ± 75.34	2.25 ± 1.45	3.18 ± 40.03	0.08 ± 0.97

201 Table 2. Sleep variables (mean ± standard deviation) in the infant (n=522) and preschool-
202 age group (n=401), and their difference computed as during-before the lockdown from BISQ
203 (infants, reported in minutes) or CSHQ (preschoolers, reported as frequency).

204
205

Eating behavior

Eating variables	Infants			Preschoolers		
	Decreased	Similar	Increased	Decreased	Similar	Increased
Meal size	68 (16.71%)	276 (67.81%)	63 (15.48%)	37 (12.67%)	204 (69.86%)	51 (17.47%)
Salty snacks	66 (19.58%)	239 (70.92%)	32 (9.5%)	56 (19.51%)	200 (69.69%)	31 (10.8%)
Sweet snacks	87 (25.82%)	222 (65.88%)	28 (8.31%)	105 (36.33%)	165 (57.09%)	19 (6.57%)
Fruits	88 (25.29%)	204 (58.62%)	56 (16.09%)	56 (19.18%)	195 (66.78%)	41 (14.04%)
Vegetables	45 (12.93%)	250 (71.84%)	53 (15.23%)	34 (11.64%)	213 (72.95%)	45 (15.41%)

Table 3. Count and percentage for eating behavior variables in infants (n=522) and
preschool children (n=401).

The eating behavior overall remained similar for the majority, ranging from 57 to 73% across
all categories in infants and preschool children (Table 3). In the remainder, meal size
increased in 15-17% and decreased in 12-17% of infants and preschool children.
The consumption of Salty snacks increased in 10-11% and decreased in 19.5% of children.
For Sweet snacks, an increase was observed in 6.5-8.3% and a decrease in 25.8-36.3% for
both age groups. Children increased their consumption of Fruits in 14-16% of cases, while
19.2-25.3% decreased it. Vegetables eating increased in 15.2-15.4% of children and
decreased in 11.6-12.9%.

Social isolation and eating behavior

We then examined whether the extent of social contact related to eating behavior in both
age groups. Overall, neither the change in meal size, nor snacks, fruits and vegetables
were related to the infants' extent of social contacts (Table 4, all p>0.05). In preschool
children, a trend was observed in the association between change in meal size and social
contact through quarantine status (p=0.07). In other words, this would indicate that
preschool children in quarantine were more likely to increase their meal size, compared to
non-quarantined preschool children (Figure 1). The other measures of social contact were
neither related to meal size nor to type of consumed food (Table 5, all p>0.05).

The control variables included in our model (age, sex and parental stress) were not related to the children's eating behavior, with the exception that older children decreased meal size to a larger extent from before to during the pandemic (b = -0.217, p < 0.001), and ate more vegetables (b = 0.214, p < 0.001).

	Infants									
	Meal size		Vegetables		Fruits		Salty snacks		Sweet snacks	
	b	p	b	p	b	p	b	p	b	p
Household size	0.057	0.195	-0.028	0.831	-0.080	0.331	0.019	0.833	-0.059	0.543
Quarantine status	0.015	0.828	-0.008	0.919	-0.090	0.525	0.023	0.833	0.072	0.543
% Social activities	0.133	0.484	-0.404	0.143	-0.218	0.525	0.251	0.559	0.161	0.543
Sex	0.054	0.484	-0.136	0.155	-0.135	0.331	0.115	0.536	-0.029	0.728
Age	0.217	**<0.001**	0.214	**<0.001**	0.144	0.192	0.020	0.833	-0.143	0.113
Parental stress	0.025	0.484	-0.014	0.857	-0.015	0.721	0.008	0.833	0.033	0.543

Table 4. Association between extent of social contact and eating behavior in infants. Unstandardized beta coefficients (b) and corrected p-values (p) are indicated from the generalized linear model. Significant associations (p < 0.05) are presented in **bold**.

Figure 1. Change in meal size for the groups of quarantined and non-quarantined preschool children. Numbers refer to percentages and counts of preschool children for whom a change in meal size was reported.

| | Preschoolers | | | | | | | | | |
| | Meal Size | | Vegetables | | Fruits | | Salty snacks | | Sweet snacks | |
	b	p	b	p	b	p	b	p	b	p
Household size	0.013	0.736	0.001	0.978	-0.038	0.623	0.010	0.848	-0.016	0.869
Quarantine status	0.193	0.071	0.153	0.305	0.108	0.623	-0.111	0.483	-0.026	0.869
% Social activities	-0.108	0.736	-0.383	0.305	-0.408	0.608	0.261	0.483	-0.039	0.869
Sex	-0.028	0.736	0.113	0.386	0.055	0.623	-0.096	0.483	-0.115	0.869
Age	-0.069	0.182	-0.009	0.978	-0.004	0.936	-0.038	0.667	0.045	0.869
Parental stress	0.065	0.093	0.014	0.963	-0.024	0.623	-0.042	0.483	-0.020	0.869

246 Table 5. Association between extent of social contact and eating behavior in preschool
247 children. Unstandardized beta coefficients (b) and corrected p-values (p) from the
248 generalized linear model.

Social isolation and sleep behavior

We then examined whether the extent of social contact was related to children's sleep
behavior. Neither in infants nor in preschool children was the extent of social contact
associated with sleep variables (*i.e.*, duration, sleep latency, bedtimes nor with the number
of nighttime awakenings; Table 6, all p>0.05). Change in sleep behavior was not related to
age, sex and parental stress, except for a link between parental stress and sleep latency in
preschool children, such that increased stress related to longer sleep latency (b = 0.130, p =
0.043).

| | Infants | | | | | | | |
| | Sleep Duration | | Sleep latency | | Bedtimes | | Nighttime awakenings | |
	b	p	b	p	b	p	b	p
Household size	5.846	0.334	-6.137	0.521	7.993	0.760	-1.705	0.895
Quarantine status	-20.355	0.107	19.213	0.222	-31.804	0.760	10.324	0.174
% Social activities	-51.058	0.107	-2.123	0.932	38.103	0.760	1.655	0.895
Sex	-4.342	0.745	1.088	0.932	-13.249	0.760	1.184	0.895
Age	13.975	0.107	-2.401	0.932	-48.104	0.088	7.735	0.095
Parental stress	5.541	0.334	-9.111	0.222	-1.543	0.908	-0.773	0.895

258
259

| | Preschoolers | | | | | | | |
| | Sleep Duration | | Sleep latency | | Bedtimes | | Nighttime awakenings | |
	b	p	b	p	b	p	b	p
Household size	0.043	0.746	0.029	0.722	0.065	0.622	0.007	0.933
Quarantine status	-0.043	0.746	-0.249	0.139	0.271	0.117	-0.049	0.933
% Social activities	0.234	0.746	-0.221	0.636	-0.222	0.622	0.491	0.848
Sex	0.042	0.746	0.039	0.722	-0.096	0.622	-0.023	0.933
Age	0.001	0.978	-0.058	0.636	-0.003	0.965	-0.076	0.848
Parental stress	0.067	0.453	0.130	**0.043**	-0.099	0.117	-0.057	0.848

261 Table 6. Extent of social contact association with sleep behavior. Unstandardized beta
262 coefficients (b) and corrected p-values (p) from the linear mixed model.
263
264 Overall, the results indicate only a negligible relationship between the extent of social
265 contact with eating behavior in infant and preschool children during the COVID-19 pandemic.
266 This was indicated by a trend suggesting that being in quarantine related to increased meal
267 size in preschoolers.

Discussion

Social isolation in adults can be related to poor sleep (Azizi-Zeinalhajlou et al., 2022;
Cacioppo et al., 2002), and altered eating behavior (Mason et al., 2016; Wright & Pritchard,
2009). Yet, the interplay between those three factors remains to be elucidated in early
childhood, which was evaluated with this observational study during the COVID-19
confinement. Specifically, parents of 522 infants and 401 preschool children provided
information before and during the confinement, rating their children's social contacts, eating
behavior and sleep. Overall, our results indicate that children's social contact is not strongly
associated with sleeping behavior, nor with eating behavior, with the exception for a trend
such that quarantined preschool children tended to have increased meal size compared to
non-quarantined ones. The nuanced findings emphasize the need to better understand
which aspect of social isolation is impactful on children's health through changes in their
sleeping and eating behavior.

Contrary to our hypotheses, meal size, as well as the consumption of snacks, fruits and
vegetables, was not related to the extent of social contact, suggesting that factors beyond
social isolation play a role in dynamics of eating behavior in infants and preschoolers. We
observed that preschoolers being in quarantine had a heightened chance of increasing meal
size in relation to the confinement, yet this association remained as a trend after adjusting
for multiple comparisons. This relationship is concordant with findings demonstrating
increased subjective isolation in obese adolescents (Mason, 2020). Interestingly, our
observation prevailed only in preschool children, yet not in infants, which could point to age-
specific differences in need for social contact. Increased meal size and more unhealthy
eating behaviors during the pandemic have been reported in children 6 years and older,
suggesting that the effect of social isolation may be dependent on age (González-Monroy et
al., 2021; Pujia et al., 2021). Age effects could also be attributed to the extent of contacts
with people outside of the immediate household. For infants, the main social contacts are
their primary caregiver(s), while preschool children are more likely to have extended
contacts with other adults. Thus, as the change in social contact with primary caregiver(s)
during the pandemic did not change, infants' changes were probably less extreme. Further,
we lacked information on other aspects of eating behavior such as the timing of meals which
could also be affected by the pandemic and is related to sleeping behavior (Mühlematter et
al., 2023). Moreover, eating behavior is the primary factor determining the composition of the
gut microbiota, and interestingly animal research observed associations between social
isolation, the gut microbiota and the brain (Dunphy-Doherty et al., 2018; Lopizzo et al.,
2021). Thus, the gut microbiota could be an interesting candidate in future research on the
interplay between social isolation, sleeping and eating behavior. Our results show that only
quarantine could have an effect on eating behavior, and only in children aged 3 to 6 years
old.

307
In line with previous studies (Markovic et al., 2021), sleep behavior shifted towards a
decreased sleep quality (i.e., shorter sleep duration, longer sleep latency, more nighttime
awakenings). Similar to eating, children's changes in sleep variables were not strongly linked
to the degree of their social contacts. Interestingly, this contrast with findings in adults.
Effects of adults' social isolation on sleep quality are well-documented, showing, for example
that subjective loneliness often accompanies subjective poor sleep quality (Bhatti & Haq,
2017; Cacioppo et al., 2002; McLay et al., 2021). Thus, the difference to children's sleep
might be attributable to other factors, such as parents engagement in mindfulness
techniques or the presence of pets, as previously demonstrated (Markovic et al., 2021).
Another possibility is that children's interactions with their parents sufficiently safeguarded
them from potential negative effects of reduced social contact on their sleep. Moreover, our
study used objective ratings of social contact, whereas studies involving adults relied on
individual perceptions of isolation (Cacioppo et al., 2002, 2010). Thus, the adverse outcomes
of social isolation might be subjectively biased in adults, making results not directly
comparable. To summarize our findings, sleep behavior change during the pandemic was
not dependent on externally reported measures of social contact.
324
Age, sex, and parental stress had a small impact on children's both eating behavior and
sleep in our dataset. Higher parental stress marginally increased the chance of increased
meal size in the preschool group. Thus, parental stress, depending on the age group,
remains an important factor to consider in the framework of children's eating behavior. In
alignment with this, stress exposure in adults can increase food intake, with specifically
increasing the preference for high-fat, high-sugar foods (Tomiyama, 2019). Future
investigations to elucidate the impact of stress on young children's eating behaviors could
inform strategies for mitigation.
333
In conclusion, this study sheds light on the association of social isolation with sleeping and
eating behavior in children up to 6 years old. We report that, in contrast to findings in
adolescents and adults, social contact beyond the family household is only negligibly
associated with young children's sleep and eating behavior. Thus, more research is needed
to gain a deeper understanding of the maturational transition and the relationships between
social contact, sleep patterns and eating behavior. This is a significant concern, particularly
in light of the rising rates of obesity (Pan et al., 2021) and sleep problems in children
observed in recent decades (Chen et al., 2021). While these issues have complex causes,
social isolation might potentially contribute to them, making it a worthwhile focus for
addressing mental, metabolic and chronobiological health.
344

Acknowledgments

We thank Océane Minot, Debora Castiglioni, Valentine Horii, and Dr. Maria Berrozpe for
assistance in data collection. Our appreciation goes to the participants who contributed to
the study. We recognize the support from the Swiss National Science Foundation (PCEFP1-
181279 to SK) and the University of Zurich, Medical Faculty (Forschungskredit FK-18-047;
Stiftung für wissenschaftliche Forschung STWF-17-008 to SK).

Disclosures

The authors have no conflicts of interest to declare.

Author Contributions

CM, SK, AM, and MB designed and conducted the study. CM performed the analyses. CM and SK wrote the manuscript. All authors discussed the results and contributed to the final manuscript.

References

Azizi-Zeinalhajlou, A., Mirghafourvand, M., Nadrian, H., Samei Sis, S., & Matlabi, H. (2022). The contribution of social isolation and loneliness to sleep disturbances among older adults : A systematic review. *Sleep and Biological Rhythms*, *20*(2), 153-163. https://doi.org/10.1007/s41105-022-00380-x

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate : A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B (Methodological)*, *57*(1), 289-300.

Berkman, L. F., Glass, T., Brissette, I., & Seeman, T. E. (2000). From social integration to health : Durkheim in the new millennium p. *Social Science*.

Bhatti, A. B., & Haq, A. U. (2017). The Pathophysiology of Perceived Social Isolation : Effects on Health and Mortality. *Cureus*, *9*(1), e994. https://doi.org/10.7759/cureus.994

Cacioppo, J. T., Hawkley, L. C., Crawford, L. E., Ernst, J. M., Burleson, M. H., Kowalewski, R. B., Malarkey, W. B., Van Cauter, E., & Berntson, G. G. (2002). Loneliness and Health : Potential Mechanisms. *Psychosomatic Medicine*, *64*(3), 407.

Cacioppo, J. T., Hawkley, L. C., & Thisted, R. A. (2010). Perceived social isolation makes me sad : 5-year cross-lagged analyses of loneliness and depressive symptomatology in the Chicago Health, Aging, and Social Relations Study. *Psychology and Aging*, *25*(2), 453-463. https://doi.org/10.1037/a0017216

Chen, X., Ke, Z. L., Chen, Y., & Lin, X. (2021). The prevalence of sleep problems among

children in mainland China : A meta-analysis and systemic-analysis. *Sleep Medicine*,

83, 248-255. https://doi.org/10.1016/j.sleep.2021.04.014

Cohen, S. (2004). Social Relationships and Health. *American Psychologist*, *59*(8), 676.

https://doi.org/10.1037/0003-066X.59.8.676

Dunphy-Doherty, F., O'Mahony, S. M., Peterson, V. L., O'Sullivan, O., Crispie, F., Cotter, P.

D., Wigmore, P., King, M. V., Cryan, J. F., & Fone, K. C. F. (2018). Post-weaning

social isolation of rats leads to long-term disruption of the gut microbiota-immune-

brain axis. *Brain, Behavior, and Immunity*, *68*, 261-273.

https://doi.org/10.1016/j.bbi.2017.10.024

Ehrler, M., Hagmann, C. F., Stoeckli, A., Kretschmar, O., Landolt, M. A., Latal, B., & Wehrle,

F. M. (2022). Mental sequelae of the COVID-19 pandemic in children with and

without complex medical histories and their parents : Well-being prior to the outbreak

and at four time-points throughout 2020 and 2021. *European Child & Adolescent*

Psychiatry. https://doi.org/10.1007/s00787-022-02014-6

González-Monroy, C., Gómez-Gómez, I., Olarte-Sánchez, C. M., & Motrico, E. (2021).

Eating Behaviour Changes during the COVID-19 Pandemic : A Systematic Review of

Longitudinal Studies. *International Journal of Environmental Research and Public*

Health, *18*(21), Article 21. https://doi.org/10.3390/ijerph182111130

Holt-Lunstad, J., Smith, T. B., Baker, M., Harris, T., & Stephenson, D. (2015). Loneliness

and social isolation as risk factors for mortality : A meta-analytic review. *Perspectives*

on Psychological Science: A Journal of the Association for Psychological Science,

10(2), 227-237. https://doi.org/10.1177/1745691614568352

Leiner, D. J. (2021). *SoSci Survey (Version 3.2.31) [Computer software]* [Logiciel]. Available

at https://www.soscisurvey.de

Li, W., Wang, Z., Syed, S., Lyu, C., Lincoln, S., O'Neil, J., Nguyen, A. D., Feng, I., & Young,

M. W. (2021). Chronic social isolation signals starvation and reduces sleep in

Drosophila. *Nature*, *597*(7875), Article 7875. https://doi.org/10.1038/s41586-021-
03837-0

Lopizzo, N., Marizzoni, M., Begni, V., Mazzelli, M., Provasi, S., Borruso, L., Riva, M. A., &
Cattaneo, A. (2021). Social isolation in adolescence and long-term changes in the
gut microbiota composition and in the hippocampal inflammation : Implications for
psychiatric disorders – Dirk Hellhammer Award Paper 2021.
Psychoneuroendocrinology, *133*, 105416.
https://doi.org/10.1016/j.psyneuen.2021.105416

Markovic, A., Mühlematter, C., Beaugrand, M., Camos, V., & Kurth, S. (2021). Severe effects
of the COVID-19 confinement on young children's sleep : A longitudinal study
identifying risk and protective factors. *Journal of Sleep Research*, e13314.
https://doi.org/10.1111/jsr.13314

Mason, T. B., Heron, K. E., Braitman, A. L., & Lewis, R. J. (2016). A daily diary study of
perceived social isolation, dietary restraint, and negative affect in binge eating.
Appetite, *97*, 94-100. https://doi.org/10.1016/j.appet.2015.11.027

McLay, L., Jamieson, H. A., France, K. G., & Schluter, P. J. (2021). Loneliness and social
isolation is associated with sleep problems among older community dwelling women
and men with complex needs. *Scientific Reports*, *11*(1), 4877.
https://doi.org/10.1038/s41598-021-83778-w

Mühlematter, C., Nielsen, D. S., Castro-Mejía, J. L., Brown, S. A., Rasch, B., Jr, K. P. W.,
Walser, J.-C., Schoch, S. F., & Kurth, S. (2023). Not simply a matter of parents—
Infants' sleep-wake patterns are associated with their regularity of eating. *PLOS*
ONE, *18*(10), e0291441. https://doi.org/10.1371/journal.pone.0291441

Pan, X.-F., Wang, L., & Pan, A. (2021). Epidemiology and determinants of obesity in China.
The Lancet. Diabetes & Endocrinology, *9*(6), 373-392. https://doi.org/10.1016/S2213-
8587(21)00045-0

Pujia, R., Ferro, Y., Maurotti, S., Khoory, J., Gazzaruso, C., Pujia, A., Montalcini, T., &
Mazza, E. (2021). The Effects of COVID-19 on the Eating Habits of Children and

Adolescents in Italy : A Pilot Survey Study. *Nutrients*, *13*(8), Article 8.

https://doi.org/10.3390/nu13082641

Qualter, P., Brown, S. L., Rotenberg, K. J., Vanhalst, J., Harris, R. A., Goossens, L.,

Bangee, M., & Munn, P. (2013). Trajectories of loneliness during childhood and

adolescence : Predictors and health outcomes. *Journal of Adolescence*, *36*(6),

1283-1293. https://doi.org/10.1016/j.adolescence.2013.01.005

Teixeira, M. T., Vitorino, R. S., da Silva, J. H., Raposo, L. M., Aquino, L. A. de, & Ribas, S.

A. (2021). Eating habits of children and adolescents during the COVID-19 pandemic :

The impact of social isolation. *Journal of Human Nutrition and Dietetics*, *34*(4),

670-678. https://doi.org/10.1111/jhn.12901

Tomiyama, A. J. (2019). Stress and Obesity. *Annual Review of Psychology*, *70*, 703-718.

https://doi.org/10.1146/annurev-psych-010418-102936

Wright, A., & Pritchard, M. E. (2009). An examination of the relation of gender, mass media

influence, and loneliness to disordered eating among college students. *Eating and*

Weight Disorders - Studies on Anorexia, Bulimia and Obesity, *14*(2), e144-e147.

https://doi.org/10.1007/BF03327813