A validated cloud-based genomic platform for co-ordinated, expedient global analysis of SARS-CoV-2 genomic epidemiology

Authors

Daniel Amoako,[1] Nguyen To Anh,[2] Jasmine Bastable,[3] Marc Brouard,[4] Constanza Campano Romero,[5] Andres Castillo Ramirez,[5] Bede Constantinides,[4] Derrick W. Crook,[3,4] Phan Manh Cuong,[6] Moussa Moise Diagne,[7] Amadou Diallo,[7] Nguyen Thanh Dung,[8] Laura Dunn,[3] Le Van Duyet,[6] Josie Everatt,[1] Katherine Fletcher,[4] Philip W. Fowler,[4] Mailie Gail,[9] Jessica Gentry,[3] Saheer Gharbia,[10] Hospital for Tropical Diseases SARS-CoV-2 testing team,[8] Nguyen Thi Thu Hong,[2] Martin Hunt,[4,12] Zam Iqbal,[4,12] Katie Jeffery,[3] Dikeledi Kekana,[1] Thomas Kesteman,[2] Jeff Knaggs,[4,12] Marcela Lopes Alves,[4] Dinh Nguyen Huy Man,[8] Amy J. Mathers,[11] Nghiem My Ngoc,[8] Sarah Oakley,[3] Hardik Parikh,[11] Tim E.A. Peto,[3,4] Phuong Quan,[4] Marcelo Rojas Herrera,[5] Nicholas Sanderson,[4] Vitali Sintchenko,[9] Jeremy Swann,[4] Junko Takata,[3,4] Nguyen Thi Tam,[2] Le Van Tan,[2] Pham Ngoc Thach,[6] Ndeye Marieme Top,[7] Nguyen Thu Trang,[2] Van Dinh Trang,[6] Robert Turner,[4] H. Rogier van Doorn,[2] Anne von Gottberg,[1,13] Jeremy Westhead,[4] Nicole Wolter,[1,13] Bernadette C. Young[3,4]

Author Affiliations

1. The National Institute for Communicable Diseases (NICD), a division of the National Health Laboratory Service, Johannesburg, Republic of South Africa
2. Oxford University Clinical Research Unit, Ho Chi Minh City, Vietnam.
3. Oxford University Hospitals NHS Foundation Trust, Oxford, United Kingdom
4. Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom
5. El Instituto de Salud Pública de Chile, Santiago, Chile

6. National Hospital for Tropical Diseases, Hanoi, Vietnam

7. The Institut Pasteur de Dakar, Dakar, Senegal

8. Hospital for Tropical Diseases, Ho Chi Minh City, Vietnam

9. Microbial Genomics Reference Laboratory, Institute of Clinical Pathology and

Medical Research, New South Wales Health Pathology, Sydney, Australia

10. Genomic Surveillance Unit, The Wellcome Sanger Institute, Hinxton, United

Kingdom

11. Division of Infectious Diseases, School of Medicine, University of Virginia,

Charlottesville, United States of America

12. European Bioinformatics Institute, Hinxton, United Kingdom

13. Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South

Africa

Corresponding author: Bernadette Young (bernadette.young@ndm.ox.ac.uk), Nuffield

Department of Medicine, University of Oxford, Oxford, OX3 9DU, 01865 220856.

Running title: Coordinated cloud-based genome analysis for SARS-CoV-2

Keywords: COVID-19, SARS-CoV-2, public health genomics, bioinformatics

Article Summary Line: A cross-sectional study of SARS-CoV-2 sequencing demonstrates that cloud-based sequencing analysis has the power to relieve bioinformatic bottlenecks, and facilitate collaboration in pathogen surveillance to enhance pandemic preparedness.

Abstract

Background

Viral sequencing has made critical contributions to our understanding of and response to the COVID-19 pandemic, but sequencing capacity and bioinformatic expertise remain limited in many settings. This proof-of-principle study aimed to demonstrate the utility of a cloud-based sequencing analysis pipeline, the Tiled Amplicon Pipeline (TAP), for rapid and collaborative SARS-CoV-2 sequencing across seven globally distributed sites.

Methods

In this cross-sectional study from July to August 2022, seven sites submitted all SARS-CoV-2 sequence data generated over a two-week period to our cloud platform. No patient identifying information was uploaded, and human reads were removed prior to upload to the cloud. Users could opt in to share sample information with collaborators via a tagging system. The pipeline performed sequence assembly, lineage identification and relatedness analysis.

Results

Seven sites contributed 5,432 sequences, of which 5,342 (98.3%) were from clinical samples and 90 (1.7%) were controls. Of the clinical samples that were correctly assembled, 3,439/4,179 (82.3%) had sufficient coverage for lineage assignment. Omicron lineages dominated, with BA.5, BA.4 and BA.2 comprising the vast majority, consistent with contemporary epidemiological observations at the time. Phylogenetic analysis demonstrated low diversity within lineages, and genotypically identical or highly similar sequences were recovered from globally disparate sites.

73 **Conclusions**

74 A cloud-based analysis platform like TAP addresses bioinformatic bottlenecks and

75 facilitates collaboration in pathogen surveillance, enhancing epidemic and pandemic

76 preparedness.

Introduction

Viral genome sequencing has proven pivotal to understanding the evolution of the SARS-CoV-2 virus during the COVID-19 pandemic and in shaping the public health response. International genomic surveillance and data sharing initiatives have together made it possible to track the emergence of variants globally [1], to demonstrate the impact of travel restrictions on viral dynamics across continents [2] or within countries [3, 4], and to identify transmission routes within hospitals [5]. Successive waves of infection driven by new variants showed that rapidly detecting new lineages is critical for understanding disease epidemiology and guiding subsequent public health responses [6], as well as informing the development of vaccines and therapeutics such as neutralising antibodies [7-9].

However, the pandemic also highlighted marked global variability in sequencing capacity and cost [10], with much of the SARS-CoV-2 sequencing undertaken at centralised reference laboratories [11]. Several important challenges remain in expanding genomic surveillance, including access to sequencing technology, availability of bioinformatic expertise, and interpretability of results generated using a plethora of different wet lab and bioinformatics protocols [12-14]. There is a need for accessible genomic surveillance infrastructure that can be used by researchers and clinicians from any location, to deliver an up-to-date global perspective of viral evolution.

One solution to both the shortage of bioinformatic expertise and the lack of global interpretability of results is web-accessible analysis infrastructure [15]. The Global Pathogen Analysis Service (GPAS) was rapidly set up in 2021 in response to the COVID-19 pandemic by the University of Oxford as a cloud-based, globally accessible web

102 platform. GPAS provided fast and secure access to a comprehensive SARS-CoV-2 genomic

103 analysis pipeline, delivering genome assembly, variant calling and lineage classification

104 from raw sequence data. Users retained control of their data but could opt in to share

105 their data to facilitate comparisons in a wider context, empowering laboratories to

106 control their own analysis. GPAS was validated to UKAS ISO 15189:2012 standards in a

107 UK tertiary hospital clinical microbiology laboratory, and was commissioned by the UK

108 Health Security Agency for over a year to support Pillar 1 national surveillance and

109 various clinical trials. GPAS provided proof-of-principle that standard microbiology

110 laboratories without bioinformatics expertise can generate outputs for local surveillance

111 and automatically submit sequences to public repositories.

113 In 2024, GPAS was upgraded as the Tiled Amplicon Pipeline (TAP) with major

114 enhancements to several software components. Since January 2025 TAP has been

115 deployed as the SARS-CoV-2 pipeline on EIT Pathogena (version 1.2.0), a multi-pipeline

116 genomic pathogen analysis platform that is free of charge for users in low- and middle-

117 income countries and accessible in all settings with internet access [16]. This service

118 demonstrates the potential of cloud-based platforms to overcome barriers to

119 democratising effective genomic surveillance, making advanced genomic analysis

120 available to resource-limited laboratories.

122 In this paper, we report the results of a collaborative, cross-sectional SARS-CoV-2

123 sequencing study across seven globally distributed sites, which demonstrate the utility

124 of a cloud-based sequencing platform in providing a quality assured, rapid, and

125 integrated global snapshot of viral diversity. We further describe the components of the

126   pipeline and discuss its potential for rapid adaptation in response to future viral

127   pandemics.

Methods

Pipeline development

TAP was deployed to a cloud platform and controlled via a Command Line Interface (CLI) tool. Full methodology is described in the Appendix (Supplementary Appendix A). At upload, a universally unique identifier (UUID) is generated and assigned to each sample. The mapping between the UUID and the user's sample identifier is downloaded and only held by the user, and FASTQ header lines are also truncated, ensuring that no potentially personally identifiable information is transmitted to the cloud platform. Tight access control to data applies within TAP, and by default data is not shared with other users unless explicitly authorised by the data owners.

Following FASTQ file upload to the cloud, processing commences automatically with no further user input required. Any reads mapping to the human genome are first removed [17], reducing the risk of any human reads being retained. Genome assembly is performed by an amplicon-aware genome assembly tool Viridian v1.3.1 [18], which scaffolds per-amplicon *de novo* assemblies into a single whole genome consensus assembly in FASTA format. Viridian was configured to use the SARS-CoV-2 reference genome Wuhan-Hu-1 (MN908947.3) for assembly and variant calling [19], and a library of seven amplicon primer schemes (AmpliSeq v1; ARTIC versions 3, 4.1, 5.3.2 (400), 5.2.0 (1200); Midnight 1200; VarSkip v1a-2b). When a primer scheme is not specified by the user, Viridan can automatically infer the most likely primer scheme from within its library.

After assembly, amino acid mutations are identified using Nextclade [20], and Pango lineages are assigned with Pangolin version 4.3.1 [21]. Aligned sequences are compared

using a novel algorithm, FindNeighbour5, which identifies single nucleotide variant

(SNV) distances between sequences without conflicting variant calls (0 SNVs) as well as

those differing by one, two and three SNVs [22, 23]. The main outputs from the pipeline

are presented in an access-restricted user interface portal and downloadable as a

summary file containing the lineage assignment, a list of related samples (according to

permissions), and metrics of the genome assembly (e.g. coverage, mean depth, amplicon

dropouts, etc). Detailed intermediate files such as VCF or FASTA files are also available

for download.

Pipeline validation

GPAS was validated for both Illumina and Oxford Nanopore Technologies (ONT)

sequencing platforms in 2021, using various datasets including a 'truth set' of cultured

SARS-CoV-2 samples sequenced with multiple platforms and library preparations,

annotated with manually curated variant calls (described in further detail in [18] and

[24]), in addition to community samples collected in Northumbria, UK. Overall, GPAS

showed negligible adjusted false call rates (less than 1/100,000 nucleotides) with respect

to the 'truth set', and had high concordance (less than 1.9/1,000,000 discordant events)

with the ARTIC assembly pipeline in use at the time.

TAP was subsequently validated against the same 'truth set' of cultured SARS-CoV-2

samples, which achieved 100% concordance with expected Pango lineages using the

same Pangolin version. Concordance of TAP and GPAS was further compared using data

from the current study, which showed 99.4% concordance in lineage calls (using the

same Pangolin and Viridian versions), and nucleotide call discordance of 0.59/1,000,000

sites. Further details of the validation and comparison processes are described in

Supplementary Appendix B.

Sample frame, sequencing, and upload

Between July and August 2022, seven sequencing centres participated in a two-week

sequencing pilot across seven countries: Senegal; Chile; South Africa; New South Wales

(NSW), Australia; Vietnam; United Kingdom (UK); and Virginia, United States (USA).

Centres were either accredited clinical microbiology or public health laboratories. All

clinical samples in which SARS-CoV-2 was detected were eligible for inclusion and

underwent in-house genomic sequencing at participating sites, with no more than one

submission from an individual patient. Sequencing platform (either Illumina or ONT) and

primer schemes were chosen by participating sites, who followed their established

sequencing protocols.

Raw sequences were uploaded in FASTQ format to GPAS, the working name of the

pipeline at the time, along with limited associated metadata (sample name, instrument

platform, sample type [clinical or control], collection date, and country). Submitting sites

verified the run quality reports of each sequencing batch including a review of positive

and negative controls, and batches were passed or failed accordingly. All passed samples

were tagged to release them to the shared data pool for subsequent aggregate analysis;

explicit permission was given from all collaborators to configure data access controls

such that all submitters could access and view each other's sequences, metadata, and

analytical outputs. For this study, primer scheme information was not provided by the

user, and automatic detection of primer scheme was enabled. All sequences were

uploaded to the European Nucleotide Archive (project accession: PRJEB70597) at the

time of the study.

Data analysis

FASTQ files from the study were redownloaded from the ENA and run through EIT

Pathogena (version 1.2.0 pre-release) on 6th November 2024. A maximum likelihood

phylogeny of aligned sequences was constructed for the three largest Pango lineages in

the sample set (BA.5, BA.4 and BA.2) using IQTree version 2.3.6, assuming a general time

reversible nucleotide substitution model with gamma rate heterogeneity, and using the

consensus tree from 1000 ultrafast bootstrapping [25]. All other analyses were

conducted in RStudio version 2023.06.0+421.

Results

Primer scheme detection

5,432 sequences were shared across seven sites (Supplementary Table 1), with date of collection ranging from April to July 2022. Of submitted samples, 90 (1.7%) were controls and 5,342 (98.3%) were clinical samples. Primer schemes were successfully auto-detected for 4,238/5,432 (78.0%) of samples, but were incorrectly inferred for all 1,194/5,432 (22.0%) samples from NSW, Australia which appropriately failed the Viridian quality control threshold for assembly. Further investigation showed that these samples were sequenced using a bespoke primer scheme that is not included in the current Viridian library, and these samples were excluded from further analysis.

Aggregate analysis of global genetic epidemiology

Among the clinical samples included in the analysis, 3,751/4,179 (89.8%) were assembled with at least 70% genome coverage (Table 1). 3,439/4,179 (82.3%) clinical samples were assembled with sufficient coverage and post-assembly quality to be assigned a lineage (Table 2). 3,412/3,439 (99.2%) were Omicron variants, with BA.5, BA.4 and BA.2 being the most common. A small number of Delta variant sequences were identified (0.4%, 15/3,439), all of which were collected prior to July 2022.

Centre	Submitted samples	Controls	<50% coverage*	50-70% coverage	>70% coverage	Earliest sample	Latest sample
Senegal	197	0	9	10	178	13th July 2022	20th July 2022
Chile	1205	20	4	6	1175	16th June 2022	22nd July 2022
South Africa	202	0	7	6	189	17th May 2022	24th June 2022
Vietnam	316	3	14	7	292	1st April 2022	14th July 2022
United Kingdom	1818	36	280	85	1417	12th July 2022	26th July 2022
Virginia, USA	500	0	0	0	500	7th June 2022	28th June 2022
Total	**4238**	**59**	**314**	**114**	**3751**		

238   **Table 1: Samples submitted by study centre, with date of collection and genome coverage.** Earliest

239   and latest sample dates exclude controls.

240   * includes samples that could not be assembled

Centre	Omicron			BA.5 and related				Delta	Alpha	Other*	Total
	BA.1	BA.2	BA.4	BA.5	BE	BF	Other (BG, BK)	AY.5, AY.57, B.1.617.2	B.1.1.7		
Senegal	0	17	24	28	2	12	0	0	0	0	83
Chile	0	136	570	302	26	63	1	0	0	5	1103
South Africa	0	3	82	77	8	3	0	0	0	3	176
Vietnam	2	228	0	6	0	0	0	15	0	0	251
United Kingdom	2	57	165	864	137	111	0	0	1	2	1339
Virginia, USA	0	289	89	84	12	9	3	0	0	1	487
Total	**4**	**730**	**930**	**1361**	**185**	**198**	**4**	**15**	**1**	**11**	**3,439**

Table 2: SARS-CoV-2 lineage of samples by study centre (where a lineage was assigned by Pangolin).

* Other includes: P.1, P.1.2, A, A.2.2, B.1, B.1.429, B.28, C.37, DE.1, XAM, XAN, XAS, XAZ, XBA, P.2

Global mixing across multiple Omicron lineages

For clinical samples collected between 1st June 2022 and 31st July 2022, the proportion

of samples assigned to the different Omicron sub-lineages varied substantially by study

site (Figure 1). Each of BA.2, BA.4 or BA.5 dominated (constituted >50% of samples from)

at least one site, while in Senegal no single lineage dominated, and BA.4 and BA.5 were

equally common in South Africa (44.3% and 46.9% respectively).

Figure 1: Omicron sub-lineages by study site. Proportion of samples assigned to Omicron sub-lineages

within each study site, where collection date was in June or July 2022.

Maximum likelihood phylogenies for each of the three most common lineages revealed

global mixing of Omicron lineages (Figure 2). There were no examples of sub-lineages

being completely geographically restricted to one site; one sub-lineage of BA.4 was found

predominantly in Chile, but an example of this sub-lineage was also identified in the

United Kingdom.

Figure 2: Maximum likelihood phylogeny of isolates from each major lineage (BA.2, BA.4, BA.5). Final branches are coloured according to the centre in which they

were sequenced.

For all three major lineages, there was evidence of genotypically identical (SNV = 0) and

highly related (SNV ≤ 3) samples being found in different study sites, with increasing

dispersal with higher SNV thresholds (Figure 3; Supplementary Table 2). The greatest

dispersal was seen in the BA.4 lineage: of 358 samples with another genotypically

identical sample, only 69.6% (249/358) were from the same site, with the rest

distributed across the remaining sites. In addition, only 10.3% (96/930) of BA.4 and

12.7% (173/1,361) of BA.5 samples did not have any related samples (SNV ≤ 3) in the

shared pool, in contrast to BA.2 samples where 51.8% (378/730) samples had a related

sample in the shared pool.

Figure 3: Percentage of unique samples for which all related samples were found only at the same site, at

each genotypic relatedness (SNV) threshold.

Cloud processing time and performance

At the time of the study, cloud processing performance on GPAS was measured by time elapsed between upload and completion of analysis. Median cloud processing time per sample was 30.6 minutes (IQR: 13.1 – 69.9); variability was observed, with several batches from Australia, UK and USA taking longer than 1000 minutes (Supplementary Figure 1). These were caused by either a data centre outage during an extreme heat event in the UK, or as a result of a software update prematurely applied to the upload portal, which was promptly resolved by improvements in the client side CLI software. Other data upload issues were identified and addressed case-by-case, including metadata file formatting errors, the availability of upload clients compatible with all required operating systems, and user interface display errors preventing batch release.

In TAP, median cloud processing time per sample for a typical batch of 100 was 14.5 minutes (IQR: 13.8 – 15.0) for Illumina and 15.6 minutes (IQR: 15.2 – 16.7) for ONT; due to parallelisation, processing a single batch of 100 samples was achievable in 19.7 minutes (Illumina) and 20.4 minutes (ONT) (Supplementary Figure 2). Processing all 5,432 samples took 766 minutes (12.7 hours). CPU time per sample is approximately 3 minutes (noting that this occurs in parallel, i.e. 2 CPUs working for 4 minutes would be reported as 8 minutes CPU time by the workflow manager, despite only taking 4 minutes); majority of analysis time is associated with other pipeline activities including network file transfers or reading data from disk into memory. Peak RAM usage for typical use of the pipeline was approximately 5GB.

Discussion

This pilot study demonstrates the potential of globally synchronous data processing and analysis in informing public health, through a unified protocol of genome assembly, variant calling and relatedness analysis. The aggregated analysis from this study conforms to contemporary observations of SARS-CoV-2 genetic epidemiology at the time [26-28], and demonstrates cosmopolitan global mixing of Omicron lineages between study sites. Even a BA.4 sub-lineage found almost exclusively in Chile was not completely restricted to that site and could be identified in a geographically distant site (UK). Similarly, between 15.3% to 30.4% of genotypically identical (SNV=0) sequences were identified in different countries over the same two-month period. Such mixing of lineages likely reflects the relaxation of travel restrictions in the participating sites at the time of the study, and captures the rapid global dispersal of Omicron with successive selective sweeps of new lineages over short time scales [29]. These observations highlight the potential of a cloud-based sequencing pipeline in facilitating data sharing and generating co-ordinated insights that could inform real-time decision-making.

While automatic primer scheme detection was used for this study, our results show the limitation in this approach, with one centre using a custom scheme not contained in the current Viridian primer scheme library. With viral evolution, it is inevitable that new and/or modified primer schemes will continue to be developed to maintain sequencing coverage of an evolving target genome. Viridian has the capability to accommodate custom primer schemes when provided with a suitable scheme definition in browser extensible data (BED) format. This capability will form the basis of future pipeline iterations and will allow the pipeline to pivot in future to assemble other viruses beyond SARS-CoV-2 that use tiled amplicon sequencing. In turn, this ability to rapidly adapt

existing infrastructure for emerging threats, rather than build bespoke solutions from

scratch each time, supports a proactive and 'Always On' approach to pandemic

preparedness [30].

Our study is limited by the opportunistic sampling frame used and is likely to be

geographically incomplete, with a preponderance for countries that have historically

contributed more to global sequencing efforts. Similarly, our study has the limitation of

lacking longitudinal data on how these findings changed over time. Despite these

shortcomings, we have demonstrated how even a simply structured sampling across time

and space is still well placed to rapidly identify replacement by new lineages. Such

limitations in sampling frame diversity and global disparity in sequencing volume were

reflected during the pandemic itself: sequencing efforts in high-income countries were

up to ten-fold higher than that achieved in low- and middle-income countries during the

first two years [11], and data was not collected evenly over the course of pandemic. Well

designed, suitably powered, and representative sampling is a key part of pandemic

preparedness planning, as exemplified by a cohort design in the UK [27].

The world is now better prepared to urgently initiate genomic pathogen surveillance. The

available sequencing platform capacity is much improved, and the achievements of global

data aggregation (as illustrated by INSDC, GISAID and Pathoplexus databases) have

alleviated a major obstacle to effective genomic surveillance and data sharing. In turn, the

integration of genomic data with disease manifestation and severity data would enable

the ready investigation of associations between genomic variation and clinical outcome

as suggested by others [31], and facilitate predictive modelling for anticipating the course

of future epidemics [32].

A remaining obstacle, especially in low-resource settings, is establishing local turnkey

bioinformatics to ensure standardised, quality-assured outputs that do not depend on in-

house bioinformatics expertise. In TAP, we have developed an exemplar of such a cloud-

based service, with simple ingestion of local sequence data and flexible privacy

protections for data sharing to facilitate local or international comparative analyses.

While factors such as internet bandwidth may limit real-world performance, TAP is

capable of rapid parallel data processing, with a run time of 15 minutes per sample and

20 minutes in total for a batch of 100 typical SARS-CoV-2 genomes. The EIT Pathogena

platform ensures sustainability of technical and infrastructural support to the pipeline

and is currently freely available to low- and middle-income countries, supporting long-

term use across research, clinical, or surveillance settings. Such a service, in turn,

contributes to the growing global landscape of genomic data sharing that will underpin

future pandemic preparedness.

Conclusions

The COVID-19 pandemic has demonstrated the multiple ways in which viral sequencing

can inform pandemic responses, but also the inequitable distribution of resources and

capabilities. A service such as TAP contributes to the democratisation of genomics,

enabling researchers and laboratories, regardless of their location or bioinformatics

expertise, to participate actively in global surveillance efforts. By providing accessible

and user-friendly tools for sequence assembly, lineage assignment, and data sharing, it

promotes inclusive collaboration, harmonises data, and ultimately enhances pandemic

preparedness.

Author contributions

Study conceptualisation: BCY, DWC

Study design: BCY, DWC

Software building and computational infrastructure: MB, BC, MH, ZI, PWF, JK, MLA, NS, JS, RT, JW

Data collection: DA, NTA, JB, CCR, ACR, PMC, MMD, AD, NTD, LD, LVD, JE, KF, MG, JG, SG, Hospital for Tropical Diseases SARS-CoV-2 testing team, NTTH, KJ, DK, TK, DNHM, AJM, NMN, SO, HP, MRH, VS, NTTa, LVT, PNT, NMT, NTTr, VDT, HRVD, AvG, NW

Data analysis: JT, BCY, BC, TEAP, PQ

Manuscript writing: JT, BCY, BC

Manuscript editing and review: all authors

Funding statement

GPAS is a non-profit organisation, and GPAS cloud infrastructure was supported by a donation from Oracle Corporation. For this study, all GPAS services were offered free of charge to all sites, and staff costs for the analysis team were met by the University of Oxford, with support from the National institute for Health Research (NIHR) Oxford Biomedical Research Centre (BRC). TAP is deployed on the EIT Pathogena platform, funded by the Ellison Institute of Technology Oxford. EIT Pathogena is free at the point of use for users in low- and middle-income countries.

Sequencing activities for NICD are supported by a conditional grant from the South African National Department of Health as part of the emergency COVID-19 response; a cooperative agreement between the National Institute for Communicable Diseases of the National Health Laboratory Service and the United States Centers for Disease Control and

Prevention (NU51IP000930); the South African Medical Research Council (SAMRC) with

funds received from the South African Department of Science and Innovation; the African

Society of Laboratory Medicine (ASLM) and Africa Centers for Disease Control and

Prevention through a sub-award from the Bill and Melinda Gates Foundation grant

number INV-018978; Africa PGI, the UK Foreign, Commonwealth and Development Office

and Wellcome (Grant no 221003/Z/20/Z); and the Department of Health and Social

Care's Fleming Fund using UK aid. NICD sequencing was also supported by The

Coronavirus Aid, Relief, and Economic Security Act (CARES ACT) through the Centers for

Disease Control and Prevention (CDC) and the COVID International Task Force (ITF)

funds through the CDC under the terms of a subcontract with the African Field

Epidemiology Network (AFENET) AF-NICD-001/2021.

Genomic surveillance conducted in Vietnam was supported by the Wellcome Trust

(222574/Z/21/Z). L.V.T. is supported by the Wellcome Trust of Great Britain

(204904/Z/16/Z and 226120/Z/22/Z).

Ethical statement

SARS-CoV-2 sequencing was performed for public health surveillance and ethical

approval for secondary analysis was not required. This determination was reviewed by

the University of Oxford Joint Research Office. No patient identifying information was

shared as part of this study. Sample identifiers remain with the submitter, and are never

kept on GPAS/TAP, or shared with other users. GPAS/TAP generates identifiers for each

sequence and writes a file linking anonymised ID to the submitted identifiers. Only the

user submitting sequence data has access to these records. The GPAS/TAP upload client

selects only SARS-CoV-2 sequence reads and discards all human sequences.

Data availability

Submitted sequencing reads (free from human reads) for all included samples are available from European Nucleotide Archive study accession PRJEB70597.

Conflicts of interest

AvG and NW have received grant funding from Sanofi and The Bill and Melinda Gates Foundation. PWF receives consultancy fees from the Ellison Institute of Technology, Oxford. JT, DWC and TEAP receive funding from the Ellison Institute of Technology, Oxford.

Acknowledgments

- Microbial Genomics Reference Laboratory, New South Wales Health Pathology, Sydney, Australia
- El Instituto de Salud Pública de Chile, Chile
- Centre for Respiratory Diseases and Meningitis (CRDM), National Institute for Communicable Diseases (NICD), a division of the National Health Laboratory Service, South Africa
- The Institut Pasteur de Dakar, Senegal
- Oxford University Hospitals NHS Trust, Oxford, United Kingdom
- University of Virginia, Charlottesville, United States of America
- Oxford University Clinical Research Unit, Vietnam
- Hospital for Tropical Diseases SARS-CoV-2 testing team, HTD Vietnam: Le Manh Hung, Nguyen Le Nhu Tung, Nguyen Thanh Phong, Vo Minh Quang, Pham Thi Ngoc Thoa, Nguyen Thanh Truong, Tran Nguyen Phuong Thao,

451   Dao Phuong Linh, Ngo Tan Tai, Ho The Bao, Vo Trong Vuong, Huynh Thi

452   Kim Nhung

453   • Oracle Global Health Business Unit

References

References

1.   Li J, Lai S, Gao GF, Shi W. The emergence, genomic diversity and global spread of SARS-CoV-2.
Nature. 2021;600(7889):408-18.

2.   Wilkinson E, Giovanetti M, Tegally H, San JE, Lessells R, Cuadros D, et al. A year of genomic
surveillance reveals how the SARS-CoV-2 pandemic unfolded in Africa. Science. 2021;374(6566):423-31.

3.   McCrone JT, Hill V, Bajaj S, Pena RE, Lambert BC, Inward R, et al. Context-specific emergence and
growth of the SARS-CoV-2 Delta variant. Nature. 2022;610(7930):154-60.

4.   Raghwani J, du Plessis L, McCrone JT, Hill SC, Parag KV, Theze J, et al. Genomic Epidemiology of
Early SARS-CoV-2 Transmission Dynamics, Gujarat, India. Emerg Infect Dis. 2022;28(4):751-8.

5.   Snell LB, Fisher CL, Taj U, Stirrup O, Merrick B, Alcolea-Medina A, et al. Combined epidemiological
and genomic analysis of nosocomial SARS-CoV-2 infection early in the pandemic and the role of
unidentified cases in transmission. Clin Microbiol Infect. 2022;28(1):93-100.

6.   Updated working definitions and primary actions for SARS-Cov-2 variants. WHO Technical
Advisory Group on Virus Evolution; 2023.

7.   Tao K, Tzou PL, Nouhin J, Gupta RK, de Oliveira T, Kosakovsky Pond SL, et al. The biological and
clinical significance of emerging SARS-CoV-2 variants. Nat Rev Genet. 2021;22(12):757-73.

8.   Tuekprakhon A, Nutalai R, Dijokaite-Guraliuc A, Zhou D, Ginn HM, Selvaraj M, et al. Antibody
escape of SARS-CoV-2 Omicron BA.4 and BA.5 from vaccine and BA.1 serum. Cell. 2022;185(14):2422-33
e13.

9.   Aggarwal A, Akerman A, Milogiannakis V, Silva MR, Walker G, Stella AO, et al. SARS-CoV-2
Omicron BA.5: Evolving tropism and evasion of potent humoral responses and resistance to clinical
immunotherapeutics relative to viral variants of concern. EBioMedicine. 2022;84:104270.

10.   Merhi G, Koweyes J, Salloum T, Khoury CA, Haidar S, Tokajian S. SARS-CoV-2 genomic
epidemiology: data and sequencing infrastructure. Future Microbiol. 2022;17:1001-7.

11.   Brito AF, Semenova E, Dudas G, Hassler GW, Kalinich CC, Kraemer MUG, et al. Global disparities in
SARS-CoV-2 genomic surveillance. Nat Commun. 2022;13(1):7003.

12.   Carter LL, Yu MA, Sacks JA, Barnadas C, Pereyaslov D, Cognat S, et al. Global genomic surveillance

strategy for pathogens with pandemic and epidemic potential 2022-2032. Bull World Health Organ.

2022;100(4):239-A.

13.   Inzaule SC, Tessema SK, Kebede Y, Ogwell Ouma AE, Nkengasong JN. Genomic-informed pathogen

surveillance in Africa: opportunities and challenges. Lancet Infect Dis. 2021;21(9):e281-e9.

14.   Chen Z, Azman AS, Chen X, Zou J, Tian Y, Sun R, et al. Global landscape of SARS-CoV-2 genomic

surveillance and data sharing. Nat Genet. 2022;54(4):499-507.

15.   Ohlsen EC, Hawksworth AW, Wong K, Guagliardo SAJ, Fuller JA, Sloan ML, et al. Determining Gaps

in Publicly Shared SARS-CoV-2 Genomic Surveillance Data by Analysis of Global Submissions. Emerg

Infect Dis. 2022;28(13):S85-S92.

16.   EIT Pathogena [Available from: https://eit-pathogena.com.

17.   Constantinides B, Hunt M, Crook DW. Hostile: accurate decontamination of microbial host

sequences. Bioinformatics. 2023;39(12).

18.   Hunt M, Hinrichs AS, Anderson D, Karim L, Dearlove BL, Knaggs J, et al. Addressing pandemic-

wide systematic errors in the SARS-CoV-2 phylogeny. bioRxiv. 2024:2024.04.29.591666.

19.   Wu F, Zhao S, Yu B, Chen YM, Wang W, Song ZG, et al. A new coronavirus associated with human

respiratory disease in China. Nature. 2020;579(7798):265-9.

20.   Aksamentov I, Roemer C, Hodcroft EB, Neher RB. Nextclade: clade assignment, mutation calling

and quality control for viral genomes. Journal of Open Source Software. 2021;6(67):3773.

21.   O'Toole A, Scher E, Underwood A, Jackson B, Hill V, McCrone JT, et al. Assignment of

epidemiological lineages in an emerging pandemic using the pangolin tool. Virus Evol.

2021;7(2):veab064.

22.   Mazariegos-Canellas O, Do T, Peto T, Eyre DW, Underwood A, Crook D, et al. BugMat and

FindNeighbour: command line and server applications for investigating bacterial relatedness. BMC

Bioinformatics. 2017;18(1):477.

23.   FindNeighbour5 [Available from: https://github.com/oxfordmmm/FN5.

24.   Constantinides B, Webster H, Rodger G, Hunt M, Supasa P, Dejnirattisai W, et al. A diverse
reference set of cultured SARS-CoV-2 genomes sequenced using various amplification methods and
instrument platforms 2024 [Available from: https://www.ebi.ac.uk/biostudies/studies/S-BSST1334.

25.   Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, et al. IQ-TREE
2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. Mol Biol Evol.
2020;37(5):1530-4.

26.   Tosta S, Moreno K, Schuab G, Fonseca V, Segovia FMC, Kashima S, et al. Global SARS-CoV-2
genomic surveillance: What we have learned (so far). Infect Genet Evol. 2023;108:105405.

27.   Foulkes S, Monk EJM, Sparkes D, Hettiarachchi N, Milligan ID, Munro K, et al. Early Warning
Surveillance for SARS-CoV-2 Omicron Variants, United Kingdom, November 2021-September 2022.
Emerg Infect Dis. 2023;29(1):184-8.

28.   Xu Y, Liu T, Li Y, Wei X, Wang Z, Fang M, et al. Transmission of SARS-CoV-2 Omicron Variant
under a Dynamic Clearance Strategy in Shandong, China. Microbiol Spectr. 2023;11(2):e0463222.

29.   Markov PV, Ghafari M, Beer M, Lythgoe K, Simmonds P, Stilianakis NI, et al. The evolution of
SARS-CoV-2. Nat Rev Microbiol. 2023;21(6):361-79.

30.   van der Westhuizen HM, Soundararajan S, Berry T, Agus D, Carmona S, Ma P, et al. A consensus
statement on dual purpose pathogen surveillance systems: The always on approach. PLOS Glob Public
Health. 2024;4(11):e0003762.

31.   Oude Munnink BB, Worp N, Nieuwenhuijse DF, Sikkema RS, Haagmans B, Fouchier RAM, et al.
The next phase of SARS-CoV-2 surveillance: real-time molecular epidemiology. Nat Med.
2021;27(9):1518-24.

32.   Volz E, Mishra S, Chand M, Barrett JC, Johnson R, Geidelberg L, et al. Assessing transmissibility of
SARS-CoV-2 lineage B.1.1.7 in England. Nature. 2021;593(7858):266-9.
529