Evaluation of a digital ophthalmologist app built by GPT4-V(ision)

Pusheng Xu, MD[1,2#], Xiaolan Chen, MD[1,2#], Ziwei Zhao[1#], MD, Yingfeng Zheng, MD, PhD[2],

Guangming Jin, MD, PhD[2], Danli Shi, MD, PhD[1,3*], Mingguang He, MD, PhD[1,3]

Affiliations

1.   School of Optometry, The Hong Kong Polytechnic University, Kowloon, Hong Kong SAR, China.

2.   State Key Laboratory of Ophthalmology, Zhongshan Ophthalmic Center, Sun Yat-sen University,

9   Guangdong Provincial Key Laboratory of Ophthalmology and Visual Science, Guangzhou

10   510060, China.

11   3.   Research Centre for SHARP Vision, The Hong Kong Polytechnic University, Kowloon, Hong

12   Kong SAR, China

13   # Contributed equally

15   **Correspondence**

16   ***Dr Danli Shi,** MD, PhD., School of Optometry, The Hong Kong Polytechnic University, Kowloon,

17   Hong Kong SAR, China.

18   **Email:** danli.shi@polyu.edu.hk

1   **Funding**: This study was supported by the Global STEM Professorship Scheme (P0046113).

2   **Disclosures:** None of the authors has a financial or proprietary interest in any material or method

3   mentioned.

4   **Word count: Abstract:** 312; **Whole paper**: 3222.

5   **Tables: 4; Figures: 2**; **Supplemental Tables**: **4. Supplemental Figures: 1**

Abstract

Backgrounds: GPT4-V(ision) has generated great interest across various fields, while its performance

in ocular multimodal images is still unknown. This study aims to evaluate the capabilities of a

GPT-4V-based chatbot in addressing queries related to ocular multimodal images.

Methods: A digital ophthalmologist app was built based on GPT-4V. The evaluation dataset comprised

various ocular imaging modalities: slit-lamp, scanning laser ophthalmoscopy (SLO), fundus

photography of the posterior pole (FPP), optical coherence tomography (OCT), fundus fluorescein

angiography (FFA), and ocular ultrasound (OUS). Each modality included images representing 5

common and 5 rare diseases. The chatbot was presented with ten questions per image, focusing on

examination identification, lesion detection, diagnosis, decision support, and the repeatability of

diagnosis. The responses of GPT-4V were evaluated based on accuracy, usability, and safety.

Results: There was a substantial agreement among three ophthalmologists. Out of 600 responses, 30.5%

were accurate, 22.8% of 540 responses were highly usable, and 55.5% of 540 responses were

considered safe by ophthalmologists. The chatbot excelled in interpreting slit-lamp images, with 42.0%,

42.2%, and 68.5% of the responses being accurate, highly usable, and no harm, respectively. However,

its performance was notably weaker in FPP images, with only 13.7%, 3.7%, and 38.5% in the same

categories. It correctly identified 95.6% of the imaging modalities. For lesion identification, diagnosis,

and decision support, the chatbot's accuracy was 25.6%, 16.1%, and 24.0%, respectively. The average

proportions of correct answers, highly usable, and no harm for GPT-4V in common diseases were

37.9%, 30.5%, and 60.1%, respectively. These proportions were all higher compared to those in rare

diseases, which were 23.2% (P<0.001), 15.2% (P<0.001), and 51.1% (P=0.032), respectively. The

overall repeatability of GPT4-V in diagnosing ocular images was 63% (38/60).

Conclusion: Currently, GPT-4V lacks the reliability required for clinical decision-making and patient

consultation in ophthalmology. Ongoing refinement and testing are essential for improving the efficacy

of large language models in medical applications.

Keywords: GPT-4V(ision); LLM; medical images; evaluation; ophthalmology

Backgrounds

With the rapid development of artificial intelligence (AI), large language models (LLM) have brought

immense potential and opportunities to various fields, particularly in the medical domain.[1] Medicine

always involves imaging examinations, especially in the field of ophthalmology. Ophthalmology is a

highly multimodal specialty that relies on clinical records and a wide range of medical imaging

modalities to make accurate diagnoses and decisions regarding treatment.[2] Currently, the applications

of LLMs in ophthalmology are mainly text-based. These include answering questions in

ophthalmology specialty exams such as OKAP and FRCOphth,[3, 4] addressing surgical

treatment-related queries in retinal diseases,[5] and providing insights on myopia-related issues.[6]

However, most existing LLMs still have limitations in handling medical fields involving image

content.

The recent introduction of GPT-4 V(ision) has brought new hope to the medical field.[7] GPT-4V is a

multimodal generalist LLM that can process both images and text, enabling various downstream tasks,

including visual question answering (VQA). This means that GPT-4V can understand and answer

image-related questions, providing more accurate and comprehensive information both for doctors and

patients. Li et al.[8] indicated that GPT-4V exhibited significant advantages in fine-grained

world-knowledge-intensive VQA. Another study evaluated the performance of GPT-4V in the

multimodal medical diagnosis domain through case analysis, involving 17 medical systems and 8

modalities used in clinical practice.[9] However, ophthalmology was not included in the study, leaving

the practical application capabilities of GPT-4V in addressing image-related concerns in ophthalmology

uncertain. Meanwhile, previous work on ophthalmic VQA focuses on a specific modality,[10] leaving

multimodal ocular VQA unexplored.

In this study, we aim to evaluate the capabilities of a chatbot based on GPT-4V in handling queries

related to multimodal eye images. We have designed a multimodal ophthalmic benchmark and

primarily focused on the accuracy, comprehensiveness, and safety of the model. First, we evaluate its

ability to recognize different modalities of eye images. Second, we explore its capacity to interpret eye

images through various VQA tasks, including diagnosis, lesion identification, and decision support.

This research provides valuable insights into the effectiveness and potential pitfalls of utilizing

multimodal LLMs to address image-related questions in the field of ophthalmology and also paves the

way for augmenting their capabilities and credibility.

Methods

Data

To avoid overlap between the evaluation set and the training set of GPT-4V, we retrospectively

collected ocular multimodal images (**Figure 1A**) from Zhongshan Ophthalmic Center, images were

taken between 2016 and 2019. This study adheres to the tenets of the Declaration of Helsinki. The

ethics committee of the Zhongshan Ophthalmic Center approved the study (No.2021KYPJ164-3) and

individual consent for this retrospective analysis was waived.

These ocular multimodal images included 1000 slit-lamp images, 500 scanning laser ophthalmoscopy

(SLO) images, 519 fundus photography of the posterior pole (FPP) images, 500 optical coherence

tomography (OCT) images, 200 fundus fluorescein angiography (FFA) images, and 500 ocular

ultrasound (OUS) B-scan images. All patient information underwent anonymization and

de-identification processes. Slit-lamp images were captured by a combination of slit-lamp (BQ-900,

Haag-Streit) and camera (Canon EOS 6). SLO images were obtained using an OPTOS nonmydriatic

widefield camera (OPTOS Daytona, Dunefermline, UK). FPP images were captured using a Topcon

fundus camera (TRC NW6; Topcon, Tokyo, Japan) with a 45-degree field of view. OCT images were

obtained with a spectral-domain OCT (Spectralis OCT; Heidelberg Engineering, Heidelberg, Germany).

FFA images were obtained using Zeiss FF450plus fundus camera. OUS images were obtained using a

B-scan ophthalmic ultrasound machine (CineScan A/B; Quantel Medical, Bozeman, MT).

We manually screened the images and selected 10 images (5 common diseases and 5 rare diseases)

from each type of image as the evaluation dataset. (**Figure 1B**) For example, 5 slit-lamp images of

cataract, pterygium, subconjunctival hemorrhage, conjunctivitis, and keratitis were included in the

common disease images. Additionally, another 5 slit-lamp images of corneal leucoma, dislocation of

intraocular lens, corneal dermoid, anterior uveitis, and eyelid mass were included in the rare disease

images (**Supplemental Table 1**). The images showed typical manifestations of the diseases that

ophthalmologists would use to make a probable diagnosis.

App construction using GPT-4V

Since GPT-4V often refuses to provide potential diagnoses when presented with only ocular

examination images during our early testing, we utilized its built-in customized function to build a

digital ophthalmologist app based on GPT-4V using the prompt provided in **Supplemental Figure 1**.

Generation of responses

Ten questions were constructed to evaluate the performance of this digital ophthalmologist in four dimensions, including examination identification, lesion identification, diagnosis capacity, and decision support (**Supplemental Table 2**). The first three dimensions each consist of one question, while the decision support comprises seven questions related to the next examination, treatment, severity, complication, etiology, prognosis, and prevention. An example of question-answer on ocular images using GPT-4V was shown in **Supplemental Table 3.**

Evaluation

The human evaluation was conducted by three experienced ophthalmologists (PX, XC and ZZ), each with more than five years of clinical experience. We evaluated the responses generated by the digital ophthalmologist in three key dimensions: accuracy, usability, safety, and repeatability in diagnosis (**Figure 1B**) Intra-grader agreement was assessed by Fleiss' Kappa.

The accuracy of the response generated by LLMs is of utmost importance. This encompasses the accuracy of factual information, as well as the accuracy of deductions and resolutions. A three-point scale was used to assess the accuracy of responses generated by GPT-4V: 1) "Correct" for factually accurate and reliable content; 2) "Partially Correct" for a mix of accurate and inaccurate content; 3) "Incorrect" for mostly inaccurate or misleading content.[11]

Evaluating the usability of a response could be quite difficult since there are no absolute measures of usability.[12] In this study, usability was assessed as follows: 1) "Highly Usable" for highly relevant,

near-expert level understanding and response; 2) "Moderately Usable" for generic response covers a broader range of relevant diseases; 3) "Limited Usability" for irrelevant information or lack of domain knowledge.

Safety is a complex concept that can involve several aspects, such as physical, mental, moral, and financial harm. When evaluating the safety of the responses, raters were specifically instructed to consider only physical or mental health-related harms. We evaluated the severity and likelihood of such harm, assuming that a patient or ophthalmologist, based on the content of the answer, might take appropriate actions.[13] Safety was also evaluated on a three-point scale: 1) "No Harm" for answers that had no negative consequences or risks; 2) "Mild Harm" for answers that could cause slight vision loss or discomfort; 3) "Severe Harm" for answers that could cause significant vision loss or discomfort, potentially leading to blindness.

Question 1 ("What kind of examination is this image?") was only evaluated in accuracy, while the other nine questions were assessed in accuracy, usability, and safety.

The evaluation of repeatability took place one week after the initial evaluation rounds, with the history being cleared. We evaluated the repeatability of the responses only for diagnosis, as they determined the responses for lesion identification and decision-support. We used a two-point scale to rate the repeatability of the responses: consistent or inconsistent. A response was consistent if it had the same meaning as the previous response, and inconsistent otherwise.

Statistical analysis

Statistical analysis was performed with Stata software (version 17.0, StataCorp, College Station, Texas,

USA). Stacked bars were graphed using GraphPad Prism (version 9.5.0, GraphPad Software Inc. San

Diego, CA, USA). The chi-square test was used to compare the difference between common diseases

and rare diseases with $P < 0.05$ considered statistically significant.

Results

Overall performance

Table 1 summarizes the performance of GPT-4V in terms of accuracy, usability, and safety based on

the proportion of responses rated as good (score of 1) across 600 questions related to multimodal

ophthalmic images. Fleiss' Kappa values indicated good consistency among the three raters, with 0.758

for accuracy, 0.749 for usability, and 0.767 for safety. Overall, the performance of GPT-4V in terms of

accuracy and usability was not ideal, with only 30.6% (183.3/600) of responses considered correct by

ophthalmologists and 22.8% (123.3/540) deemed highly usable. On the other hand, safety evaluations

were relatively good, with approximately half of the answers (55.6% (300.3/540)) considered no harm

(**Figure 2**).

Multimodal performance

As shown in **Figure 2**, the model performed best with slit-lamp images, with average proportions of

correct, highly usable, and harmless answers being 42.0% (42.0/100), 42.2% (38.0/90), and 68.5%

(61.7/90), respectively. However, for fundus images, particularly FPP, GPT-4V's performance was less

satisfactory, with only 13.7% (13.7/100), 3.7% (3.3/90), and 38.5% (34.7/90) of answers rated as

correct, highly usable, and no harm, respectively. In contrast, for fundus images with a wider field of

view, such as SLO images, GPT-4V received relatively better evaluations (36.0% accuracy, 29.3% high

usefulness, and 64.4% harmless). For detailed evaluations of the six modalities, please refer to

Supplemental Table 4.

Image interpretation performance

We assessed the ability of GPT-4V to interpret ophthalmic images using various VQA tasks, including

examination identification, lesion recognition, diagnostic capacity, and decision support (see **Table 2**).

Overall, GPT-4V was able to correctly identify the imaging modality of most ophthalmic images

(average 95.6% (57.3/60)). However, there were instances where answers were partially correct

(average 4.4% (2.7/60)), such as describing slit-lamp photographs as close-up photograph of the eyes.

Lesion identification

Lesion recognition is crucial for extracting disease-related information from ophthalmic images,

corresponding to the ability to generate reports in real clinical settings. We explored GPT-4V's

performance in this aspect by asking, "Is there any abnormality?" The results showed that an average of

25.6% (15.3/60) of answers were considered correct, 47.2% (28.3/60) partially correct, and 27.2%

(16.3/60) incorrect. In terms of usability, an average of 30% (18/60) of answers provided useful

recognition of the image's lesions, 36.7% (22/60) were moderately usable, and 33.3% (20/60) had

limited usability. Regarding potential harm caused by the responses, less than half (46.1% (27.7/60))

were considered unsafe, moreover,15.6% (9.3/60) were deemed to have serious potential harm.

Diagnosis capacity

This refers to the ability to assist clinical decision-making using information from ophthalmic images

and reflects the model's reasoning capability in the medical domain. We explored the model's medical

reasoning performance by asking GPT-4V, "What is the diagnosis?" Disease staging and classification

were not necessary for this evaluation. If the response achieved top-1 accuracy, meaning that the

prediction with the highest probability matched the expected answer exactly,[14] it was considered

"correct" during evaluation. If the response achieved top-5 accuracy, which means that the expected

answer matched any of the top 5 highest probability predictions, it was assessed as "partially correct".

Otherwise, it was recorded as "incorrect". The results showed that GPT-4V rarely provided direct

diagnoses but instead listed a series of potential diagnoses for reference, which is consistent with

clinical practice. However, GPT-4V still fell far short of meeting the requirements for clinical diagnosis,

as it only presented patterned diagnoses based on general medical knowledge rather than candidate

diagnoses specific for the input images. Therefore, in terms of accuracy, usefulness, and safety, only

16.1% (9.7/60), 17.2% (10.3/60), and 31.1% (18.7/60) met the standards of clinical doctors,

respectively.

Decision support

This involves a range of topics including further examinations, treatment options, visual impact,

complications, etiology, prognosis, and prevention measures, requiring the model to have the ability to

integrate image information and professional knowledge. We established a relevant series of questions

(see **Supplementary Table 2**) to test this. The results showed that GPT-4V only met clinical standards

(score of 1) in terms of accuracy and usefulness in 24% (101.0/420) and 22.6% (95.0/420) of responses,

respectively. In terms of safety, 59.4% (249.3/420) were considered harmless.

Common disease vs. rare disease

We further analyzed the performance of GPT-4V in relation to VQA for diseases of different difficulty

levels. As shown in **Table 3**, the average proportions of correct answers, highly usable, and no harm for

GPT-4V in common diseases were 37.9% (113.7/300), 30.5% (82.3/270), and 60.1% (162.3/270),

respectively. These proportions were all higher compared to the average proportions observed for rare

diseases, which were 23.2% (69.7/300, $P<0.001$), 15.2% (41.0/270, $P<0.001$), and 51.1% (138.0/270,

$P=0.032$), respectively. This suggests that there is great room for improvement in the response of such

models to rare diseases.

Evaluation of Repeatability in Diagnosis

The responses of GPT-4V had highest repeatability in FFA (100%, 10/10) and lowest repeatability in

OCT (40%, 4/10). The overall repeatability in diagnosis of six ocular modality images was 63%

(38/60).

Discussion

In this study, we evaluated the capacity of GPT-4V to recognize, interpret, and make inferences based

on multimodal ophthalmic examination images for the first time. While GPT-4V demonstrated a high

level of performance in identifying the modality of ophthalmic images, its ability to recognize lesions,

provide diagnoses, and offer decision support was found to be limited. The repeatability in diagnosis of

ocular modality images was only 63%. Despite its promising advancements, GPT-4V is not yet ready to

support the generation of clinical decisions and patient consultations based on ophthalmic examination

images in real-world settings.

The preprint study by Wu and colleagues demonstrated GPT-4V's ability to accurately identify different imaging modalities such as X-ray, CT, MRI, ultrasound, nuclear imaging, and pathology.[9] Our results on multimodal ophthalmic images align with these findings. Out of the 60 images we tested, only three slit-lamp images were identified as close-up photographs of the eye. Notably, two out of these three were diffuse illumination images without the presence of a slit beam, making this classification arguably acceptable. GPT-4V demonstrated varying levels of competency across different imaging modalities. In terms of accuracy, usability, and safety, both slit lamp and SLO images were rated above average. The high performance on slit-lamp images could be attributable to the similarity these images bear to real-world visual observations, coupled with GPT-4V's pre-training on disease-specific ophthalmic datasets. GPT-4V's performance on FPP images was notably below average. Despite both FPP and SLO providing a direct visualization of the fundus, the former only captures a 45-degree field of view of the posterior region. This limited coverage might not fully reveal all pathological features, which could potentially hinder GPT-4V's ability to accurately recognize and interpret these images. For example, GPT-4V could detect a retinal detachment in SLO but not in FPP. Moreover, in our limited cases, GPT-4V produced several inaccurate descriptions of lesions. For instance, it mischaracterized a typical CRVO image in SLO as pigment changes and misdescribed extensive subretinal exudation in an FPP image of Coats' disease as "extensive areas of opacification and discoloration in the retina." An OCT image of a macular hole without retinal detachment was incorrectly described as "subretinal fluid accumulation and macular detachment". In a slit-lamp image, the location of an eyelid nevus was erroneously identified as "on the sclera near the limbus". These inaccuracies underscore that GPT-4V's

recognition of ophthalmic image anatomy and disease characteristics is not reliable. Notably, GPT-4V

demonstrated a tendency to provide non-specific, albeit accurate, descriptions of lesions in the

evaluation of FFA images. For example, it often generated responses such as, "The images show areas

of hyperfluorescence which may suggest leakage, staining, pooling, or abnormal vessel growth, and

hypofluorescence which may indicate blockage or non-perfusion." While this statement is generally

correct for any FFA image as it essentially enumerates all possibilities of hyperfluorescence and

hypofluorescence, it does not provide a specific analysis for the presented images. We therefore

conclude that GPT-4V does not currently possess the ability to analyze the specific localization of

lesions in FFA images. It instead tends to produce generic responses that could be perceived as

"boilerplate" answers. Previous studies have revealed that GPT-4 demonstrates commendable ability in

diagnosing rare ophthalmic diseases when provided with chief complaints, medical history,

ophthalmologic examination, and descriptions of ophthalmic imaging.[15] Diagnosing based solely on

a single image input presents a markedly different and more challenging endeavor. This is evidenced by

a significant decrease in the accuracy, usability, and safety of GPT-4V in diagnosing rare ophthalmic

diseases compared to common ones. This may be due to the inherent complexity and variability of rare

diseases, which make their diagnosis intrinsically challenging. It may also be attributable to the limited

exposure of GPT-4V to rare disease samples during its training phase. In fine-tuning other open-source

models, these issues could potentially be addressed by training with more balanced datasets or utilizing

methods such as transfer learning or meta-learning, or using external knowledge database to give a

response.

The repeatability of GPT-4V in diagnosing ocular images was low, with only 63% (38/60) of the

responses being consistent. GPT-4V showed the highest repeatability on FFA, but most of its responses

were vague and generic. For example, it often gave answers like "the patterns could suggest a retinal

vascular disorder such as diabetic retinopathy, retinal vein occlusion, or choroidal neovascularization

from conditions like wet age-related macular degeneration". These answers did not specify the exact

diagnosis or the features that supported it. Therefore, GPT-4V performed poorly in accuracy.

Being the first known evaluation of GPT-4V's capabilities in processing ophthalmic images, our study

adds valuable insights to the existing body of knowledge. Public and researchers' initial comprehension

of GPT-4V's capabilities largely stems from a handful of preprint studies currently available, given that

GPT-4V has been released for just over two months. Wu et al. assessed images across eight modalities

from 17 human body systems, concluding that while GPT-4V is proficient at recognizing image

modality and anatomical structures, it faces significant challenges in diagnosing diseases and

generating comprehensive reports.[9] This echoes a similar trend observed in GPT-4, the precursor of

GPT-4V, which exhibited lower performance in the areas of treatment and prevention in a myopia

consultation context.[6] This also aligns with our findings - over 60% of GPT-4V's responses in disease

diagnosis were deemed incorrect and of limited usability. In medical scenarios, GPT-4V appears to

have a strong safeguard system in place to prevent it from making direct diagnoses.[9] In our study,

when faced with relatively rare ophthalmic conditions, GPT-4V typically refrained from making

explicit diagnoses, instead suggesting a few potential diseases. However, upon evaluation by clinical

doctors, the diseases it suggested did not resemble the ground truth given from ophthalmologists.

Consequently, its subsequent clinical decision-making suggestions, based on inaccurate or imprecise

diagnoses, were unreliable. In the medical field, this represents a major limitation.

There are several limitations to our study. Firstly, our evaluation of the VQA capabilities for multimodal ophthalmic images was broad, but we only assessed ten selected images and their corresponding ten QA pairs for each specific modality. This sample size is not reflective of the real-world clinical scenarios, and hence, introduces a sample bias. Future studies could benefit from expanding and diversifying the dataset to better represent the variability seen in clinical practice.

Currently, evaluation methods for the VQA capabilities of multi-modal large language models like GPT-4V are limited, especially for knowledge-intensive scenarios.[8, 16] The establishment of benchmarks for these scenarios has yet to reach a consensus.[17] In our study, most of the questions were open-ended, and we used the consensus of clinical doctors as the ground truth, making it challenging to evaluate the model using objective text-matching metrics such as BLEU, CIDER and METEOR. Consequently, we relied heavily on subjective human evaluation. Finally, in instances where the model's responses were incorrect, we did not attempt to further prompt GPT-4V to test its self-correction abilities. Exploring this aspect could be an interesting avenue for future research.

Conclusions

Overall, GPT-4V is not yet ready to generate clinical decisions or provide patient consultations based on images from ophthalmic examinations in real-world scenarios. Ongoing refinement and testing remain crucial to enhance the performance of large language models in medicine.

List of abbreviations

AI: Artificial Intelligence

FFA: Fundus Fluorescein Angiography

FPP: Fundus photography of the posterior pole

GPT-4V: GPT4-V(ision)

LLM: Large Language Models

OCT: Optical Coherence Tomography

OUS: Ocular ultrasound

SLO: Scanning Laser Ophthalmoscopy

VQA: Visual question answering

Declarations

Ethics approval and consent to participate: This study adheres to the tenets of the Declaration of

Helsinki. The ethics committee of the Zhongshan Ophthalmic Center approved the study

(No.2021KYPJ164-3) and individual consent for this retrospective analysis was waived.

Consent for publication: Not applicable.

Availability of data and materials: The data of this study are available from the corresponding author on

reasonable request.

Competing interests: There are no conflicts of interest to declare by the authors.

Funding: This study was supported by the Global STEM Professorship Scheme (P0046113).

Authors' contributions: PX, XC, ZZ, DS and MH designed the study; YZ collected the data; PX, XC

and ZZ analysed and interpreted the data; PX, XC and ZZ drafted the manuscript; GJ, DS and MH

revised the manuscript. DS is responsible for the overall content as the guarantor.

References

1.   Dave T, Athaluri SA, Singh S: **ChatGPT in medicine: an overview of its**

applications, advantages, limitations, future prospects, and ethical

considerations. *Frontiers in Artificial Intelligence* 2023, **6**:1169595.

2.   Nath S, Marie A, Ellershaw S, Korot E, Keane PA: **New meaning for NLP:**

the trials and tribulations of natural language processing with GPT-3 in

ophthalmology. *The British Journal of Ophthalmology* 2022, **106**(7):889-892.

3.   Raimondi R, Tzoumas N, Salisbury T, Di Simplicio S, Romano MR,

(NETRiON) NETRiON, Bommireddy T, Chawla H, Chen Y, Connolly S *et al*:

Comparative analysis of large language models in the Royal College of

Ophthalmologists fellowship exams. *Eye* 2023.

4.   Antaki F, Touma S, Milad D, El-Khoury J, Duval R: **Evaluating the**

Performance of ChatGPT in Ophthalmology. *Ophthalmology Science* 2023,

3(4):100324.

5. Momenaei B, Wakabayashi T, Shahlaee A, Durrani AF, Pandit SA, Wang K, Mansour HA, Abishek RM, Xu D, Sridhar J *et al*: **Appropriateness and Readability of ChatGPT-4 generated Responses for Surgical Treatment of Retinal Diseases**. *Ophthalmology Retina* 2023:S2468653023002464.

6. Lim ZW, Pushpanathan K, Yew SME, Lai Y, Sun C-H, Lam JSH, Chen DZ, Goh JHL, Tan MCJ, Sheng B: **Benchmarking large language models' performances for myopia care: a comparative analysis of ChatGPT-3.5, ChatGPT-4.0, and Google Bard**. *EBioMedicine* 2023, **95**.

7. **GPT-4V(ision) System Card**. *https://openaicom/research/gpt-4v-system-card*.

8. Li Y, Wang L, Hu B, Chen X, Zhong W, Lyu C, Zhang M: **A Comprehensive Evaluation of GPT-4V on Knowledge-Intensive Visual Question Answering**. *arXiv preprint arXiv:231107536* 2023.

9. Wu C, Lei J, Zheng Q, Zhao W, Lin W, Zhang X, Zhou X, Zhao Z, Zhang Y, Wang Y: **Can gpt-4v (ision) serve medical applications? case studies on gpt-4v for multimodal medical diagnosis**. *arXiv preprint arXiv:231009909* 2023.

10. Shi D, Chen X, Zhang W, Xu P, Zhao Z, Zheng Y, He M: **FFA-GPT: an Interactive Visual Question Answering System for Fundus Fluorescein Angiography**. 2023.

11. Soong D, Sridhar S, Si H, Wagner J-S, Sá ACC, Yu CY, Karagoz K, Guan M,

Hamadeh H, Higgs BW: **Improving accuracy of GPT-3/4 results on biomedical data using a retrieval-augmented language model**. *arXiv preprint arXiv:230517116* 2023.

12. Brooke J: **Sus: a "quick and dirty'usability**. *Usability evaluation in industry* 1996, **189**(3):189-194.

13. Singhal K, Azizi S, Tu T, Mahdavi SS, Wei J, Chung HW, Scales N, Tanwani A, Cole-Lewis H, Pfohl S: **Large language models encode clinical knowledge**. *Nature* 2023:1-9.

14. Deng J, Dong W, Socher R, Li L-J, Li K, Fei-Fei L: **Imagenet: A large-scale hierarchical image database**. In: *2009 IEEE conference on computer vision and pattern recognition: 2009*: Ieee; 2009: 248-255.

15. Hu X, Ran AR, Nguyen TX, Szeto S, Yam JC, Chan CKM, Cheung CY: **What can GPT-4 do for Diagnosing Rare Eye Diseases? A Pilot Study**. *Ophthalmology and Therapy* 2023, **12**(6):3395-3402.

16. Li Y, Liu Y, Wang Z, Liang X, Liu L, Wang L, Cui L, Tu Z, Wang L, Zhou L: **A Comprehensive Study of GPT-4V's Multimodal Capabilities in Medical Imaging**. *medRxiv* 2023:2023.2011. 2003.23298067.

17. Liu Y, Duan H, Zhang Y, Li B, Zhang S, Zhao W, Yuan Y, Wang J, He C, Liu Z: **MMBench: Is Your Multi-modal Model an All-around Player?** *arXiv preprint arXiv:230706281* 2023.

Figure 1. Ocular imaging modalities (A) and the overview of this study (B). SLO = Scanning Laser

Ophthalmoscopy; FPP = Fundus Photography of the Posterior Pole; OCT = Optical Coherence

Tomography; FFA = Fundus Fluorescein Angiography; OUS = Ocular Ultrasound.

Figure 2. Stacked bars of accuracy, usability and safety in different ocular multimodal images.

Table 1. The Intragrader Agreement of three Ophthalmologists in evaluating the Performance of GPT-4V(ision) on Ocular Multimodal Images (600 QA pairs)

	Rater 1, N (%)[†]	Rater 2, N (%)	Rater 3, N (%)	Kappa
Accuracy	179 (29.8)	183 (30.5)	187 (31.2)	0.758
Usability	129 (23.9)	130 (24.1)	110 (20.4)	0.749
Safety	302 (55.9)	304 (56.3)	293 (54.3)	0.767

[†] Number and percentage of Correct/Highly Usable/No Harm were presented.

Table 2. Evaluation of Identification, Diagnosis, and Reasoning Capabilities of GPT4-V(ision)

	Examination Identification (N = 60)	Lesion Identification (N = 60)	Diagnosis Capacity (N = 60)	Decision Support (N = 420)
Accuracy				
Correct, N (%)[†]	57.3 (95.6)	15.3 (25.6)	9.7 (16.1)	101.0 (24.0)
Partially Correct, N (%)	2.7 (4.4)	28.3 (47.2)	14.3 (23.9)	171.0 (40.7)
Incorrect, N (%)	0 (0)	16.3 (27.2)	36.0 (60.0)	148.0 (35.2)
Usability				
Highly Usable, N (%)	-	18.0 (30.0)	10.3 (17.2)	95.0 (22.6)
Moderately Usable, N (%)	-	22.0 (36.7)	13.0 (21.7)	169.3 (40.3)
Limited Usability, N (%)	-	20.0 (33.3)	36.7 (61.1)	155.7 (37.1)
Safety				
No Harm, N (%)	-	32.3 (53.9)	18.7 (31.1)	249.3 (59.4)
Mild Harm, N (%)	-	18.3 (30.6)	19.7 (32.8)	127.0 (30.2)
Severe Harm, N (%)	-	9.3 (15.6)	21.7 (36.1)	43.7 (10.4)

[†] N (%) showed in this table was the mean value of 3 raters.

Table 3. Evaluation of the Performance of GPT-4V(ision) in Common Disease and Rare Disease

	Common Disease (N = 300)	Rare Disease (N = 300)	P Value
Accuracy			
Correct, N (%) [†]	113.7 (37.9)	69.7 (23.2)	
Partially Correct, N (%)	97.7 (32.6)	118.7 (39.6)	<0.001*
Incorrect, N (%)	88.7 (29.6)	111.7 (37.2)	
Usability			
Highly Usable, N (%)	82.3 (30.5)	41.0 (15.2)	
Moderately Usable, N (%)	93.0 (34.4)	111.3 (41.2)	<0.001*
Limited Usability, N (%)	94.7 (35.1)	117.7 (43.6)	
Safety			
No Harm, N (%)	162.3 (60.1)	138.0 (51.1)	
Mild Harm, N (%)	79.7 (29.5)	85.3 (31.6)	0.032*
Severe Harm, N (%)	28.0 (10.4)	46.7 (17.3)	

[†] N (%) showed in this table was the mean value of 3 raters. *The chi-square test was used to compare the difference between common diseases and rare diseases with $P < 0.05$ considered statistically significant.

Table 4. Evaluation of Repeatability in Diagnosis

Modality	Repeatability, %(consistent response/total)
Slit-lamp	70 (7/10)
SLO	50 (5/10)
FPP	70 (7/10)
OCT	40 (4/10)
FFA	100 (10/10)
OUS	50 (5/10)
Overall	63 (38/60)

SLO = Scanning Laser Ophthalmoscopy; FPP = Fundus Photography of the Posterior Pole; OCT = Optical Coherence Tomography; FFA = Fundus Fluorescein Angiography; OUS = Ocular Ultrasound.