Molecular Analysis of Non-structural Protein 1 (*NSP1*) in Children Infected with Rotavirus in Babylon Province, Iraq: A Cross-Sectional Study

Mohammed Mohsen M. [a], Zaytoon A Al-khafaji [a], Layth Jasim Mohammed [b*] and Nima Yakhchalian[c]

[a] Department of Microbiology, Faculty of Medicine, Babylon University, College of Medicine, Iraq

[b] Department of Pharmacology, Faculty of Medicine, Babylon University, Hilla City, Babylon Governorate 51002, Iraq

[c] Department of Biology, Faculty of Science, Razi University, Kermanshah, Iran

Corresponding authors: Layth Jasim Mohammed; Laith.bi@yahoo.com; med996.layth.jasim@uobabylon.edu.iq Tel & Fax: 00964-782-5557384.

Abstract

Globally, a considerable number of infants and children younger than five are falling victim to diarrheal diseases predominantly caused by rotaviruses, which are non-enveloped, double-stranded RNA (dsRNA) viruses able to cause acute gastroeteritis and extragastrointestinal complications. Annually, human rotaviruses cause two million hospitalizations and over 500,000 deaths worldwide. Rotaviral replication, pathogenesis, and immune evasion are propagated by non-structural protein 1 (NSP1), encoded by segment five of their dsRNA genome. We examined 60 urine and stool samples from children aged 2-60 months admitted to an Obstetric and Children Hospital in Babylon over a 60-day time period with the diagnosis of acute group A rotavirus gastroenteritis. This study aimed to check the presence of NSP1 by immunochromatography assay and RT-qPCR. Immunochromatography assay detected NSP1 in 100% of urine and stool samples; however, RT-qPCR only detected it in 66.7% of urine and 50% of stool samples. RT-qPCR found 12 out of 30 urine and stool samples positive, accounting for 40% of participants. No significant correlations between RT-qPCR results and sociodemographic factors were found. Results found 73.3% of acute gastroenteritis cases were in children under two. Additionally, the urinary detection of NSP1 suggests that rotaviruses may cause extra-gastrointestinal infections, e.g., systemic infection or viremia.

Keywords: Extra-intestinal; NSP1; RT-qPCR; Rotavirus; Viremia

1. Introduction

Diarrheal disease is still recognized as the major cause of mortality and morbidity among the children less than 5 years of age globally (Vos et al. 2020). Prior scientific discoveries dating back earlier than 1973 had already found rotavirus causing diarrheal disease in animals, including mice (Adams and Kraft 1963), vervet monkeys (*Cercopithecus aethiops pygerythrus*) (Malherbe and Ulrich 1963), and cattle (Mebus et al. 1969). After these initial discoveries, rotavirus was later detected infecting members of different mammalian taxa such as the order Chiroptera (Greenberg and Estes 2009; He et al. 2013). In addition, rotaviruses have also been reported to infect avian species (Otto et al. 2012; Kindler et al. 2013). Since 1973, rotavirus has been found responsible for causing life-threatening, acute dehydrating gastroenteritis across the world, not only in infants and young children (Bishop et al. 1973; Flewett et al. 1973) who are particularly susceptible to infection, but also in older individuals in whom the infection is typically mild or even asymptomatic owing to the exposure-

induced immunity developed over time in response to repeated infections (Parashar et al. 1998). Following all aforementioned discoveries, nowadays rotaviruses are recognized as the most predominant diarrheal pathogens (Parashar et al. 1998), and rotavirus-associated diarrhea, also referred to as rotaviral diarrhea, is identified as the primary cause of viral diarrhea in the world (Chen et al. 2022).

Human rotaviruses are the major etiologic agents of infantile acute gastroenteritis, leading to roughly two million hospitalizations and more than 500,000 deaths each year worldwide (Mousavi-Nasab et al. 2020). Presently, human rotaviruses account for approximately 1,600 pediatric deaths per day globally (Angel et al. 2007). Even though much of the disease burden is concentrated in developing countries, in particular African and Asian ones (Troeger et al. 2018), due to the present socioeconomic issues (Angel et al. 2007); developed countries also face considerable impacts (Malek et al. 2006). As a case in point, it has been reported that rotaviruses are responsible for nearly 4-5% of all pediatric hospitalizations in the United States; further estimates indicate that 1 out of 67 to 1 out of 85 children are admitted to hospitals due to rotavirus-associated gastroenteritis by their fifth year of age (Charles et al. 2006). Although rotavirus has conventionally been viewed as a pathogen primarily infecting instestinal mature enterocytes, leading to the destruction of absorptive enterocytes and consequent diarrhea, this interpretation has been called into question in the light of recent evidence gathered from both patients and animal models documenting extra-intestinal rotavirus-associated pathologic impacts.

Out of the 10 species of rotaviruses (A-J) identified until now, only four of them (A, B, C and H) have been observed infecting humans (Matthijnssens et al. 2008a). Once the patient has been infected with rotavirus and the incubation period of 1-3 days has elapsed, varying clinical presentations start to appear (Bernstein 2009), including fever, tachycardia, diminished urination, abdomincal cramp, and complications of dehydration (Hoxha et al. 2015).

A multitude of pathologic circumstances in humans have been associated with rotavirus infection, including hepatobiliary disorders such as chronic and acute hepatitis and rotavirus-induced cholestasis (Ramig 2007); neurological deficits and neuropathologic manifestations, e.g. lesions in corpus callosum (Laizane et al. 2019), benign convulsions (Takanashi et al. 2010; Laizane et al. 2019), cerebellitis (Takanashi et al. 2010), encephalitis (Ushijima et al. 1986), encephalopathy (Maureen et al. 2001; Fukuda et al. 2009), and meningitis (WONG et al. 1984); type 1 diabetes mellitus (T1DM);

respiratory disorders (Goldwater et al. 1979) such as pneumonitis (Grech et al. 2001), cardiovascular diseases, e.g. hypertrophic cardiomyopathy and myocarditis (Grech et al. 2001; Cioc and Nuovo 2002), renal failure and thrombocytopenia (Ramig 2007).

Taxonomically, rotaviruses belong to the *Rotavirus* genus which falls within the *Reoviridae* family (Brunet et al. 2000). *Rotavirus* genus comprises at least seven serogroups, consisting of *Rotavirus* A to G, differentiated on the basis of the electrophoretic mobility of their genomic segments and distinct antigenic properties (Bitton 2002; Bonkoungou et al. 2010). Out of these seven serogroups, the only Groups able to infect humans are Groups A, B, and C, out of which Group G is responsible for the common, life-threatening gastroenteritis among children in the world (Richardson et al. 1998; Kang et al. 2004a). Rotaviruses are non-enveloped, icosahedral viruses, charateristically consisting of an 11-segment dsRNA genome with nearly 18,500 base pairs (Trask et al. 2012; Dian et al. 2021).

Rotavirus particles are complex and of large size (1000 Å), comprising three concentric protein capsid layers surrounding their genome (Crawford et al. 2017). The dsRNA in rotavirus contains instructions for encoding six structural proteins that make up viral proteins (VPs), also known as viral particles, including *VP1*, *VP2*, *VP3*, *VP4*, *VP6*, and *VP7*, and also six nonstructural proteins (NSPs) which are known as *NSP1*, *NSP2*, *NSP3*, *NSP4*, *NSP5*, and NSP6 (Matthijnssens et al. 2008b; Trask et al. 2012).

VPs are the chief regulators of host specificity, producing viral transcripts, cell entry, and generation of immune response (Crawford et al. 2017). With regard to NSPs, these are synthesized inside the infected host cells, and among these proteins, *NSP1*, *NSP2*, *NSP3*, *NSP5*, and *NSP6* play a significant role in the replication of viral genome, and *NSP4* regulates morphogenesis (Matthijnssens et al. 2008b). NSPs also serve as antagonists to the host's innate immunity (Crawford et al. 2017). Genomic segments in rotavirus are monocistronic, exhibiting a 1:1 correspondence between each of the segments and every single protein encoded, and the only exception is gene 11 which is bicistronic, encoding both proteins *NSP6* and *NSP5* (Matthijnssens et al. 2008b).

From a morphological standpoint, electron microscopy of the virion particle in rotaviruses exhibits a morphology closely resembling that of a wheel; therefore, they were given the name *rotavirus*, where the linguistic morpheme "rota" refers to a wheel-like shape in Latin (Cunliffe et al. 2014). They possess an extremely stable triple-layered particle (TLP) measuring nearly 85

nanometers in diameter (Long and McDonald 2017), coated with short spikes protruding outward and helping the virus attach to the target receptors on the surface of host cells. In general, the structure of rotavirus virion includes an inner layer that primarily consists of *VP2* enclosing an RNA-dependent RNA polymerase (RdRp) called *VP1*; the intermediate layer is composed of *VP6* trimers, serving as the molecular characteristic responsible for the identification of rotavirus groups, subgroups, and species; and an outer virion layer (Crawford et al. 2017). The outer layer which is able to trigger immune reponse (Li et al. 2009; Settembre et al. 2011; Papa et al. 2021) is mainly composed of capsid glycoprotein *VP7* (Estes and Kapikian 2007) and the protease-sensitive hemagglutinin spike antigen *VP4* (Crawford et al. 2001) broken into *VP5* and *VP8* following proteolysis (Jiang et al. 2023). In rotaviruses, the virion core consists of VP2, VP1, VP3, and the dsRNA genome (Papa et al. 2021). With regard to rotavirus serotypes, the classification is performed on the basis of spike protein *VP4*, the serotype which is referred to as P-type, and *VP7*, the serotype known as G-type, representing protease-sensitive *VP4* and glycoprotein *VP7*, respectively (Cunliffe et al. 2014).

With respect to the diagnosis of rotavirus infection, clinical examinations cannot distinguish rotavirus-induced diarrheal presentations from gastroenteritis caused by other infectious pathogens infecting gastrointestinal tract, such as *Salmonella spp*, *Escherichia coli*, astroviruses, and enteric adenoviruses (Crawford et al. 2017). As regards the laboratory diagnosis of rotavirus infection, a wide variety of techniques have been utilized so far, including ICT (Vontas et al. 2019) and enzyme-linked immunosorbent assay (ELISA) (Hasegawa et al. 2017) for the detection of rotavirus antigen, polyacrylamide gel electrophoresis for the detection of rotavirus dsRNA genome (Herring et al. 1982), electron microscopy (Roingeard et al. 2019), and RT-PCR assays due to higher sensitivity and the possibility for genotyping the virus (Pickering et al. 1988; Kang et al. 2004b).

NSP1 is the protein encoded in segment five of the ds-RNA genome in rotaviruses and has been found to be responsible for antagonizing host IFN signaling (Hou et al. 2021). Once NSP1 has been expressed by rotavirus-infected cells, it propagates proteasome-mediated degradation of IFN regulatory factor 3 (IRF3) (Graff et al. 2002; Barro and Patton 2005) and also promotes the degradation of other IFN regulatory factors (IRFs), specifically IFN regulatory factor 9 (IRF9), IFN regulatory factor 7 (IRF7), and IFN regulatory factor 5 (IRF5) (Arnold et al. 2013). IRFs are the transcription factors necessary for the induction of IFN, and the promotion of proteosomal

degradation of them is one of the major strategies adopted by rotaviruses to evade immune response (Barro and Patton 2005).

Since rotavirus-encoded NSP1 is considered a specifically significant protein in pathogenesis and replication of rotaviruses, this study placed its concentration on the percentage of rotavirus-infected patients who tested positive for the NSP1 protein, addressing the question of whether the detection of NSP1 protein in their urine and stool samples can be of any diagnostic value.

2. Materials and Methods

2.1. Sample Collection

A collective of 60 samples, comprising both stool and urine, were obtained from rotavirus-infected children aged 2-60 months who were admitted to an Obstetric and Children Hospital in Babylon. All samples were collected over a 60-day time period. The amount of samples collected from each patient exhibited variation, with urine samples ranging from 1 to 1.5 milliliters, and stool samples ranging from 2 to 20 grams. The samples were collected using disposable containers and were subsequently stored at temperatures of -20°C and -80°C for further analytic tests, including, Immunochromatographic test (ICT) to identify *Rotavirus* antigens in both stool and urine and RT-qPCR to detect the *Rotavirus NSP1* in both urine and stool.

2.2. Diagnostic Methods for the Detection of Rotavirus

2.2.1. Quick profile™ Adeno/Rotavirus Combo Test

The Quick profile™ Adeno/Rotavirus Combo test was used as a point-of-care. qualitative immunochromatographic assay, which is commercially available and has widely been employed for the detection of viral enteropathogens, including rotavirus and adenovirus, in human stool samples (Weitzel et al. 2007; Artiran et al. 2017). The immunochromatography assay has a number of advantages since it is neither dependent on trained laboratory technicians nor requires specific equipment (Jiang et al. 2018). However, it lacks high sensitivity (Jiang et al. 2018), which made us use RT-qPCR as another complementary molecular detection method.

2.2.2. Reverse Transcription-Quantitative Polymerase Chain Reaction (RT-qPCR)

RT-qPCR was employed to detect the *NSP1* gene in segment five of rotaviral ds-RNA genome. RT-qPCR has been extensively used for the diagnosis of group A rotaviruses (Ward et al. 2013; Anaya-

Molina et al. 2018) due to high sensitivity, possibility for the quantitative analysis of viral load, low risk of cross-contamination, and reduced chances of false-positive results.

2.3. Ethical Approval and Consent

Prior to sample collection, all participants engaged in this study were provided with detailed information, and verbal consent was obtained from each of them. This research was given ethical approval under the reference number BMS/0231/016, on behalf of the Ethics Committee at the College of Medicine, University of Babylon, Iraq.

2.4. Statistical Analysis

Statistical analysis was conducted using SPSS version 23. Categorical information was presented using frequencies and percentages, whereas continuous variables were presented using means and standard deviations (SD). The relationship between categorical variables was assessed using Fisher's exact test and Pearson's chi-square (X2). A p-value of less than 0.05 was considered to indicate statistical significance.

3. Results

According to ICT results for identification of human rotavirus in urine and stool samples (Figure 1) at a hospital for maternity and children in Babylon, all 30 samples tested positive, indicating a 100% positivity rate (Table 1).

Table 1. The Categorization of Study Participants Based on ICT Results (N = 30).

Clinical Samples	Total Samples	ICT Positive Results No. (%)	ICT Negative Results No. (%)
Stool	30	30 (100%)	0 (0%)
Urine	30	30 (100%)	0 (0%)
Total	60	60 (100%)	0(0%)



Figure 1. Identification of Human Rotavirus by the ICT in Urine and Stool Samples.

3.1. The Distribution of Study Participants According to RT-qPCR Results (Urine and Stool) Samples

Figures 2 and 3 illustrate the distribution of participants on the basis of RT-qPCR results for urine and stool samples, encompassing both negative and positive cases. According to RT-qPCR analysis for NSP1 detection, 66.7% of the study participants (N = 20) tested positive in urine samples, whereas 50.0% of the study participants (N = 15) tested positive in stool samples.



Figure 2. Distribution of Study Participants Based on RT-qPCR Results for Urine Samples.



Figure 3. Distribution of Study Participants Based on RT-qPCR Results for Stool Samples.

3.2. The Association between PCR Results and Sociodemographic Variables

Tables 2 and 3 illustrate the relationship between RT-qPCR results for urine and stool samples, respectively, and sociodemographic variables (e.g., age and gender), including both positive and negative cases. The analysis revealed that there was no significant association between RT-qPCR results and sociodemographic variables.

Table 2. Association between PCR Results (Urine Sample) and Sociodemographic Variables.

Sociodemographic characteristics	PCR (Urine)		Total	P-value	Odds ratio	95% CI
	Positive	Negative				
Age (months)				0.372	2.429	0.391-15.085
≥ 6 months	17 (85.0)	7 (70.0)	24 (80.0)			
< 6 months	3 (15.0)	3 (30.0)	6 (20.0)			
Total	20 (100.0)	10 (100.0)	30 (100.0)			
Gender				0.682	1.714	0.278-10.589
Female	6 (30.0)	2 (20.0)	8 (26.7)			
Male	14 (70.0)	8 (80.0)	22 (73.3)			
Total	20 (100.0)	10 (100.0)	30 (100.0)			

* **P value ≤ 0.05** was significant (Fisher's exact test).

Table 3. Association between PCR Results from Stool Samples and Sociodemographic Variables.

Socio-demographic characteristics	PCR (stool)		Total	P-value	Odds ratio	95% CI
	Positive	Negative				
Age (months)				1.000	1.000	0.167-5.985
≥ 6 months	12 (80.0)	12 (80.0)	24 (80.0)			
< 6 months	3 (20.0)	3 (20.0)	6 (20.0)			
Total	15 (100.0)	15 (100.0)	30 (100.0)			
Gender				0.682	2.00	0.381-10.511
Female	5 (33.3)	3 (20.0)	8 (26.7)			
Male	10 (66.7)	12 (80.0)	22 (73.3)			
Total	15 (100.0)	15 (100.0)	30 (100.0)			

* **P value ≤ 0.05** was significant (Fisher's exact test).

3.3. Rotavirus Detection by RT-qPCR

In order for the ICT results to be evaluated, RT-qPCR was carried out to detect the *NSP1* gene in segment five of rotavirus ds-RNA.

According to the RT-qPCR results shown in Table 4, out of the 30 rotavirus-infected patients from Babylon Province, 15 stool samples (50%) and 20 urine samples (66.66%) tested positive for rotavirus *NSP1*.

Table 4. Detection of Human Rotavirus in Stool and Urine Samples using RT-qPCR.

Clinical samples	Total samples	RT-qPCR (*NSP1*) Positive (%)	RT-qPCR (*NSP1*) Negative (%)
Stool	30	15 (50%)	15 (50%)
Urine	30	20 (66.66%)	10 (33.33%)
Total	**60**	**35 (58.33%)**	**25 (41.66%)**

3.4. The Runs of Real-Time PCR

In the first run, a total of 29 samples from both urine and stool, exhibited positive results, as depicted in Figure 4.



Figure 4. The First Run of Real-Time PCR for Rotavirus.

Turning to the second run, six samples were detected and exhibited positive results (Figure 5).



Figure 5. The Second Run of Real-Time PCR for Rotavirus.

3.5. RT-qPCR Efficiency

Figures 6 and 7 show the RT-qPCR standard curve and amplification plot of positive samples in the standard curve for the rotavirus *NSP1* gene, respectively. RT-qPCR exhibited an efficiency of 103%.



Figure 6. Real-Time PCR Standard Curve for the Human Rotavirus *NSP1* Gene.



Figure 7. Real-Time PCR Amplification Plot of Positive Samples in the Standard Curve for

4. Discussion

Diarrheal diseases are globally recognized to be a major cause of morbidity and mortality, specifically in developing countries where they account for 15–30% of deaths among children under the age of five (Karampatsas et al. 2018). Prior to the introduction of vaccines against rotavirus, they accounted for around 37% of diarrhea-associated deaths, leading to 450,000 fatalities worldwide, and also accounted for up to 50% of diarrhea-associated hospitalizations in developed countries (Nugent and Stewart 2023).

In the current research, the children under the age of five who presented with acute gastroenteritis were subjected to the analysis of their urine and stool samples in order for the presence of the Rotavirus A antigen to be detected.

The research conducted in Basrah city, Iraq by (Yaqoob et al. 2016), showed through the utilization of ICT that out of 300 children admitted to hospitals with gastroenteritis, 93 of them (31%) had been infected with rotavirus. This diagnostic technique has been widely employed in studies due to its simplicity, straightforward interpretation, and minimal equipment required.

Numerous studies (Ahmed et al. 2006; Khoury et al. 2011) carried out in Iraq have demonstrated a correlation between rotavirus infection and diarrhea in children below the age of five, confirming the findings of multiple studies (Ahmad Abdul AL-Kader et al. 2016; Fenjan and Jarullah 2019).

For the first time in Iraq, this research sets out to introduce a novel diagnostic approach to detecting rotavirus in urine samples. The detection of rotavirus antigen in urine proposes that rotavirus antigen can be excreted through urine, a finding that fully cocurs with other reports (Vollet III et al. 1981; Zheng et al. 1991); however, urine is not believed to serve as a viral transmission route (Anderson and Weber 2004). We still stand in need of further research to address the inquiry of whether PH and any other chemical compounds found in urine can inactivate rotavirus infectivity or virus excreted in urine can still retain its infectivity under specific curcumstances (Crawford et al. 2006).

Urine samples were collected from 30 patients with acute gastroenteritis, and RT-qPCR results tested positive for rotavirus antigen in urine samples of 20 out of 30 patients. Among 12 patients whose urine samples tested positive for rotavirus antigen, the highest levels of antigen had been detected, and this finding was consistent with those of their stool samples. The detection of rotavirus antigen in urine suggests that it is likely for the virus to be transmitted through any routes except the

typical fecal-oral way. Other studies, for example (Yokoyama et al. 2011) show that rotavirus is able to evade gastrointestinal tract and cause extra-gastrointestinal pathologic involvement.

The detection of rotavirus in 20 out of 30 urine samples obtained from children with rotaviral diarrhea indicates that antigenemia and potentially viremia can occur as a result of rotavirus infection (Crawford et al. 2006), as indicated by other findings (Blutt et al. 2003; Ramig 2004). According to a study (Li and Wang 2003), rotavirus antigen and/or RNA have been detected in multiple organs, including the central nervous system, spleen, heart, kidney, testes, and bladder, of children who have died as a result of rotavirus infection. This finding is in agreement with other postmortem examinations (Lynch et al. 2003) that detected rotavirus RNA in organs outside the gastrointestinal tract, including heart, lymphoid tissues, bladder, kidney, and testes in children who died of rotavirus infection (Lynch et al. 2003). As per another study (Blutt et al. 2003), the presence of rotavirus both antigen and RNA, in sera collected from rotavirus-infected animals has also been confirmed, which suggests the possibility that rotavirus can cause systemic infection.

These reports support the hypothesis that rotavirus can infect organs outside the gastrointestinal tract; nevertheless, the precise sites and occurrence rate of extraintestinal pathologic involvements resulting from rotavirus infection, as well as whether rotavirus can be the etiologic agent of these pathologic involvements, still require further research. In line with the aforementioned studies, there exist a variety of animal studies that have detected rotavirus antigen or RNA in extraintestinal tissues; for instance, some studies (Kraft 1958; Uhnoo et al. 1990) reported the detection of rotavirus antigen and infectious virus in the serum, nasal secretions, lungs, hepatic tissue, renal tissue, and brain of mice.

In this research, the dynamics of viremia and the extraintestinal spread of rotavirus, as well as the cell and tissue tropism of rotaviruses, were examined. In addition, the study intended to determine if rotavirus cause extra intestinal infection because the presence of rotavirus in urine indicate that the virus may escape from gastrointestinal tract and may cause viremia. This was performed subsequent to the detection of rotavirus antigenemia in children as well as in the tissues of both rotavirus-infected children and adults.

In order to confirm the presence of rotavirus infection as a urinary tract infection, urine samples were analyzed in patients with rotavirus-induced gastroenteritis. The findings were in full agreement with those reported by (Yokoyama et al. 2011), which showed that the presence of rotavirus in urine

can infect urinary sediment cells in immunocompetent children with rotavirus gastroenteritis.

5. Conclusion

Rotavirus is still a public health concern and continues to expose the lives of nearly one-third of children in Babylon Province at risk. Children younger than two years of age are more likely to be infected with rotavirus, especially males. There was a correlation between the results of RT-qPCR assay and those of rapid test, but RT-qPCR was found to show a higher diagnostic sensitivity for the detection of rotavirus antigen compared to the rapid test. The analysis of rotavirus infections revealed that an equal number of cases occured in the autumn and winter seasons. But the prevalence was higher among younger children in Babylon Province. The detection of rotavirus antigen in urine samples from diarrheal children is reported to be a useful diagnostic method and is worth attention for further laboratory applications.

6. Acknowledgments

The authors extend the sincerest appreciation to Minister Naim Abdel Yasser Al-Aboudi from the Ministry of Higher Education and Scientific Research and to Babylon University, for their valuable contributions that remarkably increased the quality of current research paper. The minister's support and constructive feedback have played a significant role in conducting and improving this research.

7. Conflict of Interest

The authors declare no conflict of interest.

References

Adams W, Kraft L (1963) Epizootic diarrhea of infant mice: identification of the etiologic agent. Science 141:359-360

Ahmad Abdul AL-Kader R, Abed Ali Hassan a, Haifa (2016) Rota virus diarrhea in children under five admitted to Basrah General Hospital. The Medical Journal of Basrah University 34:107-114 doi: 10.33762/mjbu.2016.117213

Ahmed HM et al. (2006) Molecular characterization of rotavirus gastroenteritis strains, Iraqi Kurdistan. Emerging Infectious Diseases 12:824

Anaya-Molina Y et al. (2018) A one-step real-time RT-PCR helps to identify mixed rotavirus infections in Mexico. Diagnostic Microbiology and Infectious Disease 92:288-293 doi: https://doi.org/10.1016/j.diagmicrobio.2018.06.023

Anderson EJ, Weber SG (2004) Rotavirus infection in adults. Lancet Infect Dis 4:91-99 doi: 10.1016/s1473-3099(04)00928-4

Angel J, Franco MA, Greenberg HB (2007) Rotavirus vaccines: recent developments and future considerations. Nature Reviews Microbiology 5:529-539 doi: 10.1038/nrmicro1692

Arnold MM, Barro M, Patton JT (2013) Rotavirus NSP1 mediates degradation of interferon regulatory factors through targeting of the dimerization domain. Journal of virology 87:9813-9821

Artiran S et al. (2017) Investigation of Rotavirus with Various Methods in Children with Acute Gastroenteritis and Determination of Its Molecular Epidemiology in Kayseri Province, Turkey. J Clin Lab Anal 31 doi: 10.1002/jcla.22030

Barro M, Patton JT (2005) Rotavirus nonstructural protein 1 subverts innate immune response by inducing degradation of IFN regulatory factor 3. Proceedings of the National Academy of Sciences 102:4114-4119

Bernstein DI (2009) Rotavirus overview. Pediatr Infect Dis J 28:S50-53 doi: 10.1097/INF.0b013e3181967bee

Bishop R, Davidson G, Holmes I, Ruck B (1973) Virus particles in epithelial cells of duodenal mucosa from children with acute non-bacterial gastroenteritis. The Lancet 302:1281-1283

Bitton G (2002) Encyclopedia of environmental microbiology. (No Title)

Blutt SE et al. (2003) Rotavirus antigenaemia and viraemia: a common event? Lancet 362:1445-1449 doi: 10.1016/s0140-6736(03)14687-9

Bonkoungou IJ et al. (2010) Epidemiology of rotavirus infection among young children with acute diarrhoea in Burkina Faso. BMC pediatrics 10:1-6

Brunet J-P, Cotte-Laffitte J, Linxe C, Quero A-M, Géniteau-Legendre M, Servin A (2000) Rotavirus infection induces an increase in intracellular calcium concentration in human intestinal epithelial cells: role in microvillar actin alteration. Journal of virology 74:2323-2332

Charles MD, Holman RC, Curns AT, Parashar UD, Glass RI, Bresee JS (2006) Hospitalizations associated with rotavirus gastroenteritis in the United States, 1993–2002. The Pediatric infectious disease journal 25:489-493

Chen J, Grow S, Iturriza-Gómara M, Hausdorff WP, Fix A, Kirkwood CD (2022) The Challenges and Opportunities of Next-Generation Rotavirus Vaccines: Summary of an Expert Meeting with Vaccine Developers. Viruses 14 doi: 10.3390/v14112565

Cioc AM, Nuovo GJ (2002) Histologic and in situ viral findings in the myocardium in cases of sudden,

unexpected death. Modern pathology 15:914-922
Crawford SE et al. (2001) Trypsin cleavage stabilizes the rotavirus VP4 spike. J Virol 75:6052-6061 doi: 10.1128/jvi.75.13.6052-6061.2001
Crawford SE et al. (2006) Rotavirus Viremia and Extraintestinal Viral Infection in the Neonatal Rat Model. Journal of Virology 80:4820 - 4832
Crawford SE et al. (2017) Rotavirus infection. Nat Rev Dis Primers 3:17083 doi: 10.1038/nrdp.2017.83
Cunliffe NA, Glass RI, Nakagomi O (2014) 18 - Rotavirus and Other Viral Diarrhoea. In: Farrar J, Hotez PJ, Junghanss T, Kang G, Lalloo D, White NJ (eds) Manson's Tropical Infectious Diseases (Twenty-third Edition). W.B. Saunders, London, pp 207-214.e203
Dian Z et al. (2021) Rotavirus-related systemic diseases: clinical manifestation, evidence and pathogenesis. Crit Rev Microbiol 47:580-595 doi: 10.1080/1040841x.2021.1907738
Estes M, Kapikian A (2007) Fields Virology, eds Knipe DM, Howley PM. In. Lippincott William & Wilkins, Philadelphia
Fenjan M, Jarullah B (2019) Molecular Identification And Phylogenetic Analysis Of Rotavirus In Children Suffered From Diarrhea Under Five Years Old In Thi-Qar Provience, Of Iraq. JOURNAL OF COLLEGE OF EDUCATION FOR PURE SCIENCE 9:82-91 doi: 10.32792/utq.jceps.09.01.09
Flewett T, Bryden A, Davies H (1973) Virus particles in gastroenteritis. Virus particles in gastroenteritis.
Fukuda S, Kishi K, Yasuda K, Sejima H, Yamaguchi S (2009) Rotavirus-associated encephalopathy with a reversible splenial lesion. Pediatric neurology 40:131-133
Goldwater PN, Chrystie IL, Banatvala J (1979) Rotaviruses and the respiratory tract. British medical journal 2:1551
Graff JW, Mitzel DN, Weisend CM, Flenniken ML, Hardy ME (2002) Interferon regulatory factor 3 is a cellular partner of rotavirus NSP1. Journal of virology 76:9545-9550
Grech V, Calvagna V, Falzon A, Mifsud A (2001) Fatal, rotavirus-associated myocarditis and pneumonitis in a 2-year-old boy. Annals of tropical paediatrics 21:147-148
Greenberg HB, Estes MK (2009) Rotaviruses: From Pathogenesis to Vaccination. Gastroenterology 136:1939-1951 doi: https://doi.org/10.1053/j.gastro.2009.02.076
Hasegawa M et al. (2017) Detection of rotavirus in clinical specimens using an immunosensor prototype based on the photon burst counting technique. Biomed Opt Express 8:3383-3394 doi: 10.1364/boe.8.003383
He B et al. (2013) Characterization of a novel G3P [3] rotavirus isolated from a lesser horseshoe bat: a distant relative of feline/canine rotaviruses. Journal of virology 87:12357-12366
Herring A, Inglis N, Ojeh C, Snodgrass Da, Menzies J (1982) Rapid diagnosis of rotavirus infection by direct detection of viral nucleic acid in silver-stained polyacrylamide gels. Journal of clinical Microbiology 16:473-477
Hou G, Zeng Q, Matthijnssens J, Greenberg HB, Ding S (2021) Rotavirus NSP1 Contributes to Intestinal Viral Replication, Pathogenesis, and Transmission. mBio 12:e0320821 doi: 10.1128/mBio.03208-21
Hoxha T et al. (2015) Performance of clinical signs in the diagnosis of dehydration in children with acute gastroenteritis. Med Arch 69:10-12 doi: 10.5455/medarh.2015.69.10-12
Jiang L, Tang A, Song L, Tong Y, Fan H (2023) Advances in the development of antivirals for rotavirus

infection. Front Immunol 14:1041149 doi: 10.3389/fimmu.2023.1041149
Jiang N et al. (2018) Comparison of two different combined test strips with fluorescent microspheres or colored microspheres as tracers for rotavirus and adenovirus detection. Virol J 15:44 doi: 10.1186/s12985-018-0951-5
Kang G et al. (2004a) Quantitation of group A rotavirus by real-time reverse-transcription-polymerase chain reaction: correlation with clinical severity in children in South India. J Med Virol 73:118-122 doi: 10.1002/jmv.20053
Kang G et al. (2004b) Quantitation of group A rotavirus by real-time reverse-transcription-polymerase chain reaction: Correlation with clinical severity in children in South India. Journal of medical virology 73:118-122
Karampatsas K, Osborne L, Seah M-L, Tong CY, Prendergast AJ (2018) Clinical characteristics and complications of rotavirus gastroenteritis in children in east London: A retrospective case-control study. PloS one 13:e0194009
Khoury H, Ogilvie I, El Khoury AC, Duan Y, Goetghebeur MM (2011) Burden of rotavirus gastroenteritis in the Middle Eastern and North African pediatric population. BMC Infect Dis 11:9 doi: 10.1186/1471-2334-11-9
Kindler E, Trojnar E, Heckel G, Otto PH, Johne R (2013) Analysis of rotavirus species diversity and evolution including the newly determined full-length genome sequences of rotavirus F and G. Infection, genetics and evolution 14:58-67
Kraft LM (1958) Observations on the control and natural history of epidemic diarrhea of infant mice (EDIM). Yale J Biol Med 31:121-137
Laizane G, Smane L, Nokalna I, Gardovska D, Feemster KA (2019) Rotavirus-associated seizures and reversible corpus callosum lesion. Acta medica Lituanica 26:113
Li N, Wang ZY (2003) Viremia and extraintestinal infections in infants with rotavirus diarrhea. Di Yi Jun Yi Da Xue Xue Bao 23:643-648
Li Z, Baker ML, Jiang W, Estes MK, Prasad BV (2009) Rotavirus architecture at subnanometer resolution. J Virol 83:1754-1766 doi: 10.1128/jvi.01855-08
Long CP, McDonald SM (2017) Rotavirus genome replication: Some assembly required. PLoS Pathog 13:e1006242 doi: 10.1371/journal.ppat.1006242
Lynch M et al. (2003) The pathology of rotavirus-associated deaths, using new molecular diagnostics. Clinical infectious diseases 37:1327-1333
Malek MA et al. (2006) Diarrhea-and rotavirus-associated hospitalizations among children less than 5 years of age: United States, 1997 and 2000. Pediatrics 117:1887-1892
Malherbe H, * Harwin, R.*, Ulrich M (1963) The cytopathic effects of vervet monkey viruses. South African Medical Journal 37:407-411
Matthijnssens J et al. (2008a) Full genome-based classification of rotaviruses reveals a common origin between human Wa-Like and porcine rotavirus strains and human DS-1-like and bovine rotavirus strains. Journal of virology 82:3204-3219
Matthijnssens J et al. (2008b) Full genome-based classification of rotaviruses reveals a common origin between human Wa-Like and porcine rotavirus strains and human DS-1-like and bovine rotavirus strains. J Virol 82:3204-3219 doi: 10.1128/jvi.02257-07
Maureen L et al. (2001) Rotavirus and Central Nervous System Symptoms: Cause or Contaminant? Case Reports and Review. Clinical Infectious Diseases 33:932-938 doi: 10.1086/322650
Mebus C, Underdahl N, Rhodes M, Twiehaus M (1969) Further studies on neonatal calf diarrhea virus.

In: Proceedings, annual meeting of the United States Animal Health Association, pp 97-99
Mousavi-Nasab SD et al. (2020) A Real-Time RT-PCR Assay for Genotyping of Rotavirus. Iran Biomed J 24:399-404 doi: 10.29252/ibj.24.6.394
Nugent CA, Stewart DA (2023) Rotavirus. Pediatrics In Review 44:598-600 doi: 10.1542/pir.2022-005949
Otto PH et al. (2012) Detection of avian rotaviruses of groups A, D, F and G in diseased chickens and turkeys from Europe and Bangladesh. Veterinary Microbiology 156:8-15
Papa G, Borodavka A, Desselberger U (2021) Viroplasms: Assembly and Functions of Rotavirus Replication Factories. Viruses 13 doi: 10.3390/v13071349
Parashar UD, Bresee JS, Gentsch JR, Glass RI (1998) Rotavirus. Emerg Infect Dis 4:561-570 doi: 10.3201/eid0404.980406
Pickering LK, Bartlett III AV, Reves RR, Morrow A (1988) Asymptomatic excretion of rotavirus before and after rotavirus diarrhea in children in day care centers. The Journal of pediatrics 112:361-365
Ramig RF (2004) Pathogenesis of intestinal and systemic rotavirus infection. J Virol 78:10213-10220 doi: 10.1128/jvi.78.19.10213-10220.2004
Ramig RF (2007) Systemic rotavirus infection. Expert review of anti-infective therapy 5:591-612
Richardson S, Grimwood K, Gorrell R, Palombo E, Barnes G, Bishop R (1998) Extended excretion of rotavirus after severe diarrhoea in young children. The Lancet 351:1844-1848
Roingeard P, Raynal PI, Eymieux S, Blanchard E (2019) Virus detection by transmission electron microscopy: Still useful for diagnosis and a plus for biosafety. Rev Med Virol 29:e2019 doi: 10.1002/rmv.2019
Settembre EC, Chen JZ, Dormitzer PR, Grigorieff N, Harrison SC (2011) Atomic model of an infectious rotavirus particle. Embo j 30:408-416 doi: 10.1038/emboj.2010.322
Takanashi J et al. (2010) Clinical and Radiological Features of Rotavirus Cerebellitis. American Journal of Neuroradiology 31:1591-1595 doi: 10.3174/ajnr.A2131
Trask SD, McDonald SM, Patton JT (2012) Structural insights into the coupling of virion assembly and rotavirus replication. Nature reviews microbiology 10:165-177
Troeger C et al. (2018) Rotavirus Vaccination and the Global Burden of Rotavirus Diarrhea Among Children Younger Than 5 Years. JAMA Pediatr 172:958-965 doi: 10.1001/jamapediatrics.2018.1960
Uhnoo I et al. (1990) Extramucosal spread and development of hepatitis in immunodeficient and normal mice infected with rhesus rotavirus. Journal of Virology 64:361 - 368
Ushijima H, Bosu K, Abe T, Shinozaki T (1986) Suspected rotavirus encephalitis. Archives of disease in childhood 61:692
Vollet III J, DuPont H, Pickering L (1981) Nonenteric sources of rotavirus in acute diarrhea. Journal of Infectious Diseases 144:495-495
Vontas A, Hadjicristodoulou C, Krikelis V, Petinaki E (2019) The Use of Immunochromatographic Technique for Rotavirus Detection: Experience from a Tertiary Care Hospital in Central Greece. Intervirology 62:164-168 doi: 10.1159/000502007
Vos T et al. (2020) Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. The Lancet 396:1204-1222
Ward P, Poitras E, Leblanc D, Gagnon CA, Brassard J, Houde A (2013) Comparison of different RT-

qPCR assays for the detection of human and bovine group A rotaviruses and characterization by sequences analysis of genes encoding VP4 and VP7 capsid proteins. Journal of Applied Microbiology 114:1435-1448 doi: https://doi.org/10.1111/jam.12165

Weitzel T et al. (2007) Field evaluation of a rota- and adenovirus immunochromatographic assay using stool samples from children with acute diarrhea in Ghana. J Clin Microbiol 45:2695-2697 doi: 10.1128/jcm.00562-07

WONG CJ, PRICE Z, BRUCKNER DA (1984) Aseptic meningitis in an infant with rotavirus gastroenteritis. The Pediatric Infectious Disease Journal 3:244-245

Yaqoob M, Mahdi K, Al-Hmudi H, Mohammed Ali M (2016) Detection of Rotavirus A and Escherichia coli from Diarrhea Cases in Children and Coliphage Characterization. International Journal of Current Microbiology and Applied Sciences 5:68-83 doi: 10.20546/ijcmas.2016.504.011

Yokoyama T et al. (2011) Molecular and immunohistochemical detection of rotavirus in urinary sediment cells of children with rotavirus gastroenteritis. Clinical Microbiology and Infection 17:1190-1193 doi: https://doi.org/10.1111/j.1469-0691.2011.03522.x

Zheng BJ et al. (1991) Rotavirus infection of the oropharynx and respiratory tract in young children. Journal of medical virology 34:29-37