Developing Cardiac Digital Twins at Scale: Insights from Personalised Myocardial Conduction Velocity

Shuang Qian*, Devran Ugurlu*, Elliot Fairweather*, Marina Strocchi*, Laura Dal Toso*, Yu Deng*, Reza Razavi*, Alistair Young*, Pablo Lamata*, Martin Bishop*, Steven Niederer§,‡

* Department of Biomedical Engineering, School of Imaging Sciences and Biomedical Engineering,

Kings College London, London, UK

§Alan Turing Institute, British Library, 96 Euston Rd, London, UK

‡Imperial College London, London, UK

Corresponding author: Dr Qian, 4th North Wing, St Thomas' Hospital, London, SE1 7EH

(shuang.qian@kcl.ac.uk)

Abstract

Large-cohort studies using cardiovascular imaging and diagnostic datasets have assessed cardiac anatomy, function, and outcomes, but typically do not reveal underlying biological mechanisms. Cardiac digital twins (CDTs) provide personalized physics- and physiology- constrained *in-silico* representations, enabling inference of multi-scale properties tied to these mechanisms.

We constructed 3464 anatomically-accurate CDTs using cardiac magnetic resonance images from UK biobank and personalised their myocardial conduction velocities (CVs) from electrocardiograms (ECG), through an automated framework.

We found well-known sex-specific differences in QRS duration were fully explained by myocardial anatomy, as CV remained consistent across sexes. Conversely, significant associations of CV with ageing and increased BMI suggest myocardial tissue remodelling. Novel associations were observed with left ventricular ejection fraction and mental-health phenotypes, through a phenome-wide association study, and CV was also linked with adverse clinical outcomes.

Our study highlights the utility of population-based CDTs in assessing intersubject variability and uncovering strong links with mental health.

Introduction

Large cohort multi-modality cardiovascular imaging and diagnostic data sets are increasingly available and are being used to link heart anatomy and function with physiological, lifestyle, and clinical outcomes[1–3]. While providing hypothesis-generating correlations, they have been less successful at identifying the underlying mechanisms that drive these correlations. This is in part because they are restricted to the analysis of observed attributes and do not identify the underlying physiology that may explain the observed correlations.

A strategy to alleviate this limitation is the personalization of Cardiac Digital Twins (CDTs)[4–6]. CDTs provide physics- and physiology- constrained *in-silico* representations of specific individuals. They are personalized by integrating multimodal data, enabling multi-scale structure and function to be inferred from clinical measurements. The model parameters that explain the data become the attributes that describe the underlying physiology. Early forms of CDTs have shown great potential in supporting clinical decision-making and providing tailored therapies as in prospective studies of ventricular tachycardia[7], atrial fibrillation[8] and cardiomyopathy[9].

However, creation of CDTs is associated with challenges. Complex data processing and specialist methodology and requirement of large computational resources are required, which limit their broad adoption in both industrial and clinical settings. Presently, studies are constrained to working with small patient cohorts with at most 100 patients[10], limiting their application and scalability to large population datasets.

The creation of CDTs involves two crucial steps. First is the construction of the anatomical twin, the computational replica of the anatomical structures of each subject's heart from medical images. Previously, semi-automatic workflows of heart mesh generation have been developed[11,12], demanding substantial computational resources and considerable manual interventions by trained experts. The second is to build the functional twin, i.e. identifying bespoke electrophysiological parameters that replicate clinical measurements, for example, electrocardiograms (ECGs). This step is often more challenging, requiring numerous computationally intensive simulations to calibrate the multi-scale parameters, based on the specific biophysical fidelity needed. Parameter personalization remains an ongoing challenge, with the majority of modelling studies relying on 'average' parameters derived from the literature[7,13,14].

The feasibility of generating CDTs at scale hinges on the development of a computationally and time-efficient automated workflow for both CDT creation steps. Recent advances in image segmentation, including nnU-Net[15] and Atlas-based approaches[16] provide robust and precise 3D anatomical structures within abbreviated timeframes, facilitating the rapid creation of anatomical meshes from medical images. The emergence of surrogate models as novel statistics and machine learning tools, for example, Gaussian Process Emulators (GPE)[17], provide a low-cost statistical representation of computationally expensive models. Such surrogate models enable global sensitivity analysis to identify important model parameters and therefore constrain the viable parameter space, which can reduce the number of simulations necessary for model calibration.

In this study, we developed a methodology that integrates multi-modality data from the UK biobank[18] within a CDT framework and first demonstrated the feasibility of CDT creation at scale. Within this framework, we inferred a key myocardial tissue electrical property, myocardial conduction velocity (CV), for each CDT. We then reported how CV varies across sex, body mass index and age, together with imaging and ECG-derived phenotypes (biventricular myocardial mass and QRS duration (QRSd) respectively). We also conducted a phenome-wide association study to explore

76 their relationships with other phenotypes reported in the UK Biobank, followed by assessing their
77 ability to independently predict clinical outcomes.

78 Results

79 Anatomical and functional CDT generation workflow

The CDTs were built from UK Biobank magnetic resonance imaging (MRI) and ECG data sets as
depicted in Fig. 1. Each CDT replicates the ECG QRSd and consists of a model of the heart anatomy,
the preferred myocyte orientation or fibre structure, a fast endocardial conducting layer, the
location of the activating Purkinje fascicles, the activation timing, the tissue conductivity, the degree
of anisotropy, and the location of the virtual ECG electrodes. The personalization involved the
generation of the bespoke bi-ventricular anatomy from MRI and the inference of the CV that
replicates the ECG QRSd. The anatomical personalization of a CDT, the first step in the pipeline, took
5 minutes to be built from MRI images (avg mesh resolution 0.9mm) on a desktop with 16 cores.

Fig. 1: The automated anatomical and functional (Electrophysiological) CDT generation workflow. The anatomical models
are personalised finite element meshes with physiological-detailed myocardial fibres constructed from the short-axis and
long-axis heart images in the UK biobank following the steps of segmentations, surface meshes and volumetric meshes
construction as well as myocardial fibre generation. The functional CDT workflow is to replicate the electrophysiological
activities within the anatomical models to match the QRS duration from the clinically measured 12-lead ECGs.

The functional personalization required a preliminary analysis to determine which parameters could
be inferred from the available data and which needed to be set to reference prior values. Thus, we
performed two global sensitivity analyses. In 10 representative cases sampled across sex, BMI, and
age, we calculated the sensitivity of the QRSd prediction to 20 tissue-level physiological parameters
(Fig. 2a and Supplementary Table 2) or 30 ECG electrode positions combined with the most
important tissue-level property (Fig. 2b; Supplementary table 3). Notably, the first analysis revealed
that CV has the dominant effect on QRSd, accounting for 72.7 ± 3.2 % the sensitivity, across all
electrophysiological parameters, see Fig. 2. The second analysis then studied the potential impact of
ECG electrode positions, finding that CV affects the QRSd the most (67.2 ± 10.4 %).

Given CV was identified as the only key parameter that explains most of the variation in QRSd, we
chose to calibrate the models by applying a computational-efficient bisection method to search for
the personalized CV (constrained by physiological measurements in literature [19,20]) to match with
measured QRSd from the 12-lead ECGs at rest during sinus rhythm. The CDT calibration process

assumed all other parameters were set to reference prior values taken from the literature,
introducing an estimated uncertainty of -13% to +15.6% around the 'true' CV (Supplementary Figure
2). The personalised CV parameter was inferred in 8 minutes on a desktop with 16 cores.

Fig. 2: Global sensitivity analysis. The total effects of input EP parameters explain the variance of output QRS duration in 10
subjects sampled from the cohort based on sex, age and BMI. a is for only tissue-level EP parameters and b is for the 30 ECG
electrodes' location parameters combining with the most important tissue-level parameter from a.

We used the 4,329 first participants from the UK biobank that had adequate geometrical
information[16] and reported QRSd, sex, age, BMI/weight and height information. Of these, 3945
(91.1%) and 3464 (80%) were successfully processed through the anatomical and functional
personalization workflows. Summary participant characteristics are shown in Supplementary Table
1.

Model validation

To validate the CDT workflow, we compared the QRS morphology in simulated ECGs against the
measured ECGs in the 10 representative subjects. As the ECGs were simulated from a reference
torso and heart location, we do not expect perfect matches in all leads. We have adopted a lead-to-
lead comparison approach used to compare ECGs clinically[21]. To quantify the ability of the model to
replicate the ECG morphology, we plotted the simulated 12 lead ECGs (filtered, scaled, and
temporally aligned) against measured ECG and correlation coefficients (r) were calculated.
Supplementary Figure 3 shows the comparison, with plots ordered in descending order of the
correlation coefficients. 56% of ECG leads in all subjects matched with the recordings well ($r > 0.5$).
Supplementary Figure 4 shows the averaged r of the best correlated ECG leads as the number of
leads considered increases. The maximum average r considering only the best correlated ECG leads
was 0.95, while its value dropped to 0.79 when considering the top 6 ECG leads.

To further validate our modelling approach, we investigated whether subjects with pathologically
slow conduction, for instance, caused by fascicular block or heart failure, can be differentiated
through their personalized CVs. Those subjects were identified using the summary diagnoses for
hospital inpatients in the UK Biobank with the specific disease types as shown in Supplementary
Table 5. Fig. 3 shows that subjects with fascicular block (N=46) had 16.8% lower CV (0.482± 0.11 vs
0.579±0.08 m/s, $P = 1.1 \times 10^{-14}$), and 23.3% longer QRSd (108.4±26.0 vs 87.9±12.6ms, $P =$
2.2×10^{-26}), comparing to their counterparts. Similarly, we also observed 5.2% slower CV (0.548±
0.10 vs 0.578±0.08m/s, $P = 6.6 \times 10^{-3}$) and 10.7% longer QRSd (97.6±20.8 vs 88.1±12.8ms, $P =$
2.1×10^{-8}) in subjects with heart failure (N=60), comparing to their counterparts.

Fig. 3: Boxplots of QRS duration (red) and conduction velocity (yellow) for comparing groups of participants afflicted with fascicular block and non-afflicted counterparts. The green dots indicate the mean. The corresponding P values are from the student's t-test. *P<0.05; **P<0.01; ***P<0.001; ****P<0.0001.

Comparison for different sex, BMI and age groups

We compared QRSd, CV and biventricular myocardial mass, categorized into different groups of sex, BMI and age as shown in Fig. 4. The QRSd was 9.4% longer in males (92.9±13.2 vs 84.2±11.5 ms, $P = 2.5 \times 10^{-89}$). This is consistent with males tending to have larger hearts (male: 180.9±28.5g vs 131.1±18.9g, $P = 0$). However, CVs were the same in men and women (0.576±0.09 vs 0.579±0.08m/s, P=0.25).

QRSd was longer for overweight and obese ($25 \leq BMI \leq 30$ and $BMI \geq 30$) compared to healthy ($BMI \leq 25$) groups (overweight: 88.6±12.9 and obese: 89.1±13.0, vs healthy: 87.5±13.2ms, P=0.03 and 0.01). Again, this increase in QRSd with BMI was reflected with a corresponding increase in heart size (healthy: 141.4±29.2, overweight: 158.7±33.5, obese: 170.5±36.9g, all $P < 1 \times 10^{-13}$). In contrast with sex, this increase in QRSd was also associated with a progressive increase in CVs (healthy: 0.569±0.08m/s vs overweight and obese: 0.582±0.08 and 0.587±0.09m/s, $P = 1.8 \times 10^{-5}$ and 7×10^{-6}), which suggests the compensatory mechanism for the increment of heart size (i.e. CV is increased as a mechanism to reduce QRSd when the heart needs to grow to meet the larger demand of an increased BMI).

QRSd increased with ageing, which was significant when comparing the $Age \leq 60$ group (87.3±11.6ms) with $Age \geq 65$ (87.9±12.0ms, P=0.03) and $60 \leq Age \leq 65$ (89.3±14.8ms, $P = 1.1 \times 10^{-4}$). This QRS increase is consistent with an observed CV decrease ($Age \leq 60$: 0.586±0.08 vs $60 \leq Age \leq 65$: 0.578±0.08 vs $Age \geq 65$: 0.569±0.09m/s, all P<0.02), and myocardial mass increase (141.4±29.2 vs 158.7±33.5 vs 170.5±36.9g, all $P < 1 \times 10^{-13}$). By inferring CV, it is possible to determine when changes in QRSd with age, sex or BMI are caused by changes in cardiac anatomy or biological material properties.

Fig. 4: Boxplots of QRS duration (red), conduction velocity (yellow) and myocardial mass(purple) for different groups of sex,
BMI and age. The green dots indicate the mean. The corresponding P values are from the student's t-test. *P<0.05;
**P<0.01; ***P<0.001; ****P<0.0001.

Phenome-wide association study

We performed a phenome-wide association study (PheWAS) to explore the correlations between
the multimodal and CDT-derived phenotypes and 473 UKBB reported phenotypes in categories:
pulse wave analysis, LV size & function (automatically derived from heart MRIs), abdominal
composition, medication (medical treatment received), primary demographics, early-life
information, self-reported medical conditions, lifestyle diet, alcohol, smoking, physical activity,
physical measures, education and employment, mental health, and clinical outcomes of seven
common diseases categorized from summary diagnoses in UKBB (Supplementary Table 5).

Fig. 5: Manhattan plot showing the $-log_{10}P$ (two-sided t-test) for correlations between the QRS duration, CV and
myocardial mass, and UKBB reported phenotypes. The size of the dots indicates the absolute Pearson's correlation
coefficient. The dashed horizontal lines are the Bonferroni threshold (Bonf) and the false-discovery rate (fdr) (α = 0.05). Note
the plot is clipped at 17 for better visualization for QRS duration and CV. The complete plot is in the Supplementary Figure 5.

Fig. 5 shows the Manhattan plot of the univariate correlation P values (two-sided) between M = 3
multimodal phenotypes (myocardial mass, QRSd and inferred CV) and N = 473 UKBB phenotypes for
M × N = 1419 times, with 87 correlations reaching the Bonferroni threshold for multiple comparisons
($P_{bonf} = 3.5 \times 10^{-5}$ for α = 0.05) and 146 correlations reaching the false-discovery rate (FDR)
threshold ($P_{fdr} = 0.005$ for α = 0.05). For the correlation coefficients in the Manhattan plot see
Supplementary Figure 6.

The QRSd was significantly associated with seven cardiac structural and functional phenotypes
including LV end-diastolic, end-systolic and stoke volumes, cardiac output, pulse/heart rate
($-log_{10}P \geq 2.4$, $|r| \geq 0.05$) as well as LV ejection fraction ($-log_{10}P = 3.2$, r=-0.06). In contrast, CV
was only significantly associated with LV ejection fraction ($-log_{10}P = 3.5$, r=0.06). The biventricular
myocardial mass was significantly associated with all phenotypes mentioned above ($-log_{10}P > 9.6$,
$|r| \geq 0.16$) where higher correlations were seen for structural phenotypes such as LV stroke, end
diastolic and end-systolic volumes ($-log_{10}P > 68.8$, $|r| > 0.3$).

Interestingly, the three phenotypes were all significantly associated with mental health-related
phenotypes. Both QRSd and CV were significantly associated with 'the longest period of depression'
(both: $-log_{10}P = 7.5$; QRSd: r=0.15, CV: r=-0.15) and 'Seen doctor (GP) for nerves, anxiety, tension
or depression' (QRSd:$-log_{10}P = 2.5$, r=0.05; CV: $-log_{10}P = 3.3$, r=-0.06). CV and myocardial mass
were significantly associated with 'Seen a psychiatrist for nerves, anxiety, tension or depression' (CV:
$-log_{10}P = 3.5$, r=-0.06; myocardial mass: $-log_{10}P = 2.6$, r=-0.05). Consistent with previous
studies[1], we find that myocardial mass was solely associated with the Neuroticism score, Frequency
of depressed mood in the last 2 weeks, and Happiness ($2.6 < -log_{10}P < 3.9$, r: [-0.1, -0.05]).

Association with clinical diagnoses

We investigated the associations of the multimodal and CDT-derived phenotypes with seven
common diseases, categorized using the summary diagnoses for hospital inpatients. We trained a
logistic regression model on QRSd, myocardial mass and CDT-derived CV to predict disease
occurrence, adjusted with demographics/anthropometrics factors including sex, age BMI, age*BMI
and sex*age. Fig. 6 and Supplementary Table 6 presents the odd ratios (ORs) derived from the
regression coefficients (represented as $value_{[95\% \; confidence \; interval]}$) and the corresponding P
values.

Smaller CV and greater myocardial mass were associated with a high risk of Fascicular block
(ORs=$0.43_{[0.37,0.51]}$ and $1.72_{[1.54,1.92]}$, both $P < 1 \times 10^{-16}$). In contrast, both greater QRSd and
myocardial mass were associated with a higher risk of heart failure (ORs=$1.38_{[1.20,1.58]}$ and $1.66_{[1.50,1.83]}$,
$P = 3 \times 10^{-6}$ and $P < 1 \times 10^{-16}$). Ischaemic diseases were more likely to develop with greater
QRSd (ORs=$1.34_{[1.19,1.52]}$, $P = 4 \times 10^{-6}$), while diabetes was more likely to develop with smaller CV
(ORs=$0.75_{[0.66,0.86]}$, $P = 3.6 \times 10^{-5}$). Reduced CV was a better predictor than QRSd for both neurotic
and mood disorders, (CV: ORs=$0.65_{[0.57,0.73]}$ and $0.62_{[0.54,0.72]}$, $P < 1.07 \times 10^{-11}$ vs QRSd: ORs=
$0.77_{[0.68,0.88]}$ and $0.76_{[0.66,0.87]}$ $P < 1.27 \times 10^{-4}$). The extended analysis (Supplementary Figure 7)
shows CV was significantly lower in participants having mood disorders than their counterparts
(0.564±0.085 vs 0.578± 0.085m/s, P=0.03), while QRSd and myocardial mass did not (QRSd: 88.8±
14.2 vs 88.2±13.0ms, P=0.57 and vol: 142.3± 30.5 vs 147± 32.9g, P=0.07). Hypertension was solely
associated with greater myocardial mass (OR=$1.33_{[1.18,1.49]}$, $P = 1.26 \times 10^{-6}$).

Fig. 6: Association of CV, QRS duration and myocardial mass with common diseases. (1) Odd ratios for phenotypes as input risk factors for common diseases as the outcomes. Sex, age, BMI, age*BMI and sex*age were included in the logistic regression analysis (N=3464). (2) The corresponding P values (two-sided t-test) for odd ratios. *Result reaches Bonferroni threshold ($P_{Bonf} = 6.9 \times 10^{-4}$ for α=0.05).

Discussion

This study makes three major contributions. First, we have developed a fully automated CDT generation workflow including both patient-specific anatomical and electrophysiological models, using multimodal data including both medical imaging and clinically measured ECGs. This workflow is the first of its kind to construct biophysically detailed CDTs at a large scale, benefiting from using statistical and AI tools that allow global sensitivity analysis on all known model parameters.

Second, we have showcased the capacity of utilizing the quantitative CDT-derived phenotype to unveil the underlying biological mechanisms and elucidate the variability in observable attributes such as imaging and ECG phenotypes among different populations according to sex, age and BMI. Specifically, we show that well-known sex-specific differences in QRSd can be fully explained by myocardial anatomy and that CV is the same in men and women, and that myocardial remodeling leads to changes in CV associated with ageing and increased BMI.

Third, we performed a large-scale PheWAS study and found more significant associations of the CDT-derived phenotype with cardiac and mental-health phenotypes relating to neurological disorders, including depression, compared against known ECG and imaging phenotypes. Furthermore, we found a stronger association between the decreased CV and a higher risk of neurotic and mood disorders, compared to the myocardial mass and QRSd.

These results demonstrate the usefulness of generating CDT-derived phenotypes, even from the UK biobank where the majority of participants identified as healthy. The ability to identify the electrophysiological biomarkers: myocardial CV at scale could help identify potential therapeutic targets and evaluate the therapeutic potential (or side effects) of existing drugs and heart disease medications for mental health and neurodegenerative disorders.

CDTs are rapidly advancing in recent decades, leveraging their capacity to guide, inform, monitor, diagnose and prognose therapies and surgical interventions in many current prospective clinical studies[22], paving the way for moving into industrial and clinical settings[4]. This shift requires a step change in the speed, robustness, validation, and uncertainty quantification in both anatomical and functional model creation workflow.

To date, a wide range of automated anatomical model creation workflows based on machine
learning have been proposed allowing large-scale population analysis such as on the UK Biobank
dataset[1,23,24], however, these models only account for surfaces depicting the overall heart structure,
but not high-quality volumetric meshes. There are also volumetric mesh creation workflows existing
for atria[25,26], ventricles[12] and whole hearts[11,27], but these have only been applied to smaller datasets
(<100) due to the high computational costs and manual steps required. Here we reported a fully
automated volumetric mesh generation workflow that can create personalized anatomical meshes
at scale within clinical timescales (~5 minutes/CDT).

Aside from anatomical model personalization, functional model personalization is more challenging,
where most current modelling studies were developed using 'average' material property values from
broader physiological studies with limited personalization[28,29]. Recent developments in high fidelity
biological and physiological electrophysiological CDT frameworks incorporated detailed features like
the His-Purkinje fascicles to replicate detailed QRS complex morphology [12,30]. However, only a subset
of parameters underwent personalization, yet still demanding significant computation resources and
time. Alternative computationally efficient approaches based on machine learning and statistical
methods were reported but often fall short in capturing all anatomical/functional details and
struggle to generalize effectively[31,32].

To move to population-level studies, it is of critical importance to balance biophysical fidelity,
parameter inference and computational cost. In this work, instead of making CDTs to replicate
recorded ECG morphologies that are more likely to be afflicted by subject-specific noises, we
constructed feature-specific CDTs to replicate the QRSd which can be extracted from models
robustly and computationally-efficiently[33]. Similar to previous biophysical-detailed frameworks[12], our
CDT EP framework incorporates parameters encapsulating knowledge derived from physiological
and histological/anatomical experiments in the past. Our sensitivity analysis suggests that an optimal
trade-off between model fidelity and parameter identifiability is to restrict the EP personalization to
one single parameter, CV, showing a dominating effect on QRSd (>67.2%), ensuring its
computational efficiency (~8 mins/CDT).

To truly realize the predictive capability of developed CDTs, it is important to determine the bounds
of fidelity and validity of our CDTs (i.e. which aspects of the real-world system are sufficiently
recapitulated?). We have conducted two model validations. First, we compared how well our
simulated 12-lead ECGs reproduce the recordings lead-wisely by conducting correlation tests on the
10 representative subjects. We found good correlations in 56% of lead ECGs ($r > 0.5$) from all
subjects, as we expected. We have standardized the ECG electrode locations due to data
unavailability, where such information has been previously shown to be the main driver of QRS
complex morphological difference, but not for QRSd[29,34]. Therefore, we can conclude our CDT
framework is still reliable with good accuracy for QRSd. Second, our CDT approach was tested to
identify the CVs of subjects afflicted with fascicular block or heart failure. These are two conditions
with known slow cardiac conduction substrates, and we found that their CV was indeed significantly
slower than the counterparts by 16.8% and 5.2% (Fig. 3), given we used a unified healthy conduction
system in our personalized QRSd calibration process. An additional positive validation result is the
fact that the separate association test on clinical outcomes identified that lower CV is associated
with a much higher risk of fascicular block (Fig. 6). Further, in a verification test, we quantified the
uncertainty of the inferred CVs to be <15.6% variation around the 'true' value, considering the
uncertainty in assuming all other parameters (electrophysiology and ECG electrodes positioning) as
reference prior values, which further increases the credibility of our approach.

By building CDTs, we have been able to separate the impact of conduction velocity and heart
anatomy on QRSd and quantitatively assess their relative changes across populations of different
sex, BMI and age, which unveils mechanistic insights about the underlying biological process.
Consistent with previous studies[35–38], we found that men have longer QRSd and larger myocardial
mass in comparison to women. This change in QRSd can be entirely explained by the change in
anatomy, with no discernible disparity in CVs between the sexes, suggesting no significant biological
differences in myocardium that modulate the ventricular depolarization wave propagation.
Accordingly, clinical guidelines that use QRSd as criteria for elective therapy, such as cardiac
resynchronization therapy, can develop sex-specific thresholds of decision based on heart size
differences and disregard the potential impact of CV variations. This result brings additional
evidence to support the proposal of reducing the threshold for CRT in female subjects by 9-13 ms[39].

Similar to sex, longer QRSds are observed in overweight and obese groups compared to healthy
groups, consistent with literature[37], which are primarily attributed to the increased myocardial mass.
Increased myocardial mass implies an increased length of the pathway for the ventricular activation
wave to travel through, resulting in longer QRSd, which is consistent with previous studies showing
obesity, as a common risk factor of cardiovascular diseases, can lead to structural remodelling, for
example, left ventricular hypertrophy[40]. Obesity may also lead to electrical remodeling including
conduction slowing and conduction heterogeneity, which is often observed in diseased clinical
cohorts[41,42]. We have extended our knowledge of obesity-related electrical remodelling based on a
relatively healthy cohort from the UK biobank. In contrast to earlier findings from clinical cohorts, we
have observed a concurrent small but significant elevation in CVs corresponding to the increase in
BMI. This suggests a potential adaptive response by the heart, possibly compensating for the effects
of heightened myocardial mass on QRSd within obese groups. This novel insight, initially identified in
our study, warrants further investigations to delve deeper into this adaptive mechanism and its
implications.

We also observed longer QRSd in the elderly groups, which is attributed jointly to a decreased CV
and an increased myocardial mass. This finding reaffirms previous research highlighting age-related
changes including both structural changes such as increased LV wall mass[43], possibly driven by
cardiomyocyte hypertrophy and functional changes such as the decline in diastolic function[24,44]. This
decline in the CV may be driven by well-established age-related cellular and tissue level remodelling
including impaired sodium channel function[44] (potentially though loss-of-function genetic mutations
such as SCN5A as identified in previous ECG age-delta GWAS[45] and experimental studies[46]) and the
development of myocardial fibrosis[24,47]. It may also be affected by gap junction decoupling such as
connexin 43 downregulation which is a known pathological remodelling in patients with ventricular
hypertrophy and ischaemic heart diseases[48].

PheWAS is a data-driven method to generate new hypotheses based on the identification of
correlations between exposure like genetic and environmental factors and phenotypes like diseases
and clinical outcomes. Unlike previous PheWAS studies focusing on either MRI-derived or genetic
phenotypes[1,24,44], our PheWAS study is the first to include an inferred tissue-level phenotype: CV,
uniquely estimated with our CDT approach, which can form a bridge between research findings on
the genetic, molecular/cellular level to tissue and the whole organ level.

Despite a relatively smaller sample size (N=3464), we found that biventricular myocardial mass was
highly associated with multiple structural phenotypes such as LV stroke, end diastolic and end
systolic volumes, as found in previously [1]. Interestingly, we found that compared to the QRSd
association with seven MRI-derived phenotypes, CV was only positively associated with LV ejection
fraction, clinically used as a metric of functional performance particularly in the diagnosis of heart

failure and post-myocardial infarction. Thus, CV may be used as a new biomarker for disease
evaluation and also as a potential therapeutic target for guiding the development of new drugs.

Furthermore, we also identified notable correlations with multiple mental-health factors. The
strongest correlations were observed between CV and QRSd with 'the longest period of depression'
(QRSd: r=0.15 and CV:r=-0.15). Interestingly, this was not significantly associated with myocardial
mass, although myocardial mass shows relatively weaker correlations with several other mental
health factors ($|r| < 0.1$). Furthermore, in the separate association test on clinical outcomes, we
found that reduced CV is more closely associated with an increased risk of both neurotic and mood
disorders, in contrast to QRSd, while no significant results were observed for myocardial mass.
Previous research has established a bidirectional relationship between depression and
cardiovascular diseases, while a recent study specifically shows the association between low
depression frequency with decreased risk of cardiometabolic disease using the UK Biobank data[49].
Increased evidence has linked psychological disorders with altered cardiac morphology and
functions, for example, reduced LV mass, increased myocardial fibrosis and [50,51], which were also
observed as changes in ECG metrics such as heart rate, QT interval, QRSd but not in all[52–54]. Our
results reaffirmed these previous findings on myocardial mass and extend existing knowledge by
identifying novel changes in CV and QRSd associated with neurotic and mood disorders (i.e. anxiety
and depression etc).

Overall, the CDT-derived CVs exhibit greater sensitivity to functional alternations, compared to
anatomical factors such as myocardial mass and ECG-derived metrics such as QRSd. This heightened
sensitivity could be attributed to the fact that CV is directly modulated by the physical properties of
cardiac myocytes and their interconnections. Therefore, the changes in CV may more accurately
reflect cardiac remodelling such as fibrotic alterations resulting from the decoupling of cell-cell
connections and coupling of myocytes with fibroblasts[55].

In conclusion, we provide a proof-of-concept cross-sectional study that demonstrates the potential
for applying a CDT workflow to larger cohorts. This approach may yield deeper insights into the
biological connections between changes in cardiac morphology and function with neurological
disorders. Further validation can be pursued through longitudinal studies utilizing resources such as
the UK Biobank's repeat imaging, and causal relationships can be established through large-scale
genetic studies, such as the emerging area of heart-brain connection[56].

Reference

1.   Bai, W. *et al.* A population-based phenome-wide association study of cardiac and aortic
structure and function. *Nat Med* **26**, 1654–1662 (2020).

2.   Yan, C. L. & Grazette, L. A review of biomarker and imaging monitoring to predict heart failure
recovery. *Front Cardiovasc Med* **10**, (2023).

3.   Raisi-Estabragh, Z., Harvey, N. C., Neubauer, S. & Petersen, S. E. Cardiovascular magnetic
resonance imaging in the UK Biobank: A major international health research resource. *Eur*
Heart J Cardiovasc Imaging **22**, 251–258 (2021).

4.   Niederer, S. A., Sacks, M. S., Girolami, M. & Willcox, K. Scaling digital twins from the artisanal
to the industrial. *Nat Comput Sci* **1**, 313–320 (2021).

5.   Corral-Acero, J. *et al.* The 'Digital Twin' to enable the vision of precision cardiology. *Eur Heart*
J **41**, 4556-4564B (2020).

6.   Acosta, J. N., Falcone, G. J., Rajpurkar, P. & Topol, E. J. Multimodal biomedical AI. *Nat Med* **28**,
1773–1784 (2022).

7.   Prakosa, A. *et al.* Personalized virtual-heart technology for guiding the ablation of infarct-
related ventricular tachycardia. *Nat Biomed Eng* **2**, 732–740 (2018).

8.   Patrick M. Boyle, Tarek Zghaib, Sohail Zahid, Rheeda L. Ali, Dongdong Deng, William H.
Franceschi, Joe B. Hakim, Michael J. Murphy, Adityo Prakosa, Stefan L. Zimmerman, Hiroshi
Ashikaga, Joseph E. Marine, Aravindan Kolandaivelu, Saman, N. A. T. Computationally guided
personalized targeted ablation of persistent atrial fibrillation. *Nat Biomed Eng* **3**, 870–879
(2019).

9.   Qian, S. *et al.* Additional coils mitigate elevated defibrillation threshold in right-sided
implantable cardioverter defibrillator generator placement: a simulation study. *Europace* **25**,
1–11 (2023).

10.   Niederer, S. A. *et al.* Creation and application of virtual patient cohorts of heart models:
Virtual Cohorts of Heart Models. *Philosophical Transactions of the Royal Society A:*
Mathematical, Physical and Engineering Sciences **378**, (2020).

11.   Strocchi, M. *et al.* A publicly available virtual cohort of fourchamber heart meshes for cardiac
electromechanics simulations. *PLoS One* **15**, 1–26 (2020).

12.   Gillette, K. *et al.* A Framework for the generation of digital twins of cardiac electrophysiology
from clinical 12-leads ECGs. *Med Image Anal* **71**, 102080 (2021).

13.   Arevalo, H. J. *et al.* Arrhythmia risk stratification of patients after myocardial infarction using
personalized heart models. *Nat Commun* **7**, (2016).

14.   Qian, S. *et al.* Optimization of anti-tachycardia pacing efficacy through scar-specific delivery
and minimization of re-initiation: a virtual study on a cohort of infarcted porcine hearts. *EP*
Europace **44**, 1–10 (2022).

15.   Isensee, F., Jaeger, P. F., Kohl, S. A. A., Petersen, J. & Maier-Hein, K. H. nnU-Net: a self-
configuring method for deep learning-based biomedical image segmentation. *Nat Methods*
18, 203–211 (2021).

16.   Mauger, C. *et al.* Right ventricular shape and function: Cardiovascular magnetic resonance
reference morphology and biventricular risk factor morphometrics in UK Biobank. *Journal of*
Cardiovascular Magnetic Resonance **21**, (2019).

17.   Longobardi, S. *et al.* Predicting left ventricular contractile function via Gaussian process
emulation in aortic-banded rats: Aortic banded rat heart simulations. *Philosophical*
Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences **378**,
(2020).

18.   Manolio, T. A. UK Biobank debuts as a powerful resource for genomic research. *Nat Med* **24**,
1792–1794 (2018).

19. Spear, J. F., Michelson, E. L. & Moore, E. N. Cellular electrophysiologic characteristics of
chronically infarcted myocardium in dogs susceptible to sustained ventricular
tachyarrhythmias. *J Am Coll Cardiol* **1**, 1099–1110 (1983).

20. Glukhov, A. v. *et al.* Conduction remodeling in human end-stage nonischemic left ventricular
cardiomyopathy. *Circulation* **125**, 1835–1847 (2012).

21. De Chillou, C. *et al.* Localizing the critical isthmus of postinfarct ventricular tachycardia: The
value of pace-mapping during sinus rhythm. *Heart Rhythm* **11**, 175–181 (2014).

22. Niederer, S. A., Lumens, J. & Trayanova, N. A. Computational models in cardiology. *Nat Rev
Cardiol* **16**, 100–111 (2019).

23. Xia, Y. *et al.* Automatic 3D+t four-chamber CMR quantification of the UK biobank: integrating
imaging and non-imaging data priors at scale. *Med Image Anal* **80**, 102498 (2022).

24. Thanaj, M. *et al.* Genetic and environmental determinants of diastolic heart function. *Nature
Cardiovascular Research* **1**, 361–371 (2022).

25. Nagel, C., Schuler, S., Dössel, O. & Loewe, A. A bi-atrial statistical shape model for large-scale
in silico studies of human atria: Model development and application to ECG simulations. *Med
Image Anal* **74**, 102210 (2021).

26. Solís-Lemus, J. A. *et al.* Evaluation of an open-source pipeline to create patient-specific left
atrial models: A reproducibility study. *Comput Biol Med* **162**, (2023).

27. Gillette, K. *et al.* A personalized real-time virtual model of whole heart electrophysiology.
Front Physiol **13**, 1–14 (2022).

28. Lee, A. W. C. *et al.* A rule-based method for predicting the electrical activation of the heart
with cardiac resynchronization therapy from non-invasive clinical data. *Med Image Anal* **57**,
197–213 (2019).

29. Mincholé, A., Zacur, E., Ariga, R., Grau, V. & Rodriguez, B. MRI-Based Computational
Torso/Biventricular Multiscale Models to Investigate the Impact of Anatomical Variability on
the ECG QRS Complex. *Front Physiol* **10**, (2019).

30. Camps, J. *et al.* Inference of ventricular activation properties from non-invasive
electrocardiography. *Med Image Anal* **73**, 102143 (2021).

31. Zettinig, O. *et al.* Data-driven estimation of cardiac electrical diffusivity from 12-lead ECG
signals. *Med Image Anal* **18**, 1361–1376 (2014).

32. Pezzuto, S. *et al.* Reconstruction of three-dimensional biventricular activation based on the
12-lead electrocardiogram via patient-specific modelling. *Europace* **23**, 640–647 (2021).

33. Gemmell, P. M. *et al.* A computational investigation into rate-dependant vectorcardiogram
changes due to specific fibrosis patterns in non-ischæmic dilated cardiomyopathy. *Comput
Biol Med* **123**, 103895 (2020).

34. Gillette, K., Gsell, M. A. F., Prassl, A. J. & Plank, G. Influence of Electrode Placement on the
Morphology of in Silico 12 Lead Electrocardiograms. *Comput Cardiol (2010)* **2021-Septe**, 1–4
(2021).

35. Randolph, T. C. *et al.* Race and Sex Differences in QRS Interval and Associated Outcome
Among Patients With Left Ventricular Systolic Dysfunction. *J Am Heart Assoc* **6**, 1–11 (2017).

36. Hnatkova, K., Smetana, P., Toman, O., Schmidt, G. & Malik, M. Sex and race differences in
QRS duration. *Europace* **18**, 1842–1849 (2016).

37. Rao, A. C. A. *et al.* Electrocardiographic QRS duration is influenced by body mass index and
sex. *IJC Heart and Vasculature* **37**, 100884 (2021).

38. Petersen, S. E. *et al.* Reference ranges for cardiac structure and function using cardiovascular
magnetic resonance (CMR) in Caucasians from the UK Biobank population cohort. *Journal of*
Cardiovascular Magnetic Resonance **19**, 1–19 (2017).

39. Lee, A. W. C. *et al.* Sex-Dependent QRS Guidelines for Cardiac Resynchronization Therapy
Using Computer Model Predictions. *Biophys J* **117**, 2375–2381 (2019).

40. Yao, Y., Xue, J. & Li, B. Obesity and sudden cardiac death: Prevalence, pathogenesis,
prevention and intervention. *Front Cell Dev Biol* **10**, 1–9 (2022).

41. Narayanan, K. *et al.* QRS fragmentation and sudden cardiac death in the obese and
overweight. *J Am Heart Assoc* **4**, 1–8 (2015).

42. Schram-Serban, C. *et al.* Heterogeneity in Conduction Underlies Obesity-Related Atrial
Fibrillation Vulnerability. *Circ Arrhythm Electrophysiol* **13**, (2020).

43. Forman, D. E. *et al.* Cardiovascular Biomarkers and Imaging in Older Adults: JACC Council
Perspectives. *J Am Coll Cardiol* **76**, 1577–1594 (2020).

44. Shah, M. *et al.* Environmental and genetic predictors of human cardiovascular ageing. *Nat*
Commun **14**, (2023).

45. Verkerk, A. O., Amin, A. S. & Remme, C. A. Disease Modifiers of Inherited SCN5A
Channelopathy. *Front Cardiovasc Med* **5**, (2018).

46. Maurya, S. *et al. Outlining cardiac ion channel protein interactors and their signature in the*
human electrocardiogram. *Nature Cardiovascular Research* vol. 2 (Springer US, 2023).

47. Hung, C. L. *et al.* Age- and Sex-Related Influences on Left Ventricular Mechanics in Elderly
Individuals Free of Prevalent Heart Failure: The ARIC Study (Atherosclerosis Risk in
Communities). *Circ Cardiovasc Imaging* **10**, 1–9 (2017).

48. Severs, N. J. *et al.* Gap junction alterations in human cardiac disease. *Cardiovasc Res* **62**, 368–
377 (2004).

49. Honigberg, M. C. *et al.* Low depression frequency is associated with decreased risk of
cardiometabolic disease. *Nature Cardiovascular Research* **1**, 125–131 (2022).

50. Mahmood, A. *et al.* Neuroticism personality traits are linked to adverse cardiovascular
phenotypes in the UK Biobank. *Eur Heart J Cardiovasc Imaging* **00**, 1–8 (2023).

51. Levine, G. N. *et al.* Psychological Health, Well-Being, and the Mind-Heart-Body Connection A
Scientific Statement From the American Heart Association. *Circulation* **143**, E763–E783
(2021).

52. Hu, M. X., Lamers, F., Penninx, B. W. J. H. & de Geus, E. J. C. Association between depression,
anxiety, and antidepressant use with T-Wave amplitude and QT-interval. *Front Neurosci* **12**,
(2018).

53. Tang, M., Xi, J. & Fan, X. QT interval is correlated with and can predict the comorbidity of
depression and anxiety: A cross-sectional study on outpatients with first-episode depression.
Front Cardiovasc Med **9**, (2022).

54. Yılmaz, M. & Yılmaz, S. Electrocardiographic Frontal Qrs-T Angle is Independently Associated
With Panic Disorder. *Int J Psychiatry Med* **0**, 1–15 (2023).

55. King, J. H., Huang, C. L. H. & Fraser, J. A. Determinants of myocardial conduction velocity:
Implications for arrhythmogenesis. *Front Physiol* **4 JUN**, 1–14 (2013).

56. Zhao, B. *et al.* Heart-brain connections: Phenotypic and genetic insights from magnetic
resonance images. *Science* **380**, abn6598 (2023).

Methods

Cardiac digital twin creation

The data used in this study is from the UK biobank, an open-access resource, under Application Number 88878. Ethical approval is obtained from the Northwest Research Ethics Committee (REC reference: 11/ NW/0382) and written consent is obtained from all participants. The full heart MRI protocol is described previously[1]. The present study was performed on the 4,329 first participants from the UK biobank who had adequate geometrical information[2] and reported QRS duration (QRSd), sex, age, BMI/weight and height information. The details of the selection of population sample size and quality control were described previously[2,3]. The CDT creation workflow includes anatomical mesh construction and electrophysiological model calibration, and Supplementary Table 1 shows the summary participant characteristics for subjects who successfully went through both steps with existing QRSd, sex, age, BMI/weight, and height information reported in UKBB.

Anatomical mesh generation

We used a nnU-net based architecture[4] for automatic segmentation of the LV and RV blood pools and LV myocardium, trained based on manual segmentations on short-axis heart images[3]. The end-diastolic (ED) phase was selected as the first phase of acquisition. The contours and landmarks of LV and RV derived from the segmentations on the ED frame, were fed to an atlas-based pipeline (previously validated)[2] to construct personalized biventricular surface meshes. The RV epicardium was estimated by extending the RV endocardium points normal to the surfaces by 3 mm consistent with experimental measurements[5,6]. The surface meshes were used to construct tetrahedral finite element meshes using Meshtool[7] including regions of LV myocardium, RV myocardium, aortic, tricuspid, pulmonary and mitral valves. The biventricular myocardial mass was computed from the myocardial volume using a density of $1.05\ g/mL$.

To enable automated computation for CDTs, a morphological coordinate system, known as universal ventricular coordinates (UVCs), was introduced for describing positions within ventricles based on the apical-basal (Z), transmural (ρ) (from endocardium to epicardium), rotational (φ) (anterior, anteroseptal, inferior, inferolateral, anterolateral) and chamber-wise (left ventricle and right ventricle) coordinates. Biventricular myocardial fibre structure was implemented using a rule-based approach with a transmural variation of angle α as from 60° to -60° in longitudinal fibre directions and angle β as from $-65°$ to 25° in transverse fibre directions from endocardium to epicardium.

Electrophysiological (EP) model framework

The electrophysiological simulations are performed using OpenCarp[8] (https://opencarp.org/). We used a reaction-eikonal model without diffusion to compute the sinus ventricular activation times and transmembrane potential transient over time[9]. The activation wavefront propagation in the myocardium Ω is described as:

$$\begin{cases} \sqrt{\nabla t_a^T\ \mathbf{V}\ \nabla\ t_a} = 1\ in\ \Omega \\ t_a = t_0\ in\ \Gamma \end{cases}$$

where t_a is the local activation time at any location in the myocardium, t_0 are the instants of initial activation at locations Γ and the tense field $\mathbf{V}$ encodes the spatially heterogeneous orthotropic squared conduction velocity (CV). The ventricular myocardium was treated as transversely isotropic

conductors. The ten Tusscher ionic model was used to simulate electrophysiological dynamics in the
ventricular myocytes[10]. The ventricular depolarization during sinus rhythm is initiated by the His-
Purkinje system (HPS). As direct measurement of the HPS in the cohort is not available, a fascicular-
based model was used to represent the emergent physiological features of the HPS[11]. Overall, the
electrophysiological framework consists of 20 EP parameters with uncertainty as shown in Extended
Fig. 1 and Supplementary Table 2.

Extended Fig. 1: Fascicular-based model illustration replicating the realistic ventricular activation modulating by the His-
Purkinje system. The SE layer (yellow) represents the fast conduction regions, where the fascicles are located, defined by
apical-basal coordinate Z and transmural coordinate ρ, bounded by physiological measurements from literature as shown
in Supplementary Table 2. There are also five early activation sites represented by disks with a thickness of 5% of the
ventricular wall (δ_z and δ_ρ=0.05).) and having a fixed radius of $20\mu m$, representing ~25 cells. The disks are centred at root
locations, defined by apical-basal coordinates Z and rotational coordinates φ, which are activated at specific timings. Both
coordinates and timings of the five sites are bounded by physiological measurements from literature as shown in
Supplementary Table 2.

Computing electrocardiograms (ECGs) requires information on the position of the heart within the
torso, however, this information was not available. The heart models were therefore registered to a
heart enclosed in an existing torso model[12] using the UVCs. In this torso model, the locations of
electrodes used in measuring 12-lead ECGs were identified the corresponding extracellular
potentials were simulated, and the ECGs were computed. The QRSd was computed by finding the
time points at which the spatial velocity exceeds 0.15 of the maximum spatial velocity in the
reconstructed corresponding vectorcardiogram (VCG) from 12-lead ECGs, which fuse the
information in all ECG traces[12]. This approach introduced extra uncertainty regarding the relative
locations of the ten ECG electrodes within the real torsos. To quantify this uncertainty, we
introduced another 30 parameters (Supplementary Table 3) which describe the variation of the
cartesian coordinates for the ten electrodes and assumed that each cartesian coordinate can vary
$\pm 5 \, cm$, sufficiently to take account of all possible variations in electrode locations[13].

In summary, the entire electrophysiological framework resulted in 50 parameters with uncertainty,
including 20 tissue-level characteristics (Supplementary Table 2) and 30 additional parameters for
the electrode locations (Supplementary Table 3).

Gaussian process emulator and global sensitivity analysis

To build personalized CDTs, we need to set 50 parameters. However, the clinical measurements
needed to constrain these parameters are not available and classical calibration techniques are
prohibitive due to the massive computational costs required per case and the number of cases that
we need to calibrate. We use Gaussian process emulators (GPEs) as surrogate models to accelerate
the evaluation of the effect of the input parameters on the model output of interest: QRSd. This
allows us to (1) gain important mechanistic knowledge about the input-output interactions and (2)
through a GSA exclude the parameters that have little/no effects on the output, speeding up the
personalization pipeline.

To investigate the effects of input parameters on QRSds in the cohort, we generated 10
representative samples by Latin hypercube sampling on basic characteristics including sex, age and
BMI and identifying the 10 cases, who have the closest information to those samples, as
representative subjects. Then we trained Gaussian processes emulators (GPEs) for each subject with
the output as QRSd and performed a global sensitivity analysis (GSA) using the trained GPEs, to
identify the key input parameters explaining the majority variation in the output QRSd. We train
GPEs and perform GSA, firstly on the 20 tissue-level EP parameters to exclude parameters having
small effects on the output QRSd. Then we combined the key input parameters with the other 30
parameters of ECG electrodes' locations to identify the key input parameters in the whole EP
framework, explaining the majority of variation in the output QRSd.

The training of GPEs is described previously[14] by maximizing the model log-marginal likelihood using
the GPErks emulation tool (http://github.com/stelong/GPErks). We evaluated the accuracy of GPEs
using both coefficient of determination R^2 and independent standard error ISE as:

$$R^2 = 1 - \frac{\sum_{i=1}(y_i^{true} - y_i^{mean})^2}{\sum_{i=1}(y_i^{true} - \bar{y})^2}$$

$$ISE := \frac{100}{n} \cdot \sum_{i=1}^{n} (\frac{|y_i^{true} - y_i^{mean}|}{\sqrt{y_i^{var}}} < 2)$$

where y_i^{true} is the true output, $\bar{y}$ is the mean of true outputs, y_i^{mean} and y_i^{var} is the predicted
posterior mean and variance of emulator outputs. The R^2 evaluates the error between the
predictions and the observations with close to 1 indicating a lower error. The ISE accounts for the
distance between predictions to the true values and quantifies the GPE uncertainty. If ISE is close to
100%, it means that the true values are falling in the region of the predictions within 2 standard
deviations. The R^2 and ISE of trained GPEs for the 10 subjects were shown in Supplementary Table
4.

We then performed a variance-based global sensitivity analysis (GSA)[15] on the trained GPEs to
quantify the total effect for each input parameter (S_T) on QRSd. The total effect consists of the first-
order effect and the higher-order interactions, which are computed using the Saltelli method[15] with
the SALib Python library by taking N=1000 samples from the posterior distribution of trained GPEs.
This allowed us to account for the effect of GPE uncertainty on the sensitivity indices. From the GSA,
we ranked the total effects (S_T) of the input parameters to identify the key parameters explaining

the majority variation in the output QRSd. To train each GPE, we used Latin hypercube sampling to
obtain 300 samples, and we ran EP simulations for those samples to build the training dataset. The
impact of training data set size on the GSA was tested in the supplementary Figure 1.

Personalised QRS duration calibration

To facilitate the replication of the clinically measured QRSd in the UK biobank across populations, a
calibration workflow without any manual intervention is preferred. The GSA performed above
enables us to identify the key input parameters responsible for the variation of QRSd, and allows us
to reduce the number of input parameters required for the calibration process. Here we choose to
only vary the most important parameter identified in the GSA for QRSd calibration, therefore
allowing a computationally efficient bisection method to search for the subject-specific parameter
(constrained by physiological measurements in literature) to match the corresponding QRSd.

To quantify the uncertainty brought by only varying the most important parameter to QRSd, we
have computed confidence intervals for the inferred parameter for the 10 subjects used in GPE
training. First, we randomly sampled (N=300) the other 49 parameters, assuming they are from a
normal distribution with a mean equal to the median of the physiological bounds and the upper and
lower bounds set at mean plus or minus three standard deviations (encompassing 99.72% of the
data). Then, we performed simulations using the 300 samples combined with the subject-specific
fitted parameter, to estimate the potential range of the output QRSd in each subject given the
uncertainty in the uncalibrated 49 parameters. We then used the 5th and 95th percentiles of the
output QRSd as the targets, to refit the key input parameter and then calculate their deviations from
the original subject-specific fitted parameters which represents the confidence interval of the
inferred parameter, considering the variation of other 49 parameters as in the Supplementary Figure
2.

12 lead ECGs comparison between the simulations and the recordings in UKBB

To assess the approximate errors of the simulated subject-specific 12 lead ECGs due to using a
standard torso and fixed electrode locations in the personalized QRSd calibration, we compared the
simulated 12 lead ECGs of 10 representative subjects across sex, age and BMI with their recorded
ECGs in UKBB. The simulated 12 lead ECGs were first filtered by the same filters as used in processed
recorded ECGs in the UKBB. It includes a low pass filter at 100 Hz, a high pass filter at 50 Hz and a
notch filter at 50 Hz. Then, each lead of simulated ECGs was temporally aligned to the corresponding
recorded lead ECG by matching the time points that the maximum energy was achieved (V^2) and
was also scaled in amplitude to obtain the same maximum absolute values as in the recorded ECG.
Each lead of the recorded ECG was also cropped to have the same length as the aligned and scaled
simulated lead ECG. Finally, each pair of simulated and recorded lead ECGs were compared by
computing the Pearson correlation coefficient (r). For each subject, the 12-lead ECGs were ranked
by r and the average r were calculated by considering different numbers of ranked ECG leads.

Statistical analysis

145 Statistical analysis was performed using the Python Statsmodels library. The results were presented
146 as mean ± a standard deviation unless specified. The body mass index (BMI) was calculated from
147 height and weight measures taken at the time of the MRI being taken. Codes for the UK Biobank
148 fields are included in brackets. Age was computed using the year of birth (34), month of birth (52)
149 and date of attending the assessment centre (53) to get the actual age when imaging occurred. The

Student t-test was used to compare two different groups. In the phenome-wide association (PheWAS) study, we used a similar approach as in[16]. Before computing univariate cross-correlation, effects such as age, sex, weight, and height were regressed out of the multimodal phenotypes, as they may confound with many phenotypes. The phenotypes from UKBB were normalized and then univariate cross-correlation was applied between the de-confounded multimodal phenotypes and the UKBB phenotypes. The UKBB phenotypes were categorized into 15 groups, including pulse wave analysis (128), LV size and function (133), abdominal composition (149), primary demographics (1001), early life (1002), self-reported medical conditions (1003), lifestyle diet (1004), physical measures (1006), education employment (1007), mental health (1018), summary diagnosis derived from summary diagnoses for hospital inpatient (41270), lifestyle alcohol (100051), physical activity (100054), smoking (100058), and medication. The self-reported medical condition and summary diagnosis were processed to have each column representing one disease code sorted in ascending order, before using in PheWAS. The medications group consists of all phenotypes relating to medications being used summarized from UKBB with codes including 20003, 22170, 22172, 22179, 22167, 22174, 22169, 22171, 22168, 22176, 22166, 22175, 22180, 22181, 22173, 22178, 22177, 6177, 6153, 10004, 6154, 20504, 20551, 20076, 20549, 20546, 2492. We cleaned the data before performing PheWAS by discarding phenotypes with more than 90% missing data and if two highly correlated phenotypes with correlation coefficient>0.9999, only one phenotype was kept.

Data availability

The imaging data and non-imaging participant phenotypes and clinical outcomes are available from UK Biobank via a standard application procedure at http://www.ukbiobank.ac.uk/register-apply. The model generation and simulations are performed using open-sourced software: OpenCarp, available at https://opencarp.org/.

Acknowledgements

This project is supported by the Wellcome Trust and EPSRC Centre for Medical Engineering (WT203148/Z/16/Z). S.N is also supported by NIH R01-HL152256, ERC PREDICT-HF 453 (864055), BHF (RG/20/4/34803), EPSRC (EP/X012603/1, EP/P01268X/1) and by the Technology Missions Fund under the EPSRC Grant EP/X03870X/1 & The Alan Turing Institute. M.B acknowledges BHF Project Grants PG/22/11159 and PG/22/10871.

Author contributions

S.Q, M.S, R.R, A.Y, P.L, M.B, and S.N conceived and designed the study. S.Q, D.U, E.F, L.T, and Y.D implemented the anatomical model generation pipeline and applied it to the cohort. S.Q implemented the functional twinning pipeline, applied it to the cohort, analyzed the results and wrote the manuscript. All authors contributed to the manuscript revision and approved the final submitted version.

Competing interests

The authors declare no competing interests.

Reference

1.   Petersen, S. E. *et al.* UK Biobank's cardiovascular magnetic resonance protocol. *Journal of*
Cardiovascular Magnetic Resonance **18**, (2016).

2.   Mauger, C. *et al.* Right ventricular shape and function: Cardiovascular magnetic resonance
reference morphology and biventricular risk factor morphometrics in UK Biobank. *Journal of*
Cardiovascular Magnetic Resonance **21**, (2019).

3.   Petersen, S. E. *et al.* Reference ranges for cardiac structure and function using cardiovascular
magnetic resonance (CMR) in Caucasians from the UK Biobank population cohort. *Journal of*
Cardiovascular Magnetic Resonance **19**, 1–19 (2017).

4.   Isensee, F., Jaeger, P. F., Kohl, S. A. A., Petersen, J. & Maier-Hein, K. H. nnU-Net: a self-
configuring method for deep learning-based biomedical image segmentation. *Nat Methods*
18, 203–211 (2021).

5.   Matsukubo, H., Matsuura, T., Endo, N., Asayama, J. & Watanabe, T. Echocardiographic
measurement of right ventricular wall thickness. A new application of subxiphoid
echocardiography. *Circulation* **56**, 278–284 (1977).

6.   Ho, S. Y. & Nihoyannopoulos, P. Anatomy, echocardiography, and normal right ventricular
dimensions. *Heart* **92**, 2–13 (2006).

7.   Neic, A., Gsell, M. A. F., Karabelas, E., Prassl, A. J. & Plank, G. Automating image-based mesh
generation and manipulation tasks in cardiac modeling workflows using Meshtool. *SoftwareX*
11, 100454 (2020).

8.   Plank, G. *et al.* The openCARP simulation environment for cardiac electrophysiology. *Comput*
Methods Programs Biomed **208**, 1–19 (2021).

9.   Neic, A. *et al.* Efficient computation of electrograms and ECGs in human whole heart
simulations using a reaction-eikonal model. *J Comput Phys* **346**, 191–211 (2017).

10. Ten Tusscher, K. H. W. J. & Panfilov, A. V. Alternans and spiral breakup in a human ventricular
tissue model. *Am J Physiol Heart Circ Physiol* **291**, 1088–1100 (2006).

11. Gillette, K. *et al.* Automated Framework for the Inclusion of a His–Purkinje System in Cardiac
Digital Twins of Ventricular Electrophysiology. *Ann Biomed Eng* **49**, 3143–3153 (2021).

12. Gemmell, P. M. *et al.* A computational investigation into rate-dependant vectorcardiogram
changes due to specific fibrosis patterns in non-ischæmic dilated cardiomyopathy. *Comput*
Biol Med **123**, 103895 (2020).

13. Monaci, S. *et al.* Automated Localization of Focal Ventricular Tachycardia From Simulated
Implanted Device Electrograms: A Combined Physics–AI Approach. *Front Physiol* **12**, 1–15
(2021).

14. Longobardi, S., Sher, A. & Niederer, S. A. In silico identification of potential calcium dynamics
and sarcomere targets for recovering left ventricular function in rat heart failure with
preserved ejection fraction. *PLoS Comput Biol* **17**, 1–22 (2021).

15. Saltelli, A. *et al.* Variance based sensitivity analysis of model output. Design and estimator for
the total sensitivity index. *Comput Phys Commun* **181**, 259–270 (2010).

16. Bai, W. *et al.* A population-based phenome-wide association study of cardiac and aortic
structure and function. *Nat Med* **26**, 1654–1662 (2020).

232