

# SARS-CoV-2 Infection is Associated with an Increase in New Diagnoses of Schizophrenia Spectrum and Psychotic Disorder: A Study Using the US National COVID Cohort Collaborative (N3C)

Asif Rahman<sup>1</sup>, Michael Russell<sup>2</sup>, Wanhong Zheng<sup>3</sup>, Daniel Eckrich<sup>4</sup>, Imtiaz Ahmed<sup>1\*</sup>,  
On behalf of the N3C Consortium

**1** Department of Industrial & Management Systems Engineering, West Virginia University, Morgantown, WV, USA

**2** School of Medicine, West Virginia University, Morgantown, WV, USA

**3** School of Medicine, West Virginia University, Morgantown, WV, USA

**4** Nemours Children's Health, Jacksonville, FL, USA

\* [imtiaz.ahmed@mail.wvu.edu](mailto:imtiaz.ahmed@mail.wvu.edu)

## Abstract

Amid the ongoing global repercussions of SARS-CoV-2, it's crucial to comprehend its potential long-term psychiatric effects. Several recent studies have suggested a link between COVID-19 and subsequent mental health disorders. Our investigation joins this exploration, concentrating on Schizophrenia Spectrum and Psychotic Disorders (SSPD). Different from other studies, we took acute respiratory distress syndrome (ARDS) and COVID-19 lab negative cohorts as control groups to accurately gauge the impact of COVID-19 on SSPD. Data from 19,344,698 patients, sourced from the N3C Data Enclave platform, were methodically filtered to create propensity matched cohorts: ARDS (n = 222,337), COVID-positive (n = 219,264), and COVID-negative (n = 213,183). We systematically analyzed the hazard rate of new-onset SSPD across three distinct time intervals: 0-21 days, 22-90 days, and beyond 90 days post-infection. COVID-19 positive patients consistently exhibited a heightened hazard ratio (HR) across all intervals [0-21 days (HR: 4.6; CI: 3.7-5.7), 22-90 days (HR: 2.9; CI: 2.3 -3.8), beyond 90 days (HR: 1.7; CI: 1.5-1.9)]. These are notably higher than both ARDS and COVID-19 lab-negative patients. Validations using various tests, including the Cochran Mantel Haenszel Test, Wald Test, and Log-rank Test confirmed these associations. Intriguingly, our data indicated that younger individuals face a heightened risk of SSPD after contracting COVID-19, a trend not observed in the ARDS and COVID-negative groups. These results, aligned with the known neurotropism of SARS-CoV-2 and earlier studies, accentuate the need for vigilant psychiatric assessment and support in the era of Long-COVID, especially among younger populations.

## Introduction

It has been over three years since the initial identification of SARS-CoV-2 infection (hereafter referred to as COVID-19) in the USA. Despite the development of a vaccine and efforts to combat the pandemic, there are still many unanswered questions. Particularly, the long-term effects of COVID on mental health are yet to be fully unwrapped and associated with the disease. Several preliminary studies [1-3] have suggested an increased risk of mental illness following a COVID-19 diagnosis, including

but not limited to anxiety, depression, mood disorder, post-traumatic stress disorder (PTSD), insomnia, dementia, delirium, encephalitis, psychosis, and nerve disorder. It is important to note that viral infections resulting from recent outbreaks of severe acute respiratory syndrome (SARS) in 2002 and Middle East respiratory syndrome (MERS) in 2012, both caused by coronavirus closely related to SARS-CoV-2, were also associated with neurological manifestations in some cases [3].

COVID-19 has a multi-organ pathology that includes the human brain and the central nervous system [4]. It has been detected both in the brain and cerebrospinal fluids of the diagnosed patients. The COVID-19 patients show a greater cognitive decline compared to the non-COVID patients. It has also been associated with brain structural change [5]. Recent studies have suggested that more than one-third of the infected individuals develop neurological symptoms in the acute phase of the disease, and around 34% of them show brain abnormalities [6, 7]. COVID-19 has been linked to excessive and dysregulated immune responses that can lead to systemic inflammation. Patients with severe COVID-19 have been found to have elevated levels of various inflammatory markers in their blood, such as C-reactive protein (CRP), ferritin, interleukin-6 (IL-6), and others. Among them, higher levels of CRP and IL-6 are linked to a higher risk of different neurological conditions such as major depressive disorders.

It has been well established that a bidirectional interaction exists between the central nervous system, specifically the brain, and systemic inflammation [9]. In the brain, microglia are the principal cells involved in modulating the effects of remote inflammatory stressors resulting in neuro-inflammatory manifestations of systemic processes arising from multiple causes (trauma, infection, auto-immune processes, etc.) [10]. Microglial functional and structural alterations have been found in multiple major psychiatric disorders [10–12] although presence, degree and configuration of such alterations vary between diagnoses. Therefore the etiologic and therapeutic significance of these observations remains unclear [13].

A growing body of literature has implicated systemic inflammation associated with critical illness in the development of delirium [17]. In turn, the occurrence of delirium during critical illness is associated with persistent deficits in neurocognitive function following survival [18]. The presence of pre-exposure decline in cognitive function is associated with an increased risk of post-critical illness persistent neurocognitive disability [19, 20]. Difference in epigenetic DNA methylation patterns in critically ill patients with and without delirium have recently been reported [21]. It is now well established that systemic inflammation affects brain function in critical illness and that these effects are persistent beyond the intensive care episode [22, 23]. While the exact mechanisms are still incompletely characterized, epigenetic modification of DNA directed protein transcription may play a potential role.

Major psychiatric illness is known to have a strong genetic component. However, variable penetrance suggests that environmental factors are also important in the development of clinical disease [24]. Emerging data suggests a significant association between neuroinflammatory changes and major psychiatric illness. The strongest associations to date involve the development of schizophrenia, bipolar disorder and major depression [10, 11, 25].

Given the increasing evidence demonstrating a link between systemic inflammation and neurocognitive function, along with the role of neuroinflammation in manifesting major psychiatric disorders and the systemic inflammatory effects of COVID-19, the authors sought to establish if COVID-19 could lead to a rise in the onset of significant psychiatric conditions. We decide to concentrate on the schizophrenia spectrum and psychotic disorder (SSPD), as the data linking inflammatory conditions with the emergence or progression of the disease is most compelling in this context.

We are aware that several studies [2, 3, 8] tried to establish an association between

COVID-19 and psychiatric manifestations. However, a significant portion of these studies lacked an appropriate comparison group, leading to an incomplete understanding of the incidence and prevalence of neuro-psychiatric disorders in COVID-19 patients. To address this limitation, our study incorporated a control group comprising individuals with Acute Respiratory Distress Syndrome (ARDS) and those who tested negative for COVID-19. This approach ensured a more precise correlation between COVID-19 and SSPD. Furthermore, we leveraged the N3C platform, utilizing its vast, robust, and long-term data set to effectively discern and quantify the impact of COVID-19 on SSPD.

## Materials and Methods

This study was a retrospective cohort study. All data was collected from the National COVID-19 Cohort Collaborative (N3C) Data Enclave platform. The dataset was retrieved on May 31, 2023, and we limited our analysis to include only records up to that date. Throughout the process of data collection and subsequent analysis, the authors did not have access to any information that could be used to identify individual participants.

## Study Design and Data Collection

To achieve our objectives, we initiated a systematic filtering process as depicted in Figure 1. Out of an initial dataset of 19 million patients, we categorized them into three primary groups: Acute Respiratory Distress Syndrome (ARDS), COVID-positive, and COVID-negative. We applied specific criteria to refine these cohorts. Firstly, we considered only those with a minimum of three visits. Secondly, we excluded patients with any pre-existing mental health conditions and further narrowed down our scope to individuals aged between 17 and 70 years. Notably, within the COVID-positive group, we focused on patients characterized by moderate, severe, or terminal outcomes due to the virus. After implementing these filters, our COVID-positive cohort size was finalized at 244,226. To ensure that our data from these groups could be directly compared, we also implemented a propensity score matching technique.

**Fig 1.** Final cohort selection by applying exclusion criteria

## Cohorts

Three distinct cohorts were constructed for this study: one case and two controls. The case cohort was made up of patients diagnosed with COVID-19, based on the N3C defined computable phenotype version 4.0. [50]. In order to derive meaningful insights, we limited our COVID-19 positive patients to those with moderate to severe manifestations. This categorization was determined by several factors, including the duration of inpatient hospital stays, usage of invasive ventilation, application of extracorporeal membrane oxygenation (ECMO), and even unfortunate fatalities.

For our controls, we selected patients diagnosed with ARDS post-January 1st, 2020 but without any record of a COVID-19 diagnosis during the pandemic. The third group consisted of those who tested negative for COVID-19 and had no prior history of either COVID-19 or ARDS. The starting population for our study needed to have a history of at least three medical visits spanning 365 days or more. To standardize the timeline across cohorts, index dates were determined based on the earliest date of relevant diagnosis or lab test results.

We excluded individuals below 17 years or over 70 years at the time of the index date. Furthermore, patients with the following mental health disorders prior to the index date were also removed from consideration:

- Schizophrenia Spectrum Disorders
- Bipolar Disorders
- Major Depression
- Personality Disorders
- Trauma

For homogeneity, all cohorts underwent 1:1 nearest neighbor (NN) matching on propensity scores using the R “MatchIt” package. This matching was based on the following attributes:

- Gender
- Race
- Ethnicity
- Age groups
- Prior psychiatric drug prescription or administration
- Prior Hypothyroidism diagnosis
- Prior Anxiety diagnosis
- Prior Substance Abuse diagnosis
- Prior Insomnia diagnosis

Please refer to the appendix for specific Observational Medical Outcomes Partnership (OMOP) codesets [51] used for identifying these attributes. Post-matching, the cohort sizes were 219,264 for COVID-19 positive patients, 213,183 for the lab negatives, and 222,337 for ARDS patients.

## Outcome

Once the cohorts were built, we looked at the first incident post-index date of SSPD. The code sets were developed using the OMOP concept and `concept_ancestor` tables [51] and reviewed by subject matter experts. While psychiatry continued to debate over the relationship between psychotic symptoms and mood symptoms [52,53], in this study, to focus on the hallmark thought symptoms characteristic of schizophrenia (such as delusions, hallucinations, or disorganized speech), we separated SSPD from bipolar and depressive disorders. Therefore, while acute psychotic disorder, schizophreniform, schizophrenia, schizoaffective disorders, and related ICD-10 diagnoses were included in the SSPD category, mood disorders with psychotic features were counted in the latter two categories and thus excluded from the analysis.

The data was subsequently structured to include three pivotal columns: a boolean flag denoting whether a patient was diagnosed with SSPD after the index date, the exact date of the patient’s initial SSPD diagnosis post the index date, and the duration in days between the index date and this diagnosis. This data arrangement culminated in a matrix with a single record for each patient, encompassing cohort identifiers, demographic details, other relevant covariates, and the critical outcome variable.

## Statistical Analysis

To examine the association between COVID-19 and SSPD, we performed a comparative analysis among the matched cohorts using the N3C platform in R (version 4.0). Our primary predictor variable is the disease type, categorized into three groups: COVID-positive, COVID-negative, and ARDS. The outcome variable is binary, indicating either SSPD or non-SSPD.

We utilized the Cox Proportional Hazard Model [43] to derive the hazard ratios (HR) of COVID-positive patients relative to the COVID-negative and ARDS patients. The time-to-event for patients diagnosed with SSPD was measured from their SSPD diagnosis date up to the COVID and ARDS reference dates, which include the dates of their positive COVID test, negative COVID lab test, and ARDS diagnosis. For patients without an SSPD diagnosis, this duration was taken from their most recent recorded visit to the reference date of either their COVID or ARDS diagnosis.

Before deploying the Cox model, it was imperative to test the proportional hazard assumption. In doing so, the Schoenfeld residual analysis, a conventional diagnostic tool for this purpose [44], yielded a significant p-value. This necessitated rejecting the null hypothesis of a uniform proportional hazard over the comprehensive time frame of 180 days. Consequently, we segmented the cohort into three distinct time intervals: 0-21 days, 22-90 days, and beyond 90 days. These intervals were subsequently validated for the proportional hazard assumptions, and the Cox model was then applied to each to ascertain the hazard ratio (HR).

In tandem with the Cox model, we also conducted the Cochran Mantel Haenszel Test [45], the Likelihood Ratio Test [46], the Wald Test [45], and the Log-rank Test [47] across the three time intervals. A p-value threshold of 0.05 served as the determinant for statistical significance in all these tests.

## Results

The results shed light on the potential long-term psychiatric implications of SARS-CoV-2 infection. The study found robust evidence linking SARS-CoV-2 infection to an augmented risk of Schizophrenia Spectrum and Psychotic Disorders (SSPD). Specifically, COVID-positive patients displayed almost double the incident rate (0.56%) compared to COVID-negative (0.33%) and ARDS (0.29%) patients, as showcased in Table 1.

Using the Cox model, a marked difference in the hazard ratio of new-onset psychiatric outcomes became evident between COVID-19 positive patients and the cohorts of ARDS and COVID-19 lab negative patients (Please refer to Table 2). For all time intervals considered, COVID-negative patients were the benchmark for hazard ratio computations. In the immediate 21 days following exposure, the hazard ratio for COVID-positive patients was notably high (HR: 4.6; 95% CI: 3.7 to 5.7) when contrasted with ARDS patients (HR: 0.73 CI: 0.53 to 0.99). This suggests that, during the acute phase, individuals positive for COVID-19 were significantly more likely to be diagnosed with SSPD than both their COVID-negative and ARDS counterparts.

In the subsequent interval of 22-90 days, the hazard ratio for COVID-positive patients remained elevated (HR: 2.9; 95% CI: 2.3 to 3.8), and interestingly, ARDS patients exhibited their peak hazard ratio of the study (HR:1.1, CI: 0.79 to 1.43). Beyond 90 days, the hazard ratio for the COVID-positive group experienced a reduction (HR: 1.7; 95% CI: 1.5 to 1.9) yet was consistently higher than that of the ARDS patients (HR:0.97, CI: 0.86 to 1.47). Despite the reduction in hazard ratios as time progressed, it's salient to note that COVID-19 survivors remain at a heightened risk for SSPD well beyond the immediate aftermath of their infection. The hazard ratios for

Table 1. Characteristics of Cohorts

Characteristics	ARDS		COVID Positive		COVID Negative	
	N = 222337		N = 219264		N = 213183	
	Count	Percentage	Count	Percentage	Count	Percentage
<b>Gender:</b>						
Female	105175	47.30%	103528	47.20%	105959	49.70%
Male	117162	52.70%	115736	52.80%	107196	50.30%
Other or Unknown	0	0.00%	0	0.00%	28	0.01%
<b>Age Groups:</b>						
21 and Younger	7896	3.55%	7449	3.40%	7840	3.68%
22 to 29	19885	8.94%	19160	8.74%	19644	9.21%
30 to 39	31070	14.00%	30298	13.80%	29400	13.80%
40 to 49	35871	16.10%	34407	15.70%	31594	14.80%
50 to 59	52529	23.60%	53059	24.20%	51115	24.00%
60 and Older	75086	33.80%	74891	34.20%	73590	34.50%
<b>Race:</b>						
Asian	5519	2.48%	5580	2.54%	5568	2.61%
Black/African American	48251	21.70%	47047	21.50%	45269	21.20%
White/Caucasian	132914	59.80%	126061	57.50%	127741	59.90%
Unknown	29357	13.20%	31810	14.50%	27014	12.70%
Nothing mentioned	1746	0.79%	2701	1.23%	2758	1.29%
Other	4550	2.05%	6065	2.77%	4833	2.27%
<b>Ethnicity:</b>						
Hispanic or Latino	34320	15.40%	38196	17.40%	31002	14.50%
Not Hispanic or Latino	171609	77.20%	165071	75.30%	166149	77.90%
Missing/Unknown	16408	7.38%	15997	7.30%	16032	7.52%
<b>Any Psychiatric Disease (before index date):</b>						
No	145612	65.50%	138029	63.00%	137490	64.50%
Yes	76725	34.50%	81235	37.00%	75693	35.50%
<b>Anxiety Disorder:</b>						
No	198285	89.20%	201933	92.10%	192958	90.50%
Yes	24052	10.80%	17331	7.90%	20225	9.49%
<b>Insomnia Disorder:</b>						
No	211921	95.30%	212449	96.90%	204869	96.10%
Yes	10416	4.68%	6815	3.11%	8314	3.90%
<b>Hypothyroidism Disorder:</b>						
No	206762	93.00%	204850	93.40%	198391	93.10%
Yes	15575	7.01%	14414	6.57%	14792	6.94%
<b>Hypothyroidism Disorder (Lab Test):</b>						
No	212801	95.70%	208228	95.00%	201818	94.70%
Yes	9536	4.29%	11036	5.03%	11365	5.33%
<b>Drug Abuse / Addiction Disorder:</b>						
No	189594	85.30%	190323	86.80%	188360	88.40%
Yes	32743	14.70%	28941	13.20%	24823	11.60%
<b>Drug Abuse / Addiction Disorder (Lab Test):</b>						
No	218686	98.40%	211424	96.40%	209335	98.20%
Yes	3651	1.64%	7840	3.58%	3848	1.81%
<b>All Mental Health:</b>						
No	207448	93.30%	203123	92.60%	197204	92.50%
Yes	14889	6.70%	16141	7.36%	15979	7.50%
<b>SSPD:</b>						
No	221700	99.70%	218035	99.40%	212476	99.70%
Yes	637	0.29%	1229	0.56%	707	0.33%
<b>Bipolar Disorder:</b>						
No	214506	96.50%	210281	95.90%	205014	96.20%
Yes	7831	3.52%	8983	4.10%	8169	3.83%
<b>Personality Disorder:</b>						
No	222034	99.90%	218951	99.90%	212889	99.90%
Yes	303	0.14%	313	0.14%	294	0.14%
<b>Depression:</b>						
No	210088	94.50%	205253	93.60%	199865	93.80%
Yes	12249	5.51%	14011	6.39%	13318	6.25%
<b>Trauma:</b>						
No	219432	98.70%	217107	99.00%	210308	98.70%
Yes	2905	1.31%	2157	0.98%	2875	1.35%

different time intervals are visually summarized in Figure 2.

Following the detailed hazard ratio analysis, several statistical tests (Please refer to Table 3) were conducted to further validate the findings. The Schoenfeld residuals test returned a p-value exceeding 0.05, suggesting the retention of the null hypothesis that the Hazard Ratio is consistent over time. Additionally, tests such as the Cochran Mantel Haenszel Test, Likelihood Ratio Test, Wald Test, and Log-rank Test consistently



**Table 2. Hazard Ratios with 95% Confidence Interval (CI) For SSPD In Different Time Frames.**

Time Frame (Days)	COVID Negative				COVID Positive				ARDS		
	Hazard Ratio	95 % CI			Hazard Ratio	95 % CI			Hazard Ratio	95 % CI	
		Lower	Upper			Lower	Upper			Lower	Upper
0 - 21	*	*	*		4.6	3.7	5.7		0.73	0.53	0.99
22 - 90	*	*	*		2.9	2.3	3.8		1.1	0.79	1.43
Beyond 90	*	*	*		1.7	1.5	1.9		0.97	0.86	1.47

\*COVID Negative patient group is considered as references.

**Fig 2. SSPD Hazard Ratio Comparisons.**

showed p-values less than 0.05, reinforcing a significant association between being COVID-positive and receiving an SSPD diagnosis.

In our efforts to determine potential demographic factors influencing SSPD occurrence among COVID-positive patients, several key demographics emerged as more susceptible to SSPD following a COVID-19 diagnosis. Specifically, males, individuals aged 21 or younger, those of African American descent, and non-Hispanic or Latino individuals showcased a heightened vulnerability (see Table 4 and Table 5 in the appendix for details). Intriguingly, these same demographic trends, with the exception of the age factor, were mirrored in the SSPD occurrence among both the COVID-negative and ARDS groups (refer to Table 6 and Table 7 in the appendix for further insights). While these findings undoubtedly warrant a deeper exploration, we strongly advocate for future research endeavors to prioritize this critical observation regarding the younger people being more susceptible to post-COVID SSPD.

**Table 3. P Value For Different Tests In Different Time Frames.**

P Value	Time Frame (In Days)		
	0 - 21	22 - 90	Beyond 90
Schoenfeld Residuals Test	0.24	0.75	0.15
Cochran Mantel Haenszel Test	< 0.05	< 0.05	< 0.05
Likelihood Ratio Test	< 0.05	< 0.05	< 0.05
Wald Test	< 0.05	< 0.05	< 0.05
Logrank Test	< 0.05	< 0.05	< 0.05

## Discussion

Our study indicates that the likelihood of developing SSPD after a COVID-19 infection is higher than in ARDS and COVID-19-negative patients. The significance of various demographic factors has also emerged from our results. These insights underscore the vital importance of keeping a close watch on the mental well-being of those recovering from COVID-19. Their persistent increased risk points to a wider societal concern, especially regarding severe psychiatric conditions like SSPD.

Extensive literature has accumulated since 2000 that indicates an association between various inflammatory markers, changes in the structure and function of a variety of cellular components of the brain, and the development of major psychiatric illnesses. No longer thought to be “immunologically privileged” by virtue of the blood-brain permeability barrier, it is now well established that the brain is extensively influenced by systemic inflammation and, in turn, can modulate systemic inflammation through descending [26] and biochemical [27] pathways. Inflammatory influences in the brain are structural [28] and functional [9]. Effects of maternal development on

in-utero brain development and subsequent offspring behavior have been demonstrated in animal models [29] although the relevance to mental illness in humans remains to be determined. Inflammatory influences on the developed brain have also been demonstrated and tied to clinically relevant behavior [15,30]. However, studies evaluating the relationship between various inflammatory markers and clinically relevant behavior have yielded inconsistent results [17,32–34]. At the moment, these inflammatory changes cannot be causally linked directly to specific conditions, but they do provide potential mechanistic insights and may become plausible therapeutic targets in conditions where the response to currently available medications varies widely [17,28,30].

With that background, our group sought to establish whether any association existed between SARS-CoV-2 infection and the extensive accompanying inflammatory response and new-onset psychiatric illness. Our results are consistent with the hypothesis that COVID-19 infection and the accompanying inflammatory state (“cytokine storm”) is positively associated with the new onset of SSPD. Further, these results strongly suggest a direct relationship between the development of SSPD and the severity of the disease state and presumably the intensity of the attendant inflammatory response.

The strengths of our study are underscored by the utilization of an expansive and meticulously curated national dataset, enabling an in-depth analysis of a vast number of records. Furthermore, our rigorous approach to defining exclusionary criteria and cohort matching bolster the robustness of our findings. However, the study does possess notable limitations. Foremost among these is, same as any other retrospective studies, the dependence on documented diagnoses (ICD-10 coding) to pinpoint new instances of SSPD. It is conceivable — perhaps even probable — that certain patients might have been inaccurately diagnosed, thereby skewing their categorization as determined by our data extraction methodology. The markedly elevated hazard ratio observed during the acute (0-21 day) phase, for instance, is challenging to rationalize, given that definitive diagnoses for many severe psychiatric conditions, especially SSPD, typically necessitate prolonged periods of behavioral observation. One possible reason might be that many individuals went through an early phase of unnoticed or undeclared symptoms before their clinical visit for COVID. When diagnosing SSPD, doctors likely factored in this extended course of symptoms, likely triggered by the brain inflammation caused by the coronavirus infection. There also exists a possibility that some individuals could have encountered acute and transient psychotic disorders (ATPD), which were erroneously identified as SSPD. Additionally, patients manifesting early signs hinting at SSPD may have exhibited more distinct symptoms post-COVID infection, subsequently leading to accurate diagnostic coding. Absent direct clinical assessments, pinpointing the primary influencing factor remains elusive. Nonetheless, the sustained elevated hazard ratios even beyond the initial 90 days post-infection underscore the continued influence of COVID-19 on the emergence of SSPD.

## Conclusion

In this study, we have found a substantial increase in the likelihood of being diagnosed with a schizophrenia spectrum and psychotic disorder (SSPD) after experiencing moderate to severe illness due to SARS-CoV-2 infection, in comparison to a group of individuals who had non-COVID Acute Respiratory Distress Syndrome (ARDS). Our work is consistent with the known neurotropism of the SARS-CoV-2 virus [39,40] and other reports of increased risk of major psychiatric disorders following COVID-19 infection [41–43]. Further research is required to identify specific characteristics of populations and individuals who may be at a particularly high risk of developing SSPD and potentially other significant psychiatric conditions following COVID-19 infection.



Understanding these psychiatric risks associated with COVID-19 is an essential component of our strategy to address the evolving landscape of Long-COVID.

## Acknowledgments

We would like to acknowledge West Virginia University (WVU) and WVU Clinical & Translational Research Institute (CTSI) for their support. The project described was supported by the National Institute Of General Medical Sciences, 5U54GM104942-04. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

### N3C Attribution

The analyses described in this publication were conducted with data or tools accessed through the NCATS N3C Data Enclave (<https://covid.cd2h.org>) and N3C Attribution & Publication Policy v 1.2-2020-08-25b supported by NCATS U24 TR002306. This research was possible because of the patients whose information is included within the data and the organizations (<https://ncats.nih.gov/n3c/resources/data-contribution/data-transfer-agreement-signatories>) and scientists who have contributed to the ongoing development of this community resource [<https://doi.org/10.1093/jamia/ocaa196>].

**Disclaimer** The N3C Publication Committee confirmed that this manuscript MSID:1645.616 is in accordance with N3C data use and attribution policies; however, this content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or the N3C program.

### IRB

The N3C data transfer to NCATS is performed under a Johns Hopkins University Reliance Protocol # IRB00249128 or individual site agreements with NIH. The N3C Data Enclave is managed under the authority of the NIH; information can be found at <https://ncats.nih.gov/n3c/resources>.

### Individual Acknowledgements For Core Contributors

We gratefully acknowledge the following core contributors to N3C:

Adam B. Wilcox, Adam M. Lee, Alexis Graves, Alfred (Jerrod) Anzalone, Amin Manna, Amit Saha, Amy Olex, Andrea Zhou, Andrew E. Williams, Andrew Southerland, Andrew T. Girvin, Anita Walden, Anjali A. Sharathkumar, Benjamin Amor, Benjamin Bates, Brian Hendricks, Brijesh Patel, Caleb Alexander, Carolyn Bramante, Cavin Ward-Caviness, Charisse Madlock-Brown, Christine Suver, Christopher Chute, Christopher Dillon, Chunlei Wu, Clare Schmitt, Cliff Takemoto, Dan Housman, Davera Gabriel, David A. Eichmann, Diego Mazzotti, Don Brown, Eilis Boudreau, Elaine Hill, Elizabeth Zampino, Emily Carlson Marti, Emily R. Pfaff, Evan French, Farrukh M Koraisky, Federico Mariona, Fred Prior, George Sokos, Greg Martin, Harold Lehmann, Heidi Spratt, Hemalkumar Mehta, Hongfang Liu, Hythem Sidky, J.W. Awori Hayanga, Jami Pincavitch, Jaylyn Clark, Jeremy Richard Harper, Jessica Islam, Jin Ge, Joel Gagnier, Joel H. Saltz, Joel Saltz, Johanna Loomba, John Buse, Jomol Mathew, Joni L. Rutter, Julie A. McMurphy, Justin Guinney, Justin Starren, Karen Crowley, Katie Rebecca Bradwell, Kellie M. Walters, Ken Wilkins, Kenneth R. Gersing, Kenrick Dwain Cato, Kimberly Murray, Kristin Kostka, Lavance Northington, Lee Allan Pyles, Leonie Misquitta, Lesley Cottrell, Lili Portilla, Mariam Deacy, Mark M. Bissell, Marshall Clark, Mary Emmett, Mary Morrison Saltz, Matvey B. Palchuk, Melissa A. Haendel, Meredith Adams, Meredith Temple-O'Connor, Michael G. Kurilla, Michele Morris, Nabeel Qureshi, Nasia Safdar, Nicole Garbarini, Noha Sharafeldin, Ofer Sadan, Patricia A. Francis, Penny Wung Burgoon, Peter Robinson, Philip R.O. Payne, Rafael Fuentes, Randeep Jawa, Rebecca Erwin-Cohen, Rena Patel, Richard A. Moffitt, Richard L. Zhu, Rishi Kamaleswaran, Robert Hurley, Robert T. Miller, Saiju Pyarajan,

Sam G. Michael, Samuel Bozzette, Sandeep Mallipattu, Satyanarayana Vedula, Scott Chapman, Shawn T. O'Neil, Soko Setoguchi, Stephanie S. Hong, Steve Johnson, Tellen D. Bennett, Tiffany Callahan, Umit Topaloglu, Usman Sheikh, Valery Gordon, Vignesh Subbian, Warren A. Kibbe, Wendy Hernandez, Will Beasley, Will Cooper, William Hillegass, Xiaohan Tanner Zhang. Details of contributions are available at <https://covid.cd2h.org/core-contributors>

#### Data Partners with Released Data

The following institutions whose data is released or pending:

Available: Advocate Health Care Network — UL1TR002389: The Institute for Translational Medicine (ITM) • Aurora Health Care Inc — UL1TR002373: Wisconsin Network For Health Research • Boston University Medical Campus — UL1TR001430: Boston University Clinical and Translational Science Institute • Brown University — U54GM115677: Advance Clinical Translational Research (Advance-CTR) • Carilion Clinic — UL1TR003015: iTHRIV Integrated Translational health Research Institute of Virginia • Case Western Reserve University — UL1TR002548: The Clinical & Translational Science Collaborative of Cleveland (CTSC) • Charleston Area Medical Center — U54GM104942: West Virginia Clinical and Translational Science Institute (WVCTSI) • Children's Hospital Colorado — UL1TR002535: Colorado Clinical and Translational Sciences Institute • Columbia University Irving Medical Center — UL1TR001873: Irving Institute for Clinical and Translational Research • Dartmouth College — None (Voluntary) Duke University — UL1TR002553: Duke Clinical and Translational Science Institute • George Washington Children's Research Institute — UL1TR001876: Clinical and Translational Science Institute at Children's National (CTSA-CN) • George Washington University — UL1TR001876: Clinical and Translational Science Institute at Children's National (CTSA-CN) • Harvard Medical School — UL1TR002541: Harvard Catalyst • Indiana University School of Medicine — UL1TR002529: Indiana Clinical and Translational Science Institute • Johns Hopkins University — UL1TR003098: Johns Hopkins Institute for Clinical and Translational Research • Louisiana Public Health Institute — None (Voluntary) • Loyola Medicine — Loyola University Medical Center • Loyola University Medical Center — UL1TR002389: The Institute for Translational Medicine (ITM) • Maine Medical Center — U54GM115516: Northern New England Clinical & Translational Research (NNE-CTR) Network • Mary Hitchcock Memorial Hospital & Dartmouth Hitchcock Clinic — None (Voluntary) • Massachusetts General Brigham — UL1TR002541: Harvard Catalyst • Mayo Clinic Rochester — UL1TR002377: Mayo Clinic Center for Clinical and Translational Science (CCaTS) • Medical University of South Carolina — UL1TR001450: South Carolina Clinical & Translational Research Institute (SCTR) • MITRE Corporation — None (Voluntary) • Montefiore Medical Center — UL1TR002556: Institute for Clinical and Translational Research at Einstein and Montefiore • Nemours — U54GM104941: Delaware CTR ACCEL Program • NorthShore University HealthSystem — UL1TR002389: The Institute for Translational Medicine (ITM) • Northwestern University at Chicago — UL1TR001422: Northwestern University Clinical and Translational Science Institute (NUCATS) • OCHIN — INV-018455: Bill and Melinda Gates Foundation grant to Sage Bionetworks • Oregon Health & Science University — UL1TR002369: Oregon Clinical and Translational Research Institute • Penn State Health Milton S. Hershey Medical Center — UL1TR002014: Penn State Clinical and Translational Science Institute • Rush University Medical Center — UL1TR002389: The Institute for Translational Medicine (ITM) • Rutgers, The State University of New Jersey — UL1TR003017: New Jersey Alliance for Clinical and Translational Science • Stony Brook University — U24TR002306 • The Alliance at the University of Puerto Rico, Medical Sciences Campus — U54GM133807: Hispanic Alliance for Clinical and Translational Research

(The Alliance) • The Ohio State University — UL1TR002733: Center for Clinical and Translational Science • The State University of New York at Buffalo — UL1TR001412: Clinical and Translational Science Institute • The University of Chicago — UL1TR002389: The Institute for Translational Medicine (ITM) • The University of Iowa — UL1TR002537: Institute for Clinical and Translational Science • The University of Miami Leonard M. Miller School of Medicine — UL1TR002736: University of Miami Clinical and Translational Science Institute • The University of Michigan at Ann Arbor — UL1TR002240: Michigan Institute for Clinical and Health Research • The University of Texas Health Science Center at Houston — UL1TR003167: Center for Clinical and Translational Sciences (CCTS) • The University of Texas Medical Branch at Galveston — UL1TR001439: The Institute for Translational Sciences • The University of Utah — UL1TR002538: Uhealth Center for Clinical and Translational Science • Tufts Medical Center — UL1TR002544: Tufts Clinical and Translational Science Institute • Tulane University — UL1TR003096: Center for Clinical and Translational Science • The Queens Medical Center — None (Voluntary) • University Medical Center New Orleans — U54GM104940: Louisiana Clinical and Translational Science (LA CaTS) Center • University of Alabama at Birmingham — UL1TR003096: Center for Clinical and Translational Science • University of Arkansas for Medical Sciences — UL1TR003107: UAMS Translational Research Institute • University of Cincinnati — UL1TR001425: Center for Clinical and Translational Science and Training • University of Colorado Denver, Anschutz Medical Campus — UL1TR002535: Colorado Clinical and Translational Sciences Institute • University of Illinois at Chicago — UL1TR002003: UIC Center for Clinical and Translational Science • University of Kansas Medical Center — UL1TR002366: Frontiers: University of Kansas Clinical and Translational Science Institute • University of Kentucky — UL1TR001998: UK Center for Clinical and Translational Science • University of Massachusetts Medical School Worcester — UL1TR001453: The UMass Center for Clinical and Translational Science (UMCCTS) • University Medical Center of Southern Nevada — None (voluntary) • University of Minnesota — UL1TR002494: Clinical and Translational Science Institute • University of Mississippi Medical Center — U54GM115428: Mississippi Center for Clinical and Translational Research (CCTR) • University of Nebraska Medical Center — U54GM115458: Great Plains IDeA-Clinical & Translational Research • University of North Carolina at Chapel Hill — UL1TR002489: North Carolina Translational and Clinical Science Institute • University of Oklahoma Health Sciences Center — U54GM104938: Oklahoma Clinical and Translational Science Institute (OCTSI) • University of Pittsburgh — UL1TR001857: The Clinical and Translational Science Institute (CTSI) • University of Pennsylvania — UL1TR001878: Institute for Translational Medicine and Therapeutics • University of Rochester — UL1TR002001: UR Clinical & Translational Science Institute • University of Southern California — UL1TR001855: The Southern California Clinical and Translational Science Institute (SC CTSI) • University of Vermont — U54GM115516: Northern New England Clinical & Translational Research (NNE-CTR) Network • University of Virginia — UL1TR003015: iTHRIV Integrated Translational health Research Institute of Virginia • University of Washington — UL1TR002319: Institute of Translational Health Sciences • University of Wisconsin-Madison — UL1TR002373: UW Institute for Clinical and Translational Research • Vanderbilt University Medical Center — UL1TR002243: Vanderbilt Institute for Clinical and Translational Research • Virginia Commonwealth University — UL1TR002649: C. Kenneth and Dianne Wright Center for Clinical and Translational Research • Wake Forest University Health Sciences — UL1TR001420: Wake Forest Clinical and Translational Science Institute • Washington University in St. Louis — UL1TR002345: Institute of Clinical and Translational Sciences • Weill Medical College of Cornell University — UL1TR002384: Weill Cornell Medicine Clinical and

Translational Science Center • West Virginia University — U54GM104942: West  
 Virginia Clinical and Translational Science Institute (WVCTSI) • Submitted: Icahn  
 School of Medicine at Mount Sinai — UL1TR001433: ConduITS Institute for  
 Translational Sciences • The University of Texas Health Science Center at Tyler —  
 UL1TR003167: Center for Clinical and Translational Sciences (CCTS) • University of  
 California, Davis — UL1TR001860: UC Davis Health Clinical and Translational Science  
 Center • University of California, Irvine — UL1TR001414: The UC Irvine Institute for  
 Clinical and Translational Science (ICTS) • University of California, Los Angeles —  
 UL1TR001881: UCLA Clinical Translational Science Institute • University of California,  
 San Diego — UL1TR001442: Altman Clinical and Translational Research Institute •  
 University of California, San Francisco — UL1TR001872: UCSF Clinical and  
 Translational Science Institute • Pending: Arkansas Children’s Hospital —  
 UL1TR003107: UAMS Translational Research Institute • Baylor College of Medicine —  
 None (Voluntary) • Children’s Hospital of Philadelphia — UL1TR001878: Institute for  
 Translational Medicine and Therapeutics • Cincinnati Children’s Hospital Medical  
 Center — UL1TR001425: Center for Clinical and Translational Science and Training •  
 Emory University — UL1TR002378: Georgia Clinical and Translational Science  
 Alliance • HonorHealth — None (Voluntary) • Loyola University Chicago —  
 UL1TR002389: The Institute for Translational Medicine (ITM) • Medical College of  
 Wisconsin — UL1TR001436: Clinical and Translational Science Institute of Southeast  
 Wisconsin • MedStar Health Research Institute — None (Voluntary) • Georgetown  
 University — UL1TR001409: The Georgetown-Howard Universities Center for Clinical  
 and Translational Science (GHUCCTS) • MetroHealth — None (Voluntary) • Montana  
 State University — U54GM115371: American Indian/Alaska Native CTR • NYU  
 Langone Medical Center — UL1TR001445: Langone Health’s Clinical and Translational  
 Science Institute • Ochsner Medical Center — U54GM104940: Louisiana Clinical and  
 Translational Science (LA CaTS) Center • Regenstrief Institute — UL1TR002529:  
 Indiana Clinical and Translational Science Institute • Sanford Research — None  
 (Voluntary) • Stanford University — UL1TR003142: Spectrum: The Stanford Center  
 for Clinical and Translational Research and Education • The Rockefeller University —  
 UL1TR001866: Center for Clinical and Translational Science • The Scripps Research  
 Institute — UL1TR002550: Scripps Research Translational Institute • University of  
 Florida — UL1TR001427: UF Clinical and Translational Science Institute • University  
 of New Mexico Health Sciences Center — UL1TR001449: University of New Mexico  
 Clinical and Translational Science Center • University of Texas Health Science Center  
 at San Antonio — UL1TR002645: Institute for Integration of Medicine and Science •  
 Yale New Haven Hospital — UL1TR001863: Yale Center for Clinical Investigation.

## References

1. Schou, T., Joca, S., Wegener, G. & Bay-Richter, C. Psychiatric and neuropsychiatric sequelae of COVID-19—A systematic review. *Brain, Behavior, And Immunity*. **97** pp. 328-348 (2021).
2. Huang, C., Huang, L., Wang, Y., Li, X., Ren, L., Gu, X., Kang, L., Guo, L., Liu, M., Zhou, X. & Others 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study. *The Lancet*. **397**, 220-232 (2021).
3. Taquet, M., Geddes, J., Husain, M., Luciano, S. & Harrison, P. 6-month neurological and psychiatric outcomes in 236 379 survivors of COVID-19: a retrospective cohort study using electronic health records. *The Lancet Psychiatry*. **8**, 416-427 (2021).

4. Egbert, A., Cankurtaran, S. & Karpiak, S. Brain abnormalities in COVID-19 acute/subacute phase: a rapid systematic review. *Brain, Behavior, And Immunity*. **89** pp. 543-554 (2020).
5. Douaud, G., Lee, S., Alfaro-Almagro, F., Arthofer, C., Wang, C., McCarthy, P., Lange, F., Andersson, J., Griffanti, L., Duff, E. & Others SARS-CoV-2 is associated with changes in brain structure in UK Biobank. *Nature*. **604**, 697-707 (2022).
6. Kumar, S., Veldhuis, A. & Malhotra, T. Neuropsychiatric and cognitive sequelae of COVID-19. *Frontiers In Psychology*. **12** pp. 577529 (2021).
7. Helms, J., Kremer, S., Merdji, H., Clere-Jehl, R., Schenck, M., Kummerlen, C., Collange, O., Boulay, C., Fafi-Kremer, S., Ohana, M. & Others Neurologic features in severe SARS-CoV-2 infection. *New England Journal Of Medicine*. **382**, 2268-2270 (2020).
8. Kong, X., Zheng, K., Tang, M., Kong, F., Zhou, J., Diao, L., Wu, S., Jiao, P., Su, T. & Dong, Y. Prevalence and factors associated with depression and anxiety of hospitalized patients with COVID-19. *MedRxiv*. pp. 2020-03 (2020).
9. Pavlov VA, Tracey KJ. Neural circuitry and immunity. *Immunologic research*. 2015 Dec;63:38-57.
10. Rahimian R, Wakid M, O'Leary LA, Mechawar N. The emerging tale of microglia in psychiatric disorders. *Neuroscience & Biobehavioral Reviews*. 2021 Dec 1;131:1-29.
11. Dietz AG, Goldman SA, Nedergaard M. Glial cells in schizophrenia: a unified hypothesis. *The Lancet Psychiatry*. 2020 Mar 1;7(3):272-81..
12. Trépanier MO, Hopperton KE, Mizrahi R, Mechawar N, Bazinet RP. Postmortem evidence of cerebral inflammation in schizophrenia: a systematic review. *Molecular psychiatry*. 2016 Aug;21(8):1009-26.
13. Mondelli V, Vernon AC, Turkheimer F, Dazzan P, Pariante CM. Brain microglia in psychiatric disorders. *The Lancet Psychiatry*. 2017 Jul 1;4(7):563-72.
14. Kealy J, Murray C, Griffin EW, Lopez-Rodriguez AB, Healy D, Tortorelli LS, Lowry JP, Watne LO, Cunningham C. Acute inflammation alters brain energy metabolism in mice and humans: role in suppressed spontaneous activity, impaired cognition, and delirium. *Journal of Neuroscience*. 2020 Jul 15;40(29):5681-96.
15. Girard TD, Ware LB, Bernard GR, Pandharipande PP, Thompson JL, Shintani AK, Jackson JC, Dittus RS, Ely EW. Associations of markers of inflammation and coagulation with delirium during critical illness. *Intensive care medicine*. 2012 Dec;38:1965-73.
16. McNeil JB, Hughes CG, Girard T, Ware LB, Ely EW, Chandrasekhar R, Han JH. Plasma biomarkers of inflammation, coagulation, and brain injury as predictors of delirium duration in older hospitalized patients. *PLoS One*. 2019 Dec 19;14(12):e0226412.
17. Lozano-Vicario L, García-Hermoso A, Cedeno-Veloz BA, Fernández-Irigoyen J, Santamaría E, Romero-Ortuno R, Zambom-Ferraresi F, Sáez de Asteasu ML, Muñoz-Vázquez ÁJ, Izquierdo M, Martínez-Velilla N. Biomarkers of delirium risk



- in older adults: a systematic review and meta-analysis. *Frontiers in Aging Neuroscience*. 2023 May 12;15:1174644.
18. Fong TG, Inouye SK. The inter-relationship between delirium and dementia: the importance of delirium prevention. *Nature Reviews Neurology*. 2022 Oct;18(10):579-96.
  19. Culley DJ, Flaherty D, Fahey MC, Rudolph JL, Javedan H, Huang CC, Wright J, Bader AM, Hyman BT, Blacker D, Crosby G. Poor performance on a preoperative cognitive screening test predicts postoperative complications in older orthopedic surgical patients. *Anesthesiology*. 2017 Nov 1;127(5):765-74.
  20. Davis DH, Skelly DT, Murray C, Hennessy E, Bowen J, Norton S, Brayne C, Rahkonen T, Sulkava R, Sanderson DJ, Rawlins JN. Worsening cognitive impairment and neurodegenerative pathology progressively increase risk for delirium. *The American journal of geriatric psychiatry*. 2015 Apr 1;23(4):403-15.
  21. Alexander SA, Conley YP, Girard TD, McVerry BJ, Ren D. Temporal Variability in Inflammatory Gene Methylation and Delirium in Critically Ill Patients. *American Journal of Critical Care*. 2022 Sep 1;31(5):367-74.
  22. Widmann CN, Heneka MT. Long-term cerebral consequences of sepsis. *The Lancet Neurology*. 2014 Jun 1;13(6):630-6.
  23. Rengel KF, Hayhurst CJ, Pandharipande PP, Hughes CG. Long-term cognitive and functional impairments after critical illness. *Anesthesia & Analgesia*. 2019 Apr 1;128(4):772-80.
  24. Jablensky A. Prevalence and incidence of schizophrenia spectrum disorders: implications for prevention. *Australian & New Zealand Journal of Psychiatry*. 2000 Feb;34(1 suppl):A26-34.
  25. Yirmiya R, Rimmerman N, Reshef R. Depression as a microglial disease. *Trends in neurosciences*. 2015 Oct 1;38(10):637-58.
  26. McAllen RM, McKinley MJ, Martelli D. Reflex regulation of systemic inflammation by the autonomic nervous system. *Autonomic Neuroscience*. 2022 Jan 1;237:102926.
  27. Cerejeira J, Nogueira V, Luís P, Vaz-Serra A, Mukaetova-Ladinska EB. The cholinergic system and inflammation: common pathways in delirium pathophysiology. *Journal of the American Geriatrics Society*. 2012 Apr;60(4):669-75.
  28. Salter MW, Stevens B. Microglia emerge as central players in brain disease. *Nature medicine*. 2017 Sep 1;23(9):1018-27.
  29. Kealy J, Murray C, Griffin EW, Lopez-Rodriguez AB, Healy D, Tortorelli LS, Lowry JP, Watne LO, Cunningham C. Acute inflammation alters brain energy metabolism in mice and humans: role in suppressed spontaneous activity, impaired cognition, and delirium. *Journal of Neuroscience*. 2020 Jul 15;40(29):5681-96.
  30. Smith RJ, Rabinstein AA, Cartin-Ceba R, Singh VP, Lachner C, Khatua B, Trivedi S, Gajic O. Chemokines in ICU Delirium: An Exploratory Study. *Critical Care Explorations*. 2022 Jul;4(7).



31. Schreuder L, Eggen BJ, Biber K, Schoemaker RG, Laman JD, de Rooij SE. Pathophysiological and behavioral effects of systemic inflammation in aged and diseased rodents with relevance to delirium: a systematic review. *Brain, Behavior, and Immunity*. 2017 May 1;62:362-81.
32. Knaak C, Vorderwülbecke G, Spies C, Piper SK, Hadzidiakos D, Borchers F, Brockhaus WR, Radtke FM, Lachmann G. C-reactive protein for risk prediction of post-operative delirium and post-operative neurocognitive disorder. *Acta Anaesthesiologica Scandinavica*. 2019 Nov;63(10):1282-9.
33. Brummel NE, Hughes CG, Thompson JL, Jackson JC, Pandharipande P, McNeil JB, Raman R, Orun OM, Ware LB, Bernard GR, Ely EW. Inflammation and coagulation during critical illness and long-term cognitive impairment and disability. *American journal of respiratory and critical care medicine*. 2021 Mar 15;203(6):699-706.
34. McNeil JB, Hughes CG, Girard T, Ware LB, Ely EW, Chandrasekhar R, Han JH. Plasma biomarkers of inflammation, coagulation, and brain injury as predictors of delirium duration in older hospitalized patients. *PLoS One*. 2019 Dec 19;14(12):e0226412.
35. Shanazz K, Nalloor R, Lucas R, Vazdarjanova A. Neuroinflammation is a susceptibility factor in developing a PTSD-like phenotype. *Frontiers in Behavioral Neuroscience*. 2023 Mar 29;17:1112837.
36. Speer K, Upton D, Semple S, McKune A. Systemic low-grade inflammation in post-traumatic stress disorder: a systematic review. *Journal of inflammation research*. 2018 Mar 22:111-21.
37. Zuo C, Zhuang Z, Yang P, Zhang H, Li X, Huang T, Ahluwalia TS. Dissecting the causal association between inflammation and post-traumatic stress disorder: A bidirectional Mendelian randomization study. *Journal of Affective Disorders*. 2023 Jul 15;333:436-45.
38. Herridge MS, Azoulay É. Outcomes after critical illness. *New England Journal of Medicine*. 2023 Mar 9;388(10):913-24.
39. Stein SR, Ramelli SC, Grazioli A, Chung JY, Singh M, Yinda CK, Winkler CW, Sun J, Dickey JM, Ylaya K, Ko SH. SARS-CoV-2 infection and persistence in the human body and brain at autopsy. *Nature*. 2022 Dec 22;612(7941):758-63.
40. Albornoz EA, Amarilla AA, Modhiran N, Parker S, Li XX, Wijesundara DK, Aguado J, Zamora AP, McMillan CL, Liang B, Peng NY. SARS-CoV-2 drives NLRP3 inflammasome activation in human microglia through spike protein. *Molecular psychiatry*. 2022 Nov 1:1-6.
41. Steardo Jr L, Steardo L, Verkhatsky A. Psychiatric face of COVID-19. *Translational psychiatry*. 2020 Jul 30;10(1):261.
42. Xie Y, Xu E, Al-Aly Z. Risks of mental health outcomes in people with covid-19: cohort study. *bmj*. 2022 Feb 16;376.
43. Coleman B, Casiraghi E, Blau H, Chan L, Haendel MA, Laraway B, Callahan TJ, Deer RR, Wilkins KJ, Reese J, Robinson PN. Risk of new-onset psychiatric sequelae of COVID-19 in the early and late post-acute phase. *World Psychiatry*. 2022 Jun;21(2):319.

44. Schoenfeld, D. Partial residuals for the proportional hazards regression model. *Biometrika*. **69**, 239-241 (1982).
45. Agresti A. Categorical data analysis. John Wiley & Sons; 2012 Dec 3.
46. Li B, Babu GJ. A graduate course on statistical inference. Springer New York; 2019 Aug 2.
47. Hammond F, Malec J, Nick T, Buschbacher R, editors. Handbook for clinical research: design, statistics, and implementation. Demos Medical Publishing; 2014 Aug 26.
48. Ngasa SN, Tchouda LA, Abanda C, Ngasa NC, Sanji EW, Dingana TN, Babila CS. Prevalence and factors associated with anxiety and depression amongst hospitalised COVID-19 patients in Laquintinie Hospital Douala, Cameroon. *PLoS One*. 2021 Dec 2;16(12):e0260819.
49. Mazza MG, Palladini M, De Lorenzo R, Magnaghi C, Poletti S, Furlan R, Ciceri F, Rovere-Querini P, Benedetti F, COVID-19 BioB Outpatient Clinic Study group. Persistent psychopathology and neurocognitive impairment in COVID-19 survivors: effect of inflammatory biomarkers at three-month follow-up. *Brain, behavior, and immunity*. 2021 May 1;94:138-47.
50. N3C computable phenotype: COVID-19 Phenotype Documentation, Version 4.0. [https://github.com/National-COVID-Cohort-Collaborative/Phenotype\\_Data\\_Acquisition/wiki/Latest-Phenotype](https://github.com/National-COVID-Cohort-Collaborative/Phenotype_Data_Acquisition/wiki/Latest-Phenotype). Updated March 11, 2022.9.
51. OHDSI Standardized Data. <https://www.ohdsi.org/data-standardization/>. Accessed November 10, 2022.
52. Kempf, L., Hussain, N. & Potash, J. Mood disorder with psychotic features, schizoaffective disorder, and schizophrenia with mood features: trouble at the borders. *International Review Of Psychiatry*. **17**, 9-19 (2005).
53. Lewine, R. & Hart, M. Schizophrenia spectrum and other psychotic disorders. *Handbook Of Clinical Neurology*. **175** pp. 315-333 (2020).

## A Appendix A: Tables in Appendix

Table 4. SSPD Patient Comparison Among ARDS, COVID Positive and COVID Negative Group.

	ARDS Patient N = 637		COVID Positive Patient N =1229		COVID Negative Patient = 707		P Value
	Count	Percentage	Count	Percentage	Count	Percentage	
<b>Gender:</b>							
Female	222	35%	518	42%	244	35%	<2.2e-16
Male	415	65%	711	58%	463	65%	<2.2e-16
<b>Age Groups:</b>							
21 and Younger	637	Total	1229	Total	707	Total	
#	#	#	62	5.04%	28	3.96%	4.91E-8
22 to 29	<65	<10.20%	112	9.11%	73	10.33%	0.0001715
30 to 39	94	14.76%	181	14.73%	116	16.41%	1.52E-07
40 to 49	113	17.74%	170	13.83%	112	15.84%	0.0002312
50 to 59	164	25.75%	263	21.40%	156	22.07%	1.15E-08
60 and Older	190	29.83%	441	35.88%	222	31.40%	<2.2e-16
<b>Race:</b>							
Black/African American	637	Total	1229	Total	707	Total	
Asian and White/Caucasian	246	38.62%	361	29.37%	279	39.46%	6.98E-06
	297	46.62%	711	57.85%	319	45.12%	<2.2e-16
<b>Ethnicity:</b>							
Hispanic or Latino	73	11.46%	118	9.60%	88	12.45%	0.003535
Not Hispanic or Latino	507	79.59%	1015	82.59%	571	80.76%	<2.2e-16

The (#) marked group has a count or proportion that is too small to quantitatively report

Table 5. SSPD Patient Comparison Within COVID Positive Group.

Characteristics	COVID Positive				
	All N = 219264	Only SSPD Patient = 1229	Percentage	Adjusted Value	Adjusted Percentage
	Count	Count		Adjusted Count	centage
<b>Gender:</b>					
Female	103528	518	47.20%	1097.457627	42%
Male*	115736	711	52.80%	1346.590909	58%
<b>Age Groups:</b>				2444.048536	<b>Total</b>
21 and Younger*	7449	62	3.40%	1823.529412	23.15%
22 to 29	19160	112	8.74%	1281.464531	16.27%
30 to 39	30298	181	13.80%	1311.594203	16.65%
40 to 49	34407	170	15.70%	1082.802548	13.75%
50 to 59	53059	263	24.20%	1086.77686	13.80%
60 and Older	74891	441	34.20%	1289.473684	16.37%
<b>Race:</b>				7875.641237	<b>Total</b>
Black/African American*	47047	361	21.46%	1682.451676	58.69%
Asian and White/Caucasian	131641	711	60.04%	1184.256455	41.31%
<b>Ethnicity:</b>				2866.708131	<b>Total</b>
Hispanic or Latino	38196	118	17.40%	678.1609195	33.47%
Not Hispanic or Latino*	165071	1015	75.30%	1347.941567	66.53%
				2026.102487	<b>Total</b>

The (\*) marked group has a significantly higher likelihood of having SSPD.

**Table 6.** SSPD Patient Comparison Within COVID Negative Group.

Characteristics	COVID Negative				
	All N = 213183		Only SSPD Patient = 707		Adjusted Value
	Count	Percentage	Count	Percentage	Adjusted Count
<b>Gender:</b>					
Female	105959	49.70%	244	35%	490.945674
Male*	107196	50.30%	463	65%	920.4771372
<b>Age Groups:</b>			707	<b>Total</b>	<b>Total</b>
21 and Younger	7840	3.68%	28	3.96%	760.8695652
22 to 29	19644	9.21%	73	10.33%	792.616721
30 to 39	29400	13.80%	116	16.41%	840.5797101
40 to 49	31594	14.80%	112	15.84%	756.7567568
50 to 59	51115	24.00%	156	22.07%	650
60 and Older	73590	34.50%	222	31.40%	643.4782609
<b>Race:</b>			707	<b>Total</b>	<b>Total</b>
Black/ African American*	45269	21.20%	279	39.46%	1316.037736
Asian and White/ Caucasian	133309	62.53%	319	45.12%	510.1334269
<b>Ethnicity:</b>					1826.171163
Hispanic or Latino	31002	14.50%	88	12.45%	606.8965517
Not Hispanic or Latino*	166149	77.90%	571	80.76%	732.9910141
					1339.887566

The (\*) marked group has a significantly higher likelihood of having SSPD.

Table 7. SSPD Patient Comparison Within ARDS Group.


Characteristics	ARDS					
	All N = 222337		Only SSPD Patient = 637		Adjusted Value	
	Count	Percentage	Count	Percentage	Adjusted Count	Adjusted Percentage
<b>Gender:</b>						
Female	105175	47.30%	222	35%	469.3446089	37%
Male*	117162	52.70%	415	65%	787.4762808	63%
<b>Age Groups:</b>			637	<b>Total</b>	1256.82089	<b>Total</b>
21 and Younger	7896	3.55%	#	#	422.5352113	11.31%
22 to 29	19885	8.94%	<65	< 10.20%	682.3266219	18.27%
30 to 39	31070	14.00%	94	14.76%	671.4285714	17.98%
40 to 49	35871	16.10%	113	17.74%	701.863354	18.79%
50 to 59	52529	23.60%	164	25.75%	694.9152542	18.60%
60 and Older	75086	33.80%	190	29.83%	562.1301775	15.05%
<b>Race:</b>			637	<b>Total</b>	3735.19919	<b>Total</b>
Black/African American*	48251	21.70%	246	38.62%	1133.640553	70.38%
Asian and White/Caucasian	138433	62.26%	297	46.62%	477.0111823	29.62%
<b>Ethnicity:</b>					1610.651735	<b>Total</b>
Hispanic or Latino	34320	15.40%	73	11.46%	474.025974	41.92%
Not Hispanic or Latino*	171609	77.20%	507	79.59%	656.7357513	58.08%
					1130.761725	<b>Total</b>

The (\*) marked group has a significantly higher likelihood of having SSPD.  
The (#) marked group has a count or proportion that is too small to quantitatively report.



## B Appendix B: Supporting Information

- S1 Table** Schizophrenia Spectrum and Psychotic Disorders Code Set: list of diagnosis codes used to evaluate outcomes
- S2 Table** Exclusion Code Set: list of diagnosis codes used to exclude patients if one or more codes were included in the patient record prior to the index date.
- S3 Table** Matching Criteria Code Set: diagnosis and drug exposure codes used in the R MatchIt package to build case and control cohorts.



## Hazard Ratios For Different Time Intervals

