Trait responsiveness to verbal suggestions predicts nocebo responding: A meta-analysis

Madeline V. Stein [a], Monika Heller [b], Sarah Chapman [c], James Rubin [d], & Devin B. Terhune [a]

[a] *Department of Psychology, Institute of Psychology, Psychiatry, and Neuroscience, King's College London, London, UK*

[b] *Centre for Behavioural Medicine, UCL School of Pharmacy, London, UK*

[c] *Institute of Pharmaceutical Science, King's College London, London, UK*

[d] *Department of Psychological Medicine, Institute of Psychology, Psychiatry, and Neuroscience, King's College London, London, UK*

Correspondence address:

Madeline V. Stein

madeline.stein@kcl.ac.uk

Department of Psychology

Institute of Psychiatry, Psychology & Neuroscience

King's College London

16 De Crespigny Park

London SE5 8AB, UK

Abstract

Nocebo responding involves the experience of adverse health outcomes in response to contextual cues. These deleterious responses impact numerous features of mental and physical health but are characterized by pronounced heterogeneity. Suggestion is widely recognised as a contributing factor to nocebo responding but the moderating role of trait responsiveness to verbal suggestions (suggestibility) in nocebo responding remains poorly understood. We conducted a pre-registered meta-analysis (PROSPERO registration number CRD42023425605) to quantitatively synthesize available research on the relationship between suggestibility and nocebo responding. Four electronic databases were searched for original studies involving both the assessment of suggestibility and symptom reports in response to an inactive stimulus. Of 7,729 search results, 10 articles presenting 13 correlations between suggestibility and nocebo responding were analysed. A random-effects meta-analysis revealed a significant, albeit weak, positive correlation, r=0.21 [95% CI: 0.04, 0.37], between suggestibility and nocebo responses, such that more highly suggestible individuals displayed larger responses. Sensitivity and meta-regression analyses demonstrated that studies of higher methodological quality, including those that maintained experimenter blinding, exhibited stronger effect sizes. These results corroborate proposals that trait responsiveness to verbal suggestions confers greater response to nocebos and warrants renewed attention to the role of suggestibility in symptom induction and perception.

Keywords: nocebo, suggestibility, suggestion, meta-analysis, expectation, patient–clinician communication

Introduction

Nocebo responding is a heterogenous phenomenon involving symptom induction or exacerbation

from contextual cues, such as conditioning and/or verbal suggestions, and it can occur in a diverse

array of clinical, experimental, and social contexts (Colloca & Barsky, 2020; Petrovic, 2008). Nocebo

responding prototypically presents as side effects or adverse events in response to medications (Heller

et al., 2022), inert treatments (placebos; (Bender et al., 2023) and vaccines (Haas et al., 2022). The

development of symptom expectations can occur in response to verbal or written information about

potential side effects during informed consent processes (Spiegel, 1997; Wells & Kaptchuk, 2012)

and patient education (Colloca & Miller, 2011; Hauser et al., 2012), and/or after observing another

person displaying symptoms (Meeuwis et al., 2023). The effects of nocebo responding extend beyond

the induction or exacerbation of specific symptoms. Nocebo effects can increase patient distress

(Nasiri-Dehsorkhi et al., 2023), hinder treatment adherence, lead to the premature discontinuation of

treatments and skew trial outcomes (Ciaramella et al., 2013; Colloca & Miller, 2011).

Responsiveness to nocebos is characterised by pronounced heterogeneity and it remains

controversial whether nocebo responding is a stable trait-like variable (Kern et al., 2020; Rooney et

al., 2022). Verbal suggestion, a communication for an involuntary change in experience or behaviour

(e.g., "this might hurt a bit"), has been consistently demonstrated to be efficacious in the induction of

nocebo responses (Petersen et al., 2014; Rooney et al., 2023; Webster et al., 2016). For example,

meta-analytic evidence indicates that verbal suggestions seem to induce a nocebo response of

moderate magnitude, which surpasses that of conditioning alone but falls short of the effect size

produced by the joint application of both suggestion and conditioning (Rooney et al., 2023). Nocebo

induction methods typically include direct verbal suggestions (e.g., "this procedure will increase your

experience of pain") or indirect verbal suggestions, which are more implicit (e.g., "some people might

experience pain") (Oakley et al., 2021; Polczyk, 2016). Trait responsiveness to direct verbal

suggestions (direct verbal suggestibility) is highly stable over time (Piccione et al., 1989) and is a

reliable predictor of response to hypoalgesia suggestions in both clinical and experimental contexts

(Milling et al., 2021; Thompson et al., 2019). As such, the capacity to respond to verbal suggestions

represents a promising candidate predictor of nocebo responding. Moreover, elevated suggestibility is a hallmark feature of a range of phenomena germane to nocebo responding, such as the experience of symptoms attributed to environmental factors (idiopathic environmental intolerance) (Stein et al., 2023), functional neurological disorder (Bell et al., 2011; Fiorio et al., 2022; Wieder et al., 2021), and mass psychogenic illness (Sapkota et al., 2020). Further indirect evidence for a link between suggestibility and nocebo responding comes from evidence showing that both trait empathy (Meeuwis et al., 2023) and absorption (Brascher et al., 2020) predict nocebo responding, and both have been found to correlate with suggestibility (Wickramasekera & Szlyk, 2003). Within the predictive processing framework (Hohwy, 2020), nocebo responses can be attributed to the formation of precise symptom priors (e.g., beliefs and expectations) that are overweighted relative to sensory information, yielding symptoms (Fiorio et al., 2022; Van den Bergh et al., 2017). By conferring greater responsiveness to suggestion, higher trait suggestibility may predispose individuals to be more likely to form strong or precise symptom priors and/or to be less likely to update these priors on the basis of sensory evidence (stubborn predictions). Although multiple studies have provided evidence of a positive link between suggestibility and nocebo responding (Corsi & Colloca, 2017; Khan et al., 2009; Winter & Braw, 2022), this effect hasn't been replicated in others (Zech et al., 2022; Zech et al., 2020). Previous reviews and meta-analyses have largely neglected the role of suggestibility in nocebo responding and, to our knowledge, there has not yet been an attempt to formally quantify the magnitude of this relationship.

We conducted a pre-registered meta-analysis in order to quantitatively synthesize the extant literature on the association between suggestibility and nocebo responding. Toward this end, we systematically integrated research studies examining the correlation between psychometric suggestibility measures and symptom outcomes in response to an inactive (sham) procedure. We elected to omit studies with active treatments to diminish the potential confounding effect of an active treatment. In addition to quantifying the correlation between suggestibility and the magnitude of nocebo responses, we also examined whether these correlations were moderated by different measurement variables including sample type (e.g., clinical or healthy sample), nocebo suggestion

type (e.g., direct or indirect verbal suggestion), target symptom domain, and various methodological features.

Methods

We completed this meta-analysis in accordance with Meta-analyses Of Observational Studies in Epidemiology (MOOSE) (Brooke et al., 2021) and Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (Page et al., 2021). We prospectively registered this meta-analysis on PROSPERO (registration number: CRD42023425605).

Inclusion criteria

The data subjected to analyses met the following criteria: published in English; published in academic peer-reviewed journal; use of at least one experimental manipulation for symptom induction (e.g., verbal suggestion); use of a procedure that involved either an inactive intervention or suggestion alone that was reported as a nocebo with no intervention; and inclusion of a symptom measure. We adopted an inclusive approach when defining aversive outcomes and include outcomes from placebo studies that were not uniformly aversive (e.g., intoxication). Finally, we included papers that did not include a control condition, as we anticipated the lack of a control comparator to be a limitation of many potentially relevant studies. We had broad inclusion criteria in order to obtain an accurate representation of the experimental nocebo literature.

Search strategy

PsycInfo, Embase, MEDLINE, and PubMed were searched in December 2022. The search was repeated in October 2023 and September 2024 and yielded no new articles to include. The search string (see Supplementary Materials) was augmented with manual searches of all included articles and relevant reviews. One author group was contacted to confirm that all relevant articles from their group had been captured by the initial search.

Study selection

Two reviewers (MVS and MH) independently screened titles and abstracts of all retrieved articles as part of another review. One reviewer (MVS) set aside articles that met the current review's inclusion criteria and confirmed study selection with the second reviewer (MH). Three author groups were contacted for additional data in order to be included in the analyses and were subsequently included.

Data extraction

Data extraction was performed independently by the same two reviewers on full texts. A comprehensive list of all extracted data can be found in the supplementary methods. The primary outcomes were correlation coefficient(s) between a suggestibility scale and nocebo response (operationalised as nocebo condition alone [no control], nocebo-control, or nocebo-baseline). When multiple correlations from a single suggestibility scale with subscales were reported, we only extracted the correlation for the total scale score. Additionally, studies were coded for the following criteria: suggestibility scale administration context (i.e. hypnosis-related context or not); suggestibility scale administration type (live vs. recording); scale type (direct, indirect, or not applicable); scale administration context (group vs. individual); and good psychometric properties for the suggestibility scale (yes vs. no), based on relevant reliability and validity data presented in the paper. The reviewers displayed good agreement (86%); any discrepancies were resolved through discussion with DBT.

Study methodological quality

A 15-item scale was developed to assess study methodological quality (see Supplementary Methods). Items were adapted from a previous measure (Thompson et al., 2019; Wieder et al., 2021) and included items based on Cochrane criteria and PRISMA recommendations. The two reviewers independently rated each item categorically (0=criterion not met, 1=criterion met), with any discrepancies resolved through discussion with DBT, and a total score (percentage of criteria met) was computed for each study. Agreement between reviewers was good (percentage agreement: 82%; Cohen's kappa=.62).

Data synthesis

All analyses were performed using Jamovi (The jamovi project, 2023). Individual study effect sizes included unadjusted correlation coefficients between suggestibility scale scores and nocebo responses that were transformed to *z*-scores using Fisher's *r*-to-*z* transformation (Z_r) and analysed using random-effects meta-analysis with the DerSimonian-Laird method. Alongside aggregate Z_r values, we report raw correlation coefficients for ease of interpretation. Heterogeneity was computed through I^2 and τ^2, where I^2 >50% indicates moderate or greater heterogeneity and τ^2 describes the variance of true effect sizes, with values >10% indicating greater heterogeneity. We assessed publication bias by examining funnel plots of effect sizes against standard errors for asymmetry and tested for asymmetry using Egger's bias test (Egger et al., 1997), where *p*<.10 is suggestive of asymmetry. We also estimated asymmetry-corrected cumulative effect sizes using the trim-and-fill method (Duval & Tweedie, 2000).

In order to account for heterogeneity in effect sizes, we performed subgroup analyses and a series of meta-regression analyses using binary and continuous moderators when there was a minimum of two studies per moderator level and ten studies total, respectively. Moderator variables with >2 levels were decomposed into simpler 2-level moderators. Moderators included 18 categorical variables (0=absent/no, 1=present/yes unless otherwise specified): study design (0=between-groups 1=design within-subjects), blinding of participant and/or experimenter to experimental condition, nocebo suggestion target symptom domain (pain, dyspnoea, dizziness, nausea, motor inhibition, cognitive function; each symptom coded as 0=absent vs. 1=present, e.g. no pain vs. pain), use of conditioning procedure, suggestion administration type (verbal, video, non-verbal), suggestibility scale type (0=indirect, 1=direct), nocebo administration procedure (pill, cream, procedure, nasal spray, inhalation, injection; each administration procedure coded as 0=absent vs. 1=present, e.g. no pill vs. pill), control comparator (control condition, baseline, or none), presence of hypnotic context, scale administration context (0= individual, 1=group); and one continuous variable: methodological quality percentage score.

Results

Study inclusion

A PRISMA diagram presenting study selection can be found in Fig. 1. The final sample of 10

included papers reported 13 suggestibility-nocebo response correlation pairs with three papers

reporting two correlations. Details of the included studies can be found in Table 1 and a full list of

included papers can be found in the Supplementary Results.

Figure 1. *PRISMA flowchart of the study selection process.*

1 **Table 1.** *Principal features of included papers reporting associations between suggestibility and nocebo responding (n=10).*

Source	Country	Study design	Sample	N (% female)	Age M (SD)	Symptom domain	Symptom outcome measure	Experimental condition	Control condition	Suggestibility scale
Corsi & Colloca, 2017	US	Within	Healthy volunteers	46 (52%)	27.41 (1.07)	Pain	VAS pain	Verbal suggestion; conditioning procedure	Condition: conditioned control cue	MISS
Di Stefano et al., 2022	-	Between	GI patients	40 (87%)	39 (16)	Pain, dyspnoea, nausea	VAS GI symptom scale	Verbal suggestion; sham lactose test	Condition; sham lactose test	MISS
Khan et al., 2009	-	Between	Patients experiencing seizures	47 (58%)	-	Seizure	Symptom induction	Verbal suggestion	-	HIP
Leigh et al., 2003	CA	Within	Asthma patients	17 (64%)	46.8 (14.25)	Dyspnoea	Borg Dyspnoea Scale	Verbal suggestion; saline inhalation	Baseline	CIS
Sharav et al., 2023	-	Mixed	Healthy volunteers	24 (62%)	26.65 (-)	Pain	VAS pain	Verbal suggestion; conditioning	-	SHALIT
Winter & Braw, 2021	IL	Between	Recovered COVID-19 patients	90 (75%)	Nocebo: 38.60 (1.61) Control: 39.84 (1.76)	Cognitive performance	Cognitive Failures Questionnaire	Textual suggestion	Control group; textual suggestion	SSS
Woody et al., 1997*	CA	Between	Undergraduate students	93 (44%)	20.5 (-)	Dizziness, nausea, motor inhibition, cognitive function	Subjective experience measure; Unsuggested Symptom Checklist	Verbal suggestion; sham alcohol	-	HGSHS:A
Zech et al., 2019*	DE	Within	Healthy, non-medical professionals	46 (54%)	34.3 (15.2)	Motor performance	Maximum arm muscle strength (% decrease from baseline)	Verbal suggestion	Baseline	HGSHS:A
Zech et al., 2020*	DE	Within	Elective surgery patients	45 (55%)	43.8 (15)	Motor performance	Maximum arm muscle strength (% decrease from baseline)	Verbal suggestion	Baseline	HGSHS-5
Zech et al., 2022	DE	Mixed	Healthy volunteers	50 (58%)	29.1 (12.7)	Motor performance	Spirometry	Verbal suggestion	Baseline	HGSHS-5

Notes: *= reports 2 correlation coefficients with no variation in principle features; - = Not reported; Within = within-groups design; Between = between-groups design; Mixed = mixed experimental design; US= United States; CA= Canada; IL= Israel; DE=Germany; CIS = *Creative Imagination Scale*; GI = Gastro-Intestinal; HGSHS:A = *Harvard Group Scale of Hypnotic Susceptibility: Form A*; HGSHS-5 = *Harvard Group Scale of Hypnotic Susceptibility-5 item*; HIP = *Hypnotic Induction Profile*; MISS = *Multidimensional Iowa Suggestibility Scale*; SHALIT = *Six-minute Arm Levitation Test*; SSS = *Short Suggestibility Scale*; VAS = Visual Analogue Scale

Methodological quality criteria

Methodological quality scores varied across the sample of included papers (*n*=10), with most tending

to meet around half of the criteria (*M%±SD*: 61±13; range: 40-80; see Supplementary Table 1). There

was no variability in methodological scores and ratings across studies extracted from a single paper.

All studies used a reliable and valid suggestibility scale, and the majority (90%; *n*=9) included a

referenced symptom measure. However, a little more than half of the included papers (60%; *n*=6)

ensured appropriate experimenter blinding to participants' suggestibility level or nocebo response,

whereas 50% (*n*=5) of the papers ensured participants were adequately blinded.

Meta-analysis of bivariate correlations between nocebo response and suggestibility

A random-effects meta-analysis of correlation coefficients (Fisher's *r*-to-*Z* [Z_r]) between symptom

outcomes in response to an inactive (sham) procedure and suggestibility scale scores (*k*=13; see

Figure 2) revealed a significant association, Z_r=0.22 [95% CI: 0.04, 0.38], *Z*=2.48, *p*=.013,

corresponding to a weak correlation, *r*=.21 [0.04, 0.37]. As can be seen in Figure 2, there was

considerably heterogeneity in effect sizes, I^2=78%, τ^2=.07, with correlations ranging from Z_r=-0.32 to

Z_r=0.73 (*r* range: -0.31 to 0.62). A Jackknife analysis, in which each correlation pair was sequentially

omitted and the analysis was reperformed, confirmed the aggregate effect size varied in the weak

range (Z_r range: 0.13-0.26). The cumulative effect remaining statistically significant in all of the 13

iterations, indicating that it was not driven by a single study. These results demonstrate that

suggestibility is a significant, reliable, predictor of nocebo responding.

Given the relatively low average methodological quality score (~60%), and the high level of

heterogeneity in effect sizes, we next performed a series of sensitivity analyses to determine whether

the aggregate effect size would be stable in more methodologically robust studies. Toward this end,

we repeated the central analyses on subgroups of studies with superior methodological features; we

only conducted these subgroup analyses when three or more studies could be included, resulting in

four analyses. First, we restricted the analysis to studies with higher methodological quality scores

(rating>60%; k=7), the majority of the studies included in this subgroup (k=6) were also well-powered to detect moderate-to-large effect sizes ($n \geq 46$). This resulted in a larger overall effect size, Z_r=0.41 [0.23, 0.59], Z=4.54, p<.001, equivalent to r=.39, which fell outside the 95% CIs of the original correlation, with a modest reduction in heterogeneity, I^2=67% and τ^2=.03. Similarly, studies that implemented participant condition blinding and experimenter blinding of condition, suggestibility level, and nocebo response (k=7), demonstrated a larger, and highly significant, effect size, Z_r=0.37 [0.16, 0.58], Z=3.46, p<.001, equivalent to r=.35, with only a minor decrease in heterogeneity, I^2=76% and τ^2=.05. By contrast, studies that used a measure of direct verbal suggestibility (k=10) demonstrated a comparable, significant effect to the full sample, Z_r=0.20 [0.01, 0.38], Z=2.09, p=.036, equivalent to r=.19, with a minor decrease in heterogeneity, I^2=74% and τ^2=.06. In a similar manner, studies that included a control condition (k=9) exhibited a weak effect size that was comparable to that of the total sample, albeit non-significant, Z_r=0.19 [-0.03, 0.41], Z=1.72, p=.085, equivalent to r=.18, with no reduction in heterogeneity, I^2=78% and τ^2=.08. Similarly, studies that used behavioural/physiological outcomes (k=6), yielded an effect size comparable to the overall main effect, Z_r=0.19 [-0.08, 0.46], Z=1.36, p=.17, with moderate heterogeneity, I^2=80%, τ^2=.09. Cumulatively, these analyses indicate that the utility of suggestibility as a predictor of nocebo responding is relatively stable when the data are restricted to methodologically superior studies, and indeed is substantially greater among the most methodologically rigorous studies.

Figure 2. *Forest plot of correlation coefficients (Z$_r$) [95% CIs] between suggestibility and nocebo responses (k=13).*

Notes. Marker sizes reflect study weights, with smaller and larger markers denoting smaller and larger weights, respectively.

Publication bias

Egger's test was used to assess publication bias and did not reveal significant funnel plot asymmetry, Z=-0.73, p=.46, in the total sample (Figure 3), suggesting that there was no clear evidence for publication bias. No values were calculated using the trim-and-fill method; therefore, none were applied.

Figure 3. *Funnel plot of nocebo-suggestibility correlation coefficients (Z_r) as a function of standard error (k=13). Markers denote individual study effect sizes.*

Meta-regression analyses

Owing to the moderate heterogeneity in effect sizes in the total sample and subgroups, we undertook a series of meta-regression analyses to identify the variables that moderated the magnitude of these associations (see Table 2 and Supplementary Table 3). We first considered the potential moderating effects of individual methodological quality criteria. Due to our inclusion criterion of at least two studies per moderator level (e.g., unblinded v. blinded experimenter), seven individual methodological quality items were eligible for analysis and were included as binary moderators of correlation coefficients (Table 2). Among these criteria, the only significant moderator of effect sizes was experimenter blinding to the suggestibility level or nocebo response of the participants. In particular, experimenter blinding was a highly significant, positive moderator, such that studies that ensured that the experimenter was blinded exhibited larger correlations between suggestibility and

nocebo responding. Consistent with this result, the overall methodological quality score was also a

highly significant continuous moderator; this effect was also positive in direction, although the effect

size was weak in magnitude. Both results align with the preceding results of our sensitivity analyses.

Taken together these results suggest that studies that were of higher methodological quality, in

particular those that implemented experimenter blinding, tended to display a higher correlation

between nocebo responding and suggestibility.

Table 2. *Meta-regression analyses on correlation coefficients between suggestibility and nocebo responding (k=13).*

Moderator	(ΔZ_r) [95% CIs]	Z	p	I^2
Item 1: Reliability data presented in paper	0.22 [-0.17, 0.61]	1.08	.28	77%
Item 3: Clear sample recruitment site	0.23 [-0.18, 0.65]	1.11	.26	78%
Item 4: Clear participant recruitment	0.16 [-0.31, 0.65]	0.68	.49	79%
Item 8: Clear protocol reporting	0.31 [-0.15, 0.76]	1.33	.18	76%
Item 9: Experimenter blinding to suggestibility or nocebo response	0.45 [0.18, 0.71]	3.35	<.001	61%
Item 10: Participant blinding	0.19 [-0.07, 0.47]	1.44	.15	78%
Item 11: Experimenter condition blinding	0.14 [-0.38, 0.66]	0.53	.59	78%
Total methodological quality score	0.02 [0.01, 0.03]	3.36	<.001	61%
Pain as symptom outcome	-0.32 [-0.73, 0.07]	-1.58	.11	77%
Use of conditioning	-0.22 [-0.72, 0.28]	-0.85	.39	79%
Use of control comparator	-0.07 [-0.46, 0.31]	-0.38	.69	79%
Use of direct symptom suggestion	0.04 [-0.49, 0.41]	-0.19	.85	80%

Notes: ΔZ_r= Fisher's r-to-z correlation difference; all moderators were binary except Total methodological quality score; item refers to the methodological quality criteria (see Supplementary results); for all binary moderators, 0=variable absent, 1=variable present.

We next examined whether 16 binary variables moderated correlation coefficients in the total

sample (Table 2 and Supplementary Table 3). There were no significant moderating effects in relation

to target symptom domain, suggestibility measurement context, such as within a hypnotic context,

suggestion type or suggestibility scale type. Due to the limited sample size these non-significant

results should be interpreted cautiously; however, they suggest that the association between

suggestibility and nocebo responding is not significantly contingent upon these variables.

Considering the largely unaccounted for heterogeneity, and diverse selection of suggestibility

measures, we undertook a series of exploratory meta-regression and subgroup analyses to better

understand how the choice of suggestibility scale might account for the observed variability. In

particular, we distinguished three types of suggestibility scales: established measures of direct verbal

suggestibility (k=5), Z_r=0.35 [0.13, 0.57], Z=3.12, p=.002 [r=.38], I^2=68%, τ^2=.04, abbreviated

measures of direct verbal suggestibility (k=5), Z_r=0.03 [-0.24, 0.29], Z=0.17, p=.86 [r=.02], I^2=71%,

τ^2=.07, and retrospective self-report measures of suggestibility (k=3), Z_r=0.27 [-0.19, 0.73], Z=1.14,

p=.24 [r=.25], I^2=89%, τ^2=.11. Meta-regression analyses indicated that nocebo correlations with the

established scales displayed a non-significant trend towards being greater than those with the

abbreviated scales, ΔZ_r=0.33 [-0.01, 0.67], Z=1.86, p=.062 [Δr=.32], I^2=70%, τ^2=.05. By contrast,

correlations with the established direct scales did not significantly differ from those with retrospective

scales, ΔZ_r=0.08 [-0.36, 0.53], Z=0.36, p=.71 [Δr=.08], I^2=83%, τ^2=.06, nor did they differ between

abbreviated direct scales and retrospective scales, ΔZ_r=0.25 [-0.25, 0.75], Z=0.98, p=.33 [Δr=.23],

I^2=80%, τ^2=.09. Taken together, these results imply that the observed association between nocebo

responding and suggestibility was primarily driven by established direct verbal suggestibility scales

and to a lesser extent retrospective self-report scales.

Discussion

Our meta-analysis quantified the relationship between suggestibility and nocebo responding by

synthesizing data from available empirical studies. The analyses demonstrated that suggestibility is a

reliably significant, albeit weak, positive predictor of nocebo responding. Notably, the magnitude of

this effect was greater in studies with superior methodological rigour (and larger sample sizes),

particularly those that implemented experimenter blinding. These data complement previous

systematic reviews and meta-analyses showing the clear efficacy of verbal suggestion in the induction

of nocebo responses (Petersen et al., 2014; Rooney et al., 2023; Webster et al., 2016) but bridge a gap

in the literature by illustrating the significant moderating influence of trait responsiveness to suggestion on nocebo responding. This aligns with a wealth of data showing that trait suggestibility is a reliable predictor of responses to suggestion-based interventions (Jensen et al., 2023; Milling et al., 2021; Thompson et al., 2019) but also seems to function as a risk factor for conditions and symptoms adjacent to nocebo responding including functional neurological disorder (Wieder et al., 2021), mass psychogenic illness (Sapkota et al., 2020) and symptoms associated with environmental factors (idiopathic environmental intolerance) (Stein et al., 2023).

The frequently overlooked role of suggestibility in predicting nocebo responding has multiple salient implications for the empirical study of nocebo responding, as well as the occurrence of nocebo effects in clinical practice. Despite the cumulative observed association between suggestibility and nocebo responding, this effect was characterised by moderate heterogeneity that remained mostly unaccounted for in our sensitivity and meta-regression analyses. This unaccounted-for heterogeneity is potentially partly attributable to undetected publication bias. Although publication bias as a source of the observed effect cannot be ruled out, multiple studies included here reported non-significant correlations and suggestibility within our sample was most often a secondary point of interest, thereby minimising the motivation to avoid reporting non-significant suggestibility-nocebo correlations. The inter-study variability is plausibly attributed to methodological differences across studies, the influence of which was not observed because of the overall small sample size. Some plausible moderators of this association include the characteristics of nocebo suggestions, suggestibility instruments, and target symptom domain.

Although the efficacy of verbal suggestion in nocebo responding is well established (Petersen et al., 2014; Rooney et al., 2023; Webster et al., 2016), the nuances of suggestion have received only scant attention in research on nocebo effects. For example, verbal suggestions ("your pain will reduce") and instructions ("try to reduce your pain") are frequently conflated (e.g., Rooney et al., 2023), even though they are hypothesized to reflect distinct phenomena (Brass et al., 2017; Terhune et al., 2017). Similarly, researchers have tended to neglect important differences in the phrasing of verbal suggestions, such as the distinction between direct and indirect suggestions. Using meta-

regression analyses, we aimed to disentangle the role of suggestion type, though our results should be considered preliminary as we found no significant evidence for moderating effects. In our sample of included studies, the specific verbal suggestions used to induce nocebo effects were only rarely reported, which hindered our ability to reliably code suggestion type. The moderating influence of suggestibility is plausibly contingent on the suggestion type administered. Accordingly, we recommend future studies report the exact wording of suggestions and provide specific details about when suggestions were administered during an experiment. This approach will enhance the reproducibility of experimental research in this domain and permit future analyses on the moderating influence of different features of verbal, textual, and non-verbal suggestions (see also Halligan & Oakley, 2014).

Another plausible source of heterogeneity in the association between suggestibility and nocebo responding is variability in the psychometric measurement of suggestibility. The domain of suggestibility includes responsiveness to direct and indirect verbal suggestions, often measured through work-sample instruments (e.g., Woody & Barnier, 2008). Little consideration has been devoted to salient differences between suggestibility measures, with most researchers implicitly assuming a uniform construct of suggestibility. In nocebo studies, trait suggestibility has been measured with established work-sample instruments or retrospective self-report measures. The former (direct verbal suggestibility scales) typically draw upon measures from experimental hypnosis research, in which responsiveness to a set of verbal suggestions for changes in motor control, cognition and perception are administered by an experimenter and then evaluated with brief tests (Acunzo & Terhune, 2021; Woody & Barnier, 2008). In our sample, 10 studies used direct verbal suggestibility scales, however, there was considerable variability in the number and type of suggestions included. Of note, the studies (Khan et al., 2009; Leigh et al., 2003; Woody et al., 1997; Zech et al., 2019) that used established, longer direct verbal suggestibility scales (Barber & Wilson, 1978; Shor & Orne, 1963) yielded non-significantly larger correlations than the studies (Sharav et al., 2023; Zech et al., 2022; Zech et al., 2020) that used abbreviated direct verbal suggestibility scales (Hilgard et al., 1979; Riegel et al., 2021). It remains unclear whether these differences are due to a

greater scoring range in the established scales, or their inclusion of cognitive-perceptual suggestions, which are plausibly more relevant to symptom induction than the mostly motor suggestions on the abbreviated scales. On the other hand, the use of retrospective self-report measures of suggestibility (Kotov et al., 2004) yielded more ambiguous results. Although preliminary, these results suggest that future research will benefit from the inclusion of more established suggestibility scales that include a more diverse set of suggestions and the avoidance of abbreviated suggestibility scales.

Nocebo symptoms can be reliably induced but their severity is inconsistent across symptom domains (Bagaric et al., 2022; Meeuwis et al., 2023; Rooney et al., 2023; Wolters et al., 2019). We assessed several symptom domain moderators: pain, dyspnoea, dizziness, nausea, motor inhibition, and cognitive function, all of which were found to be non-significant, which indicates that the nocebo-suggestibility association was not greater in these domains than in all others. This is likely due to the relatively small sample of included studies, such that the moderation analyses plausibly lacked sufficient power to detect weak correlation differences, especially given that each symptom domain included only 2-4 studies. Alternatively, it is plausible that the diversity in nocebo induction methods could account for the lack of moderation effects. For example, suggestibility was previously found to correlate with placebo responding when suggestion alone was used to induce placebo hypoalgesia (Parsons et al., 2021). As such, it might be that the role of suggestibility in predicting nocebo responding is dependent on suggestion-centred symptom induction techniques. Future research would benefit from clarifying how suggestibility relates to nocebo effects from different nocebo induction methods and corroborate the results of previous reviews by disentangling the moderating effects of target symptom domain within nocebo responding.

The results of this meta-analysis complement previous research demonstrating links between suggestibility and phenomena germane to nocebo responding and corroborate proposals that suggestibility confers risk for somatic symptom reporting and nocebo responding more broadly (Corsi & Colloca, 2017; Spiegel, 1997). As such, highly suggestible individuals may be more susceptible to nocebo responses due to potential side effect information, informed consent processes, or treatment misinformation. From a predictive processing perspective (Hohwy, 2020), highly suggestible

individuals may have a capacity for forming more precise symptom priors that shape perception in the direction of contextual cues (Fiorio et al., 2022; Van den Bergh et al., 2017). For example, during treatment information processes, side effect information may function as a type of indirect verbal suggestion leading to the formation of precise symptom priors, inducing premature or imitative adverse events (i.e., nocebo response). These symptom priors might compound and generalize, from treatment-specific nocebo responding, to more pronounced functional conditions depending on other individual difference factors (e.g., treatment history). Future research should explore the interplay between suggestibility, contextual cues, and the formation of precise symptom priors, investigating how individual differences and life experiences might modulate the transition from treatment-specific nocebo responses to broader functional conditions. Additionally, future research should examine suggestibility as a predictor of response to active treatments (Nitzan et al., 2015; Szigeti et al., 2024)

Limitations

The presented results should be interpreted within the context of their limitations. As in the broader experimental study of nocebo effects, the majority our sample relied on self-report measures of symptom outcomes (k=7), which are susceptible to reporting biases. However, we partially accounted for this possibility in our sensitivity analyses which suggest that the overall main effect is unlikely to be driven by the use of self-report measures. Although we observed stronger effects in higher-quality studies, the overall methodological quality across studies was relatively low with no study scoring >80%. In particular, very few of the studies included control conditions and many used baseline as a control comparator. Future research examining the role of suggestibility in nocebo responding will need to employ more rigorous designs and ensure clear reporting of experimental parameters including the precise wording of nocebo verbal suggestions. The high heterogeneity in our sample indicates that unaccounted variables impacted the association between suggestibility and nocebo responding. Our inclusion criteria resulted in a diverse sample of outcomes (e.g., pain, motor control) and nocebo induction methods (e.g., verbal suggestion, conditioning), which might have contributed to the heterogeneity and limit the generalizability of our findings. Finally, most of the studies (k=6; n range: 46-93) were only sufficiently powered to detect moderate-to-large effects, and thus were

insufficiently powered to reliably detect weak effects. This limitation may have also led to an

underestimation of the association between suggestibility and nocebo responding. Insofar as the

observed effect size is small in magnitude, this indicates that suggestibility is only one of multiple

factors contributing to variability in response to nocebos including state anxiety (Rooney et al., 2022)

and empathy (Meeuwis et al., 2023). It will be important in the future to evaluate whether these reflect

independent or interacting effects. Cumulatively, these limitations highlight the need for rigorously

designed experiments with meticulous control comparisons to further examine the apparent link

between suggestibility and nocebo responses.

Summary

This meta-analysis found a reliable association between trait suggestibility and nocebo responding.

Given the small effect size, the present results raise important questions about the practical

significance of this effect. However, owing to the relatively small number of studies, methodological

heterogeneity, and inconsistent reporting of verbal suggestions, future research is needed to better

assess the practical significance of suggestibility as a moderator of nocebo responding, particularly

outside of an experimental context. From a clinical standpoint, understanding the role of suggestibility

in nocebo responding and symptom perception more generally can pave the way for more

personalized patient care. Highly suggestible individuals might be more susceptible to negative

treatment information and thus could benefit from specific communication strategies that minimize

nocebo effects (Arrow et al., 2022; Kari et al., 2021; Spiegel, 1997; Wells & Kaptchuk, 2012).

Integrating suggestibility into existing models of nocebo responding, as has been done in accounts of

related conditions such as functional neurological disorder (Fiorio et al., 2022), could offer a more

holistic understanding of variability in nocebo responses. By doing so, we can potentially uncover

nuanced interactions between suggestibility and other established factors, such as the manner in which

side effect information is communicated to participants and patients, as well as their level of trust

Figin health care professionals. As we aim for more personalized and effective patient care and move

towards an era of precision medicine (Seyhan & Carini, 2019), understanding individual differences

factors that moderate nocebo responding is paramount for improving treatment outcomes.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or

publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this

article.

Data availability statement

All data relevant to the present meta-analysis is provided in the article. Data are freely available from

previous research studies.

References

Acunzo, D. J., & Terhune, D. B. (2021). A critical review of standardized measures of hypnotic suggestibility. *Int J Clin Exp Hypn*, *69*(1), 50-71. https://doi.org/10.1080/00207144.2021.1833209

Arrow, K., Burgoyne, L. L., & Cyna, A. M. (2022). Implications of nocebo in anaesthesia care. *Anaesthesia*, *77 Suppl 1*, 11-20. https://doi.org/10.1111/anae.15601

Bagaric, B., Jokic-Begic, N., & Sangster Jokic, C. (2022). The nocebo effect: A review of contemporary experimental research. *Int J Behav Med*, *29*(3), 255-265. https://doi.org/10.1007/s12529-021-10016-y

Barber, T. X., & Wilson, S. C. (1978). The barber suggestibility scale and the creative imagination scale: Experimental and clinical applications. *Am J Clin Hypn*, *21*(2-3), 84-108. https://doi.org/10.1080/00029157.1978.10403966

Bell, V., Oakley, D. A., Halligan, P. W., & Deeley, Q. (2011). Dissociation in hysteria and hypnosis: Evidence from cognitive neuroscience. *J Neurol Neurosurg Psychiatry*, *82*(3), 332-339. https://doi.org/10.1136/jnnp.2009.199158

Bender, F. L., Rief, W., & Wilhelm, M. (2023). Really just a little prick? A meta-analysis on adverse events in placebo control groups of seasonal influenza vaccination rcts. *Vaccine*, *41*(2), 294-303. https://doi.org/10.1016/j.vaccine.2022.11.033

Brascher, A. K., Sutterlin, S., Scheuren, R., Van den Bergh, O., & Witthoft, M. (2020). Somatic symptom perception from a predictive processing perspective: An empirical test using the thermal grill illusion. *Psychosom Med*, *82*(7), 708-714. https://doi.org/10.1097/PSY.0000000000000824

Brass, M., Liefooghe, B., Braem, S., & De Houwer, J. (2017). Following new task instructions: Evidence for a dissociation between knowing and doing. *Neuroscience & Biobehavioral Reviews*, *81*, 16-28. https://doi.org/https://doi.org/10.1016/j.neubiorev.2017.02.012

Brooke, B. S., Schwartz, T. A., & Pawlik, T. M. (2021). Moose reporting guidelines for meta-analyses of observational studies. *JAMA Surgery*, *156*(8), 787-788. https://doi.org/10.1001/jamasurg.2021.0522

Colloca, L., & Barsky, A. J. (2020). Placebo and nocebo effects. *New England Journal of Medicine*, *382*(6), 554-561. https://doi.org/10.1056/NEJMra1907805

Colloca, L., & Miller, F. G. (2011). Harnessing the placebo effect: The need for translational research. *Philos Trans R Soc Lond B Biol Sci*, *366*(1572), 1922-1930. https://doi.org/10.1098/rstb.2010.0399

Corsi, N., & Colloca, L. (2017). Placebo and nocebo effects: The advantage of measuring expectations and psychological factors. *Front Psychol*, *8*, 308. https://doi.org/10.3389/fpsyg.2017.00308

Duval, S., & Tweedie, R. (2000). Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics*, *56*(2), 455-463. https://doi.org/10.1111/j.0006-341x.2000.00455.x

Egger, M., Davey Smith, G., Schneider, M., & Minder, C. (1997). Bias in meta-analysis detected by a simple, graphical test. *BMJ*, *315*(7109), 629-634. https://doi.org/10.1136/bmj.315.7109.629

Fiorio, M., Braga, M., Marotta, A., Villa-Sanchez, B., Edwards, M. J., Tinazzi, M., & Barbiani, D. (2022). Functional neurological disorder and placebo and nocebo effects: Shared mechanisms. *Nat Rev Neurol*, *18*(10), 624-635. https://doi.org/10.1038/s41582-022-00711-z

Haas, J. W., Bender, F. L., Ballou, S., Kelley, J. M., Wilhelm, M., Miller, F. G., Rief, W., & Kaptchuk, T. J. (2022). Frequency of adverse events in the placebo arms of covid-19 vaccine trials: A systematic review and meta-analysis. *JAMA Network Open*, *5*(1), e2143955-e2143955. https://doi.org/10.1001/jamanetworkopen.2021.43955

Hauser, W., Hansen, E., & Enck, P. (2012). Nocebo phenomena in medicine: Their relevance in everyday clinical practice. *Dtsch Arztebl Int*, *109*(26), 459-465. https://doi.org/10.3238/arztebl.2012.0459

Heller, M. K., Chapman, S. C. E., & Horne, R. (2022). Beliefs about medicines predict side-effects of placebo modafinil. *Ann Behav Med*, *56*(10), 989-1001. https://doi.org/10.1093/abm/kaab112

Hilgard, E. R., Crawford, H. J., & Wert, A. (1979). The stanford hypnotic arm levitation induction and test (shalit): A six-minute hypnotic induction and measurement scale. *International Journal of Clinical and Experimental Hypnosis*, *27*(2), 111-124. https://doi.org/10.1080/00207147908407551

Hohwy, J. (2020). New directions in predictive processing. *Mind & Language*, *35*(2), 209-223. https://doi.org/10.1111/mila.12281

Jensen, M. P., Ehde, D. M., Hakimian, S., Pettet, M. W., Day, M. A., & Ciol, M. A. (2023). Who benefits the most from different psychological chronic pain treatments? An exploratory analysis of treatment moderators. *J Pain*, *24*(11), 2024-2039. https://doi.org/10.1016/j.jpain.2023.06.011

Kari, A. L., Lauren, C. H., & Alia, J. C. (2021). Changing mindsets about side effects. *BMJ Open*, *11*(2), e040134. https://doi.org/10.1136/bmjopen-2020-040134

Kern, A., Kramm, C., Witt, C. M., & Barth, J. (2020). The influence of personality traits on the placebo/nocebo response: A systematic review. *Journal of Psychosomatic Research*, *128*, 109866. https://doi.org/https://doi.org/10.1016/j.jpsychores.2019.109866

Khan, A. Y., Baade, L., Ablah, E., McNerney, V., Golewale, M. H., & Liow, K. (2009). Can hypnosis differentiate epileptic from nonepileptic events in the video/eeg monitoring unit? Data from a pilot study. *Epilepsy Behav*, *15*(3), 314-317. https://doi.org/10.1016/j.yebeh.2009.04.004

Kotov, R. I., Bellman, S. B., & Watson, D. B. (2004). Multidimensional iowa suggestibility scale (miss) brief manual *Unpublished manuscript*.

Leigh, R., MacQueen, G., Tougas, G., Hargreave, F. E., & Bienenstock, J. (2003). Change in forced expiratory volume in 1 second after sham bronchoconstrictor in suggestible but not suggestion-resistant asthmatic subjects: A pilot study. *Psychosom Med*, *65*(5), 791-795. https://doi.org/10.1097/01.psy.0000079454.48714.1b

Meeuwis, S. H., Wasylewski, M. T., Bajcar, E. A., Bieniek, H., Adamczyk, W. M., Honcharova, S., Di Nardo, M., Mazzoni, G., & Babel, P. (2023). Learning pain from others: A systematic review and meta-analysis of studies on placebo hypoalgesia and nocebo hyperalgesia induced by observational learning. *Pain*. https://doi.org/10.1097/j.pain.0000000000002943

Milling, L. S., Valentine, K. E., LoStimolo, L. M., Nett, A. M., & McCarley, H. S. (2021). Hypnosis and the alleviation of clinical pain: A comprehensive meta-analysis. *International Journal of Clinical and Experimental Hypnosis*. https://doi.org/http://dx.doi.org/10.1080/00207144.2021.1920330

Nasiri-Dehsorkhi, H., Vaziri, S., Esmaillzadeh, A., & Adibi, P. (2023). Psychological distress, perceived stress and nocebo effect (multifood adverse reaction) in irritable bowel syndrome patients. *J Educ Health Promot*, *12*, 257. https://doi.org/10.4103/jehp.jehp_221_23

Nitzan, U., Chalamish, Y., Krieger, I., Erez, H. B., Braw, Y., & Lichtenberg, P. (2015). Suggestibility as a predictor of response to antidepressants: A preliminary prospective trial. *J Affect Disord*, *185*, 8-11. https://doi.org/10.1016/j.jad.2015.06.028

Oakley, D. A., Walsh, E., Mehta, M. A., Halligan, P. W., & Deeley, Q. (2021). Direct verbal suggestibility: Measurement and significance. *Conscious Cogn*, *89*, 103036. https://doi.org/10.1016/j.concog.2020.103036

Page, M. J., Moher, D., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., Shamseer, L., Tetzlaff, J. M., Akl, E. A., Brennan, S. E., Chou, R., Glanville, J., Grimshaw, J. M., Hróbjartsson, A., Lalu, M. M., Li, T., Loder, E. W., Mayo-Wilson, E., McDonald, S., . . . McKenzie, J. E. (2021). Prisma 2020 explanation and elaboration: Updated guidance and exemplars for reporting systematic reviews. *BMJ*, *372*, n160. https://doi.org/10.1136/bmj.n160

Parsons, R. D., Bergmann, S., Wiech, K., & Terhune, D. B. (2021). Direct verbal suggestibility as a predictor of placebo hypoalgesia responsiveness. *Psychosom Med*, *83*(9), 1041-1049. https://doi.org/10.1097/psy.0000000000000977

Petersen, G. L., Finnerup, N. B., Colloca, L., Amanzio, M., Price, D. D., Jensen, T. S., & Vase, L. (2014). The magnitude of nocebo effects in pain: A meta-analysis. *Pain*, *155*(8), 1426-1434. https://doi.org/10.1016/j.pain.2014.04.016

Petrovic, P. (2008). Placebo analgesia and nocebo hyperalgesia--two sides of the same coin? *Pain*, *136*(1-2), 5-6. https://doi.org/10.1016/j.pain.2008.03.004

Polczyk, R. (2016). Factor structure of suggestibility revisited: New evidence for direct and indirect suggestibility. *Current Issues in Personality Psychology*, *4*(2), 87-96. https://doi.org/10.5114/cipp.2016.60249

Riegel, B., Tönnies, S., Hansen, E., Zech, N., Eck, S., Batra, A., & Peter, B. (2021). German norms of the harvard group scale of hypnotic susceptibility, form a (hgshs:A) and proposal of a 5-item short-version (hgshs-5:G). *International Journal of Clinical and Experimental Hypnosis*, *69*(1), 112-123. https://doi.org/10.1080/00207144.2021.1836645

Rooney, T., Sharpe, L., Todd, J., Richmond, B., & Colagiuri, B. (2022). The relationship between expectancy, anxiety, and the nocebo effect: A systematic review and meta-analysis with recommendations for future research. *Health Psychol Rev*, 1-28. https://doi.org/10.1080/17437199.2022.2125894

Rooney, T., Sharpe, L., Todd, J., Tang, B., & Colagiuri, B. (2023). The nocebo effect across health outcomes: A systematic review and meta-analysis. *Health Psychol*. https://doi.org/10.1037/hea0001326

Sapkota, R. P., Brunet, A., & Kirmayer, L. J. (2020). Characteristics of adolescents affected by mass psychogenic illness outbreaks in schools in nepal: A case-control study. *Front Psychiatry*, *11*, 493094. https://doi.org/10.3389/fpsyt.2020.493094

Seyhan, A. A., & Carini, C. (2019). Are innovation and new technologies in precision medicine paving a new era in patients centric care? *J Transl Med*, *17*(1), 114. https://doi.org/10.1186/s12967-019-1864-9

Sharav, Y., Haviv, Y., & Tal, M. (2023). Placebo or nocebo interventions as affected by hypnotic susceptibility. *Applied Sciences*, *13*(2), 931. https://www.mdpi.com/2076-3417/13/2/931

Shor, R. E., & Orne, E. C. (1963). Norms on the harvard group scale of hypnotic susceptibility, form a. *International Journal of Clinical and Experimental Hypnosis*, *11*(1), 39-47. https://doi.org/10.1080/00207146308409226

Spiegel, H. (1997). Nocebo: The power of suggestibility. *Prev Med*, *26*(5 Pt 1), 616-621. https://doi.org/10.1006/pmed.1997.0229

Stein, M. V., Holt, R., Wieder, L., & Terhune, D. B. (2023). Responsiveness to direct verbal suggestions and dissociation independently predict symptoms associated with environmental factors. *Psychopathology*, *56*(4), 324-328.

Szigeti, B., Weiss, B., Rosas, F. E., Erritzoe, D., Nutt, D., & Carhart-Harris, R. (2024). Assessing expectancy and suggestibility in a trial of escitalopram v. Psilocybin for depression. *Psychol Med*, 1-8. https://doi.org/10.1017/s0033291723003653

Terhune, D. B., Cleeremans, A., Raz, A., & Lynn, S. J. (2017). Hypnosis and top-down regulation of consciousness. *Neuroscience and Biobehavioral Reviews*, *81*(Part A), 59-74. https://doi.org/http://dx.doi.org/10.1016/j.neubiorev.2017.02.002

The jamovi project. (2023). *Jamovi (version 2.3) [computer software]. Retrieved from https://www.Jamovi.Org*.

Thompson, T., Terhune, D. B., Oram, C., Sharangparni, J., Rouf, R., Solmi, M., Veronese, N., & Stubbs, B. (2019). The effectiveness of hypnosis for pain relief: A systematic review and meta-analysis of 85 controlled experimental trials. *Neuroscience and Biobehavioral Reviews*, *99*, 298-310. https://doi.org/http://dx.doi.org/10.1016/j.neubiorev.2019.02.013

Van den Bergh, O., Witthoft, M., Petersen, S., & Brown, R. J. (2017). Symptoms and the body: Taking the inferential leap. *Neurosci Biobehav Rev*, *74*(Pt A), 185-203. https://doi.org/10.1016/j.neubiorev.2017.01.015

Webster, R. K., Weinman, J., & Rubin, G. J. (2016). A systematic review of factors that contribute to nocebo effects. *Health Psychol*, *35*(12), 1334-1355. https://doi.org/10.1037/hea0000416

Wells, R. E., & Kaptchuk, T. J. (2012). To tell the truth, the whole truth, may do patients harm: The problem of the nocebo effect for informed consent. *Am J Bioeth*, *12*(3), 22-29. https://doi.org/10.1080/15265161.2011.652798

Wickramasekera, I. E., & Szlyk, J. P. (2003). Could empathy be a predictor of hypnotic ability? *International Journal of Clinical and Experimental Hypnosis*, *51*(4), 390-399. https://doi.org/10.1076/iceh.51.4.390.16413

Wieder, L., Brown, R., Thompson, T., & Terhune, D. B. (2021). Suggestibility in functional neurological disorder: A meta-analysis. *Journal of Neurology, Neurosurgery and Psychiatry*, *92*(2), 150-157. https://doi.org/10.1136/jnnp-2020-323706

Winter, D., & Braw, Y. (2022). Covid-19: Impact of diagnosis threat and suggestibility on subjective cognitive complaints. *Int J Clin Health Psychol*, *22*(1), 100253. https://doi.org/10.1016/j.ijchp.2021.100253

Wolters, F., Peerdeman, K. J., & Evers, A. W. M. (2019). Placebo and nocebo effects across symptoms: From pain to fatigue, dyspnea, nausea, and itch. *Front Psychiatry*, *10*, 470. https://doi.org/10.3389/fpsyt.2019.00470

Woody, E. Z., & Barnier, A. J. (2008). Hypnosis scales for the twenty-first century: What do we need and how should we use them? In *The oxford handbook of hypnosis: Theory, research, and practice.* (pp. 255-280). Oxford University Press.

Woody, E. Z., Drugovic, M., & Oakman, J. M. (1997). A reexamination of the role of nonhypnotic suggestibility in hypnotic responding. *Journal of Personality and Social Psychology*, *72*(2), 399-407. https://doi.org/http://dx.doi.org/10.1037/0022-3514.72.2.399

Zech, N., Scharl, L., Seemann, M., Pfeifer, M., & Hansen, E. (2022). Nocebo effects of clinical communication and placebo effects of positive suggestions on respiratory muscle strength. *Front Psychol*, *13*, 825839. https://doi.org/10.3389/fpsyg.2022.825839

Zech, N., Schrödinger, M., Seemann, M., Zeman, F., Seyfried, T. F., & Hansen, E. (2020). Time-dependent negative effects of verbal and non-verbal suggestions in surgical patients-a study on arm muscle strength. *Front Psychol*, *11*, 1693. https://doi.org/10.3389/fpsyg.2020.01693

Zech, N., Seemann, M., Grzesiek, M., Breu, A., Seyfried, T. F., & Hansen, E. (2019). Nocebo effects on muscular performance - an experimental study about clinical situations. *Front Pharmacol*, *10*, 219. https://doi.org/10.3389/fphar.2019.00219