1   **ADHD and intelligence polygenic scores associations with developmental dimensions in children**

2   **with attention, learning and memory difficulties**

Andrea M. Santangelo[1,2], Olena Ohlei[3], Silvana Mareva[1,4], Diandra Brkic[1,5], CALM team[1], Lars

Bertram[3], Joni Holmes[1,6], Duncan Astle[1,7] and Kate Baker[1,8,9] *

[1] MRC Cognitive and Brain Sciences Unit, University of Cambridge, UK

[2] MRC Integrative Epidemiology Unit, University of Bristol, UK

[3] Lubeck Interdisciplinary Platform for Genome Analysis, University of Luebeck, Germany

[4] Psychology Department, University of Exeter, UK

[5] Brain Health Department, Nestlé Institute of Health Sciences, Switzerland

[6] School of Psychology, University of East Anglia, UK

[7] Department of Psychiatry, University of Cambridge, UK

[8] Department of Genomic Medicine, University of Cambridge, UK

[9] Department of Pathology, University of Cambridge, UK

* corresponding author: kb488@cam.ac.uk

Acknowledgments

Genotyping was carried out within the "Healthy minds 0-100 years: Optimising the use of European brain imaging cohorts (Lifebrain)" project, funded by EU Horizon2020 (Grant agreement number: 732592). We are grateful to Professor Sadaf Farooqi and her laboratory for DNA extractions and storage, and to NIHR BRC-MRC BioRepository for sample archiving.

The Centre for Attention Learning and Memory (CALM) research clinic was based at the MRC Cognition and Brain Sciences Unit, University of Cambridge, funded by the MRC through UKRI (https://calm.mrc-cbu.cam.ac.uk/). The authors wish to thank the many professionals working in children's services in the South-East and East of England for their support, and the children and their families for contributing to the project. The lead investigators are Duncan Astle, Kate Baker, Susan Gathercole, Joni Holmes, Rogier Kievit and Tom Manly. Data collection is assisted by a team of researchers and PhD students that includes Danyal Akarca, Joe Bathelt, Marc Bennett, Madalena Bettencourt, Giacomo Bignardi, Sarah Bishop, Erica Bottacin, Lara Bridge, Diandra Brkic, Annie Bryant, Sally Butterfield, Elizabeth Byrne, Gemma Crickmore, Edwin Dalmaijer, Fánchea Daly, Tina Emery, Laura Forde, Grace Franckel, Delia Furhmann, Andrew Gadie, Sara Gharooni, Jacalyn Guy, Erin Hawkins, Rebeca Ianov-Vitanov, Christian Iordanov, Agnieszka Jaroslawska, Sara Joeghan, Amy Johnson, Jonathan Jones, Silvana Mareva, Jessica Martin, Elise Ng-Cordell, Sinead O'Brien, Cliodhna O'Leary, Joseph Rennie, Andrea M. Santangelo, Ivan Simpson-Kent, Roma Siugzdaite, Tess Smith, Stepheni Uh, Maria Vedechkina, Francesca Woolgar, Natalia Zdorovtsova, Mengya Zhang.

1   **Abstract**

Common genetic variants make a significant contribution to neurodevelopmental characteristics

such as cognitive abilities and ADHD symptoms. The relevance and structure of these associations

amongst children with transdiagnostic difficulties in cognition, attention and learning has not been

explored. Polygenic scores (PGS) derived from the largest genome-wide association study (GWAS)

data at the time of this study on ADHD (38,691 individuals with ADHD and 186,843 controls) and

Intelligence (269,867 individuals) were calculated for 524 children and young people (5-18 years old)

referred to the Centre for Attention, Learning and Memory (CALM). PGS-trait associations were

assessed via linear regression analyses, for a range of cognitive and behavioural dimensional

measures, and factor scores from a hierarchical model of psychopathology. PGS associations were

explored with and without co-varying for socio-economic status (SES). Within this sample, we found

the expected positive associations between ADHD-PGS and ADHD primary symptoms, and between

Intelligence-PGS and IQ. ADHD-PGS were also associated with broader externalising behaviours and

intelligence scores, and these associations remained significant after removing ADHD-diagnosed

participants, or after covarying with SES. Intelligence-PGS showed associations with verbal and non-

verbal cognitive skills, but no significant associations with ADHD traits were detected. For the

hierarchical model of psychopathology, ADHD-PGS, but not intelligence-PGS, showed associations

with the general mental health factor, externalising factor, and social maladjustment factor, only

when SES was not included as a covariate. In summary, PGS for neurodevelopmental traits may

contribute to both general and specific cognitive and behavioural dimensions in a paediatric

transdiagnostic sample. Future studies investigating PGS associations with neural correlates, as well

as gene-by-environment interactions, will contribute to our understanding of developmental

pathways and risk-resilience mechanisms in child mental health.

Introduction

Children with neurodevelopmental difficulties often do not meet criteria for a specific

diagnosis, or they meet criteria for multiple diagnoses. Whilst case–control designs or population-

based studies have been instrumental in establishing disorder-specific and population-level genetic

associations, these approaches have shown limited ability to capture the substantial phenotypic and

genetic heterogeneity that characterises psychiatric and neurodevelopmental conditions. Individuals

with sub-diagnostic or transdiagnostic symptoms are often underrepresented: in case/control

studies with samples enriched in extreme phenotypes due to strict inclusion/exclusion criteria; and

in general population studies, with samples including unselected individuals, where the genetic

effects of sub-diagnostic symptoms may be attenuated due to low prevalence of clinical symptoms.

Recognising these complexities, investigations of neurodevelopment and mental health have

increasingly adopted a transdiagnostic framework [1]. Within this framework, it is not necessary to

meet formal diagnostic criteria to be included in a study, neither is the presence of co-occurring

difficulties an exclusion criterion i.e. better mimicking the lived experience of multi-dimensional

neurodevelopmental difficulties. Whether the underlying genetic architecture of

neurodevelopmental dimensions within transdiagnostic samples of children with heterogeneous and

co-occurring difficulties is different from diagnostic samples and general populations is still

unknown.

Attention-Deficit/Hyperactivity Disorder (ADHD) is a common neurodevelopmental disorder

characterised by symptoms in either the inattentive or hyperactive/impulsive domains, or both [2].

ADHD affects approximately 5% of children and 2.5% of adults, with a high heritability of

approximately 70-80% estimated from twin-based studies [3], and high comorbidity with other

psychiatric conditions [4]. At the time of this study, the latest genome-wide association meta-

analysis (meta-GWAS) of ADHD case-control study identified 27 genome-wide significant risk loci,

including genes enriched for expression in early brain development, with 14 % of SNP heritability

accounting for ADHD [5].

A proportion of children with symptoms of ADHD or other behavioural difficulties also have

cognitive and learning-related difficulties. The ability to think abstractly and problem solve

independently of any previously acquired knowledge is known as fluid intelligence [6]. Fluid

intelligence is highly heritable, with a general factor of intelligence (g) accounting for about 58% of

the genetic variance in cognitive traits [7]. A large-scale, meta-GWAS for the g factor of general

intelligence has identified 205 genomic loci implicating brain-expressed genes involved in nervous

system development, neuron differentiation, and synapse structure and activity [8].

Polygenic scores (PGS) are estimates of individual common genetic variation associated with

a trait or disorder and can provide insights into the genetic architecture and underlying mechanisms

contributing to complex phenotypes [9, 10]. Higher ADHD-PGS have been associated with ADHD

diagnosis in independent case-control samples [11], as well as with ADHD symptoms in population-

based studies [5, 12, 13], suggesting that an ADHD diagnosis represents the extreme expression of

one or more heritable quantitative traits. In addition, ADHD-PGS have also been negatively

correlated with cognitive abilities including general intelligence, attention, verbal and non-verbal

reasoning, and working memory [5, 14]. Conversely, genetic correlation and Mendelian

randomization analysis showed a negative causal association between intelligence-related variants

and ADHD and depression symptoms [6, 8].

Shared genetic contributions across disorders are well-recognised [15–17], with ADHD

sharing 84–98% of risk variants with other psychiatric disorders [5, 11]. Thus, PGS may influence

aspects of neurobiological and cognitive development relevant to a diverse set of educational,

behavioural and psychiatric vulnerabilities [7, 18]. Whilst examining specific PGS-behavioural

relationships has value and builds on previous literature, children with co-occurring developmental

difficulties experience a combination of cognitive, behavioural, social and emotional challenges, that

may be better understood in an integrated fashion. Hierarchical models of psychopathology, such as

the HiTOP framework [19–21], offer a structured way to examine how genetic liability relates to

mental health across multiple levels of phenotypic organization. The HiTOP model proposes a p-

factor of general psychopathology at the apex, under which more specific factors are described [22,

23]. The p factor is highly heritable (50-60% from twin studies), and the model aligns well with

current understanding of the higher order genetic structure of psychopathology. Although ADHD

polygenic risk has previously been linked to HiTOP-related factors [12, 24–26], most evidence comes

from population-based or adult samples, leaving it unclear whether these hierarchical patterns

generalize to clinically referred paediatric populations characterized by high comorbidity and

neurodevelopmental heterogeneity.

Following this approach, the Centre for Attention Learning and Memory in Cambridge

(CALM, http://calm.mrc-cbu.cam.ac.uk/) recruited a cohort of approximately 800 children

recognised by educational or clinical professionals as having difficulties in attention, learning and/or

memory [27]. This unique, transdiagnostic paediatric sample provides an opportunity to study the

factors (genetics and environmental) and mechanisms that drive diversity in neurodevelopment,

independent of categorical diagnoses and their co-occurrence. The CALM cohort is a deep-

phenotyped sample from the point of view of behavioural features [23, 28–30], dimensions of

cognitive and executive functions [31–33], dimensions of mental health [23, 34], and neuroimaging

[35, 36].

A transdiagnostic design enables the investigation of ADHD and intelligence PGS across

multiple diagnostic categories and symptom dimensions, thereby allowing the detection of shared,

divergent, and moderating genetic effects that may not be observable in diagnostically

homogeneous samples. Within a transdiagnostic sample enriched for individuals with learning

difficulties, ADHD and intelligence PGS may show associations with cognitive, behavioural, and

emotional outcomes that cut across traditional diagnostic boundaries, reflecting pleiotropic genetic

influences and dimensional symptom expression. For example, ADHD PGS may relate to attentional

or executive functioning deficits in individuals without a formal ADHD diagnosis, while intelligence

PGS may exert protective or compensatory effects on cognitive outcomes across diverse psychiatric

presentations. Such cross-disorder and moderating effects are likely to be attenuated in population

cohorts due to low symptom burden, and cannot be adequately examined in single-disorder case–

control studies. Consequently, transdiagnostic analyses provide a more mechanistically informative

framework for understanding how polygenic liability for ADHD and intelligence contributes to clinical

heterogeneity and functional outcomes across psychiatric disorders.

The current paper addresses the question of whether and how ADHD and Intelligence PGS

are associated with relevant neurodevelopmental characteristics within the transdiagnostic CALM

sample. We focused on ADHD and Intelligence PGS because these traits reflect important aspects of

behavioural and cognitive heterogeneity within the CALM cohort, and there are robust large-scale

GWAS studies to draw on for planned analyses. In addition, we conducted association analyses with

and without co-variation with socio-economic status (SES), approximated via the Index of Multiple

Deprivation (MDI), an important demographic factor that contributes to behavioural symptom

likelihood and correlates with polygenic variation [37].

Our first objective was to establish whether ADHD and Intelligence PGS showed associations

with their respective primary traits i.e. ADHD symptoms and cognitive test scores, within the CALM

sample. Our second objective was to examine whether PGS associations were specific to these traits,

or extended to additional behavioural traits (externalising and internalising problems) which would

suggest cross-over between behavioural and cognitive dimensions. Thirdly, we explored whether

PGS could contribute to our understanding of a hierarchical model of psychopathology (HiToP)

within this population. The extent to which PGS show specific or shared associations within the

CALM sample could provide insight into the inter-relationships between dimensions of

neurodevelopmental difficulty, at a behavioural or mechanistic level.

Methods

Participants

A cohort of children aged 5 to 18 years was recruited by the Centre for Attention, Learning

and Memory (CALM, http://calm.mrc-cbu.cam.ac.uk/) between February 2014 and December 2021

[27]. The cohort consisted of 800 children referred by a health or educational professional because

of difficulties related to attention, learning and/or memory. In addition, 200 children were recruited

from the same schools as controls. This paper reports analysis on the referred children only.

Comprehensive phenotyping was carried out for all children (for details see Holmes et al., 2019)

(**Table 1**), and an optional DNA sample (saliva) was provided. A total of 615 samples were submitted

from the referred children for genotyping. After QC filtering (see below), a final N=524 participants

remained, with 361 males and 163 females, and average age of 9.5 ± 2.4 years (**Table 2**). Index of

Multiple Deprivation (IMD) was used to approximate the socioeconomic status of the sample [38].

Scores for postcode areas in the United Kingdom range from 1st to 32844th (most to least deprived).

The range of IMD for the sample indicated participants came from areas with varying degrees of

deprivation, with an average ranking above the national median (**Table 2**). The decile distribution for

IMD in this cohort is shown in **Fig. S1**. Within this group of participants, 38% had received a diagnosis

of any developmental or psychiatric disorder from community services, 22.5% had received a

diagnosis of ADHD, 11.8% had been diagnosed with dyslexia or were receiving speech and language

therapy, and 7% had received a diagnosis of autism.

Phenotypic measures

The measures used in the current analysis are listed in **Table 1**, together with descriptive

data for each measure used within the CALM PGS-analysed sample.

The Conners 3 – Parent Rating Scale Short Form (CPSF) is a parent/carer-reported

questionnaire used to assess symptoms commonly associated with ADHD. It includes 45 items

describing behavioural difficulties observed over the past month, which are grouped into six

subscales: Inattention, Hyperactivity/Impulsivity, Learning Problems, Executive Function, Aggression,

and Peer Relations. We used for our analyses the subscales of Inattention and

Hyperactivity/Impulsivity.

The Strengths and Difficulties Questionnaire (SDQ) is a 25-item parent/carer-rated measure

that assesses children's emotional and behavioural functioning over the past six months. It includes

five subscales: Emotional Symptoms, Conduct Problems, Hyperactivity/Inattention, Peer

Relationship Problems, and Prosocial Behaviour. We used the 'externalising' (sum of the conduct

and hyperactivity scales) and 'internalising' (sum of the emotional and peer problems scales) scores

in our study.

The Revised Children's Anxiety and Depression Scale – Parent Version (RCADS) is a 47-item

questionnaire completed by parents/carers to measure the frequency of symptoms related to

anxiety and depression. It provides subscale scores for Separation Anxiety Disorder, Social Phobia,

Generalized Anxiety Disorder, Panic Disorder, Obsessive–Compulsive Disorder, and Major Depressive

Disorder. This questionnaire was only included within the HiToP model (see below).

HiToP factor scores were obtained according to previously published method [23]. Briefly,

higher-order dimensions of psychopathology within the CALM cohort were determined using

Goldberg's bass-ackward factor analytic method [39]. This method is based on multivariate

hierarchical models of 1 factor, 2 factors, 3 factors and so on, which enables the sequential, top-

down extraction of dimensions whilst retaining the shared variance among the measures (the model

ceases at the maximal number of factors that can be extracted). All six RCADS subscales were

included in the hierarchical dimensional model to comprehensively represent internalizing

difficulties. This approach was preferred over the SDQ Emotion subscale, which combines anxiety

and depression into a single score. Subscales from the CPSF were then added to capture

externalizing symptoms and neurodevelopmental difficulties, specifically Inattention,

Hyperactivity/Impulsivity, Executive Function, Aggression, and Peer Relations. CPSF was prioritized

over SDQ at this stage due to its broader coverage of neurodevelopmental symptoms. Finally, two

1 SDQ subscales—Conduct Problems and Prosocial Behaviour (reverse-coded to reflect low prosocial

2 behaviour)—were included to supplement the model with domains not captured by the other

3 measures. To avoid bias in the dimensional structure, only one indicator per symptom domain was

4 selected, ensuring balanced representation across all symptom types. See supplementary **Figure S2**

5 for the diagram of the HiTOP model used in this study. Because the RCADS questionnaire was

6 introduced later into the CALM study protocol, HiToP factor analysis was conducted for the

7 subgroup of participants with all measures available, for whom genetic data was available (N=218

8 individuals after QC filtering).

10 ***DNA extraction, genotyping, QC filtering, imputations, and PGS calculation***

11  For detailed methods, see Supporting Material. In brief, DNA samples were collected from

12 saliva using the Oragene® DNA self-collection kits (DNA Genotek Inc). DNA extraction was performed

13 using prepIT·L2P Kit (DNA Genotek Inc) following manufacturer instructions. Genome-wide SNP

14 genotyping was carried out using the Global Screening Array with shared custom-content (GSA,

15 Illumina, Inc., USA) on an iScan instrument. The raw intensity data was subjected to genotype calling,

16 extensive pre-imputation quality control and SNP imputation using an automated computational

17 workflow described previously, which includes post-imputation step to only retain autosomal SNPs

18 with minimac3 Rsq ≥0.3 (Hong et al., 2020, 2021). A total of 18 samples were excluded (see

19 Supporting Material for details). The LD pruned dataset was used for principal component analysis in

20 PLINK [42] along with the reference dataset of the 1000 Genome Project Consortium Phase 3 (The

21 1000 Genomes Project Consortium, 2015) to ancestry groups. Only European-descent samples were

22 used in the subsequent analyses, removing 40 non-European samples. Genotype imputation was

23 performed with MiniMac3 [43] software using the "Haplotype Reference Consortium" v1.1 reference

24 panel (McCarthy et al., 2016). Overall, this procedure resulted in 39,131,578 imputed SNPs.

25  We computed PGS as described previously (Choi et al., 2020). Allele status and effect-size

26 estimates were taken from the summary statistics of the largest ADHD [5] and Intelligence [8] meta-

GWAS published to date. Variant QC filtering was performed on both GWAS databases: duplicate

and ambiguous SNPs were removed, and only SNPs with MAF >0.05 and imputation quality r^2 >0.8

were included. The target CALM sample did not overlap with the GWAS datasets. GWAS and CALM

genotyping data were annotated on the GRCh37/hg19 genome build. Further 9 samples were

removed due to extreme heterozygosity (more than 3 SD from the mean), and 24 closely related

individuals were also removed, giving a final sample size of N=524. Variant matching and recoding

between the ADHD/Intelligence GWAS base and CALM target datasets was performed using R

(version 3.5.3), resulting in 5,583,323 and 3,994,342 overlapping SNPs for ADHD and Intelligence,

respectively. After SNP-clumping, PGS were computed for a variety of P-value thresholds (P_t) in the

primary GWAS data (0.001, 0.05, 0.10, 0.20, 0.30, 0.40, 0.50) using PLINK.

Statistical analysis

PGS-trait association analyses were conducted on standardized values using R (v4.5.0) for

each P_t. ADHD and Intelligence PGS were treated as independent variables in linear regression

models adjusting for sex, age, and the first 6 components of a PCA performed on the pruned SNP

pool dataset to account for population stratification. We performed sensitivity analysis including

index of multiple deprivation (IMD), a proxy for socioeconomic status (SES), as a covariate to reduce

potential bias arising from social stratification and gene–environment correlation in polygenic score

analyses [37]. As discussed by Abdellaoui and Verweij (2021), polygenic scores for behavioural traits

may capture not only direct genetic effects but also indirect genetic and environmental pathways

that are correlated with social background. SES is associated with both ADHD polygenic scores and

behavioural outcomes, partly because heritable traits related to education and cognition influence

family environments and social positioning. Adjusting for SES helps to improves the interpretability

of the estimated associations. We also performed sensitivity analysis removing participants that

have been diagnosed with ADHD to assess whether significant ADHD-PGS associations were driven

by these participants. The variance explained (r^2) was derived by comparing results from the full

model (outcome phenotype with PGS and covariates) vs the null model (outcome phenotype with

covariates only). P_t values that explained the highest phenotypic variance (max r^2) were selected.

Significant PGS-trait associations were determined after Benjamini-Hochberg adjustment to control

for false discovery rate [45], (Tables S1-S5).

We included index of multiple deprivation (IMD) a proxy for socioeconomic status (SES), as a

covariate to reduce potential bias arising from social stratification and gene–environment

correlation in polygenic score analyses.

As a post-hoc step to aid interpretability, statistical power for ADHD- and Intelligence-PGS

was estimated using AVENGEME R package [46] assuming a highly polygenic architecture. SNP-

heritability estimates were taken from the discovery GWAS (h^2 = 0.14 for ADHD; h^2 = 0.19 for

intelligence), with π set to 0.03 and 0.10 respectively. Target sample size was constant across traits

(N = 524 for behavioural and cognitive measures, N= 405 after removing ADHD diagnosed

participants, and N= 218 for the HiTOP factor analysis). Observed R^2 values from regression models

are reported alongside predicted power (Table S6).

Results

ADHD-PGS and Intelligence-PGS are associated with their respective primary phenotypes within

the CALM cohort

We first assessed the associations between ADHD and Intelligence PGS with their respective

behavioural and cognitive primary phenotypes within the transdiagnostic CALM cohort. We found

that ADHD-PGS were positively associated with Hyperactivity/Impulsivity scores (r^2= 0.03; Beta=0.17

SE=0.04; p=3.52e-05) and Inattention (r^2= 0.02; Beta=0.14 SE=0.04; p=1.57e-03) (**Fig. 1, Table S1**).

These associations remained significant after covariation with Index of Multiple Deprivation (IMD,

Table S2). Sensitivity analysis removing ADHD-diagnosed participants showed that the associations

between ADHD-PGS and ADHD traits were not solely observed within or biased by the presence of

participants with this diagnosis (**Table S3**).

Intelligence-PGS were positively associated with verbal (r^2= 0.05; Beta=0.22 SE=0.04;

p=4.00e-08) and non-verbal (r^2= 0.05; Beta=0.22 SE=0.04; p=7.41e-07) intelligence test scores (**Fig. 1,**

Table S1), and these associations remained significant after covariation with IMD (**Table S2**).

Fig. 1 ADHD-PGS and IQ-PGS associations with dimensional measures Forest plots showing
standardized effect size (Beta) and 95% CI (error bars) from the linear regressions between ADHD
and Intelligence-PGS and behavioural and cognitive measures within the CALM cohort. Vignette size
corresponds to the linear regression P values (Table S1)
19

ADHD-PGS are associated with behavioural and cognitive dimensions beyond ADHD-related

characteristics

We then analysed whether the impact of ADHD-PGS on behavioural traits can be observed

beyond the narrowly defined dimensions of inattention and hyperactivity. ADHD-PGS showed

positive association with externalising behaviours encompassing SDQ Conduct and

Hyperactivity/Inattention scales (r^2= 0.04; Beta=0.21 SE 0.04; p=1.06e-06), but not with internalising

behaviours encompassing Emotional and Peer problems scales (r^2= 0.01; Beta=0.09 SE=0.04;

p=5.19e-02) (**Fig. 1, Table S1**). With respect to cognitive dimensions, ADHD-PGS showed negative

associated with verbal (r^2= 0.02; Beta=-0.14 SE=0.04; p=1.02e-03), and non-verbal (r^2= 0.01; Beta=-

0.11 SE=0.04; p=1.57e-02) intelligence (**Fig 1, Table S1**). All these associations remained significant

after covariation with IMD (**Table S2**) or removing participants diagnosed for ADHD (**Table S3**).

Intelligence-PGS show no association with behavioural dimensions beyond IQ

Intelligence-PGS were not significantly associated with ADHD inattention or hyperactivity

scores, neither with externalising or internalising behavioural scores (**Fig. 1** and **Tables S1**; see **Table**

S2 for values covarying with IMD). Statistical power estimated from predicted variance explained

(see **Materials and Methods**) were similar for both ADHD and Intelligence PGS, with slightly higher

values for Intelligence-PGS (5.06% for ADHD-PGS and 5.11% for Intelligence-PGS, **Table S6**),

confirming that the lack of association between Intelligence-PGS and behavioural dimensions was

not due to reduced statistical power specifically for the intelligence-PGS.

ADHD-PGS contributes to hierarchical factors of mental health within the CALM cohort

To assess whether ADHD and Intelligence PGS were associated with mental health,

considered as a hierarchical tiered set of inter-related factors, we conducted regression analysis

using the HiTOP factor scores calculated for CALM participants with data availability (see **Materials**

and Methods for details). ADHD-PGS were positively associated with the level 1 general factor of

psychopathology (r^2= 0.03; Beta=0.17 SE=0.07; p=1.37e-02), the level 2 broad externalizing factor

(r^2= 0.03; Beta=0.17 SE=0.07; p=1.06e-02) and the level 3 social maladjustment factor (r^2= 0.03;

Beta=0.19 SE=0.07; p=6.42e-03) (**Fig. 2**, **Table S4**). However, these associations were no longer

significant when covarying with IMD (**Table S5**).

Intelligence-PGS showed no significant association with any of the HiTOP psychopathology

factors, with or without IMD as a covariate (**Fig. 2, Table S4 and S5**).

Consistent with a smaller sample size for the PGS-HiTOP scores association analyses,

statistical power estimates were slightly reduced (5.01% for ADHD-PGS and 5.02% for Intelligence-

PGS, **Table S6**). Again, power values were similar for both PGS, thus, the lack of associations

between the intelligence-PGS and HiTOP scores cannot be explained by a reduction in power

specifically for intelligence-PGS.

Fig. 2 ADHD-PGS and IQ-PGS associations with HiTOP factor scores Forest plots showing
standardized effect size (Beta) and 95% CI (error bars) from the linear regressions between ADHD
and Intelligence-PGS and psychopathology factors (HiTOP) within the CALM cohort. Vignette size
corresponds to the linear regression P values (Table S4)

Discussion

We examined polygenic contributions to behavioural and cognitive variation in the CALM

cohort, a clinically referred paediatric sample characterized by high levels of neurodevelopmental

heterogeneity, comorbidity and learning difficulties. This transdiagnostic design provides an

opportunity to test the generalizability of well-established polygenic associations for ADHD and

intelligence in a setting that more closely reflects the complexity encountered in clinical practice,

where diagnostic boundaries are often blurred and multiple domains of functioning are affected.

Using summary statistics from the largest available GWAS at the time of analysis, we derived

polygenic scores for ADHD and intelligence and evaluated their associations not only with

corresponding primary phenotypes (ADHD symptoms and IQ), but also with broader behavioural

dimensions assessed by the SDQ and with latent psychopathology factors derived from a hierarchical

model of psychopathology (HiTOP). Our approach moves beyond single-phenotype prediction and

directly examines the specificity and overlap of polygenic influences across multiple levels of

phenotypic organization within a complex, transdiagnostic clinical sample.

First, we found that ADHD-PGS were associated with hyperactivity-impulsivity and

inattention questionnaire scores, and Intelligence-PGS with verbal and non-verbal intelligence test

scores. This confirms that GWAS-derived metrics of common genomic variation are relevant to their

expected dimensions of neurodevelopmental variation despite the inclusive ascertainment criteria,

high levels of co-occurring diagnoses, and sub-diagnostic difficulties of the CALM cohort. Enrichment

for multi-dimensional behavioural and cognitive difficulties within the cohort means that PGS-trait

relationships can be observed despite a relatively small sample size. Although a high proportion of

the cohort had received a diagnosis of ADHD from community services prior to recruitment to CALM,

removing these participants from analyses did not alter results. These results provide a starting point

for future explorations of the associations between PGS, cognitive functions and neuroimaging

correlates, and longitudinal associations within the CALM cohort.

Second, we found that PGS derived from ADHD meta-GWAS statistics were associated with

neurodevelopmental dimensions beyond the core ADHD diagnostic symptoms, extending into

externalising difficulties and verbal intelligence. This is consistent with previous studies that have

found that higher ADHD-PGS are also associated with elevated neurodevelopmental, externalizing,

and depressive and anxiety symptoms in diagnostic [47] and population studies [12, 48]. There are

possible explanations for these results. The SDQ externalising score incorporates a hyperactivity

subscale, so the association between ADHD-PGS and externalising scores may be driven by this

subscale. The relationship between ADHD and language/verbal difficulties is well-documented both

in CALM and in the broader literature [49, 50]. A single dimension of behavioural difficulties that

includes both hyperactivity and inattention, was strongly and negatively associated with pragmatic

communication skills within the CALM cohort [50]. The case groups contributing to ADHD-GWAS

studies from which the ADHD-PGS is drawn likely included many individuals with ADHD plus

additional behavioural symptoms plus reduced verbal intelligence, meaning that the ADHD-PGS may

summate genomic variation contributing to each of these separate dimensions [5]. The symptoms

assessed by ADHD questionnaires (particularly for the inattention subscale) are diverse and

incorporate cognitive and behavioural features, meaning cross-over associations of the ADHD-PGS

with diverse neurodevelopmental dimensions. Alternatively, genomic variation contributing to risk

of ADHD symptoms may also contribute to risk of these additional phenotypes (a truly

transdiagnostic genetic effect), evidenced by the significant genetic correlations between ADHD risk

variants and IQ and educational attainment [11, 14]. Future studies could test these alternative

explanations.

With respect to internalizing symptoms, we observed a weak positive association with

ADHD-PGS that did not reach statistical significance. In contrast, Schlag et al. (2022) reported a

robust association between ADHD-PGS and peer problems, a specific subscale of internalizing

symptoms construct measured using the SDQ [51]. Their analyses were conducted in two large UK

population-based cohorts (ALSPAC, N = 6,174; TEDS, N = 7,112), providing substantially greater

power to detect polygenic effects than in our sample. Although non-significant, the direction of

association in our study was consistent with their findings. Importantly, the CALM cohort differs

from these population-based samples in that it is transdiagnostic and enriched for children with

learning difficulties, which may introduce greater phenotypic heterogeneity. When participants with

an ADHD diagnosis were excluded—thereby creating a sample that more closely resembles a general

population-based study with low rate of clinical symptoms—the association between ADHD-PGS and

internalizing symptoms reached statistical significance despite the reduced sample size (**Table S3**).

Taken together these findings highlight that differences in sample composition, rather than power

alone, may modulate detectability of genetic and behavioural associations, and suggest that ADHD-

PGS contribution to internalizing behaviours may be sample dependent, reflecting complex

influences that require further investigation.

12   Contrary to our transdiagnostic effects showed by the ADHD-PGS, Intelligence-PGS showed a

relationship only with verbal and non-verbal IQ test scores and did not cross-associate with any

behavioural measures within the CALM sample. Several features of the CALM sample may contribute

to this pattern. The cohort is clinically referred and enriched for learning difficulties, with high

comorbidity and potentially restricted variation in cognitive outcomes, which may limit the

detectability of cross-trait associations in a modestly sized sample. Whilst ADHD-PGS have been

shown to correlate with intelligence and educational attainment in population-based cohorts [11,

14], the reverse relationship between cognitive abilities and mental health seems to be strongly

influenced by environmental factors [52]. Additionally, cross-trait prediction in deeply phenotyped

but smaller cohorts inherently yield smaller effect sizes compared with within-trait prediction in

large GWAS. However, the lack of intelligence-PGS and ADHD-traits associations cannot be explained

by a reduction in intelligence-PGS statistical power, as both PGS showed similar power values when

estimated based on predicted variance explained using the R package AVENGEME [46] (Suppl **Table**

S6). Although the power estimates indicated limited ability to detect the expected polygenic signal in

our target sample, this reflects conservative assumptions derived from GWAS architecture and

same-trait prediction. Our analyses instead focused on cross-trait, dimensional associations in a

deeply phenotyped cohort, with an observed variance explained within the range reported for

dimensional ADHD symptom scores (~0.7–3.3%) [53]. Finally, the observed pattern of effects—

positive associations with externalizing and inattention measures and negative associations with

cognitive traits—was directionally consistent with known genetic correlations of ADHD [14],

supporting the interpretation of partially separable polygenic contributions to cognitive and

behavioural domains.

Using the transdiagnostic CALM cohort, we tested whether ADHD polygenic risk is associated

with general psychopathology, more specific externalizing or neurodevelopmental dimensions, or

across multiple hierarchical levels in childhood. In addition, by examining ADHD-PGS alongside

Intelligence-PGS within the same HiTOP framework, we directly compared the breadth and

specificity of behavioural and cognitive polygenic effects, extending prior work beyond single-liability

analyses. Here again, we found that Intelligence-PGS may not have predictive value in explaining

psychopathological variation within the sample. These findings seem not to be caused by a reduction

in power specifically for intelligence-PGS, as power values were similar for both ADHD and

Intelligence-PGS (**Table S6**). ADHD-PGS analyses in contrast, suggest that there are polygenic

contributions to an overarching p factor within CALM, indexing general susceptibility to emotional-

behavioural symptoms as has been shown previously [12, 24]. Interestingly, we did not detect any

association between ADHD-PGS and the lower-level "neurodevelopmental" factor which

encompasses inattention, hyperactivity/impulsivity and executive function. Instead, ADHD-PGS were

associated with "social maladjustment", encompassing conduct-related and peer-related problems

assessed by the SDQ. We speculate that this observed association may be explained by the high

comorbidity between ADHD and conduct disorders [2]. A within-case analysis of individuals with

comorbid ADHD and conduct disorders showed a positive linear relationship between ADHD-PGS

and conduct problems, especially aggression [54]. Alternatively, the relationship between ADHD-PGS

and verbal abilities, rather than a direct effect on behavioural characteristics, may have also

contributed to these observations. The importance of communication abilities (pragmatic skills in

particular) and their relationship with behavioural symptomatology have been emphasised in

previous CALM analyses [50], and future hypothesis-driven studies could investigate the links

between ADHD-PGS and language development. However, it is important to note that the sample

available for PGS-HiTop analysis was smaller, with the concomitant reduction in power, thus, the

specificity of associations should be treated with caution.

We found that the PGS associations with cognitive and behavioural measures were

insensitive to socioeconomic status approximated using IMD. These results perhaps reflect the

limited range of IMD within the CALM sample (Figure S1), with an overage above the national

median, and are consistent with previous work conducted in large cohort showing that genetic

influence on educational and cognitive outcomes is larger within more socioeconomically deprived

contexts [55]. Associations between ADHD-PGS and HiTop psychopathology factors were weakened

(or rendered non-significant) when co-varying for IMD. This may indicate that the observed

associations are artefacts of co-variation, or that the sample is under-powered to detect significant

associations once co-variates are accounted for, or alternatively that there are important gene x

environment interactions at play. Although strongly influential, SES is only one of the environmental

factors that contribute to behavioural and environmental outcomes [37]. Future studies could

explore gene x environment interactions within this sample by assessing the impact of other

important environmental factors, such as parental education and perinatal health, highly relevant in

neurodevelopmental outcomes [56].

In summary, we have investigated a unique transdiagnostic sample and found that PGS

contribute significantly and specifically to behavioural characteristics and cognitive abilities. This is a

first step toward investigating links between genomic variation, brain development and cognitive

processes contributing to learning and behavioural problems within transdiagnostic populations.

1   **Statements and Declarations**

Competing Interests

The authors have no relevant financial or non-financial interests to disclose.

Funding

D. Astle and K. Baker are supported by the Medical Research Council (MC_UU_00030/2;

MC_UU_00030/3). A.M. Santangelo was supported by the Isaac Newton Trust / Wellcome Trust ISSF

/ University of Cambridge Joint Research Grants scheme, and by the Medical Research Council

(MC_UU_00030/3 and MC_UU_00032/3). All research at the Department of Psychiatry in the

University of Cambridge is supported by the National Institute for Health and Care Research

Cambridge Biomedical Research Centre (NIHR203312) and the NIHR Applied Research Collaboration

East of England.

Data Availability

Behavioural and cognitive data are available online through the CALM Managed Open Access

repository: https://portal.ide-cam.org.uk/overview/1158. Genetic data are available via

collaboration with CALM.

Ethics

All procedures complied with the ethical standards of the national and institutional committees on

human experimentation and with the Helsinki Declaration of 1975, as revised in 2008. All procedures

involving human participants were approved by the National Health Service, Health Research

Authority NRES Committee East of England (REC: 13/EE/0157, IRAS 127675).

Consent to participate

Informed consent was obtained from legal guardians.

Authors' contributions

Conceptualization (KB, DA, DB, AS); Formal analysis (AS); Funding acquisition (KB); Investigation (JH,

DA, SM, CALM team); Resources (OO, LB); Writing – original draft (AS, KB); Writing – review & editing

(all authors).

References

1.   Astle DE, Holmes J, Kievit R, Gathercole SE (2022) Annual Research Review: The
transdiagnostic revolution in neurodevelopmental disorders. J Child Psychol Psychiatry
63:397–417. https://doi.org/10.1111/JCPP.13481

2.   Posner J, Polanczyk G V., Sonuga-Barke E (2020) Attention-deficit hyperactivity disorder.
Lancet 395:450–462. https://doi.org/10.1016/S0140-6736(19)33004-1

3.   Faraone S V., Perlis RH, Doyle AE, et al (2005) Molecular genetics of attention-
deficit/hyperactivity disorder. Biol Psychiatry 57:1313–1323.
https://doi.org/10.1016/J.BIOPSYCH.2004.11.024

4.   Jensen CM, Steinhausen HC (2015) Comorbid mental disorders in children and adolescents
with attention-deficit/hyperactivity disorder in a large nationwide study. Atten Defic
Hyperact Disord 7:27–38. https://doi.org/10.1007/S12402-014-0142-1

5.   Demontis D, Walters GB, Athanasiadis G, et al (2023) Genome-wide analyses of ADHD identify
27 risk loci, refine the genetic architecture and implicate several cognitive domains. Nature
Genetics 2023 55:2 55:198–208. https://doi.org/10.1038/s41588-022-01285-8

6.   Deary IJ, Cox SR, Hill WD (2021) Genetic variation, brain, and intelligence differences.
Molecular Psychiatry 2021 27:1 27:335–353. https://doi.org/10.1038/s41380-021-01027-y

7.   de la Fuente J, Davies G, Grotzinger AD, et al (2020) A general dimension of genetic sharing
across diverse cognitive traits inferred from molecular data. Nature Human Behaviour 2020
5:1 5:49–58. https://doi.org/10.1038/s41562-020-00936-2

8.   Savage JE, Jansen PR, Stringer S, et al (2018) Genome-wide association meta-analysis in
269,867 individuals identifies new genetic and functional links to intelligence. Nat Genet
50:912–919. https://doi.org/10.1038/S41588-018-0152-6

9.   Andlauer TFM, Nöthen MM (2020) Polygenic scores for psychiatric disease: From research
tool to clinical application. Medizinische Genetik 32:39–45.
https://doi.org/10.1515/MEDGEN-2020-2006/ASSET/GRAPHIC/J_MEDGEN-2020-
2006_FIG_002.JPG

10.   Choi SW, Mak TSH, O'Reilly PF (2020) Tutorial: a guide to performing polygenic risk score
analyses. Nat Protoc 15:2759–2772. https://doi.org/10.1038/S41596-020-0353-1

11.   Demontis D, Walters RK, Martin J, et al (2019) Discovery of the first genome-wide significant
risk loci for attention deficit/hyperactivity disorder. Nat Genet 51:63–75.
https://doi.org/10.1038/S41588-018-0269-7

12.   Brikell I, Larsson H, Lu Y, et al (2020) The contribution of common genetic risk variants for
ADHD to a general factor of childhood psychopathology. Mol Psychiatry 25:1809–1821.
https://doi.org/10.1038/S41380-018-0109-2

13. Martin J, Hamshere ML, Stergiakouli E, et al (2014) Genetic Risk for Attention-Deficit/Hyperactivity Disorder Contributes to Neurodevelopmental Traits in the General Population. Biol Psychiatry 76:664. https://doi.org/10.1016/J.BIOPSYCH.2014.02.013

14. van der Laan CM, Ip HF, Schipper M, et al (2025) Genome-wide association meta-analysis of childhood ADHD symptoms and diagnosis identifies new loci and potential effector genes. Nat Genet 57:2427–2435. https://doi.org/10.1038/S41588-025-02295-Y

15. Lee PH, Anttila V, Won H, et al (2019) Genomic Relationships, Novel Loci, and Pleiotropic Mechanisms across Eight Psychiatric Disorders. Cell 179:1469-1482.e11. https://doi.org/10.1016/J.CELL.2019.11.020

16. Gidziela A, Ahmadzadeh YI, Michelini G, et al (2023) A meta-analysis of genetic effects associated with neurodevelopmental disorders and co-occurring conditions. Nature Human Behaviour 2023 7:4 7:642–656. https://doi.org/10.1038/s41562-023-01530-y

17. Romero C, Werme J, Jansen PR, et al (2022) Exploring the genetic overlap between twelve psychiatric disorders. Nature Genetics 2022 54:12 54:1795–1802. https://doi.org/10.1038/s41588-022-01245-2

18. Ronald A, de Bode N, Polderman TJC (2021) Systematic Review: How the Attention-Deficit/Hyperactivity Disorder Polygenic Risk Score Adds to Our Understanding of ADHD and Associated Traits. J Am Acad Child Adolesc Psychiatry 60:1234–1277. https://doi.org/10.1016/J.JAAC.2021.01.019

19. Waszczuk MA, Eaton NR, Krueger RF, et al (2020) Redefining phenotypes to advance psychiatric genetics: Implications from hierarchical taxonomy of psychopathology. J Abnorm Psychol 129:. https://doi.org/10.1037/ABN0000486

20. Allegrini AG, Cheesman R, Rimfeld K, et al (2020) The p factor: genetic analyses support a general dimension of psychopathology in childhood and adolescence. J Child Psychol Psychiatry 61:30–39. https://doi.org/10.1111/JCPP.13113

21. Kotov R, Waszczuk MA, Krueger RF, et al (2017) The Hierarchical Taxonomy of Psychopathology (HiTOP): A dimensional alternative to traditional nosologies. J Abnorm Psychol 126:454–477. https://doi.org/10.1037/ABN0000258

22. Michelini G, Barch DM, Tian Y, et al (2019) Delineating and validating higher-order dimensions of psychopathology in the Adolescent Brain Cognitive Development (ABCD) study. Transl Psychiatry 9:. https://doi.org/10.1038/S41398-019-0593-4

23. Holmes J, Mareva S, Bennett MP, et al (2021) Higher-Order Dimensions of Psychopathology in a Neurodevelopmental Transdiagnostic Sample. J Abnorm Psychol 130:909–922. https://doi.org/10.1037/abn0000710

24. Waszczuk MA, Miao J, Docherty AR, et al (2023) General v. specific vulnerabilities: polygenic risk scores and higher-order psychopathology dimensions in the Adolescent Brain Cognitive Development (ABCD) Study. Psychol Med 53:1937–1946. https://doi.org/10.1017/S0033291721003639

25. Riglin L, Thapar AK, Leppert B, et al (2020) Using Genetics to Examine a General Liability to Childhood Psychopathology. Behav Genet 50:213–220. https://doi.org/10.1007/S10519-019-09985-4

26. Pat N, Riglin L, Anney R, et al (2022) Motivation and Cognitive Abilities as Mediators Between Polygenic Scores and Psychopathology in Children. J Am Acad Child Adolesc Psychiatry 61:782-795.e3. https://doi.org/10.1016/J.JAAC.2021.08.019

27. Holmes J, Bryant A, Gathercole SE (2019) Protocol for a transdiagnostic study of children with problems of attention, learning and memory (CALM). BMC Pediatr 19:. https://doi.org/10.1186/S12887-018-1385-3

28. Mareva S, Akarca D, Astle D, et al (2023) Transdiagnostic profiles of behaviour and communication relate to academic and socioemotional functioning and neural white matter organisation. J Child Psychol Psychiatry 64:. https://doi.org/10.1111/JCPP.13685

29. Williams KL, Holmes J, Farina F, et al (2022) Inconsistencies between Subjective Reports of Cognitive Difficulties and Performance on Cognitive Tests are Associated with Elevated Internalising and Externalising Symptoms in Children with Learning-related Problems. Res Child Adolesc Psychopathol 50:1557–1572. https://doi.org/10.1007/S10802-022-00930-4

30. Bryant A, Guy J, Holmes J (2020) The Strengths and Difficulties Questionnaire Predicts Concurrent Mental Health Difficulties in a Transdiagnostic Sample of Struggling Learners. Front Psychol 11:. https://doi.org/10.3389/FPSYG.2020.587821

31. Bathelt J, Holmes J, Astle DE, et al (2018) Data-Driven Subtyping of Executive Function-Related Behavioral Problems in Children. J Am Acad Child Adolesc Psychiatry 57:252-262.e4. https://doi.org/10.1016/J.JAAC.2018.01.014

32. Holmes J, Guy J, Kievit RA, et al (2021) Cognitive Dimensions of Learning in Children With Problems in Attention, Learning, and Memory. J Educ Psychol 113:1454–1480. https://doi.org/10.1037/edu0000644

33. Mareva S, Holmes J (2019) Transdiagnostic associations across communication, cognitive, and behavioural problems in a developmentally at-risk population: A network approach. BMC Pediatr 19:. https://doi.org/10.1186/s12887-019-1818-7

34. Guy J, Mareva S, Franckel G, Holmes J (2022) Dimensions of cognition, behaviour, and mental health in struggling learners: A spotlight on girls. JCPP advances 2:. https://doi.org/10.1002/JCV2.12082

35. Akarca D, Vértes PE, Bullmore ET, et al (2021) A generative network model of neurodevelopmental diversity in structural brain organization. Nat Commun 12:. https://doi.org/10.1038/s41467-021-24430-z

36. Siugzdaite R, Bathelt J, Holmes J, Astle DE (2020) Transdiagnostic Brain Mapping in Developmental Disorders. Current Biology 30:1245-1257.e4. https://doi.org/10.1016/j.cub.2020.01.078

37. Abdellaoui A, Verweij KJH (2021) Dissecting polygenic signals from genome-wide association studies on human behaviour. Nature Human Behaviour 2021 5:6 5:686–694. https://doi.org/10.1038/s41562-021-01110-y

38. (2019) The English Indices of Deprivation 2019

39. Goldberg LR (2006) Doing it all Bass-Ackwards: The development of hierarchical factor structures from the top down. J Res Pers 40:347–358. https://doi.org/10.1016/J.JRP.2006.01.001

40. Hong S, Dobricic V, Ohlei O, et al (2021) TMEM106B and CPOX are genetic determinants of cerebrospinal fluid Alzheimer's disease biomarker levels. Alzheimers Dement 17:1628–1640. https://doi.org/10.1002/ALZ.12330

41. Hong S, Prokopenko D, Dobricic V, et al (2020) Genome-wide association study of Alzheimer's disease CSF biomarkers in the EMIF-AD Multimodal Biomarker Discovery dataset. Transl Psychiatry 10:. https://doi.org/10.1038/S41398-020-01074-Z

42. Purcell S, Neale B, Todd-Brown K, et al (2007) PLINK: a tool set for whole-genome association and population-based linkage analyses. Am J Hum Genet 81:559–575. https://doi.org/10.1086/519795

43. Das S, Forer L, Schönherr S, et al (2016) Next-generation genotype imputation service and methods. Nat Genet 48:1284–1287. https://doi.org/10.1038/NG.3656

44. McCarthy S, Das S, Kretzschmar W, et al (2016) A reference panel of 64,976 haplotypes for genotype imputation. Nat Genet 48:1279–1283. https://doi.org/10.1038/NG.3643

45. Lee S, Lee DK (2018) What is the proper way to apply the multiple comparison test? Korean J Anesthesiol 71:353–360. https://doi.org/10.4097/KJA.D.18.00242

46. Dudbridge F (2013) Power and predictive accuracy of polygenic risk scores. PLoS Genet 9:. https://doi.org/10.1371/JOURNAL.PGEN.1003348

47. Stergiakouli E, Martin J, Hamshere ML, et al (2015) Shared genetic influences between attention-deficit/hyperactivity disorder (ADHD) traits in children and clinical ADHD. J Am Acad Child Adolesc Psychiatry 54:322–327. https://doi.org/10.1016/J.JAAC.2015.01.010

48. Du Rietz E, Coleman J, Glanville K, et al (2018) Association of Polygenic Risk for Attention-Deficit/Hyperactivity Disorder With Co-occurring Traits and Disorders. Biol Psychiatry Cogn Neurosci Neuroimaging 3:635–643. https://doi.org/10.1016/j.bpsc.2017.11.013

49. Korrel H, Mueller KL, Silk T, et al (2017) Research Review: Language problems in children with Attention-Deficit Hyperactivity Disorder - a systematic meta-analytic review. J Child Psychol Psychiatry 58:640–654. https://doi.org/10.1111/JCPP.12688

50. Hawkins E, Gathercole S, Astle D, Holmes J (2016) Language Problems and ADHD Symptoms: How Specific Are the Links? Brain Sci 6:. https://doi.org/10.3390/BRAINSCI6040050

51. Schlag F, Allegrini AG, Buitelaar J, et al (2022) Polygenic risk for mental disorder reveals distinct association profiles across social behaviour in the general population. Mol Psychiatry 27:1588–1598. https://doi.org/10.1038/S41380-021-01419-0

52. Wang Y, Anney R, Pat N (2025) The relationship between cognitive abilities and mental health as represented by cognitive abilities at the neural and genetic levels of analysis. Elife 14:. https://doi.org/10.7554/ELIFE.105537.2

53. Ronald A, de Bode N, Polderman TJC (2021) Systematic Review: How the Attention-Deficit/Hyperactivity Disorder Polygenic Risk Score Adds to Our Understanding of ADHD and Associated Traits. J Am Acad Child Adolesc Psychiatry 60:1234–1277. https://doi.org/10.1016/J.JAAC.2021.01.019

54.　Hamshere ML, Langley K, Martin J, et al (2013) High loading of polygenic risk for ADHD in
children with comorbid aggression. American Journal of Psychiatry 170:909–916.
https://doi.org/10.1176/APPI.AJP.2013.12081129/ASSET/IMAGES/LARGE/909F1.JPEG

55.　Rask-Andersen M, Karlsson T, Ek WE, Johansson Å (2021) Modification of Heritability for
Educational Attainment and Fluid Intelligence by Socioeconomic Deprivation in the UK
Biobank. Am J Psychiatry 178:625–634. https://doi.org/10.1176/APPI.AJP.2020.20040462

56.　Leffa DT, Caye A, Belangero SI, et al (2023) The synergistic effect of genetic and
environmental factors in the development of attention-deficit/hyperactivity disorder
symptoms in children and adolescents. Dev Psychopathol 1–11.
https://doi.org/10.1017/S0954579423000366

57.　Conners CK (2008) Conners parent rating scale short form, 3rd ed. Pearson, London, UK

58.　Goodman R (1997) The strengths and difficulties questionnaire: A research note. J Child
Psychol Psychiatry 38:581–586. https://doi.org/10.1111/J.1469-7610.1997.TB01545.X

59.　(2005) Wechsler Individual Achievement Test Second UK Edition (WIAT-II UK). Wechsler
Individual Achievement Test Second UK Edition (WIAT-II UK).
https://doi.org/10.53841/BPSTEST.2005.WIAT2

60.　Dunn LM, Dunn DM (2007) Peabody Picture Vocabulary Test. Pearson Education,
Minneapolis, USA.

61.　Chorpita BF, Yim L, Moffitt C, et al (2000) Assessment of symptoms of DSM-IV anxiety and
depression in children: a revised child anxiety and depression scale. Behaviour research and
therapy 38:835–855. https://doi.org/10.1016/S0005-7967(99)00130-8

Table 1: Cognitive and behavioural measures, and HiTOP factor scores, for the CALM sample included in the PGS analyses (after QC filtering)

Measures	Instrument	Variable	Mean	SD	Range
Behavioural (N=524)					
Hyperactivity/Impulsivity	Conners 3 - Parent Rating Scale Short Form[1]	Raw scores	10.98	5,59	0-18
Inattention	(CPSF)		11.77	3.44	0-16
Externalising behaviours	Strengths and Difficulties Questionnaire[2]	Sum of Conduct and Hyperactivity/Inattention scales	11.11	4.12	1-20
Internalising behaviours	(SDQ)	Sum of Emotional and Peer problems scales	7.85	4.62	0-19
Cognitive (N=524)					
Non-verbal intelligence	Wechsler Abbreviated Scales of Intelligence II[3]	Matrix Reasoning subtest-T scores	43.72	9.66	20-80
Verbal intelligence	Peabody Picture Vocabulary Test[4]	Number of correct items- percentile	48.93	29.36	0.09-99.9
HiTOP (N=218)					
Level 1:					
P factor	Conners 3 - Parent Rating Scale Short Form[1]		0.01	1.02	-2.22-3.31
Level 2:	(CPSF),				
Broad internalizing	Strengths and Difficulties Questionnaire[2]		-0.01	1.01	-1.88-3.82
Broad externalizing	(SDQ),	Factor scores[5]	0.03	1.01	-2.59-2.23
Level 3:	and Revised Children's Anxiety and				
Specific internalizing	Depression Scale Parent Version[6]		-0.01	1.01	-1.74-3.79
Social maladjustment	(RCADS-P)		-0.01	1.03	-1.70-2.57
Neurodevelopmental			0.08	1.00	-3.74-1.56

[1] [57]; [2] [58]; [3] [59]; [4] [60]; [5] [23]; [6] [61]

Table 2: Description of the CALM sample after QC filtering

Sample size	Tota	Males/Females
N	524	361/163
Diagnosis	**Percentage of total**	**N**
Any developmental or psychiatric disorder	38%	199
ADHD	22.5%	118
Dyslexia/speech therapy	11.8%	34/28
Autism	7%	37
	Mean±SD	**Range**
Age (years)	9.5±2.4	5.2-18.6
Index of Multiple Deprivation (IMD)	20328±8254	155-32803