1   **Impact of a multi-pronged cholera intervention in an endemic setting**

2   Short title: Short-term impact of OCV and WASH in a cholera endemic area

3   Alexandre Blake[1], Adam Walder[2], Ephraim Hanks[2], Placide Okitayembo Welo[3], Francisco Luquero[4], Didier

4   Bompangue[3,5], Nita Bharti[1]

5   Corresponding author: Nita Bharti, nita@psu.edu

Affiliations

1: Biology Department, Center for Infectious Disease Dynamics, Penn State University, University Park,

Pennsylvania, United States of America

2: Statistics Department, Center for Infectious Disease Dynamics, Penn State University, University Park,

Pennsylvania, United States of America

3: Programme National d'Elimination du Choléra et de lutte contre les autres Maladies Diarrhéiques,

Kinshasa, Democratic Republic of the Congo

4: Epicentre, Paris, France

5: Department of Ecology and Control of Infectious Diseases, Faculty of Medicine, University of Kinshasa,

Kinshasa, Democratic Republic of the Congo

Abstract

Cholera is a bacterial water-borne diarrheal disease that causes high morbidity in sub-Saharan Africa and

Asia. It is transmitted via the fecal-oral route and can be prevented with vaccination and Water,

Sanitation, and Hygiene (WASH) improvements. WASH improvements are resource and time intensive

but provide long lasting benefits. Vaccination campaigns can be implemented more rapidly than WASH

but provide only temporary immunity; cholera vaccines are effective in epidemic settings but their utility

in endemic settings is unclear.

The city of Kalemie in the Democratic Republic of Congo, on the shore of Lake Tanganyika, is a cholera

endemic area with annual outbreaks. Both seasonal mobility and the lake, which is a bacterial reservoir,

promote transmission, though their relative contributions are unknown. In 2013-2016 Kalemie received

a targeted vaccination campaign and WASH improvements. We assessed the short-term impact of this

intervention to guide future control strategies in endemic settings.

We fit a Susceptible-Infected-Recovered-Susceptible model with a compartment for the aquatic

bacterial population. We estimated the number of cases avoided by each arm of the intervention,

explored alternative vaccination strategies, and investigated the relative contributions of mobility and

environmentally-based transmission in local cholera dynamics.

We estimated that vaccination and WASH improvements prevented 3,484 cases (95% Credible Interval:

2,412-4,833) over a 118-week period. We showed that vaccination could prevent more cases by altering

the timing and increasing the target population size. We found that transmission was primarily

environmentally-driven and that removing environmental exposure or reducing environmental

contamination could decrease local transmission.

Together, the targeted nature of the vaccination campaign, the modest scale of WASH improvements,

and the high background immunity of the population limited the impact of the intervention. Constant

environmentally-driven force of infection maintains high levels of cholera immunity in the population

and decreases the impact of vaccination in this endemic area.

Author summary

Cholera is a major global health concern that causes high morbidity. It is a bacterial water-borne disease

that can be transmitted via the fecal-oral route or the ingestion of contaminated water. Hence, both

population mobility and environmental exposure can promote cholera persistence. The primary tools to

prevent cholera include vaccination and Water, Sanitation, and Hygiene (WASH) improvements. The

effectiveness of these interventions is well understood in epidemic settings, but their impact in endemic

settings is unclear. Achieving cholera elimination requires disentangling the contributors to

transmission, specifically population mobility and aquatic reservoirs, and assessing the impact of

interventions performed in endemic settings.

This study focuses on Kalemie, a cholera endemic city in the Democratic Republic of Congo, on shore of

a lake that serves as a potential environmental reservoir. It quantifies the short-term impact of an

intervention that used targeted vaccination and WASH. The study shows that the impact of vaccination

was dampened by very high background immunity due to constant environmental exposure. This

suggests that WASH improvements should be the primary intervention in such settings despite the time-

and resource-intensive nature of implementation.

Introduction

Cholera is a bacterial, water-borne diarrheal disease. Typical cholera symptoms include vomiting and

diarrhea with rice-water stools, potentially leading to severe dehydration. The mortality rate can reach

70% among severe cases if appropriate treatment, mainly rehydration, is not provided (1). However,

individual symptoms can vary widely depending on age and prior exposures, ranging from asymptomatic

infections, to mild infections with symptoms that are hardly distinguishable from other diarrheal

diseases, to the characteristic severe watery diarrhea (2). As many as 80% of infections can be

asymptomatic in endemic areas (1), resulting in underestimates of cholera burden.

Cholera is transmitted through the fecal-oral route. The causal agent, *Vibrio cholerae* (*V. cholerae*),

survives in aquatic environments and is present in the excreta (stools and vomit) of infected individuals.

Infection is acquired by ingesting a sufficient bacterial load from the environment (indirect transmission)

or contact with infectious excreta (direct transmission). *V. cholerae* abundance in aquatic reservoirs

varies through interactions with biotic and abiotic factors. Aquatic flora, phytoplankton, and fauna,

including copepods, have been described to modulate *V. cholerae* aquatic populations. In parallel,

environmental parameters including water temperature and salinity also influence the *V. cholerae* life

cycle in its aquatic reservoir (3–8). Viable *V. cholerae* can persist in the environment in suboptimal

conditions for over 15 months in a non-culturable state (7), from which it can revert to a culturable state

in favorable conditions. There are over 200 serogroups of *V. cholerae*, but only the serogroups O1 and

O139 can cause the disease cholera (9). They are often introduced in natural or manmade water

reservoirs through inappropriate waste management (6,10,11). Consuming contaminated water can

then trigger outbreaks, potentially leading to additional contamination of water reservoirs. In non-

endemic areas, the environmental contribution to cholera transmission is often low, but in endemic

areas the relative contribution of direct and indirect transmission routes is unknown. This could make

the benefits we would expect from interventions, as traditionally implemented in outbreak response,

less clear in such settings because they do not necessarily target the dominant transmission route.

The 7th cholera pandemic has been ongoing since 1961. In addition to its traditional endemic area in the

Bengal region, cholera established endemic areas in sub-Saharan Africa (SSA) following its introduction

in 1970 (12). SSA now experiences the greatest annual global morbidity and mortality of cholera (13),

excluding major epidemic events that occurred in Haiti and Yemen. The Global Task Force on Cholera

Control (GTFCC) has set a road map to eliminate cholera in 20 endemic countries by 2030 (14), defining

SSA as an important target. Generally, diseases or pathogens are considered 'endemic' when they

display persistent local transmission for an extended period of time in a geographic area. For cholera

specifically, the World Health Organization (WHO) defines a nation as endemic when local transmission

caused cases in the previous three years. This definition encompasses a wide variety of transmission

patterns.

Cholera transmission can be prevented by improving water and sanitation infrastructures and with

vaccination. Water, sanitation, and hygiene (WASH) improvements have historically been the primary

tool to prevent cholera. WASH improvements are resource intensive and implementation requires time

to prevent cholera (15). They are extremely effective, as waste management and water infrastructures

have largely prevented cholera transmission in high income countries (16). Cholera cannot be eliminated

from an area without WASH improvements (17), however resource scarcities limit such improvements in

the countries carrying most of the global cholera burden. Sub-Saharan African nations have some of the

poorest access to clean water and improved toilets in the world as well as the greatest disparities in

infrastructures between rural and urban areas (18). The scale of the WASH improvements necessary to

address cholera in the SSA region is vast.

In comparison, implementing a vaccination campaign is fast and can reduce cholera transmission

quickly. The empirical results of the reactive use of oral cholera vaccines (OCV) in 2012 in Guinea and

theoretical results from modeling studies demonstrated the potential utility of vaccination as a tool for

preventing cholera quickly (19,20). A quick vaccine rollout leads to a fast increase in population

immunity that can mitigate cholera transmission, but this is a short term benefit because the acquired

protection declines after about 2-3 years (21). The increasing stockpile of OCV allowed for more

frequent use of vaccines in outbreak response and its novel use in endemic areas (22,23).

Both OCV and WASH improvements are important components of the multisectoral interventions

required to achieve cholera control or elimination in areas with high cholera burden (14,23). However,

the benefit of OCV is straightforward only in epidemic settings (22,24–26). The short-term impact of

OCV correlates with the resulting increase of the population immunity, but this benefit might be narrow

in an endemic setting where exposure to cholera is frequent. As such, quantifying the impact of

interventions using OCV in endemic settings will provide valuable information to inform control

strategies and achieve the ambitious goals set by the GTFCC.

Within SSA, Cholera is endemic in the African Great Lakes region. The Democratic Republic of Congo

(DRC) has consistently carried one of the highest cholera burdens in this region (13). Cholera is endemic

in the city of Kalemie in Tanganyika Province in DRC; the city lies on the shore of Lake Tanganyika (Figure

1A and Figure 1B). The area displays annual peaks of cholera cases, typically during rainy seasons (Figure

1C), and reports suspected cholera cases all year. Lake Tanganyika is a potential environmental reservoir

that likely promotes cholera persistence. In parallel, the local population is highly mobile with 24.7% of

the residents of the Tanganyika Province reporting travelling at least once in the previous 12 months for

a duration of at least 1 month (27). There is also strong fishing activity associated with the lake, with

fishermen moving seasonally and experiencing exposure to the lake and low sanitation conditions (28).

These population factors could promote cholera persistence through metapopulation dynamics (28).

The city of Kalemie received an intervention in 2013-2016 that included both an OCV campaign and

WASH improvements. The health system in DRC is organized in several levels in increasingly smaller

geographical units: Provinces, health zones, and health areas. Public health interventions are often

organized and implemented at health zone level. The vaccination campaign in Kalemie intended to

target about 120,000 people with two doses of OCV, Shanchol™, and focused on the four health areas

where the attack rates had historically been the highest as of November 2013. The vaccination campaign

was interrupted after 3 days due to security issues and resumed in July 2014 after redefining the target

population. The expiration of vaccine doses ultimately led to reducing the target to about 52,000 people

in two health areas (29). The WASH intervention focused on improving access to clean water. Doctors

Without Borders (*Médecins Sans Frontières*, MSF) extended access to tap water in the northern part of

the city and laid pipes, built water reservoirs, distributed water filters, and set up public drinking

fountains in collaboration with *Solidarites International*. In addition, sand filters were installed on paths

where people draw water from the lake, and chlorination activities were performed during outbreaks.

The WASH intervention incurred delays in the aftermath of the security issue, but was completed. We fit

a compartmental model including both interhuman and environmental cholera transmission to assess

the short-term impact of this multi-pronged intervention in Kalemie. We considered the potential

influence of seasonal migration and environmental drivers and their relative contributions to local

transmission.

Figure 1: Overview of the location and the seasonality of cholera cases in the study area. (A) Map of the DRC with

population density (with log10(population+1) transformation), the boundaries of the health zones in white, major

roads in light green, a red box around the health zones of Kalemie and Nyemba (seen in B), a red circle on the city of

Kalemie, and Lake Tanganyika in blue. (B) Detailed map of the city of Kalemie (red circle), which sits across the two

health zones of Kalemie and Nyemba (dark red boundaries), and population density (low in grey and high in red). (C)

Boxplot of the total number of reported suspected cholera cases by week (based on the International Organization for

Standardization (ISO) system) in the health zones of Kalemie and Nyemba from 2002 to 2014 (30), with week number

and the typical rainy season weeks shaded in grey.

Results

The compartmental model with interhuman and environmental transmission reproduced the reported

weekly cholera cases reasonably well, although it struggled to capture the magnitude of the peaks in

cases (Figure 2A). The model indicated high local immunity, fluctuating between 93.2% and 99.5%

(Figure 2B). This high immunity is the likely consequence of annual outbreaks and persistent

environmental exposure, which we explain further below. The targeted vaccinations occurred at times

when population immunity was at 98.2% (95% credible interval (CrI): 97.6-98.7) in November 2013,

93.4% (95% CrI: 91.3-95.4) in July 2014, and 93.4% (95% CrI: 91.3-95.3) during the catch-up in August

2014.

Figure 2: Incident cases and model fit and variation of the percentage of infected, recovered, and susceptible over

time based on the model. (A) Weekly reported suspected cholera cases residing in the city of Kalemie (empty circles)

from November 2013 to February 2016 and mean prediction of the reported weekly cholera cases by the model (dark

line) and its 95% credible interval (grey envelope) (B) Mean prediction of the percent of the population infected

(prevalence), recovered, and susceptible (dark lines) and their 95% credible interval (grey envelopes) by the model

from November 2013 to February 2016. Typical rainy seasons are shaded in grey and the timing of the distribution of

vaccine doses in vertical dashed grey lines.

All the scenarios omitting vaccination (WASH only, and no WASH and no vaccination) visibly lacked a

reduction in the susceptible proportion of the population in July 2014 (Figure 3A). Over this period of

118 weeks, we estimated an average number of cases avoided of: 2,623 (mean: 2,626.6, 95% CrI:

1,910.2-3,391.6) cases avoided by vaccination alone (scenario with WASH only), 881 (mean: 881.3, 95%

CrI: 94.9-1,764.1) cases avoided by WASH alone (scenario with vaccination only), and 3,484 (mean:

3,484.4, 95% CrI: 2,411.7-4,832.8) cases avoided by implementing both vaccination and WASH (scenario

with no vaccination and no WASH improvements) (Figure 3B).

Figure 3: Estimated impact of the components of the intervention and impact of alternative vaccination strategies.

(A) Model predictions of the mean percent of the population that was infected, recovered, and susceptible considering:

the intervention as it happened of WASH and vaccination (dark blue), WASH only (blue), vaccination only (green),

and no intervention of either vaccination or WASH (yellow) from November 2013 to February 2016. Typical rainy

seasons are shaded in grey and the timing of the distribution of vaccine doses is shown in dashed purple lines. (B)

Violin plots of numbers of additional cholera cases at the end of the study period with WASH only (blue), vaccination

only (green), or no intervention (no WASH and no vaccination) (yellow) compared to the intervention as it happened

of WASH and vaccination. The error bars show the 95% credible interval, the filled circles show the median values,

and the horizontal bars show the mean values. (C) Heatmap of the number of cases avoided by changing the timing

and the coverage of a vaccination campaign compared to a scenario without intervention. The timing of the distribution

of vaccine doses for the intervention as it happened are in dashed grey lines in A and C.

Alternative vaccination scenarios revealed that small-scale interventions, or interventions with small

target population sizes, had a very limited impact in this population with high immunity (Figure 3C).

However, the timing of a pulse of vaccination substantially influenced its potential impact. Specifically,

timing the vaccination to occur during the lowest point of population immunity and before an outbreak

began increased its impact. The best performing vaccination scenario avoided 6,858 cases (mean:

6,858.5, 95%CrI: 4,809.7-8,600.8) over 118 weeks for 200,000 vaccinated people, which is more than

twice as many cases that were avoided by vaccination alone in the intervention that occurred. Despite

this, the high level of local immunity resulted in vaccinating a large proportion of immune individuals,

reducing the impact of the vaccination.

Figure 4: Contributions of seasonal mobility and the environmental reservoir in cholera transmission. (A) Model

predictions of the mean percent of infected, recovered, and susceptible individuals in the population with seasonal

migration and environmental exposure and contamination (light blue), environmental exposure and contamination (no

seasonal mobility) (pink), seasonal mobility and environmental contamination (no environmental exposure) (beige),

and seasonal mobility and environmental exposure (no environmental contamination) (grey) from November 2013 to

February 2016. Typical rainy seasons are shaded in grey. (B) Violin plots of numbers of cholera cases avoided by the

end of the study period with environmental exposure and contamination (no seasonal mobility) (pink), seasonal

migration and environmental contamination (no environmental exposure) (beige), or seasonal migration and

environmental exposure (no environmental contamination) (grey) compared to a scenario with seasonal migration,

environmental exposure, and environmental contamination. The error bars, the filled circles, and the horizontal bars

indicate the 95% credible interval, the median, and the mean values, respectively. All the scenarios considered in A

and B assume that no intervention occurred (no WASH and no vaccination). (C) Mean prediction of the variation of

the environmental (light orange line) and interhuman (light green line) components of the force of infection and their

95% credible interval (light orange and light green envelopes) from November 2013 to February 2016. The light

orange and light green dashed lines indicate the values of the environmental exposure rate ($\boldsymbol{\beta_e}$) and interhuman

transmission rate ($\boldsymbol{\beta_h}$), respectively. (D) Left: Mean prediction of the variation of the net bacterial growth ($\boldsymbol{\varphi_t - \varepsilon}$)

(dark line) in the environment and its 95% credible interval (grey envelope) from November 2013 to February 2016.

Right: Violin plot and boxplot of the distribution of the mean prediction of the net bacterial growth rate from

November 2013 to February 2016. The dashed black line indicates 0: values below 0 show net bacterial decay and

values above 0 show net bacterial growth.

Regarding the relative contributions of mobility and environmental reservoirs to cholera transmission,

the model suggested that seasonal migration had a minimal influence on the observed cholera dynamics

relative to the environmental reservoir (Figure 4A); a scenario with no seasonal migration did not

substantially alter the number of cases. However, removing environmental exposure avoided 93,896

cases (mean: 93,896.5, 95% CrI: 72,842.6-122,297.5) and preventing environmental contamination from

infectious human excreta avoided 91,519 cases (mean: 91,519.0, 95% CrI: 68,381.1-124,373.1). Each of

these interventions virtually interrupted local cholera transmission (Figure 4B). Environmental

contamination with infectious excreta appeared necessary to maintain a bacterial load sufficient to

support environmentally-driven transmission because the fluctuation of *V. cholerae* population

averaged towards net decay (Figure 4D, right).

The high immunity inferred by the model was maintained through annual flare-ups in Kalemie and

constant environmental exposure. The environmental component of the force of infection ($\Phi_e = \beta_e \frac{B}{\kappa + B}$

$(1 + \lambda_e f(rain_t))$) was consistently greater than the interhuman transmission component ($\Phi_h = \frac{\beta_h I}{N}$)

(Figure 4C). The latter, Φ_h, remained low because epidemic flare-ups did not lead to a high prevalence of

infection as they would in a mostly susceptible population. Conversely, Φ_e strongly increased with pulses

of net bacterial growth due to environmental drivers, despite an overall trend favoring net decay (Figure

4D left and right).

Discussion

The impact of the intervention performed in Kalemie was modest when measured by cases avoided,

which we estimate around 3,484 (mean: 3,484.4, 95% CrI: 2,411.7-4,832.8) for both intervention arms

combined. Following the interruption of planned vaccination activities, the reduction of the target

population size after resuming activities, the narrow scale of the WASH improvements, and the high

level of immunity in the population likely all contributed to mitigating the impact of the intervention.

Benefitting from vaccination in endemic cholera settings, as defined by WHO, requires an understanding

of dominant local transmission routes. Our model suggests that in settings where an environmental

reservoir provides consistent exposure and maintains high immunity in the population, the impact of

vaccination is minimal. The practical limitation of not being able to identify and target the population of

susceptible individuals leads to vaccinating a majority of immune individuals. Achieving very high

vaccine coverage would immunize a greater number of susceptible individuals, but at the cost of giving

many additional doses to immune individuals. This cost could be reduced slightly by targeting the age

group most represented among susceptibles or by guiding vaccination with serosurveys. The age profile

of the suspected cholera cases residing in Kalemie (median age of 15 years, and interquartile range (IQR)

of 3-34 years) during this period would support restricting the maximum age of the target population to

increase the impact of the vaccination campaign. However, applying an age cut-off would require high

resolution epidemiological data. Similarly, using serosurveys to guide vaccination efforts to target

susceptibles would incur a substantial additional financial cost in addition to the difficulty of applying a

binary interpretation to serosurvey results.

Our findings are consistent with previous analyses suggesting that allocation of OCV doses should

prioritize reactive vaccination in epidemic settings over proactive vaccination in endemic settings (31).

However, because endemicity is more nuanced than the current WHO definition suggests, OCV could

still play an important role in some endemic settings. Ongoing studies are assessing the impact of recent

vaccination campaigns performed in other endemic settings in DRC: in Uvira (32), which is also on the

shore of Lake Tanganyika, and in Goma (33), which is on the shore of Lake Kivu, a Great Lake that sits

just north of Lake Tanganyika . These studies will provide valuable insights on the impact of OCV with

alternative strategies.

Our model suggests that a well-timed large-scale vaccination could improve the impact of vaccination in

Kalemie city, potentially avoiding an average of 6,858.5 cases (95%CrI: 4,809.7-8,600.8) for 200,000

vaccinated individuals. However, this requires implementing a very large vaccination campaign with

precise timing. Such vaccination campaigns would be difficult to implement due to logistical challenges

and high costs, and they would still achieve only short term and small-scale benefits. On the other hand,

our findings suggest that WASH improvements on a scale large enough to prevent environmental

exposure and contamination for the whole population could have a dramatic impact. Although we

estimated that the WASH improvements in Kalemie prevented a modest number of cases, this is likely

partially due to the short period of time considered to assess the impact of an intervention. The main

components of this WASH intervention consisted of extending the pipe network and building a water

reservoir, which were completed incrementally during that 118-week period. While extending the

access to the pipe network is an important step, it does not guarantee reliable and consistent access to

chlorinated tap water (34). The immense magnitude of the improvements required to ensure both

access to safe water and efficient waste management, not only in Kalemie but throughout the cholera-

affected nation of DRC, appears necessary to control cholera. Implementing WASH improvements

should be considered a priority not only to control cholera, but also to prevent the transmission of other

water-borne and fecal-oral pathogens that contribute to the disease burden in DRC (35). This approach

will also help achieve the 6th goal of the Sustainable Development Goals (36), which is to ensure

availability and sustainable management of water and sanitation for all, in a country where WASH

improvements are critically needed (18).

Kalemie is not unique regarding strong environmentally driven cholera transmission. Substantial

environmental contribution for cholera cases has been reported in Haiti and Zimbabwe (20).

Environmental drivers are also important drivers in other endemic settings like Bangladesh and India,

where they act differently: flooding in the early and late phase of the monsoon are strongly associated

with higher cholera incidence (37,38), while the peak of the monsoon is associated with a cholera lull

due the "dilution" of *V. cholerae* in its reservoir (39). Although mobility does not play a role in local

persistence within the city of Kalemie, movement allows Kalemie to be a source of cholera that can seed

outbreaks in surrounding areas where exposure is less frequent. The older age of the suspected cholera

cases residing outside of Kalemie (median age of 24.5 years, and IQR: 5.75-39.25 years) is consistent

with lower exposure rates and source-sink dynamics.

Our estimates indicating a major role of environmentally driven transmission in Kalemie's local cholera

dynamics appear plausible. Confirming our estimates of population immunity and the source of bacterial

infection would require serological data and substantial microbiological monitoring of the lake water in

the area. Evidence of environmental presence of toxigenic *V. cholerae* is scarce but it has been found in

Lake Tanganyika on the Tanzanian shore in fish and water samples with some evidence of increased

positive samples during rainy seasons (40–42). Evidence of fecal contamination in coastal water would

support the importance of the strong amplification of environmental presence through environmental

contamination during outbreaks (43).

The natural variation in abundance of the *V. cholerae* population in the lake leans in favor of net decay.

Previous modeling studies assumed bacterial growth rates to consistently be in favor of net decay,

whether it varied over time or it was constant negative growth (44,45). Our model allowed

environmental bacterial abundance to vary over time based on environmental inputs, allowing

temporary switches to net bacterial growth. It played an important role in creating pulses of a high,

environmentally-driven force of infection. The overall net bacterial decay in the lake also highlights that

regularly replenishing local bacterial population through environmental contamination is a critical

component of local persistence. This emphasizes the potential compounded benefits of comprehensive

improvements to sanitary infrastructures and access to clean water.

Although our model is not designed to assess the long-term dynamics of cholera in Kalemie city, the

importance of water contamination is consistent with observations in other modern outbreaks. The

cholera outbreak that occurred in Haiti in 2010 following its introduction by United Nations

peacekeepers was primarily transmitted along the Artibonite river in its early phase (46). Fecal

contamination of water has been commonly found in Haiti (47), where access to safe drinking water and

sanitation is among the lowest in the world (18). This provides conditions conducive to environmentally

driven transmission for cholera. In contrast, the neighboring country on Hispaniola, the Dominican

Republic, has good access to safe drinking water and strong sanitation infrastructures (18) and

experienced little cholera transmission during the massive outbreak in Haiti. The impact of improving

the quality of consumed water (reducing environmental exposure) in our simulation is impressive but

the impact of removing environmental contamination appears almost as effective. However, this would

be difficult to compare in practice since a comprehensive WASH intervention would aim to both

improve the quality of consumed water and minimize the contamination of the local water reservoir.

We did not consider cholera-induced mortality because of the low number of cholera-induced deaths in

this population and the local experience in managing cholera. However, there is evidence that a

substantial portion of cholera mortality has previously occurred in the community (48) so we cannot rule

out that some cholera-induced mortality is not captured in the reported data. The absence of dedicated

mortality surveys in Kalemie prevents us from estimating the number of deaths avoided by the multi-

pronged intervention. Our model did not consider the potential immunological impact of vaccination on

already immune individuals, which could extend the period during which they are protected against

cholera infection. This could have led us to slightly underestimate the duration of immunity but it is

unlikely to have substantial impact on our estimates considering the short study period (118 weeks,

which is a little more than 2 years) compared to our estimated average immunity period ($1/\gamma$= 6.2 years,

95% CrI: 4.6- 8.0 years). We also assumed that immunity waning for susceptible individuals who were

successfully vaccinated was similar to the immunity waning after a natural infection, although it is likely

shorter (49). Again, this would have little impact on our estimates considering the small proportion of

susceptible individuals in the population when doses were distributed in our model as well as the

previously mentioned reasons (50).

We included the potential impact of the WASH intervention in a simplistic way, assuming a linear

variation of the environmental transmission rate due to the WASH intervention. In the absence of more

detailed information, this method required the fewest additional assumptions. To estimate

environmental drivers, we used measurements of chlorophyll-a levels and surface water temperature in

the lake in addition to the influence of rain. The association between *V. cholerae* and other elements of

its aquatic reservoir is vaguely understood, but the details remain unclear (7,51). We cannot assess how

accurately we captured the main fluctuations of the environmental bacterial population in the absence

of thorough environmental sampling in the area. However, the environmental drivers we considered

have all been shown to modulate *V. cholerae* environmental abundance or exposure to the

environmental reservoir. Phytoplankton growth, indirectly measured through chlorophyll-a, has been

associated with cholera outbreaks in several studies, and cyanobacteria more specifically are a credible

reservoir for *V. cholerae* (5,7). Water temperature influences phytoplankton growth (7), and the

consequence of rainfall on environmental exposure and environmental contamination to/from *V.*

cholerae is credible in this setting along a lake with low access to water and sanitation infrastructures

(52). Finally, the complex combination of parameters to estimate critical metrics including the number

of infected individuals, the net bacterial growth, and its consequence on the environmental component

of the force of infection leads to some loss of identifiability. However, we reached convergence on the

critical metrics they helped estimate, and our model captured true values of parameters in 95% CrI using

simulated data (see Supplementary Information (SI)). We remain confident in our main conclusions.

We explicitly included a direct proxy of human presence: anthropogenic nighttime radiance. Nighttime

radiance is a reliable indicator of human presence and has previously been used to infer population

mobility in both high and low-income countries (53–56). We are confident in our finding regarding the

low contribution of seasonal mobility. We also explicitly included relevant environmental drivers making

the environmental component of our model robust, though it might be limited by the spatiotemporal

resolution and availability of remote sensing data, particularly for chlorophyll-a.

There is a scarcity of impact assessments of cholera interventions in endemic settings beyond estimates

of vaccine effectiveness and vaccine coverage. Studies like this one are crucial to guide cholera

elimination. OCV and WASH improvements are core components of the toolbox to control or eliminate

cholera. However, the value of OCV in reactive vaccination in epidemic settings has not been clear in

areas with various patterns of endemicities. The assumption that most of the target population is

susceptible becomes less accurate as transmission is increasingly environmentally driven. Reducing

cholera transmission in endemic areas will require a location-specific understanding of the transmission

routes to tailor a strategy; a "one size fits all" approach is unlikely to achieve satisfying results.

Coordinated geographically-specific strategies might also be necessary to achieve regional cholera

control.

Materials and methods

We fit a Susceptible-Infected-Recovered-Susceptible model that included a compartment, B, for the *V.*

cholerae population in the environmental reservoir, Lake Tanganyika (57). Our SIRB model explicitly

considered the influence of environmental factors on bacterial growth in the lake and the potential

consequences of seasonal population mobility.

We fit the SIRB model to the reported suspected cholera cases presenting at the only cholera treatment

center (CTC) in the city of Kalemie from November 2013 to February 2016. Detailed surveillance data

were gathered in an electronic register with support from MSF. This level of resolution is available only

for this period of time. Only residents of Kalemie city were included in this analysis. We estimated the

number of vaccinated individuals from vaccine coverage data collected in a survey performed by MSF

(29) and the associated population size estimates (see SI).

Seasonal population mobility (58) can influence local cholera dynamics through regular reintroductions

from areas with ongoing transmission. We included this connectivity by quantifying the seasonal

variation of contemporaneous anthropogenic nighttime radiance extracted from Visible Infrared

Imaging Radiometer Suite (VIIRS) data (59) (see SI).

We considered the influence of water temperature and phytoplankton, such as cyanobacteria, as

environmental drivers on aquatic bacterial growth (7). Water temperature and phytoplankton were

included in the model through lake surface temperature (SST_t) and chlorophyll-a ($chlor_t$). These values

were extracted from Moderate Resolution Imaging Spectroradiometer data (60) (see SI). Precipitation

($rain_t$) was also considered as a potential factor that could increase exposure to environmental reservoir

and the environmental contamination with infectious human excreta (52) ($\lambda_e f(rain_t)$) by contaminating

drinking water sources and flooding defecation sites ($\lambda_c f(rain_t)$). We extracted precipitation estimates

from meteorological forcings data (61).

The structure of the model is as follows:

405
$$\frac{dS}{dt} = \delta R - \beta_h \frac{SI}{N} - \beta_e \frac{B}{\kappa + B}(1 + \lambda_e f(rain_t))S - \eta S + f_S(rad_t)$$

406
$$\frac{dI}{dt} = \beta_h \frac{SI}{N} + \beta_e \frac{B}{\kappa + B}(1 + \lambda_e f(rain_t))S - \gamma I + f_I(rad_t)$$

407
$$\frac{dR}{dt} = \gamma I - \delta R + \eta S$$

408
$$\frac{dB}{dt} = \mu(1 + \lambda_c f(rain_t))I - B(\varepsilon - \varphi_t)$$

with

$f_S(rad_t) = \alpha_1 rad_t$

$f_I(rad_t) = \alpha_I rad_t$

$ratio_{S/I} = \dfrac{\alpha_1}{\alpha_I}$

$\varphi_t = e^{\alpha_2 + \alpha_3 sst_t + \alpha_4 chlor_t}$

$\eta = (\sigma_1 VC_{1t} + \sigma_2 VC_{2t})\square$

$f(rain_t) = \dfrac{rain_t}{max(rain_t)}$

We used a negative binomial process to link the predicted number of weekly incident cases (C_t) and the

weekly reported number of suspected cases (A_t): $A_t \sim NegBinom(C_t r, C_t \psi)$, with r, a constant reporting

rate, and $C_t \psi$, an overdispersion parameter that scaled with the predicted number of new cases. The

negative binomial distribution can handle overdispersion and its scaling overdispersion parameter

allows variance estimates to better scale in situations with fast and large variations of the incidence, like

in epidemics.

Susceptible individuals become infected through exposure to the environmental reservoir, β_e or through

interhuman transmission, β_h. The WASH intervention decreased the environmental exposure rate β_e by

a quantity β_{WASH} by the end of the study period. β_e was assumed to decrease linearly from β_e to β_e-

β_{WASH} from the time the first WASH improvement was completed (ISO week 40 in 2014). The model did

not allow the environmental contamination to vary because the intervention did not target human

waste management. The infection probability from an exposure to the environment followed a dose-

effect relationship, with κ being the half saturation constant. Infectious individuals transitioned to the

recovered compartment at rate γ. Susceptible individuals could gain immunity through vaccination, η, 1

week after receiving the vaccine (19). This was included in the model through a step function of the

number of people who received 1 or 2 doses (VC_{1t} and VC_{2t}) of ShancholTM. The model presented here

assumed a vaccine effectiveness of 80% (σ_1) and 85% (σ_2) respectively for one and two dose regimens.

We considered a range of alternative values for those two parameters, including estimates from studies

done in the aftermath of reactive vaccination campaigns performed in Zambia and Guinea (19,62) (see

SI). Considering the wide age range of the target population (everyone older than 1 year), we assumed

that the proportions of susceptible, infected, and recovered among the vaccinated individuals were the

same as in the general population at the time the doses were distributed. Immunity wanes at rate δ,

returning immune individuals to the susceptible compartment. We assumed the vaccine had no effect

on individuals who were immune, whether immunity was acquired through prior infection or

vaccination. We also assumed that vaccination had no impact on those who were infected at the time of

the vaccination. We added a penalty term (ϴ) to account for the spatially targeted nature of the

vaccination campaign, which focused on areas with historically high attack rates where residents

experience more cholera exposures, resulting in a low proportion of susceptibles. We considered a

range of possible values for ϴ over several models (between 0.7 and 1) (see SI).

This model did not include the births, deaths, and the age structure of the host population because of

the short time period of 118 weeks that was considered. We also assumed no cholera-induced mortality

due to case management; only 5 deaths were reported among the 1634 resident suspected cholera

cases, or 0.3%, during the study period (see SI).

Population size was allowed to vary through seasonal migration ($f_S(rad_t)$ and $f_I(rad_t)$). We assumed

that the net migration flow varied linearly with the first derivative of the nighttime radiance data in the

area (rad_t) (53). We first fit a generalized additive model with a cyclical spline to the radiance data and

then extracted its first derivative (see SI). We considered the migration of susceptible and infected

individuals but not immune individuals, who do not actively contribute to transmission. We considered a

range of values for the ratio of susceptible and infectious individuals among the mobile population (

$ratio_{S/I}$) over several models (between 10 and 100) (see SI). We explored alternative model structures

including allowing only susceptible individuals to be mobile ($f_I(rad_t) = 0$) as well as excluding all

seasonal mobility as a sensitivity analysis ($f_S(rad_t) = 0 \ and \ f_I(rad_t) = 0$) (see SI).

The bacterial population in the environment increased with contamination of the lake from the excreta

of infected individuals (μ), and a time dependent bacterial growth rate (φ_t) that varied with

environmental drivers. Conversely, it decreased through bacterial decay (ε). To minimize identifiability

issues between ε and the components of φ_t we clipped the parameter space of the decay rate using

previously reported values (44,63) (see SI).

We estimated the parameters β_h, β_e, β_{WASH}, α_1, α_2, α_3, α_4, λ_e, λ_c, δ, μ, ε, r, and ψ, and the initial

conditions S_0, I_0, R_0, B_0 through Markov chain Monte Carlo sampling using the Metropolis-Hastings

algorithm and fit a total of 64 models with every combination of the values considered for σ_1, σ_2, ⬚, and

$ratio_{S/I}$ in addition to 32 models with alternative structures for sensitivity analyses (see SI). The model

presented here minimized the deviance information criterion (DIC). All the estimates presented are the

mean values over the posterior distribution and their 95% CrI).

We assessed the short-term impact of each arm of the intervention separately and both arms together

by estimating the number of additional cases in their absence. We fixed η to 0 while keeping β_{WASH}

unchanged (simulating WASH improvements without vaccination), did not allow β_e to decrease (β_{WASH}

=0) while keeping η unchanged (simulating vaccination without WASH improvements), and then fixed

both η and β_{WASH} to 0 (simulating an intervention with no vaccination and no WASH improvements).

We sampled 10,000 sets of parameters from the posterior distribution and calculated the number of

cases in each of the alternative scenarios and compared them to the intervention as it happened.

We explored alternative vaccination strategies by varying the timing and the size of the target

population, between 50,000 to 200,000, of the vaccination campaign, assuming a campaign that

occurred only once during the 118-week period with a two-dose regimen. The maximum target

population size considered in these alternatives strategies is within the MSF vaccination capacity

observed in other vaccination campaigns against cholera (62,64). We estimated the number of cases

avoided for each scenario by calculating the difference in the number of cholera cases between a

simulation using the alternative vaccination scenario and a simulation assuming no intervention (no

vaccination and no WASH) for every set of parameters sampled from the posterior distribution. We

considered 84 combinations of alternative timing and target population size. We sampled 500 sets of

parameters for each combination, computational intensity prohibited more. As before, we assumed that

the proportion of susceptible, infected, and recovered among the vaccine-targeted individuals was the

same as in the general population at the time the doses were distributed. For these scenarios we

considered a non-spatially targeted vaccination campaign (see SI).

We investigated the relative contributions of population mobility and environmental exposure in a

context without intervention (η=0 and β_{WASH}=0, meaning no vaccination and no WASH improvements).

We simulated alternative scenarios with no seasonal mobility ($f_S(rad_t)$=0 and $f_I(rad_t)$=0), or no

environmental exposure (β_e=0), or no environmental contamination (μ=0) and calculated the

cumulative difference between these and a scenario with seasonal mobility, environmental exposure

and contamination ($f_S(rad_t)$, $f_I(rad_t)$, β_e, and μ unchanged) for each of 10,000 sets of parameters

sampled from the posterior distribution.

Ethical considerations

The Ethical Review Board of the University of Lubumbashi approved the study protocol to assess the

impact of the vaccination campaign (study protocol ethical number: UNILU/CEM/028/2013) and its

extension (study protocol ethical number: UNILU/CEM/050/2015). Individuals provided informed

consent to be part of the vaccine coverage survey. Pennsylvania State University's Institutional Review

Board determined the post-intervention handling and analyses of these anonymized data was Not

Human Research (STUDY00015621).

Acknowledgements

We would like to acknowledge support and feedback from Nathan Wikle on statistical analyses.

Author contribution

Conceptualization: AB, FL, DB, NB

Data curation: AB, AW, FL

Formal analysis, AB, AW, FL

Investigation: AB, FL, DB

Methodology: AB, AW, FL, EH, NB

Software: AB, AW

Visualization: AB, NB

Writing – Original draft preparation: AB, NB

Writing - Review and Editing: AB, AW, FL, DB, POW, EH, NB

References

1. Harris JB, LaRocque RC, Qadri F, Ryan ET, Calderwood SB. Cholera. The Lancet. 2012 Jun

30;379(9835):2466–76.

2. Nelson EJ, Harris JB, Morris JG, Calderwood SB, Camilli A. Cholera transmission: the host, pathogen

and bacteriophage dynamic. Nat Rev Microbiol [Internet]. 2009 Oct [cited 2020 Apr 8];7(10).

Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3842031/

3. Jutla A, Whitcombe E, Hasan N, Haley B, Akanda A, Huq A, et al. Environmental Factors Influencing

Epidemic Cholera. Am J Trop Med Hyg. 2013 Sep 4;89(3):597–607.

4.   Constantin de Magny G, Colwell RR. Cholera and climate: a demonstrated relationship. Trans Am

Clin Climatol Assoc. 2009;120:119–28.

5.   Constantin de Magny G, Murtugudde R, Sapiano MRP, Nizam A, Brown CW, Busalacchi AJ, et al.

Environmental signatures associated with cholera epidemics. Proc Natl Acad Sci. 2008 Nov

18;105(46):17676–81.

6.   Zerbo A, Castro Delgado R, González PA. A review of the risk of cholera outbreaks and urbanization

in sub-Saharan Africa. J Biosaf Biosecurity. 2020 Dec 1;2(2):71–6.

7.   Islam MS, Zaman MH, Islam MS, Ahmed N, Clemens JD. Environmental reservoirs of Vibrio cholerae.

Vaccine. 2020 Feb 29;38:A52–62.

8.   Vezzulli L, Pruzzo C, Huq A, Colwell RR. Environmental reservoirs of Vibrio cholerae and their role in

cholera. Environ Microbiol Rep. 2010 Feb;2(1):27–33.

9.   Finkelstein RA. Cholera, Vibrio cholerae O1 and O139, and Other Pathogenic Vibrios. In: Baron S,

editor. Medical Microbiology [Internet]. 4th ed. Galveston (TX): University of Texas Medical Branch

at Galveston; 1996 [cited 2020 Apr 8]. Available from:

http://www.ncbi.nlm.nih.gov/books/NBK8407/

10. Birmingham ME, Lee LA, Ndayimirije N, Nkurikiye S, Hersh BS, Wells JG, et al. Epidemic cholera in

Burundi: patterns of transmission in the Great Rift Valley Lake region. The Lancet. 1997 Apr

5;349(9057):981–5.

11. Frerichs RR, Keim PS, Barrais R, Piarroux R. Nepalese origin of cholera epidemic in Haiti. Clin

Microbiol Infect Off Publ Eur Soc Clin Microbiol Infect Dis. 2012 Jun;18(6):E158-163.

12. Sow S, Antonio M, Oundo JO, Mandomando I, Ramamurthy T. Endemic and Epidemic Cholera in

Africa. In: Ramamurthy T, Bhattacharya SK, editors. Epidemiological and Molecular Aspects on

Cholera [Internet]. New York, NY: Springer; 2011 [cited 2022 Dec 1]. p. 31–50. (Infectious Disease).

Available from: https://doi.org/10.1007/978-1-60327-265-0_3

13. Ali M, Nelson AR, Lopez AL, Sack DA. Updated Global Burden of Cholera in Endemic Countries. PLoS

Negl Trop Dis. 2015 Jun 4;9(6):e0003832.

14. Organization WH. Ending cholera a global roadmap to 2030. In: Ending cholera a global roadmap to

2030. 2017. p. 32–32.

15. Waldman RJ, Mintz ED, Papowitz HE. The Cure for Cholera — Improving Access to Safe Water and

Sanitation. N Engl J Med. 2013 Feb 14;368(7):592–4.

16. Luby SP, Davis J, Brown RR, Gorelick SM, Wong THF. Broad approaches to cholera control in Asia:

Water, sanitation and handwashing. Vaccine. 2020 Feb 29;38:A110–7.

17. Montgomery M, Jones MW, Kabole I, Johnston R, Gordon B. No end to cholera without basic water,

sanitation and hygiene. Bull World Health Organ. 2018 Jun 1;96(6):371-371A.

18. Organization WH. Progress on household drinking water, sanitation and hygiene 2000-2017: special

focus on inequalities. World Health Organization; 2019.

19. Luquero FJ, Grout L, Ciglenecki I, Sakoba K, Traore B, Heile M, et al. Use of Vibrio cholerae vaccine in

an outbreak in Guinea. N Engl J Med. 2014 May 29;370(22):2111–20.

20. Mukandavire Z, Morris JG. Modeling the Epidemiology of Cholera to Prevent Disease Transmission

in Developing Countries. Microbiol Spectr. 2015 Jun 19;3(3):3.3.21.

21. World Health Organization. Cholera vaccine: WHO position paper, August 2017 –

Recommendations. Vaccine. 2018 Jun 7;36(24):3418–20.

22. Pezzoli L. Global oral cholera vaccine use, 2013–2018. Vaccine. 2020 Feb;38:A132–40.

23. Shaikh H, Lynch J, Kim J, Excler JL. Current and future cholera vaccines. Vaccine. 2020 Feb

29;38:A118–26.

24. Ali M, Debes AK, Luquero FJ, Kim DR, Park JY, Digilio L, et al. Potential for Controlling Cholera Using a

Ring Vaccination Strategy: Re-analysis of Data from a Cluster-Randomized Clinical Trial. PLOS Med.

2016 Sep 13;13(9):e1002120.

25. Leung T, Eaton J, Matrajt L. Optimizing one-dose and two-dose cholera vaccine allocation in

outbreak settings: A modeling study. PLoS Negl Trop Dis. 2022 Apr 20;16(4):e0010358.

26. Finger F, Bertuzzo E, Luquero FJ, Naibei N, Touré B, Allan M, et al. The potential impact of case-area

targeted interventions in response to cholera outbreaks: A modeling study. PLOS Med. 2018 Feb

27;15(2):e1002509.

27. Congo Democratic Republic DHS, 2013-14 - Final Report (French).

28. Bompangue Nkoko D, Giraudoux P, Plisnier PD, Tinda AM, Piarroux M, Sudre B, et al. Dynamics of

cholera outbreaks in Great Lakes region of Africa, 1978-2008. Emerg Infect Dis. 2011

Nov;17(11):2026–34.

29. Massing LA, Aboubakar S, Blake A, Page AL, Cohuet S, Ngandwe A, et al. Highly targeted cholera

vaccination campaigns in urban setting are feasible: The experience in Kalemie, Democratic Republic

of Congo. PLoS Negl Trop Dis. 2018 May 7;12(5):e0006369.

30. Kayembe HCN, Bompangue D, Linard C, Muwonga J, Moutschen M, Situakibanza H, et al. Modalities

and preferred routes of geographic spread of cholera from endemic areas in eastern Democratic

Republic of the Congo. PLOS ONE. 2022 Feb 7;17(2):e0263160.

31. Moore SM, Lessler J. Optimal allocation of the limited oral cholera vaccine supply between endemic

and epidemic settings. J R Soc Interface. 2015 Oct 6;12(111):20150703.

32. The impact of mass cholera vaccination in Uvira, Democratic Republic of the Congo – Global Task

Force on Cholera Control [Internet]. [cited 2023 Apr 12]. Available from:

https://www.gtfcc.org/research/the-impact-of-mass-cholera-vaccination-in-uvira-democratic-

republic-of-the-congo/

33. Impact evaluation of OCV preventive campaigns, DRC – Global Task Force on Cholera Control

[Internet]. [cited 2023 Apr 13]. Available from: https://www.gtfcc.org/research/impact-evaluation-

of-ocv-preventive-campaigns-drc/

34. Jeandron A, Saidi JM, Kapama A, Burhole M, Birembano F, Vandevelde T, et al. Water Supply

Interruptions and Suspected Cholera Incidence: A Time-Series Regression in the Democratic

Republic of the Congo. Brocklehurst C, editor. PLOS Med. 2015 Oct 27;12(10):e1001893.

35. Reiner RC, Graetz N, Casey DC, Troeger C, Garcia GM, Mosser JF, et al. Variation in Childhood

Diarrheal Morbidity and Mortality in Africa, 2000–2015. N Engl J Med. 2018 Sep 20;379(12):1128–

38.

36. THE 17 GOALS | Sustainable Development [Internet]. [cited 2022 Dec 13]. Available from:

https://sdgs.un.org/goals

37. Carrel M, Emch M, Streatfield PK, Yunus M. Spatio-temporal clustering of cholera: the impact of

flood control in Matlab, Bangladesh, 1983-2003. Health Place. 2009 Sep;15(3):741–52.

38. Cash BA, Rodó X, Emch M, Yunus M, Faruque ASG, Pascual M. Cholera and Shigellosis: Different

Epidemiology but Similar Responses to Climate Variability. PLOS ONE. 2014 Sep 17;9(9):e107223.

39. Shackleton D, Memon FA, Nichols G, Phalkey R, Chen AS. Mechanisms of cholera transmission via

environment in India and Bangladesh: state of the science review. Rev Environ Health [Internet].

2023 Jan 16 [cited 2023 Jun 23]; Available from:

https://www.degruyter.com/document/doi/10.1515/reveh-2022-0201/html

40. Hounmanou YMG, Njamkepo E, Rauzier J, Gallandat K, Jeandron A, Kamwiziku G, et al. Genomic

Microevolution of Vibrio cholerae O1, Lake Tanganyika Basin, Africa - Volume 29, Number 1—

January 2023 - Emerging Infectious Diseases journal - CDC. [cited 2023 Apr 13]; Available from:

https://wwwnc.cdc.gov/eid/article/29/1/22-0641_article

41. Michael M. Detection and antibiotic susceptibility of vibrio cholerae In oreochromis tanganicae

(Tilapia) and water in Lake Tanganyika, Kigoma-Tanzania [Internet] [Thesis]. Sokoine University of

Agriculture; 2019 [cited 2023 Apr 13]. Available from:

http://www.suaire.sua.ac.tz/handle/123456789/3905

42. Nyambuli S. Prevalence, pathogenic markers and antibiotic susceptibility of Vibrio cholerae in

sardines, water and phytoplankton in lake Tanganyika, Tanzania. Sokoine University of Agriculture;

2017.

43. Plisnier P, Denis, Cornet Y, Cocquyt C. Cholera outbreaks at Lake Tanganyika induced by Climate

Change?(CHOLTIC) : final report [Internet]. Brussels : Belgian Scientific Policy, 2015; 2016 Jun [cited

2023 Apr 13]. Available from: https://orfeo.belnet.be/handle/internal/4240

44. Codeço CT. Endemic and epidemic dynamics of cholera: the role of the aquatic reservoir. BMC Infect

Dis. 2001 Feb 2;1(1):1.

45. Pasetto D, Finger F, Camacho A, Grandesso F, Cohuet S, Lemaitre JC, et al. Near real-time

forecasting for cholera decision making in Haiti after Hurricane Matthew. PLOS Comput Biol. 2018

May 16;14(5):e1006127.

46. Piarroux R, Barrais R, Faucher B, Haus R, Piarroux M, Gaudart J, et al. Understanding the cholera

epidemic, Haiti. Emerg Infect Dis. 2011 Jul;17(7):1161–8.

47. Widmer JM, Weppelmann TA, Alam MT, Morrissey BD, Redden E, Rashid MH, et al. Water-Related

Infrastructure in a Region of Post-Earthquake Haiti: High Levels of Fecal Contamination and Need for

Ongoing Monitoring. Am J Trop Med Hyg. 2014 Oct 1;91(4):790–7.

48. Morof D, Cookson ST, Laver S, Chirundu D, Desai S, Mathenge P, et al. Community Mortality from

Cholera: Urban and Rural Districts in Zimbabwe. Am J Trop Med Hyg. 2013 Apr 3;88(4):645–50.

49. Alam MM, Riyadh MA, Fatema K, Rahman MA, Akhtar N, Ahmed T, et al. Antigen-Specific Memory

B-Cell Responses in Bangladeshi Adults after One- or Two-Dose Oral Killed Cholera Vaccination and

Comparison with Responses in Patients with Naturally Acquired Cholera. Clin Vaccine Immunol.

2011 May;18(5):844–50.

50. Franke MF, Ternier R, Jerome JG, Matias WR, Harris JB, Ivers LC. Long-term effectiveness of one and

two doses of a killed, bivalent, whole-cell oral cholera vaccine in Haiti: an extended case-control

study. Lancet Glob Health. 2018 Sep 1;6(9):e1028–35.

51. ALMAGRO-MORENO S, TAYLOR RK. Cholera: Environmental Reservoirs and Impact on Disease

Transmission. Microbiol Spectr [Internet]. 2013 Dec [cited 2021 May 6];1(2). Available from:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4321695/

52. Lemaitre J, Pasetto D, Perez-Saez J, Sciarra C, Wamala JF, Rinaldo A. Rainfall as a driver of epidemic

cholera: Comparative model assessments of the effect of intra-seasonal precipitation events. Acta

Trop. 2019 Feb 1;190:235–43.

53. Bharti N, Tatem AJ, Ferrari MJ, Grais RF, Djibo A, Grenfell BT. Explaining Seasonal Fluctuations of

Measles in Niger Using Nighttime Lights Imagery. Science. 2011 Dec 9;334(6061):1424–7.

54. Stathakis D, Baltas P. Seasonal population estimates based on night-time lights. Comput Environ

Urban Syst. 2018 Mar 1;68:133–41.

55. Bharti N, Tatem AJ. Fluctuations in anthropogenic nighttime lights from satellite imagery for five

cities in Niger and Nigeria. Sci Data [Internet]. 2018 Nov 13 [cited 2019 Jul 30];5. Available from:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6233255/

56. Sutton P, Roberts D, Elvidge C, Baugh K. Census from Heaven: An estimate of the global human

population using night-time satellite imagery. Int J Remote Sens. 2001 Jan 1;22(16):3061–76.

57. Akman O, Corby MR, Schaefer E. Examination of models for cholera: Lett Biomath. 2016 Dec

1;3(1):93-118-93–118.

58. Mberu BU. African Migration and Population Distribution: Recent Trends. Int Handb Migr Popul
Distrib. 2015;6:245.

59. Elvidge CD, Baugh K, Zhizhin M, Hsu FC, Ghosh T. VIIRS night-time lights. Int J Remote Sens. 2017
Nov 2;38(21):5860–79.

60. MODIS-Aqua M. NASA goddard space flight center, ocean ecology laboratory, ocean biology
processing group. Moderate-Resolut Imaging Spectroradiometer MODIS Aqua L0 Data NASA OB
DAAC Greenbelt MD USA. 2018;

61. Sheffield J, Goteti G, Wood EF. Development of a 50-Year High-Resolution Global Dataset of
Meteorological Forcings for Land Surface Modeling. J Clim. 2006 Jul 1;19(13):3088–111.

62. Ferreras E, Chizema-Kawesha E, Blake A, Chewe O, Mwaba J, Zulu G, et al. Single-Dose Cholera
Vaccine in Response to an Outbreak in Zambia. N Engl J Med. 2018 08;378(6):577–9.

63. Dimitrov DT, Troeger C, Halloran ME, Longini IM, Chao DL. Comparative Effectiveness of Different
Strategies of Oral Cholera Vaccination in Bangladesh: A Modeling Study. PLoS Negl Trop Dis. 2014
Dec 4;8(12):e3343.

64. Ciglenecki I, Sakoba K, Luquero FJ, Heile M, Itama C, Mengel M, et al. Feasibility of Mass Vaccination
Campaign with Oral Cholera Vaccines in Response to an Outbreak in Guinea. PLoS Med. 2013 Sep
10;10(9):e1001512.

Supporting Information

S1 File

Supporting Information captions

S1 File. Supplementary material providing additional details on the methods

Figure 1

Figure 2

Figure 3

Figure 4