

Low *IGFBP7* expression identifies a subset of breast cancers with favorable prognosis and sensitivity to IGF-1 receptor targeting with ganitumab: Data from I-SPY2 and SCAN-B

Christopher Godina¹, Michael N Pollak², Helena Jernström¹.

¹ Division of Oncology, Department of Clinical Sciences in Lund, Lund University and Skåne University Hospital, Barngatan 4, SE-221 85 Lund, Sweden

² Lady Davis Institute for Medical Research, Jewish General Hospital and Department of Oncology, McGill University, Montreal, QC, Canada

Corresponding authors: Helena Jernström and Christopher Godina Division of Oncology, Department of Clinical Sciences in Lund, Lund University and Skåne University Hospital, Barngatan 4, SE-221 85 Lund, Sweden

E-mail:

christopher.godina@med.lu.se,

michael.pollak@McGill.Ca,

helena.jernstrom@med.lu.se

Running title: *IGFBP7* gene expression in breast cancer

Keywords: *IGFBP7*, breast cancer, molecular profiling, ganitumab, pCR, clinical outcome, I-SPY2, SCAN-B.

Abstract word count: 277

Manuscript word count: 3010

Figures: 4

Tables: 4

Supplements: 4

References: 30

Abbreviations

DEGs - Differentially expressed genes

DGE - Differential gene expression

DMFI - Distant metastasis-free interval

ER - Estrogen receptor

EMT - Epithelial-mesenchymal transition

FC - Fold change

FDR - False discovery rate

FPKM - Fragments per kilobase of exon per million mapped reads

GEO - Gene Expression Omnibus

GEX - Gene expression profile

GSEA - Gene set enrichment analysis

HER2 - Human epidermal growth factor receptor 2

HR - Hazard ratio

HR - Hormone receptor

IGF-1 - Insulin-like growth factor-1

IGF-2 - Insulin-like growth factor-2

IGFBP7 - Insulin-like growth factor receptor binding protein-7

IGF-1R - Insulin-like growth factor-1 receptor

I-SPY2 - Investigation of Serial Studies to Predict Your Therapeutic Response With Imaging And moLecular Analysis 2

OR- Odds ratio

OS - Overall survival

pCR - pathological complete response

PR - Progesterone receptor

RFI - Recurrence-free interval

RNA-seq - Massive parallel paired-end sequencing of mRNA

SCAN-B - Swedish Cancerome Atlas Network - Breast

TME - Tumor microenvironment

TNBC - Triple-negative breast cancer

Q – Quartile

Summary

There has been a long-standing interest in targeting the Insulin-like growth factor-1 receptor (IGF-1R) signaling system in breast cancer due to its key role in growth, proliferation, and survival. To date, no IGF-1R targeting agent has shown substantial clinical benefit in controlled trials despite promising preclinical data. Treatment predictive biomarkers for IGF-1R targeting agents are lacking. IGFBP7 is an atypical insulin-like growth factor binding protein as it associates with the IGF-1R, reducing its capacity to bind IGF-1 or IGF-2. We report that low *IGFBP7* gene expression identified a subset of breast cancers for which the addition of ganitumab (an anti-IGF-1R monoclonal antibody) with metformin to chemotherapy substantially improved the pathological complete response rate compared to neoadjuvant chemotherapy alone. Furthermore, high *IGFBP7* expression predicted increased distant metastasis risk. *IGFBP7* expression deserves further evaluation as a predictor of therapeutic benefit of IGF-1R targeting agents and of breast cancer prognosis.

Abstract

To date, no IGF-1R targeting agent has shown clinical benefit in patients despite promising preclinical data. We hypothesized that Insulin-like growth factor binding protein-7 (*IGFBP7*) expression predicts response to Insulin-like growth factor-1 receptor (IGF-1R) targeting agents, for which there are no biomarkers predictive of efficacy. High *IGFBP7* expression was previously associated with breast cancer progression. Therefore, the predictive value of *IGFBP7* expression was interrogated in I-SPY2 (NCT01042379) for pathological complete response (pCR) to neoadjuvant treatment and in SCAN-B (NCT02306096) for clinical outcome. In I-SPY2, *IGFBP7* expression was examined as a predictor of pCR to neoadjuvant chemotherapy alone and separately to the combination of chemotherapy plus ganitumab (anti-IGF-1R antibody) and metformin. Odds ratios (OR) and hazard ratios (HR) with 95% confidence interval (CI) were calculated with logistic and Cox regression, respectively. Higher *IGFBP7* expression conferred lower odds of achieving pCR in the ganitumab/metformin plus chemotherapy arm, $OR_{adj} 0.38$ (95% CI 0.17–0.80) but not in the chemotherapy-alone arm, adjusted OR 1.23 (95% CI 0.63–2.45; $P_{interaction}=0.016$). In the ganitumab/metformin plus chemotherapy arm, 46.9% of patients with tumors in the lowest quartile of *IGFBP7* expression achieved pCR compared to compared to only 5.6% in the highest quartile. In SCAN-B, higher *IGFBP7* expression was associated with distant metastasis risk $HR_{adj} 1.41$ (95% CI 1.16–1.73). In conclusion, low *IGFBP7* gene expression identifies a subset of breast cancer patients for whom the addition of ganitumab and metformin to chemotherapy results in a significantly improved pCR rate compared to neoadjuvant chemotherapy alone. Furthermore, we add to prior evidence that high *IGFBP7* expression is predictive of poor outcome.

Introduction

A significant portion of breast cancer patients relapse despite receiving optimal treatment adjuvant or neoadjuvant according to current guidelines¹. This fact highlights the need for new treatments and new biomarkers to better guide patient selection.

Dysregulation of insulin-like growth factor-1 receptor (IGF-1R) signaling has long been correlated with breast cancer growth, proliferation, and survival^{2,3}. Many agents have been developed to target IGF-1R⁴. To date, no studies have demonstrated a substantial clinical benefit of adding IGF-1R targeting agents to existing treatments⁴. Recently, the Investigation of Serial Studies to Predict Your Therapeutic Response With Imaging And moLecular Analysis 2 (I-SPY2) trial showed a small increase in pCR rate when the anti IGF-1R monoclonal antibody ganitumab (AMG479) and metformin were added to chemotherapy, but the effect size failed to reach the prespecified threshold for a positive result⁵. Further, eleven putative IGF-1R signaling axis biomarkers were tested, but none were predictive of ganitumab benefit⁵.

I-SPY2 has been able to identify treatment predictive biomarkers for emerging treatments due to its unique design⁶⁻⁹, and to evaluate efficacy of new treatments¹⁰⁻¹³. To further this goal, the I-SPY2-990 Data Resource has been made publicly available¹⁰.

It has previously been shown¹⁴ that unlike other IGFBPs, IGFBP7 can bind to the IGF-1R receptor (as well as to IGF-I and IGF-II), but the *in vivo* consequences of the IGFBP7 - IGF-1R interaction are unclear. There are also gaps in knowledge concerning complex interactions between IGFBP7 and IGF-1R in the presence of anti-IGF-1R antibodies such as Ganitumab. IGFBP7 binds insulin-like growth factor-1 (IGF-1) and the IGF-1R in a mutually exclusive manner¹⁴. Compared to other IGFBPs, IGFBP7 binds insulin with higher affinity than IGFs^{14,15}. The binding of IGFBP7 decreases activation and internalization of IGF-1R in response to IGF-1/2 but at the same sensitizes IGF-1R to insulin stimulation^{14,15}. It also has been shown that IGFBP7 promotes the persistence of IGF-1R at the cell surface, prolonging insulin/IGF stimulation, and enhancing Akt activation that exerts mitogenic and pro-survival effects^{15,16}.

Although we have presented prior evidence that IGFBP7 expression in breast cancer or circulating levels of this protein are related to prognosis^{17,18}, this protein was not previously studied in relation to efficacy of IGF-1R targeted therapies. We previously reported that the

prognostic value of circulating IGFBP7 levels were influenced by tumor IGF-1R status¹⁸, which suggests the idea that the interplay between IGFBP7 and IGF-1R merits further investigation. We hypothesized that IGFBP7 might compete with IGF-1R monoclonal antibody binding to IGF-1R, decreasing its efficacy and at the same time promote tumor growth and metastatic potential.

Materials and methods

I-SPY2

The I-SPY2 trial (NCT01042379) is an open-label, multicenter, adaptively randomized phase 2 trial of neoadjuvant therapy, evaluating multiple investigational arms in parallel. Neoadjuvant paclitaxel followed by doxorubicin/cyclophosphamide (with Trastuzumab in HER2+ disease) serves as the common control arm^{12,13}. Investigational agent(s) are combined with this regimen^{12,13}. The primary endpoint is pathological complete response (pCR; ypT0/is, ypN0)^{12,13}. Eligibility for I-SPY2 includes age 18 years or older, stage II or III breast cancer and primary tumor size >2.5 cm by clinical examination or >2.0 cm by imaging^{12,13}. Patients with hormone receptor (HR)+ cancers are eligible only if they have a poor prognosis estimated by the MammaPrint (MP) 70-gene based prognostic signature¹⁰. Subsequent adjuvant endocrine treatment was at the discretion of the treating physician. Ganitumab was tested in combination with metformin since ganitumab can induce hyperglycemia⁵. In the I-SPY2 study of the IGF-IR blocking antibody ganitumab, patients with HER2+ tumors were not included since there is no safety data concerning the combination of ganitumab and trastuzumab⁵. Gene expression profiling of core needle biopsies taken from the primary tumor before treatment was performed using Agilent 44K expression arrays¹⁰. Transcriptomic and clinical data of patients enrolled in the I-SPY2 trial were obtained from the Gene Expression Omnibus (GEO) database (GSE194040). For the analysis reported here, we studied patients from the ganitumab/metformin plus chemotherapy arm and the chemotherapy-alone control arm, excluding patients with HER2+ disease (Figure 1).

SCAN-B

The Swedish Cancerome Analysis Network – Breast (SCAN-B; NCT02306096) is an ongoing population-based cohort. SCAN-B prospectively includes breast cancer patients diagnosed and treated at nine Swedish hospitals^{19,20}. All newly diagnosed breast cancer

patients are invited to participate²⁰. Tumor specimens or core needle biopsies (in case of neoadjuvant treatment) from the patients' tumors are obtained in conjunction with routine clinical sampling^{19,20}. The samples are subject to gene expression profiling using RNA-seq according to custom SCAN-B workflow¹⁹⁻²¹. Gene expression levels were expressed in fragments per kilobase of exon per million mapped reads (FPKM)²¹. Clinical data was collected from the Swedish National Quality Registry for Breast Cancer¹⁹⁻²¹. Curated RNA-seq and clinical data were accessed from Staaf *et al.*²¹. A subset of 5326 patients with available follow-up for distant metastasis, no bilateral cancer, and gene expression profiles (GEXs) from a primary invasive breast cancer was analyzed (Supplementary Figure 1). If multiple GEXs for a single patient were available, the GEX with the highest RNA concentration was chosen²¹, leaving one GEX per patient for analysis.

Data analysis

In both I-SPY2 and SCAN-B, eight gene expression modules representing different biological functions in breast cancer were calculated as previously described²². The correlations between *IGFBP7* gene expression, gene expression of 15 other proteins involved in the IGF/Insulin pathway, and the eight modules were calculated using Pearson's correlation.

In the regression analyses, *IGFBP7* gene expression was modelled both as a continuous variable and categorized as quartiles (quartile 1 (Q1), quartile 2 (Q2), quartile 3 (Q3), and quartile 4 (Q4)) to allow for non-linear effects. The lowest expression of *IGFBP7* (Q1) was used as reference. Quartiles were created separately for I-SPY2 and SCAN-B based on the entire datasets.

Crude and adjusted odds ratios (OR) with 95% confidence intervals (CI) for pCR in I-SPY2 was estimated using logistic regression. The multivariable models were adjusted for HR+, HER2+, trial arm (control as reference), PAM50 subtype (Luminal A as reference), MP ultra-high risk (MP2), ISPY-2 immune (Immune+), and DNA repair deficiency (DRD+) signatures¹⁰. To investigate if there was any interaction between *IGFBP7* gene expression and efficacy of ganitumab/metformin plus chemotherapy treatment in achieving pCR, interaction analyses were performed in the ganitumab/metformin plus chemotherapy arm and the HER2 negative subset of the control arm. The interaction was tested in both crude and adjusted models by including an interaction term and comparing the models with or without the interaction term using the Likelihood ratio (LR) test. The *P*-values for the interaction term and LR test are

reported. Due to sparse data, PAM50 subtype could not be included in the interaction analyses.

The endpoints used for survival analyses in SCAN-B were recurrence-free interval (RFI), distant metastasis-free interval (DMFI), and overall survival (OS)²¹. Cox proportional hazards regression was used to estimate crude and adjusted Hazard ratios (HRs) with 95% CI. The multivariable models were adjusted for age (binned in 5-year intervals), tumor characteristics; lymph node status (pN1/2/3), tumor size (pT2/3/4), Grade (III), estrogen receptor (ER)+, progesterone receptor (PR)+, HER2+, PAM50 subtype (Luminal A as reference), PAM50 ROR category (High vs Low/Intermediate); and (neo)adjuvant treatments; endocrine treatment, chemotherapy, and trastuzumab.

Differential gene expression (DGE) analysis was performed using the ‘Limma-Voom’ package²³. The criteria used to define differentially expressed genes (DEGs) between the QT4 and Q1 of *IGFBP7* expression was false discovery rate (FDR) of ≤ 0.05 and \log_2 fold change (\log_2FC) ≥ 1.5 for up-regulated genes and $\log_2FC \leq -1.5$ for down-regulated genes. The ‘clusterprofiler’²⁴ package was used to perform gene set enrichment analysis (GSEA). Gene sets were grouped according to Hallmark Signatures²⁵. The results were visualized using ‘EnhancedVolcano’, ‘pHeatmap’ and ‘clusterprofiler’²⁴ packages.

All data analyses were conducted in R version 4.2.2. *P*-values < 0.05 was considered statistically significant. All *P*-values were two-tailed. This study followed the Reporting Recommendations for Tumor Marker Prognostic Studies (REMARK) criteria²⁶.

Results

In both ISPY-2 and SCAN-B, *IGFBP7* expression was correlated with *IGFBP3-6* and *IGF1* and *IGF2* expression (Figure 2 A-B). *IGFBP7* expression was highest in Normal-like followed by Luminal A subtype in both the ISPY-2 trial and SCAN-B cohort (Figure 2 C-D). Likewise, the correlations between *IGFBP7* expression and the eight gene modules were similar in ISPY-2 and SCAN-B. *IGFBP7* expression was positively correlated with stroma, lipid, and early response to growth signaling modules and negatively correlated with mitotic checkpoint and progression modules (Figure 2 E-F). The correlations suggest an association with aggressive tumor microenvironment (TME) and growth factor signaling. *IGFBP7* expression was stable across HR-positive and HER2-positive subtypes (Figure 2 G-H).

Descriptive statistics for clinicopathological factors are presented in Supplementary Table 1 for I-SPY2 and Supplementary Table 2 for SCAN-B.

In all patients (across all treatment arms), *IGFBP7* (continuous and in quartiles) expression was not associated with odds of achieving pCR in the multivariable model (Supplementary Table 3). There was an interaction between *IGFBP7* expression (continuous and in quartiles) and efficacy of ganitumab/metformin plus chemotherapy treatment in achieving pCR in both the crude (LR $P=0.011$ for both continuous and quartiles) and multivariable models (LR $P=0.017$ for continuous and LR $P=0.015$ for quartiles) (Table 1). In the ganitumab/metformin plus chemotherapy arm 22.6% of patients achieved pCR compared to 16.8% in the chemotherapy alone control arm. However, higher *IGFBP7* expression conferred lower odds of achieving pCR in the ganitumab/metformin plus chemotherapy arm, adjusted OR 0.38 (95% CI 0.17–0.80) but not in the chemotherapy alone control arm, adjusted OR 1.23 (95% CI 0.63–2.45; $P_{\text{interaction}}=0.016$) (Figures 1, 3 and Table 1). Likewise, Q2-4 of *IGFBP7* expression compared to Q1 also conferred lower odds of achieving pCR in the ganitumab/metformin plus chemotherapy arm (Q4 adjusted OR 0.09 (95% CI 0.00–0.58)) but not in the chemotherapy-alone control arm (adjusted OR 1.38 (95% CI 0.41–4.73; $P_{\text{interaction}}=0.031$)). The crude and adjusted ORs for Q2-3 and other clinicopathological factors are provided in Figure 1 and Table 1. When stratified by quartiles, in the ganitumab/metformin plus chemotherapy arm 46.9% of *IGFBP7* Q1 tumors achieved pCR compared to 17.2% of *IGFBP7* Q2 tumors, 11.5% of *IGFBP7* Q3 tumors, and only 5.6% of *IGFBP7* Q4 tumors (Figure 3B and Table 1). In the chemotherapy alone control arm, 14.0% *IGFBP7* Q1 tumors, 20.0% of *IGFBP7* Q2 tumors, 16.7% *IGFBP7* Q3 tumors, and 16.3% of *IGFBP7* Q4 tumors achieved pCR (Figure 1, Figure 3C and Table 1). When divided by breast cancer subtype (high risk HR+HER2- vs. TNBC), the ability of *IGFBP7* expression to identify breast cancers more likely to respond to ganitumab/metformin plus chemotherapy than to chemotherapy alone, was more apparent in TNBC, where 66.7% of *IGFBP7* Q1 tumors and no *IGFBP7* Q4 tumors achieved pCR, but the numbers are too small to draw definitive conclusions (Table 2). In an exploratory analysis, stratified by *IGFBP7* expression quartiles, ganitumab/metformin plus chemotherapy treatment conferred higher odds of achieving pCR in tumors with low *IGFBP7* expression (Q1-2) (adjusted OR 2.64 (95% CI 1.16–5.49)) but not in tumors with high *IGFBP7* expression (Q3-4), (adjusted OR 0.44 (95% CI 0.11–1.40)). The improved efficacy of ganitumab/metformin plus chemotherapy treatment compared to standard chemotherapy in achieving pCR was confined to the $\approx 25\%$

of patients in the lowest quartile of *IGFBP7* expression, where the pCR rate was 46.9% in the ganitumab/metformin plus chemotherapy group, compared to 14% in the chemotherapy alone group (Figure 1 and Table 3).

In SCAN-B, the median follow-up for the 4158 patients still at risk was 5.45 (inter quartile range 5.07–8.15) years. In the univariable survival analyses of *IGFBP7* expression in the SCAN-B cohort, *IGFBP7* expression Q4 compared to Q1 was not associated with increased risk of recurrence (HR 0.96 (95% CI 0.76–1.22)) or distant metastasis (HR 1.00 (95% CI 0.77–1.32)) (Supplementary Table 4). However, after adjustment for age, clinicopathological factors and treatment in the multivariable models, high *IGFBP7* expression (Q4 compared to Q1) was associated with significantly increased risk of recurrence (HR 1.37 (95% CI 1.04–1.82)) and distant metastasis (HR 1.60 (95% CI 1.15–2.22)) (Supplementary Table 4). When modeled as a continuous variable, *IGFBP7* expression showed similar associations with clinical outcome (Supplementary Table 5). Most importantly, *IGFBP7* (continuous) gene expression in the adjusted model also conferred a significantly increased risk of recurrence HR (1.29 (95% CI 1.09–1.53)) and distant metastasis (HR 1.41 (95% CI 1.16–1.73)) (Supplementary Table 5).

Differential gene expression (DGE) analyses were performed comparing *IGFBP7* Q4 versus Q1 tumors in I-SPY2 and SCAN-B. Notably, higher expression of several genes coding for proteins involved in endothelial cell regulation and extracellular matrix remodeling, e.g. *PDGFRB*, *COMP*, *FAP*, *COL1A1*, *OGN*, *LUM*, *COL3A1*, *SPOCK1*, *COL1A2*, *DCN*, *ANGPTL1*, and *SPON1* were seen in *IGFBP7* Q4 tumors, supporting a potential association with an active TME (Supplementary Figure 2 and Tables 6-7). Significantly enriched gene sets in *IGFBP7* Q4 tumors included EMT, TGF- β signaling, coagulation, angiogenesis and downregulated gene sets included MYC response among others (Supplementary Figure 2 and Supplementary Tables 8-9).

Discussion

We report, using data from the I-SPY trial, that breast cancer patients with tumors showing low *IGFBP7* gene expression had more than double the probability of achieving a pCR with neoadjuvant treatment using the combination of ganitumab, metformin, and chemotherapy than using neoadjuvant chemotherapy alone. In contrast, *IGFBP7* expression was unrelated to the probability of chemotherapy alone achieving a pCR. To our knowledge, this is the first

study to date that identifies a specific biomarker related to efficacy of IGF-1R targeting treatment. By the use of *IGFBP7* as a predictor of ganitumab efficacy, we identified a substantial proportion of HER2 negative breast cancers including both poor prognosis HR⁺HER⁻ cancers (54.7% of patients) and TNBC (45.3% of patients) where the addition of ganitumab/metformin plus chemotherapy substantially improves pCR rates. Furthermore, we observed that high *IGFBP7* expression was associated with worse clinical outcome in SCAN-B, a large contemporary population-based breast cancer cohort.

Our finding that low *IGFBP7* expression defines a subset of patients where the addition of ganitumab to chemotherapy approximately doubles the pCR rate, shows the utility of the I-SPY2 trial design. Furthermore, our results demonstrate that detailed examination of the publicly available I-SPY2-990 Data Resource, either hypothesis-driven (as in this case), or agnostically, can provide opportunities to test hypotheses that were not proposed before or during the execution of the trial. The initial assessment of IGF signaling axis in the I-SPY2 ganitumab arm⁵ was negative but failed to include IGFBP7, a non-canonical IGFBP. Had *IGFBP7* been included in the initial assessment of predictive biomarkers in I-SPY2, it would have identified the distinct subgroup where ganitumab showed benefit. Our findings add to the list of important studies^{8,10-13} where the I-SPY2 trial detected activity of novel agents^{7,8,12,13}, even if the benefit is shown to be confined to a biomarker-defined subset of patients.

The molecular features of tumors with high *IGFBP7* expression suggest an aggressive TME that facilitates metastasis. This is consistent with previous studies showing that both systemic and tumor-specific IGFBP7 protein levels were poor prognostic markers in breast cancer^{17,18}, and are also in line with our results from SCAN-B. Despite early studies¹⁴⁻¹⁸, there are many gaps in knowledge concerning the manner by which IGFBP7 modulates signaling by the IGF-1R family, and potentially the efficacy of anti-IGF-1R antibodies. Our findings provide a strong rationale for further experimental work to elucidate how IGFBP7 diminishes the effect of ganitumab and biological influence of IGFBP7 on the TME.

IGFBP7 merits further study as a treatment predictive biomarker in other cancer types such as colon, ovarian and prostate cancer, as well as sarcomas, where IGF-1R targeting agents have been investigated^{4,27}. We emphasize that the lack of data concerning HER2-positive breast cancers does not imply lack of activity in these patients; they simply were not eligible for the ganitumab arm of the I-SPY2 study because of non-availability of clinical safety data

regarding co-administration of ganitumab and HER2 targeting agents. In fact, prior literature²⁸ has suggested benefit of co-targeting IGF-IR and HER2 signaling.

The strengths of this study include the use of contemporary data from both the I-SPY2 trial and SCAN-B cohort. SCAN-B allows for the evaluation of biomarkers in a contemporary real-world setting due to the large-scale RNAseq analysis of consecutively enrolled breast cancer patients¹⁹⁻²¹. Further, I-SPY2 evaluates ganitumab in a randomized controlled setting, providing an ideal platform for investigation of treatment specific biomarkers for ganitumab. The *IGFBP7* expression showed stable associations with clinicopathological factors and molecular features across SCAN-B and ISPY-2.

The current study, while hypothesis driven, is a retrospective analysis of existing datasets and thus the results require further validation in prospective studies. Future work should include investigation of circulating IGFBP7 in relation to ganitumab response, as we did not have access to ISPY-2 blood samples. Another limitation is the lack of data on long-term outcome in I-SPY2, but, it has been previously shown that pCR has a strong association with distant metastasis-free survival²⁹. Due to the combination of ganitumab and metformin plus chemotherapy in the treatment arm, it is not possible to determine with certainty whether *IGFBP7* expression identifies tumors sensitive to ganitumab alone, metformin alone or the combination. However, the MA.32 randomized clinical trial did not show benefit of adjuvant metformin compared to standard therapy in early-stage breast cancer³⁰, suggesting that the observed effect in I-SPY2 was driven by ganitumab.

There are few treatment options beyond chemotherapy in TNBC. Recently PARP inhibitors and immune checkpoint inhibitors have been incorporated in clinical practice^{7,8,13,31-33}. Ganitumab plus chemotherapy combinations have a fairly benign side effect profile⁵, and may improve outcomes for TNBC and other cancers, provided that IGFBP7 is validated as predictive biomarker..

In conclusion, low *IGFBP7* gene expression identifies a subset of breast cancers for whom the addition of ganitumab and metformin to neoadjuvant chemotherapy results in a significantly improved pCR rate compared to neoadjuvant chemotherapy alone. This justifies laboratory studies to address gaps in knowledge concerning roles of IGFBP7 in neoplasia, and the relevance of this protein to IGF-1R targeting agents. Such research, together with the results presented here, may lead to a review of decisions to halt the development of IGF-1

targeting drugs, which were based on disappointing results of prior trials that did not use predictive biomarkers.

Data and code availability

Clinical and RNA-seq data from SCAN-B is accessible from Staaf *et al.* ²¹, available at Mendeley Data. The clinical and transcriptomic data from I-SPY2 are available from the GEO database (GSE194040). This paper does not report original code. Any additional information required to reanalyze the data reported in this paper is available upon request.

Ethics statement

Ethical approvals for I-SPY2 and SCAN-B were obtained in relation to the primary projects and publications ^{10,12,13,19-21}. No other separate approval was obtained for this specific study since it is based on previously published data. All participants signed written informed consent. The study was conducted in accordance with the ethical principles of the Declaration of Helsinki.

Authors' contributions

C Godina: Conceptualization, Data curation, Formal analysis, Methodology, Software, Resources, Visualization, Writing – original draft. **MN Pollak**: Conceptualization, Supervision, Writing – review & editing. **H Jernström**: Conceptualization, Funding acquisition, Formal analysis, Project administration, Supervision, Writing – original draft, Writing – review & editing. All authors have read and agreed to the published version of the manuscript.

Acknowledgments

We thank the I-SPY2 consortium for generously making their data publicly available. We would also like to thank the I-SPY2 data and safety monitoring committee, trial coordinators, project oversight committee, and investigators, and all the patients who volunteered to participate in I-SPY2. Furthermore, we would like to acknowledge the patients who provided consent to participate in the SCAN-B study, the central SCAN-B laboratory at the Division of Oncology, Lund University for sample processing and RNA sequencing, the Swedish National Quality Register for Breast Cancer for clinical and histopathological data, Regional Cancer Center South, and the South Swedish Breast Cancer Group.

This study was funded by the Swedish Cancer Society (CAN 20 0763 and CAN 23 2952), the Faculty of Medicine at Lund University, the Mrs Berta Kamprad Foundation, and the South Swedish Health Care Region (Region Skåne ALF 40620). Dr. Pollak acknowledges funding from the Terry Fox Foundation. The funders had no role in study design and conduct of the study, data collection and analysis, data interpretation, or manuscript preparation and decision to submit the manuscript for publication.

Declaration of interests

The authors declare no competing interests.

Figure legends

Figure 1. Flowchart of included and excluded patients in I-SPY2. pCR rates in the ganitumab/metformin plus chemotherapy arm and chemotherapy control arm, according to *IGFBP7* expression categories.




Figure 2. *IGFBP7* expression in relation to molecular features

Pearson correlations of *IGFBP7* gene expression (continuous) and 15 gene in the IGF/Insulin pathway (INS, INSR, IRS1, IRS2, IGF1, IGF2, IGFALS, IGF1R, IGF2R, IGFBP1, IGFBP2, IGFBP3, IGFBP4, IGFBP5, and IGFBP6) in (A) I-SPY2, and (B) SCAN-B. Pearson correlations of *IGFBP7* gene expression (continuous) and the eight gene modules (Stroma, Lipid, Immune Response, Mitotic Checkpoint, Mitotic Progression, Basal, Early Response, Steroid Response) in (C) I-SPY2, and (D) SCAN-B. *IGFBP7* expression (continuous) by PAM50 subtype in (E) I-SPY2 and (F) SCAN-B. SCAN-B. *IGFBP7* expression (continuous) by receptor subtype in (G) I-SPY2 and (H) SCAN-B.

Figure 2.





Figure 3. *IGFBP7* expression in relation to pCR

Heatmap of differentially expressed genes (DEG) in *IGFBP7* Q4 compared to in *IGFBP7* Q1 tumors in (A) I-SPY2. Patients are along columns and biomarkers are along the rows. Red indicates higher expression and blue lower. Annotation tracks reflects pCR (purple), Receptor Subtype, PAM50 subtype, and trial arm (light red: ganitumab/metformin plus chemotherapy). *IGFBP7* expression (continuous) by pCR in (B) the ganitumab/metformin plus chemotherapy arm and in (C) the control arm. pCR rate by *IGFBP7* quartiles in (D) the ganitumab/metformin plus chemotherapy arm and in (E) the control arm.

Figure 3.



Tables

Table 1. Odds ratio of achieving a pCR in relation *IGFBP7* expression in control and ganitumab+metformin arms

Variable	IGFBP7 Continuous				Interaction	IGFBP7 Quartiles				Interaction
	Control arm HER2 negative		Ganitumab + metformin			Control arm HER2 negative		Ganitumab + metformin		
	OR ¹	95% CI ¹	OR ¹	95% CI ¹		OR ¹	95% CI ¹	OR ¹	95% CI ¹	
Crude										
IGFBP7 Continuous	1.05	0.58, 1.95	0.32	0.15, 0.64	LR ² <i>P</i> = 0.011 <i>P</i> _{interaction} = 0.014					LR ² <i>P</i> = 0.011 <i>P</i> _{interaction} = 0.024 <i>P</i> _{interaction} = 0.021 <i>P</i> _{interaction} = 0.019
IGFBP7 Quartiles	pCR/Total		pCR/Total							
Q1	7/50		15/32			Ref.	Ref.			
Q2	8/40		5/29			1.54	0.50, 4.80	0.24	0.07, 0.74	
Q3	7/42		3/26			1.23	0.39, 3.91	0.15	0.03, 0.53	
Q4	7/43		1/18		1.19	0.38, 3.80	0.07	0.00, 0.39		
Multivariable										
IGFBP7 Continuous	1.23	0.63, 2.45	0.38	0.17, 0.80	LR ² <i>P</i> = 0.017 <i>P</i> _{interaction} = 0.016					LR ² <i>P</i> = 0.015 <i>P</i> _{interaction} = 0.033 <i>P</i> _{interaction} = 0.013 <i>P</i> _{interaction} = 0.031
IGFBP7 Quartiles										
Q1						Ref.	Ref.			
Q2						1.62	0.51, 5.28	0.27	0.07, 0.92	
Q3						1.86	0.53, 6.65	0.17	0.03, 0.69	
Q4						1.38	0.41, 4.73	0.09	0.00, 0.58	
HR+	1.44	0.57, 3.64	0.64	0.16, 2.38		1.38	0.54, 3.56	0.56	0.14, 2.15	
MP2	1.85	0.65, 5.70	1.40	0.20, 10.0		1.81	0.63, 5.57	1.27	0.19, 9.02	
Immune+	1.00	0.35, 2.82	1.88	0.58, 6.33		1.06	0.37, 3.00	1.77	0.52, 6.19	
DRD+	2.35	0.67, 7.84	1.84	0.43, 8.32		2.43	0.76, 8.24	1.91	0.43, 8.91	

¹OR = Odds Ratio, CI = Confidence Interval

²LR = Likelihood ratio

Table 2. pCR in relation *IGFBP7* expression in the chemotherapy control arm and chemotherapy/ganitumab/metformin arm stratified by receptor subtype

	HR+HER2-		TNBC	
	Control arm	Ganitumab + metformin	Control arm	Ganitumab + metformin
<i>IGFBP7</i> Quartiles	pCR/Total (%)	pCR/Total (%)	pCR/Total (%)	pCR/Total (%)
Q1	5/21 (23.8%)	3/14 (21.4%)	2/29 (6.9%)	12/18 (66.7%)
Q2	3/21 (14.3%)	2/16 (12.5%)	5/19 (26.1%)	3/13 (23.1%)
Q3	2/31 (6.4%)	2/15 (13.3%)	5/11 (45.5%)	1/10 (10%)
Q4	4/19 (21.0%)	1/12 (8.3%)	3/24 (12.5%)	0/5 (0%)

Table 3. Odds ratio of achieving a pCR in the ganitumab+metformin arm compared to the HER2-negative control arm, stratified by quartile and median of *IGFBP7* gene expression, I-SPY2

				<i>IGFBP7</i> expression							
All patients				Q1		Q2		Q3		Q4	
Variable		OR ¹	95% CI ¹	OR ¹	95% CI ¹	OR ¹	95% CI ¹	OR ¹	95% CI ¹	OR ¹	95% CI ¹
Crude	pCR/Total										
Trial Arm											
Control	30/179	Ref.		Ref.		Ref.		Ref.		Ref.	
Ganitumab + metformin	24/106	1.45	0.79, 2.65	5.42	1.94, 16.5	0.83	0.23, 2.82	0.65	0.13, 2.61	0.30	0.02, 1.90
Multivariable											
Trial Arm											
Control	30/179	Ref.		Ref.		Ref.		Ref.		Ref.	
Ganitumab + metformin	24/106	1.50	0.79, 2.81	6.17	2.01, 21.3	0.84	0.22, 3.05	0.35	0.06, 1.69	0.25	0.01, 2.55
HR+	22/152	1.07	0.52, 2.18	0.74	0.22, 2.42	1.00	0.23, 4.11	0.00		8.05	1.25, 75.3
MP2	40/147	2.12	0.92, 5.14	0.85	0.20, 4.13	4.00	0.57, 38.1	0.00		1.74	0.19, 20.8
Immune+	33/133	1.22	0.58, 2.57	0.77	0.16, 3.30	0.89	0.19, 4.08	12.7	1.46, 285	0.56	0.06, 4.74
DRD+	36/123	2.09	0.90, 4.99	4.59	1.02, 25.0	1.44	0.26, 8.58	0.04	0.00, 0.71	23.0	1.43, 900
				Low (Q1-2)				High (Q3-4)			
				OR¹	95% CI¹	OR¹	95% CI¹	OR¹	95% CI¹	OR¹	95% CI¹
Crude											
Trial Arm											
Control	30/179			Ref.				Ref.			
Ganitumab + Metformin	24/106			2.50		1.16, 5.49		0.49		0.13, 1.49	
Multivariable											
Trial Arm											
Control	30/179			Ref.				Ref.			
Ganitumab + Metformin	24/106			2.64		1.17, 6.10		0.44		0.11, 1.40	
HR+	22/152			0.83		0.33, 2.01		1.61		0.41, 6.60	
MP2	40/147			1.72		0.55, 6.03		3.23		0.74, 16.6	
Immune+	33/133			0.92		0.33, 2.53		2.57		0.72, 9.69	
DRD+	36/123			2.50		0.85, 7.81		1.10		0.24, 5.01	

¹OR = Odds Ratio, CI = Confidence Interval

Supplementary figure legends

Supplementary Figure 1. Flowchart of included and excluded patients in SCAN-B

Supplementary Figure 2. Molecular analyses of *IGFBP7* expression in I-SPY2 and SCAN-B

Volcano plots showing significantly up- and downregulated genes in *IGFBP7* Q4 compared to in *IGFBP7* Q1 tumors in (A) I-SPY2, and (B) SCAN-B. Dot plots showing activated and suppressed Hallmark Signatures in *IGFBP7* Q4 compared to in *IGFBP7* Q1 tumors in (C) I-SPY2, and (D) SCAN-B. (D).

References

1. Harbeck, N., *et al.* Breast cancer. *Nat Rev Dis Primers* **5**, 66 (2019).
2. Pollak, M. The insulin and insulin-like growth factor receptor family in neoplasia: an update. *Nat Rev Cancer* **12**, 159-169 (2012).
3. Sachdev, D. & Yee, D. Disrupting insulin-like growth factor signaling as a potential cancer therapy. *Mol Cancer Ther* **6**, 1-12 (2007).
4. Ekyalongo, R.C. & Yee, D. Revisiting the IGF-1R as a breast cancer target. *NPJ Precis Oncol* **1**(2017).
5. Yee, D., *et al.* Ganitumab and metformin plus standard neoadjuvant therapy in stage 2/3 breast cancer. *NPJ Breast Cancer* **7**, 131 (2021).
6. Du, L., *et al.* Predicted sensitivity to endocrine therapy for stage II-III hormone receptor-positive and HER2-negative (HR+/HER2-) breast cancer before chemohormone therapy. *Ann Oncol* **32**, 642-651 (2021).
7. Pusztai, L., *et al.* Durvalumab with olaparib and paclitaxel for high-risk HER2-negative stage II/III breast cancer: Results from the adaptively randomized I-SPY2 trial. *Cancer Cell* **39**, 989-998.e985 (2021).
8. Nanda, R., *et al.* Effect of Pembrolizumab Plus Neoadjuvant Chemotherapy on Pathologic Complete Response in Women With Early-Stage Breast Cancer: An Analysis of the Ongoing Phase 2 Adaptively Randomized I-SPY2 Trial. *JAMA oncology* **6**, 676-684 (2020).
9. Wulfkuhle, J.D., *et al.* Evaluation of the HER/PI3K/AKT Family Signaling Network as a Predictive Biomarker of Pathologic Complete Response for Patients With Breast Cancer Treated With Neratinib in the I-SPY 2 TRIAL. *JCO Precis Oncol* **2**(2018).
10. Wolf, D.M., *et al.* Redefining breast cancer subtypes to guide treatment prioritization and maximize response: Predictive biomarkers across 10 cancer therapies. *Cancer Cell* **40**, 609-623.e606 (2022).
11. Magbanua, M.J.M., *et al.* Clinical significance and biology of circulating tumor DNA in high-risk early-stage HER2-negative breast cancer receiving neoadjuvant chemotherapy. *Cancer Cell* **41**, 1091-1102.e1094 (2023).
12. Park, J.W., *et al.* Adaptive Randomization of Neratinib in Early Breast Cancer. *The New England journal of medicine* **375**, 11-22 (2016).
13. Rugo, H.S., *et al.* Adaptive Randomization of Veliparib-Carboplatin Treatment in Breast Cancer. *The New England journal of medicine* **375**, 23-34 (2016).
14. Evdokimova, V., *et al.* IGFBP7 binds to the IGF-1 receptor and blocks its activation by insulin-like growth factors. *Sci Signal* **5**, ra92 (2012).

15. Artico, L.L., *et al.* Physiologic IGFBP7 levels prolong IGF1R activation in acute lymphoblastic leukemia. *Blood Adv* **5**, 3633-3646 (2021).
16. Artico, L.L., *et al.* IGFBP7 Fuels the Glycolytic Metabolism in B-Cell Precursor Acute Lymphoblastic Leukemia by Sustaining Activation of the IGF1R-Akt-GLUT1 Axis. *Int J Mol Sci* **24**(2023).
17. Godina, C., *et al.* Prognostic impact of tumor-specific insulin-like growth factor binding protein 7 (IGFBP7) levels in breast cancer: a prospective cohort study. *Carcinogenesis* **42**, 1314-1325 (2021).
18. Rosendahl, A.H., *et al.* Pre- and Postoperative Circulating IGF-I, IGFBP-3, and IGFBP-7 Levels in Relation to Endocrine Treatment and Breast Cancer Recurrence: A Nested Case-Control Study. *Frontiers in Oncology* **11**(2021).
19. Saal, L.H., *et al.* The Sweden Cancerome Analysis Network - Breast (SCAN-B) Initiative: a large-scale multicenter infrastructure towards implementation of breast cancer genomic analyses in the clinical routine. *Genome Med* **7**, 20 (2015).
20. Rydén, L., *et al.* Minimizing inequality in access to precision medicine in breast cancer by real-time population-based molecular analysis in the SCAN-B initiative. *Br J Surg* **105**, e158-e168 (2018).
21. Staaf, J., *et al.* RNA sequencing-based single sample predictors of molecular subtype and risk of recurrence for clinical assessment of early-stage breast cancer. *NPJ Breast Cancer* **8**, 94 (2022).
22. Fredlund, E., *et al.* The gene expression landscape of breast cancer is shaped by tumor protein p53 status and epithelial-mesenchymal transition. *Breast cancer research* **14**, R113 (2012).
23. Ritchie, M.E., *et al.* limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* **43**, e47 (2015).
24. Wu, T., *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *Innovation (Camb)* **2**, 100141 (2021).
25. Liberzon, A., *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst* **1**, 417-425 (2015).
26. McShane, L.M., *et al.* Reporting Recommendations for Tumor Marker Prognostic Studies (REMARK). *Journal of the National Cancer Institute* **97**, 1180-1184 (2005).
27. Akshintala, S., *et al.* Phase I trial of Ganitumab plus Dasatinib to Cotarget the Insulin-Like Growth Factor 1 Receptor and Src Family Kinase YES in Rhabdomyosarcoma. *Clinical cancer research* **29**, 3329-3339 (2023).
28. Lu, Y., Zi, X., Zhao, Y., Mascarenhas, D. & Pollak, M. Insulin-like growth factor-I receptor signaling and resistance to trastuzumab (Herceptin). *Journal of the National Cancer Institute* **93**, 1852-1857 (2001).
29. Yee, D., *et al.* Association of Event-Free and Distant Recurrence-Free Survival With Individual-Level Pathologic Complete Response in Neoadjuvant Treatment of Stages 2 and 3 Breast Cancer: Three-Year Follow-up Analysis for the I-SPY2 Adaptively Randomized Clinical Trial. *JAMA oncology* **6**, 1355-1362 (2020).
30. Goodwin, P.J., *et al.* Effect of Metformin vs Placebo on Invasive Disease-Free Survival in Patients With Breast Cancer: The MA.32 Randomized Clinical Trial. *Jama* **327**, 1963-1973 (2022).
31. Schmid, P., *et al.* Event-free Survival with Pembrolizumab in Early Triple-Negative Breast Cancer. *The New England journal of medicine* **386**, 556-567 (2022).
32. Bianchini, G., De Angelis, C., Licata, L. & Gianni, L. Treatment landscape of triple-negative breast cancer - expanded options, evolving needs. *Nat Rev Clin Oncol* **19**, 91-113 (2022).

33. Tutt, A.N.J., *et al.* Adjuvant Olaparib for Patients with BRCA1- or BRCA2-Mutated Breast Cancer. *The New England journal of medicine* **384**, 2394-2405 (2021).