Persistent Hyposmia as Surrogate for α-Synuclein-Linked Brain Pathology

Brit Mollenhauer[1,2,6,7]*, Juan Li[3,4,6]*, Sebastian Schade[2], Sandrina Weber[1], Claudia Trenkwalder[2],

Luis Concha-Marambio[5], Julianna J. Tomlinson[3,6,7], aSCENT-PD Investigators[6,7], Michael G.

Schlossmacher[3,6,7#]

[1]Department of Neurology, University of Goettingen, Kassel

[2]Paracelsus-Elena-Klinik, Kassel, Germany

[3]Neuroscience Program, [4]Clinical Epidemiology Program, Ottawa Hospital Research Institute,

Ottawa, ON., Canada

[5]Amprion, Inc.; San Diego, CA; USA

[6]Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase,

MD. 20815

[7]aSCENT-PD Investigators include: Ben Arenkiel, Zhandong Liu, Brit Mollenhauer, Josef

Penninger, Maxime Rousseaux, Armen Saghatelyan, Natalina Salmaso, Michael G. Schlossmacher,

Christine Stadelmann-Nessler, Julianna J. Tomlinson, John M. Woulfe

*These authors contributed equally
#The corresponding author

Michael G. Schlossmacher, MD, FRCPC
Ottawa Hospital Research Institute; Ottawa, Canada
mschlossmacher@toh.ca
Phone: +1 613-562-5462

ABSTRACT

Impaired olfaction can be associated with neurodegenerative disorders. We examined odor

identification in newly diagnosed patients with parkinsonism and those at increased risk, measured

olfactory performances longitudinally, and juxtaposed results to cerebrospinal fluid (CSF) values.

Using Sniffin'-Sticks-Identification Tests (SST-ID), we examined 312 age-matched individuals at

a German center, including: 126 with Parkinson disease (PD), 109 healthy controls, 25 with other

neurodegenerative disorders and 52 with a REM-sleep behavior disorder (RBD). As expected, PD

patients had significantly lower SST-ID scores than controls. Scent identification by subjects with

other neurodegenerative diseases fell between those with PD and healthy individuals. Those with

isolated RBD, who subsequently converted to PD or dementia, had lower baseline scores than non-

converters. When monitoring olfaction in participants up to a decade, we saw small group

differences in progression rates for hyposmia. However, these variations were insignificant after

controlling for age, sex and length of intervals between testing. When analyzing participants' sense

of smell versus several CSF biomarkers linked to neurodegeneration, we found no correlation with

SST-ID scores. However, the means for normalized concentrations of α-synuclein, total tau,

phosphorylated tau and amyloid-β peptide$_{42}$ were reduced in PD. We also identified significant

age- and sex-linked differences in CSF values. Finally, we compared olfaction to the results of a

validated α-synuclein 'Seed Amplification Assay' (SAA) using CSF. We found that hyposmia

strongly correlated with a positive CSF α-synuclein SAA-test. We conclude that chronically

impaired olfaction in older adults is strongly associated with a positive α-synuclein SAA-test from

CSF but not with the concentrations of several, neuropathologically relevant CSF markers. We

posit that simple-to-administer, quantitative smell tests could serve as inexpensive screening tools

in future population studies for the identification of α-synuclein-related brain disorders, including

Parkinson's during its premotor phase.

MANUSCRIPT

Impaired olfaction can be associated with neurodegenerative disorders.[1] Its pathogenesis,

progression over time and any sex differences remain unknown. We examined odor identification

in newly diagnosed patients with parkinsonism and those at increased risk, measured olfactory

performances longitudinally, and juxtaposed results to cerebrospinal fluid (CSF) values.

Using Sniffin'-Sticks-Identification Tests (SST-ID; scale, 0-16), we examined 312 age-matched

individuals at a single center[2], including: 126 with Parkinson disease (PD), 109 healthy controls,

25 with other neurodegenerative disorders and 52 with a REM-sleep behavior disorder (RBD). In

16 RBD subjects, a neurodegenerative disorder[3] was subsequently diagnosed (**Suppl. TableS1**).

The participants' baseline SST-ID scores are shown in **Fig.1a**, with reduced olfaction defined as a

score below the 10^{th} percentile for each sex and age group.[4] As expected,[1] PD patients had lower

SST-ID scores than controls (P<2e-16). Scent identification by subjects with other

neurodegenerative diseases fell between those with PD (P=6.6e-05) and healthy individuals

(P=0.018). Those with isolated RBD, who subsequently converted to PD or dementia, had lower

baseline scores than non-converters (P=0.0021). There was no sex difference after controlling for

age (**Fig.1a**).

When monitoring olfaction in participants up to a decade, we saw small group differences in

progression rates for hyposmia (**Fig.1b**; **Suppl**. **Fig.S1**). However, these variations were

insignificant after controlling for age, sex and length of intervals between testing.

When analyzing participants' sense of smell versus several CSF biomarkers[2] linked to

neurodegeneration, we found no correlation with SST-ID scores. However, the means for

normalized concentrations of α-synuclein, total tau, phosphorylated tau and amyloid-β peptide$_{42}$

were reduced in PD. We also identified significant age- and sex-linked differences in CSF values

(**Suppl**. **Fig.S2**).

Finally, we compared olfaction to the results of a validated α-synuclein 'Seed Amplification Assay'

(SAA) using CSF. Positivity in this binary test has been shown to reflect α-synuclein-linked brain

pathology.[5] We found that hyposmia strongly correlated with a positive CSF α-synuclein SAA-

test [R: 0.63; R^2: 0.4 (P<2.2e-16); **Fig.1c**]. There was no age or sex difference in its outcome.

We conclude that chronically impaired olfaction in older adults is strongly associated with a

positive α-synuclein SAA-test from CSF but not with the concentrations of several,

neuropathologically relevant CSF markers. Future studies will determine at what age hyposmia

and SAA-test positivity begin, and why. We posit that simple-to-administer, quantitative smell

tests could serve as inexpensive screening tools in future population studies for the identification

of α-synuclein-related brain disorders, including Parkinson's during its premotor phase.

REFERENCES

1. Haehner A, Hummel T, Reichmann H. (2011) Olfactory loss in Parkinson's disease.
Parkinsons Dis. 2011:450939.

2. Mollenhauer B, Trautmann E, Sixel-Döring F, Wicke T, Ebentheuer J, Schaumburg M,
Lang E, Focke NK, Kumar KR, Lohmann K, Klein C, Schlossmacher MG, Kohnen R,
Friede T, Trenkwalder C; DeNoPa Study Group. (2013) Nonmotor and diagnostic findings
in subjects with de novo Parkinson disease of the DeNoPa cohort. *Neurology* 81(14): 1226-
34

3. Postuma RB, Iranzo A, Hu M, Högl B, Boeve BF, Manni R, Oertel WH, Arnulf I, Ferini-
Strambi L, Puligheddu M, Antelmi E, Cochen De Cock V, Arnaldi D, Mollenhauer B,
Videnovic A, Sonka K, Jung KY, Kunz D, Dauvilliers Y, Provini F, Lewis SJ, Buskova J,
Pavlova M, Heidbreder A, Montplaisir JY, Santamaria J, Barber TR, Stefani A, St Louis
EK, Terzaghi M, Janzen A, Leu-Semenescu S, Plazzi G, Nobili F, Sixel-Doering F, Dusek
P, Bes F, Cortelli P, Ehgoetz Martens K, Gagnon JF, Gaig C, Zucconi M, Trenkwalder C,
Gan-Or Z, Lo C, Rolinski M, Mahlknecht P, Holzknecht E, Boeve AR, Teigen LN, Toscano
G, Mayer G, Morbelli S, Dawson B, Pelletier A. (2019) Risk and predictors of dementia
and parkinsonism in idiopathic REM sleep behaviour disorder: a multicentre study. *Brain*.
142(3):744-759.

4. Oleszkiewicz A, Schriever VA, Croy I, Hähner A, Hummel T. (2019) Updated Sniffin'
Sticks normative data based on an extended sample of 9139 subjects. *Eur Arch*
Otorhinolaryngol. 276(3):719-728.

5. Arnold MR, Coughlin DG, Brumbach BH, Smirnov DS, Concha-Marambio L, Farris CM,
Ma Y, Kim Y, Wilson EN, Kaye JA, Hiniker A, Woltjer RL, Galasko DR, Quinn JF. (2022)
α-Synuclein Seed Amplification in CSF and Brain from Patients with Different Brain
Distributions of Pathological α-Synuclein in the Context of Co-Pathology and Non-LBD
Diagnoses. *Ann Neurol*. 92(4):650-662.

FIGURE

Figure 1: Association of chronically reduced olfaction with α-synuclein-linked brain diseases

(A) Relationship between baseline SST-ID scores (y-axis) and disease group (columns), sex (rows),
and age at baseline enrolment (x-axis). Each square represents a participant. Classification of
olfaction (square colors) was defined by SST-ID scores at or below the 10th percentile for each sex
and age group, as published.[4] The "reduced olfaction" group includes both hyposmia and anosmia.

(B) Changes in SST-ID scores over time. The legend underneath the graph summarizes the color
representation and corresponding number of participants in each group. Each square represents a
study subject at a given time point, with overlap seen due to integer scores and number of months
tracked. Each thin line tracks the changes of an individual participant, whereas thicker linear
regression lines summarize the overall trend.

(C) A Cummings estimation plot was used to illustrate CSF α-synuclein SAA-test results *versus*
SST-ID score distributions in each group at study enrolment. Each data point in the upper panel
represents the score of each participant, with colors representing CSF α-synuclein SAA-status as
either positive or negative. The vertical lines represent the conventional mean ± standard deviation
error bars. Lower panels show the mean group difference (i.e., effect size) and its 95% confidence
interval (CI) estimated by bias-corrected and accelerated bootstrapping, using healthy controls
(HC) as the reference group. SST-ID score and CSF α-synuclein SAA-status distinguish PD
patients from age- and sex-matched healthy controls with AUC values of 0.89 (95% CI 0.85-0.93)
and 0.95 (95% CI 0.92-0.98), respectively.

Abbreviations denote SST-ID as Sniffin'-Sticks-Identification test; DeNoPa, *De Novo* Parkinson
Study; RBD, isolated REM sleep behavior disorder; PD, Parkinson disease; HC, healthy control;
OND as other neurological diseases; CSF, cerebrospinal fluid; SAA, seed amplification assay.

SUPPLEMENTARY APPENDIX

Supplement to: Mollenhauer B, Li J *et al*. Persistent Hyposmia as Surrogate of α-Synuclein-

Linked Brain Pathology

This appendix has been provided by the authors to give readers additional information about the

work.

TABLE OF CONTENTS

SUPPLEMENTARY INFORMATION: METHODS

Source of data and participants

We used de-identified data from the 'DeNoPa cohort', an ongoing, single-center study based at

the Paracelsus-Elena Klinik in Kassel, Germany. Data from this study were downloaded on May

16th, 2023. The cohort[2] represents an observational, longitudinal study of patients with a newly

established ("*de novo*") diagnosis of PD (UK Brain Bank Criteria[2]) that were naïve to L-DOPA

therapy at baseline, and of age- as well as sex-matched, neurologically healthy controls (HC).

Exclusion criteria included previously known or subsequently detected brain conditions, such as

normal pressure hydrocephalus, cerebrovascular disease, early cognitive impairment, features of

atypical parkinsonism, and medication-induced parkinsonism. HC subjects were recruited through

relatives and friends of enrolled PD subjects, other patients of the hospital-based clinic as well as

through newspaper advertisements in 2009. Control subjects had to be without any active, known

or previously treated condition of their central nervous system. Diagnostic accuracies for study

participants were ensured by ongoing follow-up visits every two years (as of 2023, 10th year

follow-up visits are underway). Twenty-seven patients classified as PD at baseline were later re-

grouped as patients with "other neurological diagnoses" (OND).

The 'DeNoPa RBD cohort' is a prospective study that follows patients with clinically diagnosed

isolated RBD, using an inpatient polysomnogram and related questionnaires, with up to 8 years of

follow-up. Both recruitment and follow-up visits are currently continuing. Conversion to PD and

OND was confirmed based on the UK Brain Bank Criteria and other diagnostic criteria.[3] Of note,

there is no overlap in participants between the two cohorts, and participants' olfactory function

had no influence on their recruitment into either of the two cohorts.

The launch and ongoing analyses of these cohorts were approved by the Investigational Review

Board of Landesaerztekammer Hesse in Frankfurt, Germany (FF89/2008) and by the Regulatory

Ethics Board of The Ottawa Hospital in Ottawa, Canada (20180010-01H).

Procedures and statistical analyses

Sniffin' Sticks test (SST): SST comprises a supervised test for one's sense of smell using pen-like

odor dispensing devices and is done in the clinic.[4] This study focused on its Identification subtest

(SST-ID) containing 16 items, in which subjects choose the odor in each stick from four options

(one correct, three distractors). The German language version of the SST-ID test was completed

by DeNoPa PD and HC participants at baseline and serially during follow-up visits, as well as by

RBD participants at baseline and during follow-up visits (see **Figure 1.b** for details).

An in vitro α-synuclein-directed seed amplification assay (SAA): Methodological details of the

SAA test, which was performed at Amprion Inc., San Diego, CA. used in this study have been

described elsewhere.[5] The SAA data were available for a subset of DeNoPa (HC: 63; PD: 92; OND:

17) and RBD (n = 27) participants and downloaded on August 15th, 2023.

Biomarkers measured in cerebrospinal fluid (CSF): Methodological details of the CSF biomarkers

quantification used in this study have been described elsewhere.[6-8] The CSF data at baseline were

available for a subset of DeNoPa (HC: 61; PD: 98; OND: 19) participants and downloaded on May

10th, 2019. Five biomarkers, namely the concentrations of total α-synuclein, total tau,

phosphorylated tau, amyloid-β peptide$_{42}$, and total protein, were studied here (**Supplementary**

Figure S1 a-e). The first four readouts were then further normalized using CSF total protein

concentrations measurement in each person (**Supplementary Figure S1 f-i**).

Statistical analyses: Demographic and diagnostic characteristics of the study cohorts are

summarised using n (%) or median (the interquartile range (IQR)). The reported p-values represent

the significance from corresponding Chi-squared test, Welch Two Sample t-test, or ANOVA with

Bonferroni or Dunnett post-hoc test corrections, using the PD group as the reference group.

Coefficients reported in **Supplementary Figures S1, S2** were calculated using multivariate linear

regression. Results with p-values smaller than 0.05 were considered significant. Statistical

analyses were performed using 'R' (version 4.2.0). Cummings estimation plots were generated

using 'dabestr',[9] all other plots were generated using 'ggplot2'.[10] The package 'pROC'[11] was used

for generating ROC and AUC.

CODE AND DATA AVAILABILITY

The code for data analyses and figures is publicly accessible in GitHub
(https://github.com/JuanLiOHRI/SAA). Data can be accessed via zenodo
(https://zenodo.org/records/10222976).

TABLE S1

Demographic characteristics of adults enrolled in two single-centre cohorts from Germany

Characteristic	DeNoPa			RBD[2]
	PD	HC	OND[1]	
N	126	109	25	52
Sex: Male	82 (65.1)	67 (61.5) [p = 0.57]	15 (60) [p = 0.63]	35 (67.3)
Age at baseline in years				
All	66 (13.8)	65 (10) [p = 0.98]	71 (12.0) [p = 0.01]	68 (11.2)
Male	66 (14.8)	67 (8) [p = 0.81]	70 (12) [p = 0.20]	68 (10.5)
Female	66 (12.8)	64 (9.8) [p = 0.90]	74 (9.5) [p = 0.03]	68 (12)
Parkinsonism duration at baseline in months	14.5 (14.75)	n.a.	15 (25) [p = 0.25]	n.a.
Follow-up time in months	118 (48.5)	121 (4.0) [p = 0.05]	72 (69) [p = 0.03]	48.5 (57)

[1] Detailed diagnosis in the OND group: DLB = 3, ET = 6, MSA = 4, PSP = 5, RLS = 1, DD PD =

2, Vascular PD = 4.

[2] Detailed diagnoses in the converted cases in the RBD cohort: PD = 10, DLB = 5, DD PD = 1.

Values are n (%) or median (IQR). The reported p-values in square brackets represent the

significance from corresponding Pearson's Chi-squared test, Welch two sample t-test, or ANOVA

with Dunnett post-hoc test, using the PD group as the reference group.

DeNoPa = *De Novo* Parkinson Study. RBD = isolated REM sleep behavior disorder. IQR = the

interquartile range. PD = Parkinson disease. HC = healthy control. OND = other neurological

diseases. DLB = dementia with Lewy bodies. ET = essential tremor. MSA = multiple system

atrophy. PSP = progressive supranuclear palsy. RLS = restless leg syndrome. DD PD = uncertain

neurological diagnosis. n.a. = not available

LEGENDS / SUPPLEMENTARY FIGURES

<u>Suppl. Figure S1:</u> Rate of changes of SST-ID scores over time.

The rate of change was calculated by dividing the score difference of two successive visits with the time difference in months. The legend underneath the graph summarizes the color representation and corresponding number of participants in each group. Each square represents a study subject at a given time point. Each thin line tracks the changes of an individual participant, whereas thicker linear regression lines summarize the overall trend.

The table lists the coefficients and p-values of linear regression using "rate ~ month + group + age + sex" as the formula.

<u>**Suppl. Figure S2:**</u> **Relationship between select CSF biomarkers and disease group and sex.**

(A-I) Cummings estimation plots were used to illustrate and compare select CSF biomarker

distributions, as indicated, in each study group. Each point represents a participant. Classification

of olfaction (point colors) was defined as SST-ID scores below the 10th percentile in each sex and

age group. The vertical lines represent the conventional mean ± standard deviation error bars. The

lower panels show the mean group difference (the effect size) and its 95% confidence interval (CI)

estimated by bias-corrected and accelerated bootstrap, using HC in each sex as the reference group.

The table in each individual panel **(A-I)** lists the coefficients, p-values, and significance levels of

linear regression using "biomarker ~ group + age + sex + olfaction" as the formula.

SST-ID = Sniffin' Sticks Identification test. DeNoPa = *De Novo* Parkinson Study. PD = Parkinson

disease. HC = healthy control. OND = other neurological diseases. CSF = cerebrospinal fluid.

249
250

SUPPLEMENTARY REFERENCES

1.   Mollenhauer B, Trautmann E, Sixel-Döring F, Wicke T, Ebentheuer J, Schaumburg M, Lang
E, Focke NK, Kumar KR, Lohmann K, Klein C, Schlossmacher MG, Kohnen R, Friede T,
Trenkwalder C; DeNoPa Study Group. (2013) Nonmotor and diagnostic findings in subjects
with de novo Parkinson disease of the DeNoPa cohort. *Neurology* 81(14): 1226-34

2.   Hughes, A. J., Daniel, S. E., Kilford, L. & Lees, A. J. Accuracy of clinical diagnosis of
idiopathic Parkinson's disease. A clinico-pathological study of 100 cases. *J. Neurol.*
Neurosurg. Psychiatry 55, 181–184 (1992).

3.   Postuma RB, Berg D, Stern M, Poewe W, Olanow CW, Oertel W, Obeso J, Marek K, Litvan
I, Lang AE, Halliday G, Goetz CG, Gasser T, Dubois B, Chan P, Bloem BR, Adler CH,
Deuschl G. MDS clinical diagnostic criteria for Parkinson's disease. *Mov Disord.* 2015
Oct;30(12):1591-601.

4.   Hummel T, Sekinger B, Wolf SR, Pauli E, Kobal G. (1997) 'Sniffin' sticks': olfactory
performance assessed by the combined testing of odor identification, odor discrimination and
olfactory threshold. *Chem Senses*. 22(1):39-52.

5.   Concha-Marambio L, Weber S, Farris CM, Dakna M, Lang E, Wicke T, Ma Y, Starke M,
Ebentheuer J, Sixel-Döring F, Muntean ML, Schade S, Trenkwalder C, Soto C, Mollenhauer
B. (2023) Accurate Detection of α-Synuclein Seeds in Cerebrospinal Fluid from Isolated
Rapid Eye Movement Sleep Behavior Disorder and Patients with Parkinson's Disease in the
DeNovo Parkinson (DeNoPa) Cohort. *Mov Disord.* 2023 Apr;38(4):567-578.

6.   Mollenhauer B, Cullen V, Kahn I, Krastins B, Outeiro TF, Pepivani I, Ng J, Schulz-Schaeffer
W, Kretzschmar HA, McLean PJ, Trenkwalder C, Sarracino DA, Vonsattel JP, Locascio JJ,
El-Agnaf OM, Schlossmacher MG. (2008) Direct quantification of CSF alpha-synuclein by
ELISA and first cross-sectional study in patients with neurodegeneration. *Exp Neurol.*
213(2):315-25.

7.   Mollenhauer B, Locascio JJ, Schulz-Schaeffer W, Sixel-Döring F, Trenkwalder C,
Schlossmacher MG. (2011) α-Synuclein and tau concentrations in cerebrospinal fluid of
patients presenting with parkinsonism: a cohort study. *Lancet Neurol*. 2011 Mar;10(3):230-
40. doi: 10.1016/S1474-4422(11)70014-X. Erratum in: *Lancet Neurol*. 2011 Apr;10(4):297.

8.   Mollenhauer B, Trautmann E, Taylor P, Manninger P, Sixel-Döring F, Ebentheuer J,
Trenkwalder C, Schlossmacher MG. (2013) Total CSF α-synuclein is lower in de novo
Parkinson patients than in healthy subjects. *Neurosci Lett.* 532:44-8.
9.   Ho J.W., Tumkaya T., (2020). *dabestr: Data Analysis using Bootstrap-Coupled Estimation.*
https://cran.r-project.org/web/packages/dabestr/index.html
10.   Wickham H (2016). *ggplot2: Elegant Graphics for Data Analysis.* Springer-Verlag New York.
ISBN 978-3-319-24277-4, https://ggplot2.tidyverse.org.
11.   Robin X., et al. (2023). *pROC: Display and Analyze ROC Curves*. https://cran.r-
project.org/web/packages/pROC/index.html

ACKNOWLEDGEMENTS

This research was funded in part by Aligning Science Across Parkinson's [Grant ID: ASAP-

020625] administered by the Michael J. Fox Foundation for Parkinson's Research (MJFF) and by

the Parkinson Research Consortium Ottawa.

A

B

304 **C**

306 **Figure 1: Association of chronically reduced olfaction with α-synuclein-linked brain diseases**

	Coefficient	p-value
(Intercept)	0.02151	0.287
Month	-9.23E-05	0.188
Group: PD	-0.004101	0.466
Group: RBD_Not converted	-0.008664	0.351
Group: RBD_Converted	-0.01197	0.276
Age	-0.000252	0.4
Sex: Male	-0.001891	0.708

312 **<u>Suppl. Figure S1:</u> Rate of changes of SST-ID scores over time.**

A: Total α-synuclein in CSF by group and sex

	Coefficient	p-value	
(Intercept)	785.093	0.048464	*
Group: PD	-88.454	0.50093	
Group: OND	-53.941	0.755369	
Age	16.607	0.00432	**
Sex: Male	-370.055	0.000371	***
Olfaction: Reduced	-28.366	0.822528	

B: Total tau in CSF by group and sex

	Coefficient	p-value	
(Intercept)	39.256	0.599	
Group: PD	-30.589	0.22	
Group: OND	12.407	0.699	
Age	4.324	0.0000982	***
Sex: Male	-79.644	0.0000584	***
Olfaction: Reduced	8.801	0.712	

C: Phosphorylated tau in CSF by group and sex

	Coefficient	p-value	
(Intercept)	21.6646	0.039964	*
Group: PD	-3.9124	0.258921	
Group: OND	-1.2608	0.789619	
Age	0.4845	0.001663	**
Sex: Male	-9.8005	0.000468	***
Olfaction: Reduced	1.0947	0.742111	

D: Ab42 in CSF by group and sex

	Estimate	Pr(>|t|)	
(Intercept)	894.6618	1.82E-10	***
group: PD	-66.8745	0.1294	
group: OND	-94.3713	0.0976	.
age	-0.1737	0.9279	
sex: Male	-5.0841	0.8818	
olfaction: Reduced	69.7542	0.0994	.

E: Total protein content of CSF by group and sex

	Coefficient	p-value	
(Intercept)	589.587	0.00277	**
Group: PD	121.162	0.06337	.
Group: OND	166.213	0.0487	*
Age	-2.386	0.39993	
Sex: Male	93.304	0.06529	.
Olfaction: Reduced	-48.689	0.43644	

F: Normalized total α-synuclein in CSF by group and sex

	Coefficient	p-value	
(Intercept)	2.35543	0.041	*
Group: PD	-0.70624	0.0646	.
Group: OND	-0.61559	0.2204	
Age	0.03662	0.0289	*
Sex: Male	-1.44577	2.20E-06	***
Olfaction: Reduced	0.0792	0.8288	

G: Normalized total tau in CSF by group and sex

	Coefficient	p-value	
(Intercept)	0.203781	0.32572	
Group: PD	-0.145039	0.03651	*
Group: OND	-0.055866	0.53023	
Age	0.009416	0.00202	**
Sex: Male	-0.284432	3.33E-07	***
Olfaction: Reduced	0.034067	0.60652	

H: Normalized phosphorylated tau in CSF by group and sex

	Coefficient	p-value	
(Intercept)	0.0713408	0.0289	*
Group: PD	-0.021574	0.0451	*
Group: OND	-0.0179481	0.2205	
Age	0.0010134	0.032	*
Sex: Male	-0.0413419	2.64E-06	***
Olfaction: Reduced	0.0052111	0.6126	

I: Normalized Ab42 in CSF by group and sex

	Estimate	Pr(>|t|)	
(Intercept)	2.180539	1.4E-05	***
group: PD	-0.418569	0.01084	*
group: OND	-0.524225	0.01338	*
age	0.001771	0.80309	
sex: Male	-0.396461	0.00202	**
olfaction: Reduced	0.206328	0.18744	

314   **Suppl. Figure S2: Relationship between select CSF biomarkers and disease group and sex.**