A Increased trunk flexion in standing up is related to muscle weakness rather than pain avoidance in individuals with unilateral knee pain; a simulation study.

Eline van der Kruk[1*], Thomas Geijtenbeek[1,2]

[1]Department of Biomechanical Engineering, Faculty of Mechanical Engineering (3me), Delft
University of Technology, the Netherlands

[2]GoatStream, the Netherlands

* Correspondence:
Corresponding Author
e.vanderkruk@tudelft.nl

Keywords: *predictive simulation, neuromuscular model, knee osteoarthritis, ageing, sit-to-walk,*
timed-up-and-go

Abstract

The 'Timed Up and Go' test (TUG) is a widely used clinical tool for assessing gait and balance,
relying primarily on timing as a measure. However, there are more observable biomechanical
compensation strategies within TUG that are indicative of underlying neuromuscular issues and
movement priorities. In individuals with unilateral knee osteoarthritis, an increased trunk flexion
during TUG is a common phenomenon, often attributed to muscle weakness and/or pain avoidance.
Unfortunately, differentiating between these underlying causes is difficult to accomplish using
experimental studies. This study aimed to distinguish between muscle weakness and pain avoidance
as contributing factors, using predictive neuromuscular simulations of the sit-to-walk movement.
Muscle weakness was simulated by reducing the maximum isometric force of the vasti muscles
(ranging from 20% to 60%), while pain avoidance was integrated as a movement objective, ensuring
that peak knee load did not exceed predefined thresholds (2-4 times body weight). The simulations
demonstrate that a decrease in muscular capacity led to greater trunk flexion. Pain avoidance led to
slower movement speeds and altered muscle recruitments, but not to greater trunk flexion. The
predictive simulations thus indicate that increased trunk flexion is more likely the result of lack of
muscular reserve rather than pain avoidance. These findings align with reported differences in
kinematics and muscle activations between moderate and severe knee osteoarthritis patients,
emphasizing the impact of severe muscle weakness in those with advanced knee osteoarthritis. The
simulations offer valuable insights into the mechanisms behind altered movement strategies,
potentially guiding more targeted treatment.

1   Introduction

Quality of healthcare is at risk due to an increase of age-related health issues and a shortage of
healthcare workers in the near future. The primary focus of most governments is to prolong the
independence of older adults in their own homes. A crucial daily activity that supports this
independence is the act of rising from a seated position [1]. This seemingly simple task becomes
increasingly challenging as individuals age and is closely linked to a higher risk of falls [2].

In Western societies, a significant percentage of individuals over the age of 65 suffer from symptomatic osteoarthritis (OA), particularly affecting the knee joint [3]. The 'Timed Up and Go' test (TUG) is a clinical evaluation of gait and balance, serving as a diagnostic tool in these individuals [4]. It involves a sit-to-walk exercise where individuals rise from a chair, walk a distance of 3 meters, turn, walk back to the chair, and sit down. In a clinical setting, the time taken for this exercise is measured, if the exercise takes more than 30 seconds the patient is indicated with an high risk of falling.

Recent experimental research has demonstrated that beyond timing, biomechanical compensation strategies are indicative of the underlying age-related changes in the neuromuscular system and movement priorities in sit-to-walk [5]. Individuals with unilateral OA often exhibit asymmetrical movements while standing up, bearing additional weight on the unaffected side by leaning the trunk or by using asymmetric arm movements [6]–[10]. Additionally, they display increased maximum trunk flexion [6], [11]. Recent systematic reviews suggest that these altered trunk flexion patterns may be related to muscle weakness and/or pain avoidance [12].

Whereas experimental studies cannot distinguish between these underlying causes [13], predictive simulations are a promising tool to determine the interconnectivity and interdependency of the neuromuscular capacity, reinforcement schemes, sensory integration, and adaptation strategies in standing up. We recently published the validation of a planar neuromusculoskeletal model with reflex-based muscle control that simulates the sit-to-walk movement [14]. This communication aims to leverage this predictive neuromuscular framework to determine whether the observed alterations in trunk flexion strategies in individuals with unilateral knee osteoarthritis are more likely attributed to muscle weakness or to pain avoidance.

2   Methods

2.1   Model and controller

The design and validation of the musculoskeletal model and controller have been previously published [14]. We employed the H1120 model, featuring 11 degrees of freedom and 20 Hill-type muscles, which was initially developed as an OpenSim3 model and translated into Hyfydy (Fig. 1) [15]. Peak isometric forces in this model are based on a lower limb model by Delp et al. (1990) (G2392) [16].

Contact forces between the feet and the ground and between the buttocks and the chair were modelled with a Hunt Crossley force spheres and box, respectively. The simulation of the sit-to-walk movement involved the utilization of a standing-up controller [14] followed by a gait controller [17]. The standing-up controller operates on a reflex control principle, featuring two distinct states, each with its own set of control parameters. The reflex controller is based on monosynaptic and antagonistic proprioceptive feedback from the muscles and vestibular feedback linked to the pelvis tilt. The timing of the transition between the states is part of the optimization problem. To account for neural latencies, we incorporated data from previous studies.

States calculated from the model were muscle length (l), muscle velocity (v), muscle force (F), and pelvis tilt orientation (θ) and velocity ($\dot{\theta}$). The optimization problem encompassed a total of 551 free parameters, comprising controller gains (KC, KL+, KF±, Kp, and Kv), muscle length feedback offsets (lo), proportional feedback of θ (θo), proportional feedback of the lumbar and thoracic joints, transition times between the controllers, and stance load threshold for the gait state controller.

Figure 1 - The musculoskeletal model (H1120) has 11 degrees of freedom and is actuated using 20
Hill-type muscle-tendon units. The H1120 model is available on [15].

[Here Figure 1]

2.2   Objective Function

We used the objective measures from the published sit-to-walk controller [14]. These measures
encompassed the following criteria: a gait velocity measure with a minimum threshold of 0.8 m/s,
range penalties designed to replicate joint limitations, including lumbar extension (-50° to 0°), thorax
extension (-15° to 15°), pelvis tilt (-50° to 30°), and ankle angle (-60° to 60°), a constraint on knee
limit force (500 N/m), and the minimization of head acceleration (1m/s^2). Additionally, we included
an energy estimate aimed at reducing energy consumption during the stand-up phase and minimizing
the cost of transport during gait. The measure consisted of metabolic energy expenditure based on
[18], and was complemented by the minimization of cubed muscle activations, along with torque
minimization at the lumbar and thoracic joints to serve as a proxy for the absence of trunk muscles.
During the optimization process, this measure eventually became the sole non-zero term within the
objective function.

2.3   Optimization algorithm

The optimization process was conducted using the OpenSource software SCONE with the
Covariance Matrix Adaptation Evolutionary Strategy (CMA-ES) [19]. Simulations were performed
using the Hyfydy simulation engine [20]. The population size of each generation was 10, and the
number of iterations varied depending on the simulation duration. Multiple parallel optimizations
were executed with the same initial guesses, and the best set was subsequently employed as the
starting point for the next set of optimizations.

2.4   Conditions

2.4.1 Muscle weakness (VAS)

Muscular capacity progressively declines with age, with the largest relative decline in knee flexion-
extension. Comparing studies on PCSA in older and young adults, the most significant relative PCSA
decrease between these groups is in the vastus lateralis [21], [22] (scaling to the soleus which has the
largest PCSA; in young adults VASINT is 83% of soleus max. force, in older adults 53% of soleus
max. force). Individuals with knee osteoarthritis also have pronounced quadriceps strength decline
[23]–[25].

To simulate reduced muscular capacity in knee extension, we systematically bilaterally decreased the
vasti (VAS) maximum isometric force by 20%, 30%, 40%, 50%, and 60%. We initiated the model
with the optimized sit-to-walk controller for the standard seat [14] (neutral condition). The best set of
the reduced model was then employed as the starting point for the subsequent optimization,
replicating a progressive capacity decline scenario.

2.4.1 Pain avoidance

To simulate pain avoidance, we incorporated a cost function requiring the simulation to minimize the
peak unilateral knee load of the stepping leg to less than 4, 3, and 2 times the body weight (BW).
Individuals with symptomatic unilateral knee OA prefer to step with the affected leg first. Therefore,
we implemented a cost function that becomes zero once the peak knee load on the stepping leg (left
leg in simulations) was below the set threshold.

2.5 Measures

We analysed the trunk flexion, calculated as the angle between the global vertical and the trunk by
adding the pelvis tilt, lumbar extension, and thoracic extension. Additionally, we assessed differences
in timing, muscle activation, joint loading, and kinematics. The outcomes under various conditions
were visualized using the OpenSim software [26].

Results

3.1 Reduced VAS muscle strength

Reducing the maximum isometric strength VAS led to an increased bilateral activation of the Vasti
muscle (Fig. 2). The simulation compensates with increased hip extensor activity (HAM, GMAX)
and increased plantarflexor muscle activity (GAS, SOL) in the stepping leg. This aligns with findings
from experimental studies on sit-to-stand movements in older adults [27].

We found a bilateral rise in hip loading at the point of seat-off when VAS strength was reduced.
Additionally, there was a bilateral increase of ankle joint loading, particularly on the stepping side
which doubled in the -50% condition. Reduced VAS muscle strength thus resulted in elevated hip
and ankle loads. When VAS strength was further reduced to -60%, the model could not find a
solution to rise from the seat unaided.

A minor increase in trunk flexion was observed when VAS strength was reduced (Fig. 3). Trunk
flexion at seat-off reached 42.8 degrees in the -50% condition versus 38.3 degrees in the -20%
condition (40.8 degrees in the neutral condition).

Figure 2 Muscle activation (a) and joint loading (b) for conditions in which VAS strength is reduced

[Here Figure 2]

Figure 3 a) trunk flexion from start to seat-off in sit-to-walk simulations b) conditions shown in color
for seat-off, first toe off, and third toe off. There is a mild increase in trunk flexion when the VAS
strength was reduced. When the knee is unloaded there is a slower movement speed.

[Here Figure 3]

3.2 Unloading of the Knee

In the neutral condition, the peak knee loads (stance leg: 6.8 BW, stepping leg: 5.6 BW) and hip loads (stance leg: 3.6 BW, stepping leg: 3.4 BW) manifest just after seat-off. The peak ankle load in the stepping leg occurs during the double stance (rising) phase (stepping leg: 1.3 BW), and for stance leg, it arises at the end of the single stance phase (stance leg: 2.4 BW), consistent with experimental data [14].

Unloading of the knee is accomplished by reducing ipsilateral activation of VAS and hamstrings (HAM) (Fig. 4). Following Toe-off, there is an increased ipsilateral activation of the tibialis anterior muscle (TA), resulting in increased dorsiflexion, which compensates for reduced foot-ground clearance. Unloading the knee (ranging from -29% to -59%) led to reduced load on the ipsilateral hip (ranging from -23% to -39%) and ankle (ranging from -37% to -40%). However, the ipsilateral ankle did not exhibit proportional unloading, unlike the other joints. Loading on the contralateral knee and hip increased, ranging from +5% to +11% and 0% to +5%, respectively. As for the contralateral ankle, there was a load decrease in the <4BW condition (-4%) and an increase in the other conditions (14% and 29%).

Interestingly, during knee unloading, the simulation yielded a smaller, rather than a greater, trunk angle at seat-off compared to the neutral condition (Fig 4).

Figure 4 *Muscle activation (a) and joint loading (b) for conditions in which the knee joint of the*
stepping leg is unloaded.

167   [Here Figure 4]
168

Discussion

Our simulations reveal increased trunk flexion in response to a decrease in maximum isometric force
in VAS (representing reduced muscular capacity). This result aligns with the notion that an increased
task demand, such as reduced seat height, leads to greater trunk flexion before seat-off [12], [14].
Pain avoidance, simulated with the objective of reducing knee load, resulted in altered muscle
recruitments, characterized by reduced ipsilateral activation of VAS and HAM, and slower
movement speeds. However, it did not yield greater trunk flexion strategies. The predictive
simulations support the idea that increased trunk flexion is more likely a consequence of insufficient
muscular reserve than a pain avoidance mechanism.

The increase in time to perform the sit-to-walk movement is consistent with findings in the sit-to-
stand (STS) task. Previous studies have reported that individuals with knee OA take significantly
more time to complete the STS task and TUG compared to older adults [6], [28], [29] . Various
authors have postulated that this time difference may be attributed to increased forward body bending
[28]or quadriceps muscle weakness [30]. However, the simulations show that reducing movement
velocity alone results in decreased peak knee loading, and might therefore be the result of a pain
reduction strategy.

While the Timed-Up-and-Go test is a commonly employed clinical assessment, only the timing of the
exercise is typically reported in clinical and literature contexts. There is a scarcity of biomechanical
data pertaining to OA patients' performance in this test, particularly regarding which leg they use to
step out first. Our choice to reduce the peak load in the stepping leg was informed by clinical
observations. Given that in the sit-to-walk movement, individuals must unload one leg to initiate gait,
this assumption appears valid from a motor control perspective. We did conduct simulations where
the stance leg was unloaded (available online [31]). Those simulations demonstrate that the energy
objective costs were higher when the stance leg was unloaded compared to the stepping leg,
providing further support for the idea that unloading the stepping leg is a more natural movement.

When examining published experimental data, greater trunk flexion was only observed in individuals
with severe knee OA. In contrast, individuals with moderate knee OA exhibited a similar movement
trajectory to that of healthy individuals but with altered muscle recruitment patterns [12]. This
discrepancy can be attributed to variations in muscle strength. It is plausible that individuals with
severe OA, due to reduced daily activity, possess less muscular reserve compared to individuals with
moderate OA. Consequently, severe OA patients may resort to increased trunk flexion during the
process of standing up as a compensatory mechanism to counteract their strength limitations.

Taking this line of reasoning further, it's conceivable that in individuals with early and moderate
symptomatic knee OA, unloading the knee to alleviate pain could potentially contribute to the
development of muscular weakness in specific muscle groups, such as the vastus lateralis, over time.
Humans stand up over 60 times a day and the simulations indicate that unloading the knee during this
action is accomplished through reduced activation of the VAS. Since it is reasonable to infer that this
compensation mechanism occurs in various daily activities, we can postulate the following
hypothesis (Fig. 5): individuals with early and moderate OA may unload the knee to mitigate pain,
inadvertently leading to the underuse of specific muscle groups, particularly the vasti muscles. Over
time, this underuse could result in a decline in vasti strength, resulting in increased trunk flexion, as
observed in individuals with severe knee OA. This underscores the significance of monitoring and
addressing pain in individuals with early and moderate OA.

Limitations:

•   The simulations are conducted with a planar model. In reality, human movements during sit-
to-walk involve lateral trunk motions and arm support [5]. Individuals with unilateral
symptomatic knee osteoarthritis can use this to redistribute the load [12]. However, this
aspect could not be tested in the current model. Additionally, individuals with knee OA
exhibit increased antagonist muscle activation [8], [29], [32], [33], which is linked to the
medial-lateral and internal-external rotational stability of the knee. These complexities were
not captured by the simplified planar joint in the model.

•   In the simulations we only tested the effect of reduced VAS isometric strength. In reality, OA
patients experience muscle weakness in multiple muscle groups, which could contribute to
greater trunk flexion. Furthermore, regarding the musculoskeletal model, the maximum
isometric forces are based on (23), where VAS muscles are relatively weak for a healthy
adult, as indicated by the maximum activation of this muscle during chair rising [14].

•   The H1120 model represents a male adult with a height of 1.80m and a mass of 75 kg. Body
height and mass influence movement strategies in sit-to-walk. It has been demonstrated that
lower seat heights also lead to increased trunk flexion [14]. To ensure direct comparisons
with experimental datasets, it would be beneficial to diversify the musculoskeletal models and
simulations.

Figure 5 Hypothesized sequence of adaptive strategies in Knee OA patients.

References

[1]   P. M. Dall and A. Kerr, "Frequency of the sit to stand task: an observational study of free-
living adults," *Appl Ergon*, vol. 41, no. 1, pp. 58–61, 2010.

[2]   G. F. Fuller, "Falls in the elderly," *Am Fam Physician*, vol. 61, no. 7, pp. 2159–2168, 2000.

[3]   P. E. D. Eysink, M. Poos, R. Gijsen, G. J. Kommer, and C. H. van Gool, "Epidemiologische
data van Ziekten van het botspierstelsel en bindweefsel: Achtergrondrapport voor Programma
Zinnige Zorg," 2019.

[4]   A. Alghadir, S. Anwer, and J.-M. Brismée, "The reliability and minimal detectable change of
Timed Up and Go test in individuals with grade 1–3 knee osteoarthritis," *BMC Musculoskelet
Disord*, vol. 16, pp. 1–7, 2015.

[5] E. van der Kruk, P. Strutton, L. J. Koizia, M. Fertleman, P. Reilly, and A. M. J. Bull, "Why do
older adults stand-up differently to young adults?: investigation of compensatory movement
strategies in sit-to-walk," *npj Aging*, vol. 8, no. 1, p. 13, 2022.

[6] K. Turcot, S. Armand, D. Fritschy, P. Hoffmeyer, and D. Suvà, "Sit-to-stand alterations in
advanced knee osteoarthritis," *Gait Posture*, vol. 36, no. 1, pp. 68–72, 2012.

[7] J. E. Naili, E. W. Broström, E. M. Gutierrez-Farewik, and M. H. Schwartz, "The centre of
mass trajectory is a sensitive and responsive measure of functional compensations in
individuals with knee osteoarthritis performing the five times sit-to-stand test," *Gait Posture*,
vol. 62, pp. 140–145, 2018.

[8] B. S. Davidson, D. L. Judd, A. C. Thomas, R. L. Mizner, D. G. Eckhoff, and J. E. Stevens-
Lapsley, "Muscle activation and coactivation during five-time-sit-to-stand movement in
patients undergoing total knee arthroplasty," *Journal of Electromyography and Kinesiology*,
vol. 23, no. 6, pp. 1485–1493, 2013.

[9] I. Eitzen, L. Fernandes, L. Nordsletten, L. Snyder-Mackler, and M. A. Risberg, "Weight-
bearing asymmetries during Sit-To-Stand in patients with mild-to-moderate hip osteoarthritis,"
Gait Posture, vol. 39, no. 2, pp. 683–688, 2014.

[10] M. Lamontagne, M. L. Beaulieu, D. Varin, and P. E. Beaulé, "Lower⬜limb joint mechanics
after total hip arthroplasty during sitting and standing tasks," *Journal of Orthopaedic
Research*, vol. 30, no. 10, pp. 1611–1617, 2012.

[11] Y. Sagawa, A. Bonnefoy-Mazure, S. Armand, P. Hoffmeyer, D. Suva, and K. Turcot,
"Individuals with knee osteoarthritis exhibit altered movement patterns during the sit-to-stand
task," *Movement & Sport Sciences-Science & Motricité*, no. 98, pp. 39–49, 2017.

[12] E. van Der Kruk, A. K. Silverman, P. Reilly, and A. M. J. Bull, "Compensation due to age-
related decline in sit-to-stand and sit-to-walk," *J Biomech*, vol. 122, p. 110411, 2021.

[13] E. van der Kruk, A. K. Silverman, L. Koizia, P. Reilly, M. Fertleman, and A. M. J. Bull, "Age-
related compensation: Neuromusculoskeletal capacity, reserve & movement objectives," *J
Biomech*, vol. 122, p. 110385, 2021.

[14] E. van der Kruk and T. Geijtenbeek, "A planar neuromuscular controller to simulate age-
related adaptation strategies in the sit-to-walk movement.," *bioRxiv*, pp. 2011–2023, 2023.

[15] E. van der Kruk and T. Geijtenbeek, "Repository sit-to-walk simulations,"
https://github.com/BODIES-Lab-TU-Delft/Sit-to-walk-predictive-simulations. 2023.

[16] S. L. Delp, J. P. Loan, M. G. Hoy, F. E. Zajac, E. L. Topp, and J. M. Rosen, "An interactive
graphics-based model of the lower extremity to study orthopaedic surgical procedures," *IEEE
Trans Biomed Eng*, vol. 37, no. 8, pp. 757–767, 1990.

[17] H. Geyer and H. Herr, "A muscle-reflex model that encodes principles of legged mechanics
produces human walking dynamics and muscle activities," *IEEE Transactions on neural
systems and rehabilitation engineering*, vol. 18, no. 3, pp. 263–273, 2010.

[18] J. M. Wang, S. R. Hamner, S. L. Delp, and V. Koltun, "Optimizing locomotion controllers using biologically-based actuators and objectives," *ACM Transactions on Graphics (TOG)*, vol. 31, no. 4, pp. 1–11, 2012.

[19] T. Geijtenbeek, "Scone: Open source software for predictive simulation of biological motion," *J Open Source Softw*, vol. 4, no. 38, p. 1421, 2019.

[20] T. Geijtenbeek, "The {Hyfydy} Simulation Software," https://hyfydy.com.

[21] E. M. Arnold, S. R. Ward, R. L. Lieber, and S. L. Delp, "A model of the lower limb for analysis of human movement," *Ann Biomed Eng*, vol. 38, pp. 269–279, 2010.

[22] A. Rajagopal, C. L. Dembia, M. S. DeMers, D. D. Delp, J. L. Hicks, and S. L. Delp, "Full-body musculoskeletal model for muscle-driven simulation of human gait," *IEEE Trans Biomed Eng*, vol. 63, no. 10, pp. 2068–2079, 2016.

[23] C. Slemenda *et al.*, "Quadriceps weakness and osteoarthritis of the knee," *Ann Intern Med*, vol. 127, no. 2, pp. 97–104, 1997.

[24] T. E. McAlindon, C. Cooper, J. R. Kirwan, and P. A. Dieppe, "Determinants of disability in osteoarthritis of the knee.," *Ann Rheum Dis*, vol. 52, no. 4, pp. 258–262, 1993.

[25] M. V Hurley, D. L. Scott, J. Rees, and D. J. Newham, "Sensorimotor changes and functional performance in patients with knee osteoarthritis," *Ann Rheum Dis*, vol. 56, no. 11, pp. 641–648, 1997.

[26] S. L. Delp *et al.*, "OpenSim: open-source software to create and analyze dynamic simulations of movement," *IEEE Trans Biomed Eng*, vol. 54, no. 11, pp. 1940–1950, 2007.

[27] S. H. L. Smith, P. Reilly, and A. M. J. Bull, "A musculoskeletal modelling approach to explain sit-to-stand difficulties in older people due to changes in muscle recruitment and movement strategies," *J Biomech*, vol. 98, p. 109451, 2020.

[28] F. C. Su, K. A. Lai, and W. H. Hong, "Rising from chair after total knee arthroplasty," *Clinical biomechanics*, vol. 13, no. 3, pp. 176–181, 1998.

[29] G. Patsika, E. Kellis, and I. G. Amiridis, "Neuromuscular efficiency during sit to stand movement in women with knee osteoarthritis," *Journal of Electromyography and Kinesiology*, vol. 21, no. 5, pp. 689–694, 2011.

[30] A. Alghadir, S. Anwer, and J.-M. Brismée, "The reliability and minimal detectable change of Timed Up and Go test in individuals with grade 1–3 knee osteoarthritis," *BMC Musculoskelet Disord*, vol. 16, pp. 1–7, 2015.

[31] E. van der Kruk and T. Geijtenbeek, "Repository case study simulations," https://github.com/BODIES-Lab-TU-Delft/STW-case-study-unilateral-pain.git.

[32] G. Bouchouras, G. Patsika, V. Hatzitaki, and E. Kellis, "Kinematics and knee muscle activation during sit-to-stand movement in women with knee osteoarthritis," *Clinical biomechanics*, vol. 30, no. 6, pp. 599–607, 2015.

318 [33] G. Bouchouras, G. Sofianidis, G. Patsika, E. Kellis, and V. Hatzitaki, "Women with
319   knee osteoarthritis increase knee muscle co-contraction to perform stand to sit," *Aging*
320   *Clin Exp Res*, vol. 32, pp. 655–662, 2020.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Author Contributions

Eline van der Kruk – Conceptualization, Data Curation, Methodology, Formal Analysis, Funding, Validation, Acquisition, Supervision, Writing – Original Draft Preparation

Thomas Geijtenbeek - Software, Writing – Review & Editing

Funding

This study was funded by NWO-TTW VENI Grant 18145(2021). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Data Availability Statement

The source code and data for the predictive framework and the results presented in this manuscript are available from Git repositories: [15], [31]

Figure captions

Figure 1 The musculoskeletal model (H1120) has 11 degrees of freedom and is actuated using 20 Hill-type muscle-tendon units. The H1120 model is available on [15].

Figure 2 Muscle activation (A) and joint loading (B) for conditions in which VAS strength is reduced

Figure 3 A) trunk flexion from start to seat-off in sit-to-walk simulations B) conditions shown in color for seat-off, first toe off, and third toe off. There is a mild increase in trunk flexion when the VAS strength was reduced. When the knee is unloaded there is a slower movement speed.

Figure 4 Muscle activation (A) and joint loading (B) for conditions in which the knee joint of the stepping leg is unloaded.

Figure 5 Hypothesized sequence of adaptive strategies in Knee OA patients