

1 **Dengue virus transmission in Italy: surveillance and epidemiological trends** 2 **up to 2023**

3
4 Francesco Branda¹, Taishi Nakase², Antonello Maruotti³, Fabio Scarpa⁴, Alessandra Ciccozzi¹, Chiara
5 Romano¹, Simone Peletto⁵, Ana Maria Bispo de Filippis⁶, Luiz Carlos Junior Alcantara^{7,8}, Alessandro
6 Marcello⁹, Massimo Ciccozzi^{1^}, José Lourenço^{10,11^}, Marta Giovanetti^{7,8,12^}

- 7
- 8 1. Unit of Medical Statistics and Molecular Epidemiology, Università Campus Bio-Medico di Roma, Rome,
- 9 Italy;
- 10 2. Department of Epidemiology & Population Health, Stanford University School of Medicine, Stanford, CA,
- 11 USA
- 12 3. Department GEPLI, Libera Università Ss Maria Assunta, 00193 Rome, Italy;
- 13 4. Department of Biomedical Sciences, University of Sassari, 07100 Sassari, Italy;
- 14 5. Istituto Zooprofilattico Sperimentale del Piemonte, Liguria e Valle d'Aosta, Turin, Italy
- 15 6. Laboratório de Flavivírus, Instituto Oswaldo Cruz, Fundação Oswaldo Cruz, Rio de Janeiro, Brazil;
- 16 7. Instituto René Rachou, Fundação Oswaldo Cruz, Minas Gerais, Brazil;
- 17 8. Climate Amplified Diseases And Epidemics (CLIMADE), Brazil, Americas;
- 18 9. The Laboratory of Molecular Virology, International Centre for Genetic Engineering and Biotechnology
- 19 (ICGEB), 34149, Trieste, Italy;
- 20 10. Universidade Católica Portuguesa, Biomedical Research Center, Lisboa, Portugal;
- 21 11. Climate Amplified Diseases And Epidemics (CLIMADE), Portugal, Europe;
- 22 12. Sciences and Technologies for Sustainable Development and One Health, Università Campus Bio-Medico
- 23 di Roma, Rome, Italy.

24 **Correspondence:** jolourenco@ucp.pt

25 ^ Co-senior authors

26
27
28
29
30
31
32
33
34
35

36 **Abstract**

37 Dengue virus circulation is on the rise globally with increased epidemic activity in previously
38 unaffected countries, including within the European region. In 2023, global dengue activity
39 peaked, and Italy reported the highest number of dengue cases and local chains of transmission
40 to date. By curating several sources of information, we introduce a novel data repository focused
41 on dengue reporting in Italy. We integrate geographic, epidemiological, genomic and climatic
42 spatio-temporal data to present an overview of transmission patterns of the past eight years
43 related to circulating viral lineages, geographic distribution, hotspots of reporting, and the
44 theoretical contribution of local climate. This study contributes to a better understanding of the
45 evolving scenario in Italy, with the potential to inform reassessment and planning of adequate
46 national and European public health strategies to manage the emergence of dengue.

47

48

49 **KEYWORDS:** Dengue, public health, virus surveillance, epidemiology, climate change.

50

51

52

53

54 Introduction

55 Dengue fever is a mosquito-borne viral infection caused by the Dengue virus (DENV), which
56 belongs to the *Flaviviridae* family. Common symptoms include high fever, intense headaches,
57 and significant muscle pain [1]. Transmission primarily occurs through the bites of *Aedes spp.*
58 infected mosquitoes [1]. In recent years, there has been a significant global increase in incidence,
59 leading to impacts on and concerns for public health systems [2]. One of the primary drivers of
60 this surge is climate change [3,4]. Ongoing climatic changes, including temperature fluctuations
61 and altered rainfall patterns, have not only contributed to the increased incidence but also
62 facilitated geographic spread, notably towards higher altitudes [3]. Historically, Europe has
63 experienced mostly imported cases of dengue, with sporadic autochthonous transmission [2].
64 However, autochthonous cases have been surging in the past five years, with significant reports
65 from Spain [5], Croazia [6], France [7] and Italy [8]. In 2023, Italy reported its highest ever
66 number of locally transmitted dengue [9].

67
68 In the present study, we integrate several sources of geographic, genetic, epidemiological and
69 climate data, presenting a comprehensive overview of the spatiotemporal patterns and possible
70 drivers of dengue activity in Italy.

71 Methods

72 To assess the trends of DENV in Italy, we collected epidemiological data from the weekly
73 reported cases published by the Istituto Superiore di Sanità (ISS) on the EpiCentro website,
74 which can be accessed at [9]. The process started with downloading and pre-processing all
75 bulletins from the EpiCentro website into CSV files. These are now stored in a publicly
76 accessible GitHub repository, which is regularly updated with new Italian data as they become
77 available. The repository can be found at <https://github.com/fbranda/arbovirus>, and a detailed
78 synopsis of the file structure is provided in **Table S1**.

79

Files and format	Description
Bulletins PDF	These files provide updates on the Dengue situation throughout Italy
Dengue-ita-yyyy*.csv Comma-separated values file (.csv, UTF-8 encoded)	Number of Reported Dengue Cases Nationwide, Categorized by Sex and Type of Infection (Autochthonous and Imported)
Dengue-ita-age-yyyy*.csv Comma-separated values file (.csv, UTF-8 encoded)	Number of Dengue Cases Reported by Age and Sex
Dengue-ita-location-exposure-yyyy*.csv Comma-separated values file (.csv, UTF-8 encoded)	Percentage of Dengue Cases Relative to Total Cases by Location
Dengue-ita-regions-yyyy*.csv Comma-separated values file (.csv, UTF-8 encoded)	Reported Dengue Cases by Region

Dengue-ita-summary-cases.csv Comma-separated values file (.csv, UTF-8 encoded)	Time Series of Nationwide Reported Dengue Cases by Year, Month, and Type of Infection (Autochthonous and Imported)
Dengue-ita-summary-cases-regions.csv Comma-separated values file (.csv, UTF-8 encoded)	Time Series of Regionally Reported Dengue Cases Categorized by Year, Month, and Type of Infection (Autochthonous and Imported)

80 **Table S1.** Overview of File Structure for the Italian Dengue virus Data Repository
81 *yyyy is the year of the monitoring
82

83 **Reporting risk model**

84 We modeled reported counts per area i as Y_i , ($i = 1$ to n) using a Poisson distribution with mean
85 $E_i \times \theta_i$, where E_i is the expected counts and θ_i is the relative risk in area i . Then, the log risks are
86 modeled with a sum of an intercept to model the overall disease risk level and random effects
87 that account for extra-Poisson variability in the observed data [10]. Areas with relative risks $\theta_i >$
88 1 and $\theta_i < 1$ are areas with high and low risks, respectively. Areas with $\theta_i = 1$ has the same risk
89 as expected from the standard population. In this work, the model for disease mapping is
90 expressed as:

$$Y_i \sim Po(E_i \times \theta_i), i = 1, \dots, n,$$
$$\log \log(\theta_i) = \alpha + u_i + v_i,$$

91
92 where α denotes the overall risk level, u_i is a spatial structured random effect that models the
93 spatial dependence between the relative risks, and v_i is an unstructured exchangeable random
94 effect that models uncorrelated noise.
95

96 **Detection of clusters**

97 To identify spatio-temporal clusters (i.e., geographic areas in which the number of observed
98 incidence events is significantly different from that predicted by a random distribution) we used
99 Kulldorff's scan statistics [11]. This method uses a moving window, which can take various
100 shapes, such as circles or ellipses, and different sizes, running systematically over the entire
101 study area. For each position and size of the window, the scan statistic calculates a probability
102 index based on the ratio of the number of events observed within the window to the number of
103 events expected under a null hypothesis of random distribution (i.e. the events are randomly
104 distributed across the entire study area). The Kulldorff's scan statistics then compares the actual
105 data collected with this random distribution for each window, and calculates the likelihood ratio
106 (LR) to determine whether the observed cluster is statistically significant or could be due to
107 chance:

$$LR = \frac{L(\text{Observed}, \theta)}{L(\text{Expected}, \theta)}$$

108 where $L(Observerd, \theta)$ represents the likelihood of the observed data within the window,
109 considering the parameters θ of the alternative hypothesis, and $L(Expected, \theta)$ represents the
110 likelihood of the expected data within the window, also considering the parameters θ of the
111 alternative hypothesis. An LR greater than 1 indicates that there are more observed cases within
112 the window than expected.

113
114 To determine the statistical significance of a cluster, LR is compared to a threshold value. This
115 threshold is often represented as a $-\log(p\text{-value})$. The p-value quantifies the probability of
116 observing a cluster as extreme as, or more extreme than, the one identified under the null
117 hypothesis that cases are randomly distributed. A lower p-value indicates a more significant
118 cluster. More specifically, the $-\log(p\text{-value})$ is compared to the critical values from a chi-squared
119 distribution (χ^2). If the $\log(p\text{-value})$ is greater than or equal to the critical value from the chi-
120 squared distribution at a specified significance level (often set at 0.05 or 0.01), the cluster is
121 deemed statistically significant, and the null hypothesis of random distribution within the spatial
122 window is rejected. The outcome of this significance calculation guides the interpretation of the
123 spatial window or cluster. A statistically significant cluster suggests that the observed pattern of
124 events is unlikely to be the result of random chance and may indicate the presence of a true
125 cluster. In contrast, a non-significant cluster implies that the observed pattern can be reasonably
126 explained by random variation. All analyses were performed using R (v4.3.1) [4] and we used
127 SaTScan software (v10.1.2) (<https://www.satscan.org/>) [5] to perform geographic disease
128 surveillance for early detection of disease outbreaks and evaluate the statistical significance of
129 disease cluster alarms.

130 **Dated phylogenetics**

131 We analyzed all available genome sequences from GISAID (DENV 1 n=7, DENV 3 n=3) along
132 with reference strains (n=95 for DENV1 and n=140 for DENV3 respectively). These sequences
133 were aligned using MAFFT [12] and manually refined in Aliview [13] to eliminate anomalies.
134 The GTR model, deemed the best fit by ModelFinder in IQ-TREE2 [14], was employed for the
135 initial estimation of maximum likelihood phylogenetic trees, with robustness validated through
136 1,000 bootstrap replicates. TempEst [15] aided in detecting temporal signals, while BEAST
137 facilitated the inference of time-scaled phylogenetic trees. A comprehensive model selection
138 process, incorporating path-sampling and steppingstone methods, identified the uncorrelated
139 relaxed molecular clock model, utilizing the SRD06 model and Bayesian Skygrid coalescent
140 model, as optimal for Bayesian analysis [15,16]. We conducted phylogeographic analysis to map
141 the virus' spatial diffusion, using discrete sampling locations (countries) and an asymmetric
142 model of location transition with BSSVS [17]. To ensure thoroughness, MCMC runs were
143 performed in duplicate over 100 million iterations, achieving an effective sample size (ESS) of
144 over 200. The final maximum clade trees, post 10% burn-in exclusion, were compiled using
145 TreeAnnotator and displayed in FigTree v1.4.4.

146 **Analysis of long-term temperature trends**

147 Long-term trends in local temperature were estimated using a linear regression model on each
148 geographical pixel of monthly satellite temperature time series from 1979 (January) to 2023
149 (September) [18]. In the main text we report the estimated slopes (all had p -value < 0.0001).

150 **Climate-based transmission suitability**


151 We estimated theoretical, dengue virus climate-based transmission suitability, as per Nakase et al
152 [3] for each geographical pixel available in satellite climate data from Copernicus.eu [18]. This
153 suitability measure (index P) measures the reproductive, transmission potential of a single adult
154 female mosquito during its lifetime in a fully susceptible host population [19]. Since satellite
155 data was only available up to September 2023, all outputs in the main text related to transmission
156 suitability were summarized per year using climate data between January and September of each
157 year.

158 **Results**

159 **Spatio-temporal Dynamics of Dengue in Italy**

160
161 The spatio-temporal dynamics of reported dengue in Italy are summarized in **Figure 1**. Between
162 2015 and 20 November 2023, there have been a total of 1028 reported dengue cases nationally,
163 including local and imported cases. Of these 253 were reported between 2015 and 2017, for
164 which there was no available geolocation (**Figure 1A**). Between 2018 and 2023, a total of 775
165 cases were reported, with the Northeast, Northwest and Center macroregions each experiencing
166 about 33% of cases (N=269, 256, 225, respectively), and the South/Insular experiencing only
167 ~3% of cases (N=25).

168
169 During 2023, the year with highest ever reported cases (N=327, ~31%), there were 75 cases in
170 the Northeast, 134 in the Northwest, 111 in the Center and only 7 in the South/Insular
171 macroregions (**Figure 1B**). Among these, 82 were confirmed as autochthonous and attributable
172 to four distinct chains of transmission within the regions of Lodi in the Northwest, and Latina,
173 Rome (including metropolitan areas), Anzio, all in the Center macroregion.




174
175
176
177
178
179

Figure 1. Spatiotemporal Dynamics of Dengue in Italy. A) Map of Italy with regions (black boundaries) and macroregions (color legend). B) Time series of reported dengue cases. Reports between 2015-2017 had no geolocation, while those after 2018 are aggregated by macroregion. C) Yearly count of available DENV genome sequences by serotype in the GISAID repository (January 2015 to November 20, 2023). Missing years had no genomes.

180 Genomic data, derived from the complete genome sequences available in the public repository
181 (GISAID), revealed that three dengue serotypes have circulated in Italy in the last eight years
182 (Figure 1C). In 2015, DENV1, DENV2, and DENV3 were reported as circulating, with DENV1
183 being the predominant serotype. Between 2018 and 2020, DENV1 was the only serotype present
184 in the database, while in 2023 both DENV1 and DENV3 were detected. This recent trend in Italy
185 aligns with the global trends observed since early 2023, where there has been an increase in the
186 circulation of DENV1 and DENV3, particularly across Latin American countries [20, 21].

187 The 2023 Epidemiological Dynamics of Dengue in Italy

188
189 By November 20, 2023, active dengue virus transmission was reported in 15 of Italy's 21
190 regions. This wide geographical range has prompted a nationwide movement, aimed at
191 comprehensively assessing and identifying gaps in the public health response and preparedness.
192 Figure 2A delineates regions with high Dengue incidence in 2023, categorized based on weekly
193 case reports: Zero/Low includes regions like Abruzzo and Aosta Valley with up to five cases;
194 Medium covers regions from Campania to Veneto with six to fifteen cases; and High represents
195 Emilia Romagna, Lazio, Lombardy, and Piedmont, reporting over fifteen cases. Notably,
196 Lombardy and Lazio represent 33% and 25% of the total cases, respectively, highlighting
197 significant reporting concentrations.



198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221


222

223

Figure 2. The 2023 Epidemiological Dynamics of Dengue in Italy. **A)** Identification of reporting hotspots across Italian regions in 2023, including a summary of key characteristics within the identified clusters. **B1)** Breakdown of reported cases by age group. In red the proportion of cases per age group, and in grey the proportion of the national population per age group. **B2)** Breakdown of reported cases by gender. In blue the proportion of cases reported as female per age group, and in yellow the proportion of cases reported as male per age group. For the period 2015-2022: grey dashed line is the monthly reporting mean; grey area marks the limits of the minimum and maximum reported cases over the months; pink area marks the typical time window of peak reporting. For 2023, the full purple line is the total cases per month. The y-axis is square-root transformed for visualization.

Two major clusters were identified: the first extended across a diagonal axis within the Northwest and Northeast macroregions, from Piedmont to Emilia Romagna, covering over 20 million people; the second was located in Lazio, with a population of 5 million. The relative risk (RR) of dengue infection in these clusters was 2.48 in the first and 3.05 in the second. The median age of individuals was 37 years, with a slight male predominance of 51.68% (**Figure 2B**). In general, cases were more frequently reported among the 20-29 and 30-39 age groups, contrasting with the national age profile of the population (**Figure 2B1**). Between genders, cases were also concentrated among the 20-29 and 30-39 age groups for females, but were more uniformly distributed for males older than 20 years (**Figure 2B2**). Compared to the monthly average trends in reported cases with the period 2015-2022, the year 2023 was different in both seasonal dynamics and size (**Figure 2C**). Specifically, reports for 2023 were consistently higher than the historical average across the months (with the exception of March). Peak reporting in 2023 was in August (N=116) well within the historical time window of peak reporting, but peaking much higher (ratio = 9.6) and only approaching average historical levels late in November.

224



225
226
227
228
229
230
231
232

Figure 3. Phylogenetic analysis of DENV1-V and DENV3-III in Italy. A) Time-scaled phylogeographic tree of DENV1-V (including all Italian strains currently available $n=7$ plus $n=95$ GenBank sequences). Colors represent different sampling locations. B) Time-scaled phylogeographic tree of DENV3-III (including all Italian strains currently available $n=3$ plus $n=140$ GenBank sequences). Colors represent different sampling locations; C) Map displaying the countries involved, with virus transmission paths marked by counter-clockwise lines colored according to the viral source. DENV1-V is shown with a black dashed line, and DENV3-III is represented by a light green dashed line.


233 **Phylogenetic Analysis of co-circulating DENV1 and DENV3 in 2023**

234 We combined all available partial and complete DENV genome sequences from Italy, primarily
235 generated during the 2023 epidemic, with other DENV1 and DENV3 genomes from public
236 repositories. Our phylogenetic analysis revealed that the novel 2023 isolates belonged to DENV1
237 genotype V (DENV1-V) and DENV3 genotype III (DENV3-III). These are currently the
238 predominant genotypes circulating in Latin American countries [20, 21].

239

240 Further analysis of DENV1-V indicated two major introductions into Italy, estimated to have
241 occurred in early and late December 2022 (High Posterior Density interval ranging from July 24
242 2022, to June 10 2023) (**Figure 3A**). Discrete phylogeographic reconstruction also suggested that
243 South America, primarily represented by Brazilian sequences, was a central hub for multiple
244 viral introductions into Italy (**Figure 3B**).

245 We further traced the introduction of DENV3-III to late November 2022 (High Posterior Density
246 interval ranging from August 22 2022, to June 28 2023) (**Figure 3C**). Our findings suggested
247 that the Caribbean (including Cuba), which saw a rise in DENV3-III cases since 2022 [20],
248 might have been the source of this introduction (**Figure 3D**).
249
250



251
252 **Figure 4. Climate-based Suitability of Dengue in Italy.** **A1)** Yearly mean climate-based transmission suitability (index P) per
253 geographical pixel of the climate satellite data; presented are a subset of years. **A2)** Distributions of suitability values per year.
254 The light red horizontal line marks the historical mean (2015-2023). **B1)** Yearly number of months in which climate-based
255 transmission suitability (index P) per geographical pixel of the climate satellite data was above 1; presented are a subset of years.
256 **B2)** Proportion of land (of geographical pixels available in climate data) that present suitability above 1 for at least 1 month of the
257 year. Presented are the historical period (2015-2023) and 2023 only. North/Center is the aggregation of the Northwest, Northeast
258 and Center macroregions. **C)** Map of Italy with estimates of rate of increase (linear slope) in mean yearly temperature (color
259 legend). Black boundaries are macroregions. **D)** Time series of mean monthly temperature (1979-2022) for two geolocations as
260 marked on panel D. Line coloring follows the corresponding macroregion. **E)** Distributions of rates of increase (linear slopes) for
261 each of the macroregions (as presented in panel D).

262 Climate-based Transmission Suitability of Dengue in Italy

263 We next explored whether climate, an established driver of arboviral transmission potential [3,
264 4], could shed light on patterns of Dengue virus reporting across Italy. Using climate satellite
265 data, we estimated the theoretical transmission suitability of dengue between 2015 and 2023.
266 Suitability was found to vary across space (**Figure 4A1**) and time (**Figure 4A2**). The far north

267 and the latitude-longitude diagonal across the country, rich in high altitude land, consistently
268 presented the lowest suitability. In contrast, the island of Sicily, the continental coasts and some
269 areas of the northern macroregions consistently presented intermediate-to-high suitability.
270 Notably, these areas also showed high suitability (suitability > 1) for a few consecutive months
271 of the year (**Figure 4B1**). A suitability of 1 is theoretically relevant because it marks the
272 threshold above which a single infected female mosquito could transmit the virus to more than
273 one host during their lifetime.

274
275 The year 2023, in which an exceptionally high number of autochthonous cases were reported,
276 presented transmission suitability only slightly above the historical average (**Figure 4B1**).
277 However, 19% of total land in Italy showed suitability above 1 for at least one month in 2023. -
278 In contrast to historical trends but following the geographical distribution of reported
279 autochthonous cases, suitability above 1 occurred mostly in the Northwest, Northeast and Center
280 macroregions. (**Figure 4B2**).

281
282 Finally, we quantified historical trends in local climate by estimating rates of change in
283 temperature. Our estimates showed a universal, yearly increase in mean temperature across Italy
284 (**Figure 4C**). The highest rates of increase were found in the Northeast macroregion (**Figure**
285 **4D**), followed by the Northwest, Center and South/Insular macroregions (**Figure 4DE**).

286 **Discussion**

287 This study summarizes reported dengue infections in Italy of the last eight years, introducing a
288 new data repository of incidence reports into the future. The repository is designed to improve
289 accessibility to information by the research and education communities, thereby supporting the
290 generation of knowledge and awareness related to the emergence of dengue in the country.

291
292 We explored the potential contribution of local climate to the transmission potential of dengue
293 across Italy by estimating a commonly used climate-based suitability measure. We found that
294 suitability is spatially heterogeneous and generally much lower than that estimated for endemic
295 countries such as Brazil or the Dominican Republic (see e.g. [3,22]). Although not directly
296 quantified, suitability was generally lower at higher altitudes. Reasonable suitability was found
297 along continental coastal areas, the island of Sicily and interior regions of the Northern
298 macroregions. Coincidentally, with the exception of Sicily, these areas were involved in the
299 exceptional emergence of chains of transmission during 2023. However, our analyses also reveal
300 that the majority of land area rarely shows high suitability, with less than 20% with high
301 suitability for at least one month of the year. This suggests weak potential for long-term
302 transmission and sheds light the potential current role of local climate in limiting persistence.
303 Nonetheless, we also uncover that mean yearly temperature has been increasing rapidly,
304 especially in the Northern macroregions, conspicuously matching the areas where West Nile

305 virus and Usutu virus have emerged as public health threats in recent years [23, 24]. While
306 transmission suitability for dengue does not seem to have increased at a similar pace, the effect
307 of rising temperatures is likely to have positive effects on the life cycle and reproduction of
308 mosquito-species generally involved in arboviral transmission. Together, these results underline
309 the need for adaptable mosquito control policies due to environmental changes.

310
311 We note historical asymmetries in dengue reporting across Italy. Over the years, a highly
312 variable number of dengue cases have been reported countrywise, with specific relevance
313 regarding higher latitudes (including the official Northwest, Northeast and Center macroregions
314 of Italy). The pandemic initial years of 2020 and 2021 presented the lowest reporting. While
315 disruption to health services and awareness of infections beyond COVID-19 may have faltered
316 during this period, we also uncovered that such years presented some of the lowest climate-based
317 transmission suitability. Higher reporting was observed for 2019 and 2023, which did present
318 higher than normal suitability, but that more likely reflected the success of dengue epidemic
319 activity elsewhere. For example, 2019 was a year of resurgence in South America and 2023 is
320 now a record year for incidence across the globe. Over the years, three of the four dengue
321 serotypes have been reported to circulate in Italy. Nonetheless, their success in being introduced
322 into the country should also reflect epidemic expansion elsewhere, specifically in light of the
323 absence of herd-immunity in Italy.

324
325 The year of 2023 in Italy was exceptional, not just due to the total number of reported cases, but
326 also for the uncommon co-occurrence of a few autochthonous transmission chains of more than
327 one serotype. We summarize that such chains occurred mostly within two spatial clusters, one in
328 the Northern and another in the Center macroregions. Age profiles for 2023 were available and
329 revealed a higher incidence in the 20-39 age group. The latter contrasted with the national age
330 pyramid of the country, suggesting that active age individuals were more exposed to infection.
331 This effect was particularly strong for females, while males presented a more uniform incidence
332 above 20 years of age. The reasons for this asymmetric exposure are unknown and should be the
333 focus of future epidemiological studies.

334
335 Despite the limited number of available dengue genomes related to infections reported in Italy,
336 we were able to recover a few observations of interest. We noted, for example, that viral lineages
337 of the 2023 transmission clusters belonged to DENV1 genotype V and DENV3 genotype III.
338 Notably, circulation of DENV1 resulted from two independent introductions into the country,
339 while DENV3 resulted from a single introduction. The origin of DENV1 was suggested to be
340 from South America, while the origin of DENV3 is likely to have been the Caribbean. Once
341 again, we noted that the co-circulation of DENV1 and DENV3 in Italy in 2023 conspicuously
342 occurred while the specific genotypes experienced large expansion in Latin American countries
343 [20, 21]. It should be noted however, that the number of genomes available limit definite

344 conclusions, and as such the results presented in this study are amenable to change upon
345 inclusion of further genomic data.

346
347 The information summarized in this study highlights the necessity of a strong genomic
348 surveillance network in Italy for tracking and identifying introductions and local chains of
349 transmission, primarily facilitated by the movement of viremic travelers from elsewhere. A shift
350 from passive to active surveillance would improve responsiveness to short-term transmission
351 chains similar to those experienced during 2023. In parallel, active surveillance strategies should
352 take into account factors such as the spatio-temporal variation of climate suitable for local
353 transmission and human mobility from endemic countries. Together, this could help ensure a
354 comprehensive approach to monitoring and containing dengue introductions and arboviruses
355 more generally.

356 **Conflict of Interest**

357 The authors declare that there are no conflicts of interest.
358

359 **Author Contribution**

360 **Conception and design:** F.B., M.C., J.L., M.G.; **Investigations:** F.B., T.N., M.G., J.L.; **Data**
361 **Analysis:** F.B., T.N., J.L., M.G.; **Visualization:** F.B., J.L., M.G.; **Writing – Original:** F.B., J.L.,
362 M.G.; **Writing – Revision:** F.B., A.M., T.N., F.S., A.C., L.C.J.A., A.M.B.d.F., A.M., J.L., M.C.,
363 M.G.

364 **Acknowledgments**

365 M. Giovanetti's funding is provided by PON "Ricerca e Innovazione " 2014-2020 and by the
366 CRP-ICGEB RESEARCH GRANT 2020 Project CRP/BRA20-03, Contract CRP/20/03.

367 **Data Availability**

368 The data that support the findings of this study are openly available at the following link:
369 <https://github.com/fbranda/arbovirus>.

370 **References**

- 371 1. Adelino TÉR, Giovanetti M, Fonseca V, Xavier J, de Abreu ÁS, do Nascimento VA, et al. Field and
372 classroom initiatives for portable sequence-based monitoring of dengue virus in Brazil. *Nat*
373 *Commun.* 2021;12: 1–12.
- 374 2. Dengue worldwide overview. In: European Centre for Disease Prevention and Control [Internet]. 15
375 Nov 2023 [cited 7 Dec 2023]. Available: <https://www.ecdc.europa.eu/en/dengue-monthly>
- 376 3. Nakase T, Giovanetti M, Obolski U, Lourenço J. Global transmission suitability maps for dengue
377 virus transmitted by *Aedes aegypti* from 1981 to 2019. *Scientific Data.* 2023;10: 1–11.
- 378 4. Nakase T, Giovanetti M, Obolski U, Lourenço J. A retrospective analysis of climate-based dengue
379 virus transmission suitability and demographic changes over the last four decades. *medRxiv.* 2023.
380 p. 2023.11.21.23298791. doi:10.1101/2023.11.21.23298791
- 381 5. Herrero-Martínez J-M, Sánchez-Ledesma M, Ramos-Rincón J-M. Imported and autochthonous
382 dengue in Spain. *Rev Clin Esp.* 2023;223: 510–519.
- 383 6. Pem-Novosel I, Vilibic-Cavlek T, Gjenero-Margan I, Kaic B, Babic-Erceg A, Merdic E, Medic A,
384 Ljubic M, Pahor D, Erceg M. Dengue virus infection in Croatia: seroprevalence and entomological
385 study. *New Microbiol.* 2015 Jan;38(1):97-100. Epub 2015 Jan 1. PMID: 25821943.
- 386 7. Cochet A, Calba C, Jourdain F, Grard G, Durand GA, Guinard A, et al. Autochthonous dengue in
387 mainland France, 2022: geographical extension and incidence increase. *Euro Surveill.* 2022;27.
388 doi:10.2807/1560-7917.ES.2022.27.44.2200818
- 389 8. Cassaniti I, Ferrari G, Senatore S, Rossetti E, Defilippo F, Maffeo M, et al. Preliminary results on an
390 autochthonous dengue outbreak in Lombardy Region, Italy, August 2023. *Euro Surveill.* 2023;28.
391 doi:10.2807/1560-7917.ES.2023.28.37.2300471
- 392 9. EpiCentro. Febbre dengue. [cited 7 Dec 2023]. Available: [https://www.epicentro.iss.it/febbre-](https://www.epicentro.iss.it/febbre-dengue/aggiornamenti)
393 [dengue/aggiornamenti](https://www.epicentro.iss.it/febbre-dengue/aggiornamenti)
- 394 10. Lawson AB. *Bayesian Disease Mapping.* 3rd Edition. Chapman and Hall/CRC; 2018.
- 395 11. Kulldorff M. A spatial scan statistic. *Communications in Statistics - Theory and Methods.* 1997
396 [cited 7 Dec 2023]. doi:10.1080/03610929708831995
- 397 12. Katoh K, Rozewicki J, Yamada KD. MAFFT online service: multiple sequence alignment,
398 interactive sequence choice and visualization. *Brief Bioinform.* 2019;20: 1160–1166.
- 399 13. Larsson A. AliView: a fast and lightweight alignment viewer and editor for large datasets.
400 *Bioinformatics.* 2014;30: 3276–3278.
- 401 14. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective stochastic
402 algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol.* 2015;32: 268–274.
- 403 15. Rambaut A, Lam TT, Max Carvalho L, Pybus OG. Exploring the temporal structure of
404 heterochronous sequences using TempEst (formerly Path-O-Gen). *Virus Evol.* 2016;2: vew007.
- 405 16. Baele G, Li WLS, Drummond AJ, Suchard MA, Lemey P. Accurate model selection of relaxed

- 406 molecular clocks in bayesian phylogenetics. *Mol Biol Evol.* 2013;30: 239–243.
- 407 17. Lemey P, Rambaut A, Welch JJ, Suchard MA. Phylogeography takes a relaxed random walk in
408 continuous space and time. *Mol Biol Evol.* 2010;27: 1877–1885.
- 409 18. Copernicus Climate Data Store. [cited 7 Dec 2023]. Available:
410 <https://cds.climate.copernicus.eu/cdsapp#!/dataset/ecv-for-climate-change?tab=overview>
- 411 19. Obolski U, Perez PN, Villabona-Arenas CJ, Thézé J, Faria NR, Lourenço J. MVSE: An R-package
412 that estimates a climate-driven mosquito-borne viral suitability index. *Methods Ecol Evol.* 2019;10:
413 1357–1370.
- 414 20. Adelino, T.É.R., Giovanetti, M., Fonseca, V. et al. Field and classroom initiatives for portable
415 sequence-based monitoring of dengue virus in Brazil. *Nat Commun* 12, 2296 (2021).
416 <https://doi.org/10.1038/s41467-021-22607-0>
- 417 21. Taylor-Salmon E, Hill V, Paul LM, Koch RT, Breban MI, Chaguza C, et al. Travel surveillance
418 uncovers dengue virus dynamics and introductions in the Caribbean. *medRxiv.* 2023. p.
419 2023.11.11.23298412. doi:10.1101/2023.11.11.23298412
- 420 22. Petrone ME, Earnest R, Lourenço J, Kraemer MUG, Paulino-Ramirez R, Grubaugh ND, et al.
421 Asynchronicity of endemic and emerging mosquito-borne disease outbreaks in the Dominican
422 Republic. *Nat Commun.* 2021;12: 1–12.
- 423 23. Riccardo F, Bella A, Monaco F, Ferraro F, Petrone D, Mateo-Urdiales A, et al. Rapid increase in
424 neuroinvasive West Nile virus infections in humans, Italy, July 2022. *Eurosurveillance.* 2022;27:
425 2200653.
- 426 24. Caracciolo I, Mora-Cardenas E, Aloise C, Carletti T, Segat L, Burali MS, Chiarvesio A, Totis V,
427 Avšič-Županc T, Mastrangelo E, Manfroni G, D'Agaro P, Marcello A. Comprehensive response to
428 Usutu virus following first isolation in blood donors in the Friuli Venezia Giulia region of Italy:
429 Development of recombinant NS1-based serology and sensitivity to antiviral drugs. *PLoS Negl Trop*
430 *Dis.* 2020 Mar 30;14(3):003.