Druggable causal genes of bipolar disorder identified through Mendelian Randomization analysis offer a route to intervention in integrated stress response

Zhai Ting[1,2*]

[1] School of Humanities, Southeast University, Nanjing, 211189, China

[2]Key Laboratory of Child Development and Learning Science (Southeast University), Ministry of
Education, and Institute of Child Development and Education, Southeast University, Nanjing
210096, China

* Correspondence:
Zhai Ting
matthewzhai@foxmail.com

Keywords: bipolar disorder, integrated stress response, summary data-based Mendelian
randomization, genome-wide association study (GWAS), colocalization analysis.

Abstract

Protein misfolding and aggregation, mitochondrial dysfunction, and oxidative stress are all common
features of age-associated proteinopathies. As a consequence, protein synthesis, folding, and
degradation are altered in these disorders. The ISR acts as a central regulator of protein homeostasis.
Genome wide association studies (GWASes) on bipolar disorder (BD) have been conducted;
however, as a causative mechanism underlying the cognitive deficits and neurodegeneration in a
wide broad range of brain disorders, the impact of ISR on BD is understudied, particularly for current
GWASes of molecular phenotypes have limited sample sizes and limited protein coverages. And
many genes and gene products are not easily druggable, so some potentially important causal genes
may not offer an obvious route to intervention. We aimed to prioritize BD associated genes for
functional follow-up studies using the summary data-based Mendelian Randomization (SMR) and
Bayesian colocalization (COLOC) methods to integrate the summary-level data of the GWAS on BD
and the expression quantitative trait locus (eQTL) study in blood. We utilized the GWAS data
including 41917 BD cases and 371549 controls from the Psychiatric Genomics Consortium (PGC)
and the eQTL data from 31,684 participants of predominantly European ancestry from the eQTLGen
consortium. Plus, we use Graph DB to alt drug target. The SMR analysis identified EIF2B5 gene that
were associated with DB due to not linkage but pleiotropy or causality. The COLOC analysis
strongly suggested that EIF2B5 and the trait of BD were affected by shared causal variants, and thus
were colocalized. By means of data in EpiGraphDB we find other putative causal BD genes
(EIF2AK4 and GSK3B) to prioritise potential alternative drug targets.

1   Introduction

Bipolar disorders(In this article, the abbreviation BD will be used to refer to Bipolar disorders) are
described by the *Diagnostic and statistical manual of mental disorders: DSM-5™, 5th ed* (2013) as a
group of brain disorders that cause extreme fluctuation in a person's mood, energy, and ability to
function. People who live with bipolar disorder experience periods of great excitement, overactivity,
delusions, and euphoria (known as mania) and other periods of feeling sad and hopeless (known as

depression). The prevalence of bipolar disorders was consistent across diverse cultures and ethnic
groups, with an aggregate lifetime prevalence of 0.6% for bipolar I disorder, 0.4% for bipolar II
disorder, 1.4% for subthreshold bipolar disorder, and 2.4% for the bipolar disorder spectrum(Grande
et al., 2016). According to statistics, the total cost of BD in the United States is estimated to be
$81,559 per patient per year (Cloutier et al., 2018). Bipolar disorder (BD) has a higher lifetime rate of
suicide attempts (SA) than other psychiatric disorders(Brown et al., 2000; Sani et al., 2011; Takizawa
et al., 2008). Previous studies have reported that individuals with BD experience more negative and
stressful life events than the general population(Brown et al., 2000; Jacobs & Brewer, 2004; Sani et
al., 2011). A better understanding of the underlying pathophysiology may aid in improving
diagnostic accuracy and patient treatment stratification(Vieta et al., 2018); however, the
pathophysiology mechanism of BD remains unclear.

The integrated stress response (ISR) is an evolutionarily conserved intracellular signaling network
that helps the cell, tissue, and organism to adapt to a variable environment and maintain health. In
response to different environmental and pathological conditions, including protein homeostasis
(proteostasis) defects, nutrient deprivation, viral infection, and oxidative stress, the ISR restores
balance by reprogramming gene expression. The various stresses are sensed by four specialized
kinases (PERK, GCN2, PKR and HRI) that converge on phosphorylation of a single serine on the
eukaryotic translation initiation factor eIF2. eIF2 phosphorylation blocks the action of eIF2's guanine
nucleotide exchange factor termed eIF2B, resulting in a general reduction in protein synthesis.
However, if the stress cannot be mitigated, the ISR triggers apoptosis to eliminate the damaged
cell(Costa-Mattioli & Walter, 2020). Extensive research has shown that the ISR is activated in a wide
range of disorders of the brain. This activation is evidenced by detection of eIF2-P and
phosphorylation of PKR, PERK, and GCN2 in post-mortem brains from individuals and animal
models of cognitive and neurodegenerative disorders, including Alzheimer's disease, Parkinson's
disease, Huntington disease, amyotrophic lateral sclerosis (ALS), traumatic brain injury, Down
syndrome, and Charcot-Marie-Tooth disease (Atkin et al., 2008; Buffington et al., 2014; Chang et al.,
2002; Hoozemans et al., 2007; Ishimura et al., 2016; Moon et al., 2018; Smith & Mallucci, 2016).
Notably, ISR activation causes cognitive defects in mouse models of traumatic brain injury (Chou et
al., 2017; Sen et al., 2017), and Alzheimer's disease (Hwang et al., 2017; Tible et al., 2019). Whilst
some research has been carried out on how stress will affect mood disorder, there have been few
empirical investigations into if any change in ISR would bring patients mood disorder.

The approach to empirical research adopted for this study was Mendelian Randomization(MR)
analysis, which is a reliable method to determine causal effects between risk factors and outcomes
using genetic data, which can effectively eliminate the interference of confounding factors (Emdin et
al., 2017). In addition, colocalization is an essential analytical method for exploring the common
causal molecular mechanism among different diseases and disease-related intermediate phenotypes
(Wu et al., 2019). The specific objective of this study was to describe the association between ISR
and BD. As a result, we prioritized several eIF2-associated genes for follow-up functional studies in
future to elucidate the underlying disease mechanisms. we systematically applied summary data-
based Mendelian randomization analysis (SMR) and colocalization (COLOC) to prioritize putative
causal genes using publicly available eQTL datasets. Furthermore, we try to explore potential
alternative drug targets by searching candidate genes available from EpiGraphDB. Then, using MR
analysis to enhance our confidence in identifying potential viable drug targets.

2   Methods

2.1   Study design and data resources

Figure 1 describes the design of this study.

84   Figure 1 Study overview.

For the summary-level data of the GWAS on BD, we used summary statistics of 41917 BD cases and
371549 controls from the Psychiatric Genomics Consortium (Sullivan, 2023). Tissue cis-QTL data
were from the Genotype-Tissue Expression (GTEx) project (Consortium et al., 2020). We
downloaded lite version of the GTEx V8 eqtl data (only SNPs with P < 1e-5 are included). This is a
set of cis-eQTL summary data across 49 human tissues from the GTEx project. Only SNPs within
1Mb of the transcription start site are available. Blood eQTL summary statistics of eIF2 genes were
obtained from eQTLGen including the genetic data of blood gene expression in 31,684 individuals
derived from 37 datasets (Võsa et al., 2021).

2.2   SMR analysis

We conducted SMR and heterogeneity in dependent instruments (HEIDI) tests in cis regions using
the SMR software tool version 1.3.1. Detailed methods for SMR analysis were described in the
original SMR paper (Zhu et al., 2016). The SMR software tool was originally developed to
implement the SMR & HEIDI methods to test for pleiotropic association between the expression
level of a gene and a complex trait of interest using summary-level data from GWAS and expression
quantitative trait loci (eQTL) studies. The SMR & HEIDI methodology can be interpreted as an
analysis to test if the effect size of a SNP on the phenotype is mediated by gene expression. This tool
can therefore be used to prioritize genes underlying GWAS hits for follow-up functional studies.

The default p-value threshold to select the top associated eQTL for the SMR test is 5.0e-8. By
default, we only run the SMR analysis in the cis regions. The default threshold of eQTL p-value to

select eQTLs for the HEIDI test is 1.57e-3, which is equivalent to a chi-squared value (df=1) of 10.
The window centred around the probe to select cis-eQTLs (passing a p-value threshold) for the SMR
analysis is 2000Kb.

2.3 COLOC analysis

We conducted COLOC analysis using the coloc package (Giambartolomei et al., 2014) in R software
(version 4.3.2). The coloc package can be used to perform genetic colocalisation analysis of two
potentially related phenotypes, to ask whether they share common genetic causal variant(s) in a given
region. This package(version 5) supercedes previously published version 4 by introducing use of the
SuSiE approach to deal with multiple causal variants rather than conditioning or masking(Wang et
al., 2020).

COLOC analysis calculates posterior probabilities (PPs) of the five hypotheses: 1) H0; no association
with either gene expression or phenotype; 2) H1; association with gene expression, not with the
phenotype; 3) H2; association with the phenotype, not with gene expression; 4) H3; association with
gene expression and phenotype by independent SNVs; and 5) H4; association with gene expression
and phenotype by shared causal SNVs. A large PP for H4 (PP.H4 above 0.95) strongly supports
shared causal variants affecting both gene expression and phenotype.

2.4 PPI networks & Mendelian randomization

Systematic MR of molecular phenotypes such as proteins and expression of transcript levels offer
enormous potential to prioritise drug targets for further investigation. However, many genes and gene
products are not easily druggable, so some potentially important causal genes may not offer an
obvious route to intervention.

A parallel problem is that current GWASes of molecular phenotypes have limited sample sizes and
limited protein coverages. A potential way to address both these problems is to use protein-protein
interaction information to identify druggable targets which are linked to a non-druggable, but
robustly causal target. Their relationship to the causal target increases our confidence in their
potential causal role even if the initial evidence of effect is below our multiple-testing threshold.

Here, we use data in EpiGraphDB to prioritise potential alternative drug targets in the same PPI
network, as follows:

For an existing drug target of interests, we use PPI networks to search for its directly interacting
genes that are evidenced to be druggable.(Liu et al., 2020). We then examine the causal evidence of
these candidate genes on the disease by using two-sample Mendelian randomization (MR) (Hemani
et al., 2018).

3 Results

3.1 SMR and COLOC analyses using blood eQTL datas

138 The overall results are presented in Table 1.

Table 1 SMR/HEIDI results using the GWAS data on BD.

ProbeID	PROBECHR	GENE	TOPSNP	TOPSNP_CHR	A1	A2	FREQ	B_GWAS	SE_GWAS	P_GWAS	B_EQTL	SE_EQTL	P_EQTL	B_SMR	SE_SMR	P_SMR	P_HEIDI	NSNP_HEIDI	FDR
ENSG00000070785	1	EIF2B3	rs399628	1	A	C	0.375746	-1.0029	0.0095	0.7613	0.143125	0.0200113	8.5382E-13	-7.00716	0.981964	9.6179E-13	3.708E-10	16	1.1541E-12
ENSG00000143486	1	EIF2D	rs66928859	1	G	A	0.425447	0.99541	0.0095	0.6305	0.252463	0.0246545	1.3118E-24	3.9428	0.386871	2.1642E-24	2.8087E-25	16	5.1941E-24
ENSG00000055332	2	EIF2AK2	rs4648160	2	T	C	0.0427435	-0.97356	0.022	0.2236	-0.492181	0.0593344	1.0855E-16	1.97805	0.242615	3.5488E-16	3.6422E-15	20	6.0836E-16
ENSG00000172071	2	EIF2AK3	rs1020616	2	C	T	0.0656183	-1.02798	0.0176	0.1165	0.380813	0.0658245	7.2394E-09	-2.69944	0.468888	8.557E-09	4.0199E-13	7	9.2539E-09
ENSG00000144895	3	EIF2A	rs2868946	3	C	T	0.856859	-1.0008	0.013	0.9495	0.682109	0.0346387	2.5199E-86	-1.46721	0.0769066	3.8571E-81	2.9752E-14	20	4.6286E-80
ENSG00000145191	3	EIF2B5	rs843360	3	A	T	0.442346	-1.01217	0.0094	0.1974	0.11791	0.0204781	8.8164E-09	-8.59293	1.49602	9.2539E-09	0.3175938	8	9.2539E-09
ENSG00000086232	7	EIF2AK1	rs4724770	7	G	A	0.828032	-0.97404	0.0119	0.2699	-0.24618	0.0270822	9.8949E-20	3.95662	0.437943	1.6467E-19	8.539E-55	20	3.2934E-19
ENSG00000236493	10	EIF2S2P3	rs11187107	10	A	G	0.361829	-1.00854	0.0098	0.3851	0.341224	0.0424696	9.3928E-16	-2.95565	0.368988	1.1455E-15	4.0277E-13	20	1.7183E-15
ENSG00000111361	12	EIF2B1	rs112286578	12	A	C	0.163022	-0.99015	0.0129	0.4399	0.316611	0.0314287	5.0136E-26	-2.98588	0.28865	1.4206E-25	2.1631E-17	20	4.2619E-25
ENSG00000134001	14	EIF2S1	rs12590956	14	G	A	0.0715706	-1.04227	0.0195	0.03332	-0.190429	0.0240201	2.2285E-15	5.47327	0.697933	4.43E-15	0.0008309	20	5.9066E-15
ENSG00000119718	14	EIF2B2	rs175040	14	G	A	0.461233	-1.00361	0.0093	0.6958	-0.500686	0.0324789	1.2834E-53	2.00447	0.131348	1.3954E-52	1.9281E-31	20	8.3726E-52
ENSG00000128829	15	EIF2AK4	rs72731415	15	G	A	0.333002	-0.98857	0.0099	0.2496	0.36058	0.0289515	1.3195E-35	-2.74161	0.221833	4.3623E-35	1.0534E-25	20	1.7449E-34

First, we conducted the SMR analysis to integrate the GWAS and blood eQTL data to identify the
most relevant genes whose expression in blood was significantly associated with the trait of DB.
Only one gene passed the SMR test. In chromosome 3, EIF2B5 gene passed both the SMR and
HEIDI tests and thus were significantly associated with the trait of DB because of pleiotropy or
causality.

Next, we conducted the COLOC analysis to integrate the GWAS and blood eQTL data of the genes
that passed the SMR test and assess whether the genes were colocalized with the trait of DB. TIt can
be seen from the data in Table 2 that there is strong support for colocalization between the trait and
the EIF2B5 gene that passed both SMR and HEIDI tests.

Table 2 COLOC results of the genes that passed the SMR test

snp	pvalues.df1	MAF.df1	N.df1	V.df1	z.df1	r.df1	lABF.df1	pvalues.df2	MAF.df2	N.df2	V.df2	z.df2	r.df2	lABF.df2	internal.sum.lABF	SNP.PP.H4
rs843360	0.1974	0.510834	50000	0.000181	1.2889945	0.99549542	-1.8743189	4.2643E-25	0.510834	10000	0.00020009	10.3481992	0.99118533	50.7049839	48.830665	1

3.2 Alt Drug Target

3.2.1 Using PPI networks for alternative drug targets search

The assumption here is that the most likely alternative targets are either directly interacting with
EIF2B5 or somewhere in the same PPI network. In this example, we consider only genes that were
found to interact with EIF2B5 via direct protein-protein interactions, and require that those
interacting proteins should also be druggable.

The thousands of genes are classified with regard to their druggability by (Finan et al., 2017), where
the Tier 1 category refers to approved drugs or those in clinical testing while for other tier categories
the druggability confidence drops in order Tier 2 and then Tier 3.

Table 3 shows the results obtained from data on the druggable alternative genes.

Table 3 Druggable alternative genes

g1.name	p1.uniprot_id	p2.uniprot_id	g2.name	g2.druggability_tier
EIF2B5	Q13144	Q9P2K8	EIF2AK4	Tier 1
EIF2B5	Q13144	P49841	GSK3B	Tier 1
EIF2B5	Q13144	Q9H1Y3	OPN3	Tier 3B

3.2.2 Distinguishing vertical and horizontal pleiotropy for SNP-protein associations

A key MR assumption is that the genetic variant (SNP) is only related to the outcome of interest
through the exposure under study (the "exclusion restriction" assumption). This assumption is
potentially violated under horizontal pleiotropy, where a SNP is associated with multiple phenotypes

(e.g. proteins) independently of the exposure of interest. In contrast, vertical pleiotropy, where a SNP
is associated with multiple phenotypes on the causal pathway to the outcome, does not violate the
"exclusion restriction criterion" of MR.

Here, by integrating SNP-protein associations with biological pathway and protein-protein
interaction (PPI) information retrieved from EpiGraphDB, we have developed an approach to assess
potential horizontal pleiotropy. For a SNP associated with a group of proteins, we check the number
of biological pathways and PPIs that are shared across this group of proteins. If these proteins are
mapped to the same biological pathway and/or a PPI exists between them, then the SNP is more
likely to act through vertical pleiotropy and therefore be a valid instrument for MR.

175   Figure 2 pathway_protein_graph

176 From the graph above we can see that P49841 and Q9P2K8 have same biological pathways, the table
177 below illustrates the specific pathways that how they interact.

178 Table 4 Specific pathways that how proteins interact

node.name	node.id	node.url
Cellular responses to external stimuli	R-HSA-8953897	https://reactome.org/PathwayBrowser/#/R-HSA-8953897
Cellular responses to stress	R-HSA-2262752	https://reactome.org/PathwayBrowser/#/R-HSA-2262752

179 Then we complement the information given by pathway ontologies with data of interactions between
180 proteins. For each pair of proteins, we retrieve shared protein-protein interactions, either a direct
181 interaction or an interaction where there's one mediator protein in the middle. As can be seen from
182 the figure3, for each pair of proteins, there are shared protein-protein interactions.

Figure 3 ppi_protein_graph

185   Overall, these results indicate that our study will not violate under horizontal pleiotropy.

186   **3.2.3 Using Mendelian randomization results for causal effect estimation**

187   The next step is to find out whether any of these genes have a comparable and statistically plausible
188   effect on DB. For further analysis we select the gene of interest (EIF2B5) as well as its interacting
189   genes with Tier 1 druggability.

191   Figure 4 Scatter plots of the association of EIF2B5 and BD

193   Figure 5 Scatter plots of the association of EIF2AK4 and BD

195   Figure 6 Scatter plots of the association of GSK3B and BD

196   It is worth noting that there were also differences in the ratios of MR test. As noted by Burgess et al.
197   (2023); Burgess and Thompson (2017), due to the heterogeneity test highlighting the existence of
198   heterogeneity (P < 0.05), we give priority to using a random effect model of IVW and/or Weighted
199   median method. The results, as shown in Table 5, indicate that a causal association was found
200   between EIF2B5, EIF2AK4, GSK3B and BD.

Table 5 MR analysis of putative causal BD genes and bipolar disorder.

exposure	outcome	method	nsnp	b	se	pval	lo_ci	up_ci	or	or_lci95	or_uci95	Q	Q_df	Q_pval	egger_intercept	se	pval
							MR analyses						*Heterogeneity*			*Pleiotropy*	
EIF2B5 (clump_r2=0.18, clump_kb=1000)	BD	MR Egger	65	-5.21E-01	1.20E+00	6.65E-01	-2.87E+00	1.82E+00	5.94E-01	6.20E-02	6.20E+00	202058.3	63	0	3.98E-01	2.31E-01	9.08E-02
		Weighted median	65	1.58E+00	9.13E-02	5.37E-67	1.40E+00	1.76E+00	4.85E+00	4.05E+00	5.80E+00						
		Inverse variance weighted	65	1.25E+00	6.19E-01	4.37E-02	3.54E-02	2.46E+00	3.48E+00	1.04E+00	1.17E+01	211521	64	0			
		Simple mode	65	4.25E+00	1.84E-01	8.77E-33	3.89E+00	4.61E+00	7.03E+01	4.90E+01	1.01E+02						
		Weighted mode	65	1.50E+00	2.45E-01	6.76E-08	1.02E+00	1.98E+00	4.47E+00	2.76E+00	7.23E+00						
EIF24K4		MR Egger	290	1.72E+00	6.70E-01	1.09E-02	4.04E-01	3.03E+00	5.57E+00	1.50E+00	2.07E+01	1570106	288	0	-1.54E-01	1.15E-01	1.81E-01
		Weighted median	290	2.31E+00	6.09E-02	0.00E+00	2.19E+00	2.43E+00	1.00E+01	8.92E+00	1.13E+01						
		Inverse variance weighted	290	9.42E-01	3.39E-01	5.46E-03	2.77E-01	1.61E+00	2.57E+00	1.32E+00	4.99E+00	1579904	289	0			
		Simple mode	290	5.59E+00	8.68E-02	2.37E-173	5.41E+00	5.76E+00	2.66E+02	2.25E+02	3.16E+02						
		Weighted mode	290	3.19E+00	8.52E-02	7.01E-113	3.02E+00	3.36E+00	2.43E+01	2.05E+01	2.87E+01						
GSK3B (clump_r2=0.05, clump_kb=100)		MR Egger	19	-4.98E-01	2.78E+00	8.60E-01	-5.95E+00	4.95E+00	6.08E-01	2.62E-03	1.41E+02	37775.24	17	0	5.82E-01	4.31E-01	1.95E-01
		Weighted median	19	1.81E+00	1.21E-01	1.70E-50	1.57E+00	2.04E+00	6.09E+00	4.81E+00	7.72E+00						
		Inverse variance weighted	19	2.80E+00	1.36E+00	3.93E-02	1.38E-01	5.46E+00	1.64E+01	1.15E+00	2.35E+02	41822.66	18	0			
		Simple mode	19	5.59E+00	4.32E-01	1.50E-10	4.74E+00	6.44E+00	2.68E+02	1.15E+02	6.25E+02						
		Weighted mode	19	2.16E+00	1.07E-01	8.31E-14	1.95E+00	2.36E+00	8.63E+00	7.00E+00	1.06E+01						

4   Discussion

The results of this study show that EIF2B5 is the potential disease-causing gene of BD, EIF2AK4 and GSK3B may become alternative drug targets for this disease.

Perhaps the most interesting finding is early research regarding EIF2B5 mainly aimed to study the roles of its expression in few cancerous tissue(Hou et al., 2020; Jiao et al., 2018; Xue et al., 2014; Yang et al., 2014), multiple sclerosis(Zahoor et al., 2017), and vanishing white matter (VWM) (Dietrich et al., 2005)syndrome. Importantly, the eIF2B complex plays a central role in the cellular integrated stress response (ISR). Stress-dependent kinase activation leads to phosphorylation of eIF2, which binds eIF2B more tightly and reduces overall protein synthesis. Previous research has established that white matter microstructure alterations in bipolar disorder(Adler et al., 2004; Bellani & Brambilla, 2011; Bellani et al., 2012; López-Larson et al., 2002; Yurgelun-Todd et al., 2007). However, eukaryotic translation initiation factor 2B (EIF2B) gene family encoding the five subunits of eIF2B complex-α, β, γ, δ and ε respectively. This finding supports previous research into this brain area which links EIF2B5 and BD.

The ISR responds to a variety of different stress conditions that lead to alterations in cellular homeostasis. In metazoans, these stresses are sensed by at least four different kinases that each phosphorylate Ser51 in eIF2a to activate the ISR(Wek, 2018). One of them, GCN2(gene name EIF2AK4) contains a regulatory domain homologous with histidyl-tRNA synthetase and when amino acids are scarce, binding of deacylated His-tRNA triggers its activation (Harding et al., 2000; Hinnebusch, 2005). However, the precise mechanism(s) underlying these vastly different activation modalities remain unknown. But it is well known that the HPA axis is closely related to stress and AgRP neurons(Cleber Gama de Barcellos Filho et al., 2021; Perry et al., 2015; Perry et al., 2019). A possible explanation for this might further support the idea of Yuan et al. (2023): hypothalamic AgRP neurons are involved in the process of leucine deficiency improving depressive-like behavior induced by CRS, and the amino acid sensor general control non-derepressible 2 (GCN2) mediates this behavioral effect of AgRP neurons.

When it comes to GSK3B, Chen et al. (2014) showed that published data on the association between GSK3B -50C/T (rs334558) and bipolar disorder are inconclusive. This differs from the findings presented here. It could be argued that the positive results were due to the relatively much bigger sample size and a multiple SNPs-based approach can provide a more objective evaluation. Despite its exploratory nature, this study offers some insight into GSK3B, which is not only implicated in the pathophysiology of Alzheimer disease (Mendes et al., 2009), Parkinson disease (Wider et al., 2011), schizophrenia (Chen et al., 2020), and depressive disorders (Levchenko et al., 2018).

The generalisability of these results is subject to certain limitations. Firstly, the SMR method cannot distinguish pleiotropic genes from the causative genes. The association between prioritized genes and BD needs to be validated by follow-up functional studies. Moreover, BD cohorts in our data were collected in Europe, North America and Australia, so these findings cannot be extrapolated to all patients.

5   Conclusion

This study set out to examine the relationship between ISR and BD, also, assess the feasibility of some genes' druggability. The most obvious finding to emerge from this study is that we established causal relationships between genetic variants, candidate genes, and BD risk. The insights gained from this study may be of assistance to cure this disease.

6   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

7   Author Contributions

ZT designed this study, analyzed data, and wrote the draft of the manuscript. Prof.Deng huihua discussed and reviewed the manuscript critically. All authors contributed to the article and approved the submitted version.

8   Acknowledgments

The authors thank Yang Lab for the user-friendly SMR software. We also appreciate consortia such as the PGC, GTEx and eQTLGen for providing publicly available data.

9   Data Availability Statement

The summary-level data of the GWAS on BD was publicly available from the Psychiatric Genomics Consortium (https://figshare.com/articles/dataset/bip2021_noUKBB/22564402/1). The eQTL dataset was publicly available from the eQTLGen consortium (https://www.eqtlgen.org/). V8 release of the GTEx eQTL/sQTL summary data can be downloaded from https://yanglab.westlake.edu.cn/software/smr/#DataResource.

10   Reference

Adler, C. M., Holland, S. K., Schmithorst, V., Wilke, M., Weiss, K. L., Pan, H., & Strakowski, S. M. (2004). Abnormal frontal white matter tracts in bipolar disorder: a diffusion tensor imaging study. *Bipolar Disorders*, *6*(3), 197-203. https://doi.org/10.1111/j.1399-5618.2004.00108.x

Atkin, J. D., Farg, M. A., Walker, A. K., McLean, C., Tomas, D., & Horne, M. K. (2008). Endoplasmic reticulum stress and induction of the unfolded protein response in human sporadic amyotrophic lateral sclerosis. *Neurobiology of Disease*, *30*(3), 400-407. https://doi.org/10.1016/j.nbd.2008.02.009

Bellani, M., & Brambilla, P. (2011). Diffusion imaging studies of white matter integrity in bipolar disorder. *Epidemiology and Psychiatric Sciences*, *20*(2), 137-140. https://doi.org/10.1017/s2045796011000229

Bellani, M., Perlini, C., Ferro, A., Cerruti, S., Rambaldelli, G., Isola, M., Cerini, R., Dusi, N., Andreone, N., Balestrieri, M., Mucelli, R. P., Tansella, M., & Brambilla, P. (2012). White matter microstructure alterations in bipolar disorder. *Functional Neurology*, *27*(1), 29-34. https://pubmed.ncbi.nlm.nih.gov/22687164/

Brown, G. K., Beck, A. T., Steer, R. A., & Grisham, J. R. (2000). Risk factors for suicide in psychiatric outpatients: a 20-year prospective study. *Journal of consulting and clinical psychology*, *68*(3), 371-377. https://doi.org/10.1037/0022-006x.68.3.371

Buffington, S. A., Huang, W., & Costa-Mattioli, M. (2014). Translational Control in Synaptic Plasticity and Cognitive Dysfunction. In S. E. Hyman (Ed.), *Annual Review of Neuroscience, Vol 37* (Vol. 37, pp. 17-38). Annual Reviews. https://doi.org/10.1146/annurev-neuro-071013-014100

Burgess, S., Davey Smith, G., Davies, N., Dudbridge, F., Gill, D., Glymour, M., Hartwig, F., Kutalik,
Z., Holmes, M., Minelli, C., Morrison, J., Pan, W., Relton, C., & Theodoratou, E. (2023).
Guidelines for performing Mendelian randomization investigations: update for summer 2023
[version 3; peer review: 2 approved]. *Wellcome Open Research*, *4*(186).
https://doi.org/10.12688/wellcomeopenres.15555.3

Burgess, S., & Thompson, S. G. (2017). Interpreting findings from Mendelian randomization using
the MR-Egger method. *European Journal of Epidemiology*, *32*(5), 377-389.
https://doi.org/10.1007/s10654-017-0255-x

Chang, R. C. C., Wong, A. K. Y., Ng, H.-K., & Hugon, J. (2002). Phosphorylation of eukaryotic
initiation factor-2α (eIF2α) is associated with neuronal degeneration in Alzheimer's disease.
NeuroReport, *13*(18), 2429-2432. https://doi.org/10.1097/00001756-200212200-00011

Chen, G. D., Tang, J., Yu, G. W., Chen, Y. P., Wang, L. C., & Zhang, Y. (2014). Meta-analysis
demonstrates lack of association of the *GSK3B*-50C/T polymorphism with risk of
bipolar disorder. *Molecular Biology Reports*, *41*(9), 5711-5718.
https://doi.org/10.1007/s11033-014-3441-x

Chen, Y., Hua, S., Wang, W. P., Fan, W. X., Tang, W., Zhang, Y., & Zhang, C. (2020). A
comprehensive analysis of *GSK3B* variation for schizophrenia in Han Chinese
individuals. *Asian Journal of Psychiatry*, *47*, Article 101832.
https://doi.org/10.1016/j.ajp.2019.10.012

Chou, A., Krukowski, K., Jopson, T., Zhu, P. J., Costa-Mattioli, M., Walter, P., & Rosi, S. (2017).
Inhibition of the integrated stress response reverses cognitive deficits after traumatic brain
injury. *Proceedings of the National Academy of Sciences of the United States of America*,
114(31), E6420-E6426. https://doi.org/10.1073/pnas.1707661114

Cleber Gama de Barcellos Filho, P., Campos Zanelatto, L., Amélia Aparecida Santana, B., Calado,
R. T., & Rodrigues Franci, C. (2021). Effects chronic administration of corticosterone and
estrogen on HPA axis activity and telomere length in brain areas of female rats. *Brain
Research*, *1750*, 147152. https://doi.org/10.1016/j.brainres.2020.147152

Cloutier, M., Greene, M., Guerin, A., Touya, M., & Wu, E. (2018). The economic burden of bipolar I
disorder in the United States in 2015. *Journal of Affective Disorders*, *226*, 45-51.
https://doi.org/10.1016/j.jad.2017.09.011

Consortium, T. G., Aguet, F., Anand, S., Ardlie, K. G., Gabriel, S., Getz, G. A., Graubert, A.,
Hadley, K., Handsaker, R. E., Huang, K. H., Kashin, S., Li, X., MacArthur, D. G., Meier, S.
R., Nedzel, J. L., Nguyen, D. T., Segrè, A. V., Todres, E., Balliu, B., . . . Volpi, S. (2020).
The GTEx Consortium atlas of genetic regulatory effects across human tissues. *Science*,
369(6509), 1318-1330. https://doi.org/10.1126/science.aaz1776

Costa-Mattioli, M., & Walter, P. (2020). The integrated stress response: From mechanism to disease.
Science, *368*(6489), 384-+, Article eaat5314. https://doi.org/10.1126/science.aat5314

Diagnostic and statistical manual of mental disorders: DSM-5™, 5th ed. (2013). American
Psychiatric Publishing, Inc. https://doi.org/10.1176/appi.books.9780890425596

Dietrich, J., Lacagnina, M., Gass, D., Richfield, E., Mayer-Pröschel, M., Noble, M., Torres, C., &
Pröschel, C. (2005). *EIF2B5* mutations compromise GFAP+ astrocyte
generation in vanishing white matter leukodystrophy. *Nature Medicine*, *11*(3), 277-283.
https://doi.org/10.1038/nm1195

Emdin, C. A., Khera, A. V., & Kathiresan, S. (2017). Mendelian Randomization. *JAMA*, *318*(19), 1925-1926. https://doi.org/10.1001/jama.2017.17219

Finan, C., Gaulton, A., Kruger, F. A., Lumbers, R. T., Shah, T., Engmann, J., Galver, L., Kelley, R., Karlsson, A., Santos, R., Overington, J. P., Hingorani, A. D., & Casas, J. P. (2017). The druggable genome and support for target identification and validation in drug development. *Science Translational Medicine*, *9*(383), eaag1166. https://doi.org/10.1126/scitranslmed.aag1166

Giambartolomei, C., Vukcevic, D., Schadt, E. E., Franke, L., Hingorani, A. D., Wallace, C., & Plagnol, V. (2014). Bayesian Test for Colocalisation between Pairs of Genetic Association Studies Using Summary Statistics. *Plos Genetics*, *10*(5), Article e1004383. https://doi.org/10.1371/journal.pgen.1004383

Grande, I., Berk, M., Birmaher, B., & Vieta, E. (2016). Bipolar disorder [Review]. *Lancet*, *387*(10027), 1561-1572. https://doi.org/10.1016/s0140-6736(15)00241-x

Harding, H. P., Novoa, I., Zhang, Y., Zeng, H., Wek, R., Schapira, M., & Ron, D. (2000). Regulated Translation Initiation Controls Stress-Induced Gene Expression in Mammalian Cells. *Molecular Cell*, *6*(5), 1099-1108. https://doi.org/10.1016/S1097-2765(00)00108-8

Hemani, G., Zheng, J., Elsworth, B., Wade, K. H., Haberland, V., Baird, D., Laurin, C., Burgess, S., Bowden, J., Langdon, R., Tan, V. Y., Yarmolinsky, J., Shihab, H. A., Timpson, N. J., Evans, D. M., Relton, C., Martin, R. M., Davey Smith, G., Gaunt, T. R., & Haycock, P. C. (2018). The MR-Base platform supports systematic causal inference across the human phenome. *eLife*, *7*, e34408. https://doi.org/10.7554/eLife.34408

Hinnebusch, A. G. (2005). TRANSLATIONAL REGULATION OF GCN4 AND THE GENERAL AMINO ACID CONTROL OF YEAST. *Annual Review of Microbiology*, *59*(1), 407-450. https://doi.org/10.1146/annurev.micro.59.031805.133833

Hoozemans, J. J. M., van Haastert, E. S., Eikelenboom, P., de Vos, R. A. I., Rozemuller, J. M., & Scheper, W. (2007). Activation of the unfolded protein response in Parkinson's disease. *Biochemical and Biophysical Research Communications*, *354*(3), 707-711. https://doi.org/10.1016/j.bbrc.2007.01.043

Hou, L., Jiao, Y., Li, Y. Q., Luo, Z. P., Zhang, X. Y., Pan, G. Q., Zhao, Y. C., Yang, Z. Y., & He, M. (2020). Low <i>EIF2B5<i> expression predicts poor prognosis in ovarian cancer. *Medicine*, *99*(5), 6, Article e18666. https://doi.org/10.1097/md.0000000000018666

Hwang, K. D., Bak, M. S., Kim, S. J., Rhee, S., & Lee, Y. S. (2017). Restoring synaptic plasticity and memory in mouse models of Alzheimer's disease by PKR inhibition. *Molecular Brain*, *10*, 10, Article 57. https://doi.org/10.1186/s13041-017-0338-3

Ishimura, R., Nagy, G., Dotu, I., Chuang, J. H., & Ackerman, S. L. (2016). Activation of GCN2 kinase by ribosome stalling links translation elongation with translation initiation. *eLife*, *5*, 22, Article e14295. https://doi.org/10.7554/eLife.14295

Jacobs, D., & Brewer, M. (2004). APA practice guideline provides recommendations for assessing and treating patients with suicidal behaviors. In (Vol. 34, pp. 373-380): SLACK Incorporated Thorofare, NJ.

Jiao, Y., Fu, Z., Li, Y. Q., Meng, L. Y., & Liu, Y. H. (2018). High EIF2B5 mRNA expression and its prognostic significance in liver cancer: a study based on the TCGA and GEO database. *Cancer Management and Research*, *10*, 6003-6014. https://doi.org/10.2147/cmar.S185459

Levchenko, A., Losenkov, I. S., Vyalova, N. M., Simutkin, G. G., Bokhan, N. A., Wilffert, B.,
Loonen, A. J. M., & Ivanova, S. A. (2018). The functional variant rs334558 of GSK3B is
associated with remission in patients with depressive disorders. *Pharmacogenomics &*
Personalized Medicine, *11*, 121-126. https://doi.org/10.2147/pgpm.S171423

Liu, Y., Elsworth, B., Erola, P., Haberland, V., Hemani, G., Lyon, M., Zheng, J., Lloyd, O.,
Vabistsevits, M., & Gaunt, T. R. (2020). EpiGraphDB: a database and data mining platform
for health data science. *Bioinformatics*, *37*(9), 1304-1311.
https://doi.org/10.1093/bioinformatics/btaa961

López-Larson, M. P., DelBello, M. P., Zimmerman, M. E., Schwiers, M. L., & Strakowski, S. M.
(2002). Regional prefrontal gray and white matter abnormalities in bipolar disorder.
Biological Psychiatry, *52*(2), 93-100, Article Pii s0006-3223(02)01350-1.
https://doi.org/10.1016/s0006-3223(02)01350-1

Mendes, C. T., Mury, F. B., Moreira, E. D., Alberto, F. L., Forlenza, O. V., Dias-Neto, E., & Gattaz,
W. F. (2009). Lithium reduces Gsk3b mRNA levels: implications for Alzheimer Disease.
European Archives of Psychiatry and Clinical Neuroscience, *259*(1), 16-22.
https://doi.org/10.1007/s00406-008-0828-5

Moon, S. L., Sonenberg, N., & Parker, R. (2018). Neuronal Regulation of eIF2α Function in Health
and Neurological Disorders. *Trends in Molecular Medicine*, *24*(6), 575-589.
https://doi.org/10.1016/j.molmed.2018.04.001

Perry, R. J., Lee, S., Ma, L., Zhang, D., Schlessinger, J., & Shulman, G. I. (2015). FGF1 and FGF19
reverse diabetes by suppression of the hypothalamic–pituitary–adrenal axis. *Nature*
communications, *6*(1), 6980. https://doi.org/10.1038/ncomms7980

Perry, R. J., Resch, J. M., Douglass, A. M., Madara, J. C., Rabin-Court, A., Kucukdereli, H., Wu, C.,
Song, J. D., Lowell, B. B., & Shulman, G. I. (2019). Leptin's hunger-suppressing effects are
mediated by the hypothalamic–pituitary–adrenocortical axis in rodents. *Proceedings of the*
National Academy of Sciences, *116*(27), 13670-13679.
https://doi.org/doi:10.1073/pnas.1901795116

Sani, G., Tondo, L., Koukopoulos, A., Reginaldi, D., Kotzalidis, G. D., Koukopoulos, A. E.,
Manfredi, G., Mazzarini, L., Pacchiarotti, I., Simonetti, A., Ambrosi, E., Angeletti, G.,
Girardi, P., & Tatarelli, R. (2011). Suicide in a large population of former psychiatric
inpatients. *Psychiatry and Clinical Neurosciences*, *65*(3), 286-295.
https://doi.org/10.1111/j.1440-1819.2011.02205.x

Sen, T., Gupta, R., Kaiser, H., & Sen, N. (2017). Activation of PERK Elicits Memory Impairment
through Inactivation of CREB and Downregulation of PSD95 After Traumatic Brain Injury.
Journal of Neuroscience, *37*(24), 5900-5911. https://doi.org/10.1523/jneurosci.2343-16.2017

Smith, H. L., & Mallucci, G. R. (2016). The unfolded protein response: mechanisms and therapy of
neurodegeneration. *Brain*, *139*, 2113-2121. https://doi.org/10.1093/brain/aww101

Sullivan, P. (2023). *bip2021_noUKBB. figshare*. https://doi.org/10.6084/m9.figshare.22564402.v1

Takizawa, R., Kasai, K., Kawakubo, Y., Marumo, K., Kawasaki, S., Yamasue, H., & Fukuda, M.
(2008). Reduced frontopolar activation during verbal fluency task in schizophrenia: a multi-
channel near-infrared spectroscopy study. *Schizophr Res*, *99*(1-3), 250-262.
https://doi.org/10.1016/j.schres.2007.10.025

Tible, M., Liger, F. M., Schmitt, J., Giralt, A., Farid, K., Thomasseau, S., Gourmaud, S., Paquet, C., Reig, L. R., Meurs, E., Girault, J. A., & Hugon, J. (2019). PKR knockout in the 5xFAD model of Alzheimer's disease reveals beneficial effects on spatial memory and brain lesions. *Aging Cell*, *18*(3), 11, Article e12887. https://doi.org/10.1111/acel.12887

Vieta, E., Salagre, E., Grande, I., Carvalho, A. F., Fernandes, B. S., Berk, M., Birmaher, B., Tohen, M., & Suppes, T. (2018). Early Intervention in Bipolar Disorder. *American Journal of Psychiatry*, *175*(5), 411-426. https://doi.org/10.1176/appi.ajp.2017.17090972

Võsa, U., Claringbould, A., Westra, H.-J., Bonder, M. J., Deelen, P., Zeng, B., Kirsten, H., Saha, A., Kreuzhuber, R., Yazar, S., Brugge, H., Oelen, R., de Vries, D. H., van der Wijst, M. G. P., Kasela, S., Pervjakova, N., Alves, I., Favé, M.-J., Agbessi, M., . . . i, Q. T. L. C. (2021). Large-scale cis- and trans-eQTL analyses identify thousands of genetic loci and polygenic scores that regulate blood gene expression. *Nature Genetics*, *53*(9), 1300-1310. https://doi.org/10.1038/s41588-021-00913-z

Wang, G., Sarkar, A., Carbonetto, P., & Stephens, M. (2020). A Simple New Approach to Variable Selection in Regression, with Application to Genetic Fine Mapping. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, *82*(5), 1273-1300. https://doi.org/10.1111/rssb.12388

Wek, R. C. (2018). Role of eIF2α kinases in translational control and adaptation to cellular stress. *Cold Spring Harbor perspectives in biology*, *10*(7), a032870. https://doi.org/10.1101/cshperspect.a032870

Wider, C., Vilariño-Güell, C., Heckman, M. G., Jasinska-Myga, B., Ortolaza-Soto, A. I., Diehl, N. N., Crook, J. E., Cobb, S. A., Bacon, J. A., Aasly, J. O., Gibson, J. M., Lynch, T., Uitti, R. J., Wszolek, Z. K., Farrer, M. J., & Ross, O. A. (2011). *SNCA*, *MAPT*, and *GSK3B* in Parkinson disease: a gene-gene interaction study. *European Journal of Neurology*, *18*(6), 876-881. https://doi.org/10.1111/j.1468-1331.2010.03297.x

Wu, Y., Broadaway, K. A., Raulerson, C. K., Scott, L. J., Pan, C., Ko, A., He, A., Tilford, C., Fuchsberger, C., Locke, A. E., Stringham, H. M., Jackson, A. U., Narisu, N., Kuusisto, J., Pajukanta, P., Collins, F. S., Boehnke, M., Laakso, M., Lusis, A. J., . . . Mohlke, K. L. (2019). Colocalization of GWAS and eQTL signals at loci with multiple signals identifies additional candidate genes for body fat distribution. *Human Molecular Genetics*, *28*(24), 4161-4172. https://doi.org/10.1093/hmg/ddz263

Xue, D., Lu, M., Gao, B., Qiao, X., & Zhang, Y. (2014). Screening for transcription factors and their regulatory small molecules involved in regulating the functions of CL1-5 cancer cells under the effects of macrophage-conditioned medium. *Oncol Rep*, *31*(3), 1323-1333. https://doi.org/10.3892/or.2013.2937

Yang, S., Zhang, H., Guo, L., Zhao, Y., & Chen, F. (2014). Reconstructing the coding and non-coding RNA regulatory networks of miRNAs and mRNAs in breast cancer. *Gene*, *548*(1), 6-13. https://doi.org/10.1016/j.gene.2014.06.010

Yuan, F., Wu, S., Zhou, Z., Jiao, F., Yin, H., Niu, Y., Jiang, H., Chen, S., & Guo, F. (2023). Leucine deprivation results in antidepressant effects via GCN2 in AgRP neurons. *Life Metabolism*, *2*(1). https://doi.org/10.1093/lifemeta/load004

Yurgelun-Todd, D. A., Silveri, M. M., Gruber, S. A., Rohan, M. L., & Pimentel, P. J. (2007). White matter abnormalities observed in bipolar disorder: a diffusion tensor imaging study. *Bipolar Disorders*, *9*(5), 504-512. https://doi.org/10.1111/j.1399-5618.2007.00395.x

Zahoor, I., Haq, E., & Asimi, R. (2017). Multiple Sclerosis and EIF2B5: A Paradox or a Missing
Link. In A. A. A. Asea, F. Geraci, & P. Kaur (Eds.), *Multiple Sclerosis: Bench to Bedside:*
Global Perspectives on a Silent Killer (Vol. 958, pp. 57-64). Springer International
Publishing Ag. https://doi.org/10.1007/978-3-319-47861-6_5

Zhu, Z., Zhang, F., Hu, H., Bakshi, A., Robinson, M. R., Powell, J. E., Montgomery, G. W.,
Goddard, M. E., Wray, N. R., Visscher, P. M., & Yang, J. (2016). Integration of summary
data from GWAS and eQTL studies predicts complex trait gene targets. *Nature Genetics*,
48(5), 481-487. https://doi.org/10.1038/ng.3538