Identifying direct risk factors in UK Biobank with simultaneous Bayesian-frequentist model-averaged hypothesis testing using Doublethink

Nicolas Arning[1], Helen R. Fryer[1], Daniel J. Wilson[1,2,*]

1. Big Data Institute, Oxford Population Health, University of Oxford
2. Oxford University Department for Continuing Education
* Address for correspondence: Li Ka Shing Centre for Health Information and Discovery, Old Road Campus, Oxford, OX3 7LF, United Kingdom. Email daniel.wilson@bdi.ox.ac.uk

21 May 2025

Abstract

Big data approaches to discovering non-genetic risk factors have lagged behind genome-wide association studies that routinely uncover novel genetic risk factors for diverse diseases. Instead, epidemiology typically focuses on candidate risk factors. Since modern biobanks contain thousands of potential risk factors, candidate approaches may introduce bias, inadequately control for multiple testing, and overlook important signals. Doublethink, a novel model-averaged hypothesis testing approach, offers a solution that simultaneously controls the Bayesian false discovery rate (FDR) and frequentist familywise error rate (FWER) while accounting for uncertainty in variable selection. Here we investigate direct risk factors for COVID-19 hospitalization from among 1,912 variables in 201,917 UK Biobank participants by implementing a Doublethink-based exposome-wide association study using Markov Chain Monte Carlo. Focusing on the 2020 outbreak, we find nine individual variables and six groups of variables exposome-wide significant at 9% FDR and 0.05% FWER. We identify significant direct effects among relatively overlooked risk factors including psychiatric disorders, dementia and prior infection, which we evaluate in relation to studies of other populations. We detect significant direct effects among some commonly reported risk factors like age, sex and obesity, but not others like diabetes, cardiovascular disease, hypertension, which may be mediated instead through variables representing general comorbidity. Doublethink produces interchangeable posterior odds and *p*-values for individual variables and arbitrary groups, facilitating flexible and powerful *post-hoc* hypothesis testing. We discuss the potential for impact and limitations of joint Bayesian-frequentist hypothesis testing, including the benefits of an agnostic exposome-wide approach to discovery.

Significance

Understanding what causes disease is key to improving its treatment and prevention. Large health studies like UK Biobank measure thousands of possible causes of disease. Traditionally, scientists have studied possible causes (like smoking or exercise) one-at-a-time, in depth. For greater perspective, we could study them altogether to test which have any effect. We recently introduced Doublethink, which combines the advantages of two major statistical approaches to testing. Here we use Doublethink to test 1,912 possible causes of COVID-19 hospitalization in UK Biobank. We found strong evidence for relatively overlooked causes: psychiatric conditions, dementia and previous infections. Findings from other health studies support these causes, highlighting the need to re-evaluate them and showing how our approach can reveal valuable insights.

Keywords: UK Biobank, COVID-19 hospitalization, exposome-wide association studies, Bayesian model-averaging, FDR, FWER

Introduction

The big data era has seen the advent of biobank-scale scans for genetic determinants of diverse health outcomes in cohorts like UK Biobank (1, 2). But similar data-driven identification of non-genetic determinants, termed risk factors, has not become commonplace. Instead, current epidemiology typically reports on candidate risk factors. Studies aim to address the question: What is the total effect of a variable on the outcome? Is it non-zero? For instance, more than 100 published studies have analysed dozens of candidate risk factors for COVID-19 outcomes in UK Biobank (Table S1). Synthesizing these findings is difficult because: (i) Other, more important, risk factors that were not analysed may exist among the thousands measured. (ii) It is unclear how to appropriately limit false positives caused by multiple testing. (iii) The processes of selecting candidate risk factors and deciding to publish are vulnerable to bias. The experience of candidate gene studies, largely superseded by genome-wide association studies (GWAS), raises further questions about strength of evidence and reproducibility in candidate risk factor studies (3, 4, 5).

A major complication for systematic studies of non-genetic risk factors, compared to GWAS, is the problem of mediation (6). Mediation occurs when the total effect of a variable (e.g. age) on an outcome (e.g. COVID-19 severity) is wholly or partially caused through another variable (e.g. prior pneumonia). This conceptually divides the total effect into direct and indirect effects. Mediation is ignored in GWAS because genetic variables are coinherited at conception; they cannot generally cause one another. So the question in GWAS is effectively: What is the direct effect of a variable on the outcome? Is it non-zero? Artefactual signals generated by confounding are instead the major concern. Controlling for other variables helps avoid confounding (7), but controlling for mediating variables alters the scientific question by shifting attention from total to direct effects. Unfortunately, direct effects can differ in direction and magnitude to total effects, a source of bias known as the Table 2 fallacy (8). Further pitfalls include reverse causation and collider bias (9).

Nevertheless, the demand for GWAS-inspired exposome-wide association studies (10) presents an opportunity, which has been partly filled by machine learning (11, 12). Machine learning offers a data-driven agnostic approach. A major advantage is its ability to analyse high dimensional data with minimal intervention, even in the presence of collinearity and widespread correlation between variables. But the question in machine learning is different: What is the contribution of a variable to predicting the outcome? Usually there is no formal test. More importantly, a variable can be valuable for prediction due to confounding (13). Machine learning is therefore problematic for risk factor identification. Other concerns have been raised with artificial intelligence approaches in healthcare, particularly in terms of often difficult-to-achieve interpretability and equity (14).

Bayesian methodology offers a solution to the question of identifying direct effects in biobank-scale data while controlling for confounding (15). An important advantage is the ability to account for uncertainty in model choice by averaging over the inclusion or exclusion of other variables when estimating or testing the direct effect of each variable. This uncertainty can strongly influence conclusions. The question is therefore: What is the explanatory value of each variable, over and above all the other variables? Is it non-zero? With a careful approach to feature engineering to mitigate issues around mediation, reverse causality and collider bias, and with independent replication of discoveries, Bayesian model averaging (BMA) offers a powerful approach. But Bayesian approaches are seldom used in current epidemiology: none of 127 published studies of risk factors for COVID-19 outcomes in UK Biobank was Bayesian (Table S1). This might be explained by several issues, including lack of familiarity among researchers, high computational requirements, and difficulties specifying prior distributions (16). Many practitioners worry about the role of the prior in Bayesian hypothesis testing, which can lead different researchers to different conclusions from the same data (17).

Doublethink (18) is a new approach that aims to address these concerns by facilitating joint Bayesian-frequentist model-averaged hypothesis testing while simultaneously controlling the Bayesian false discovery rate (FDR) and the frequentist familywise error rate (FWER). By implementing Doublethink using a Markov Chain Monte Carlo (MCMC) algorithm, we are able to test for direct risk factors among thousands of individual variables, and arbitrary groups of those variables, while accounting for uncertainty in variable selection. We apply Doublethink to investigate direct risk factors for COVID-19 hospitalization in UK Biobank among 1,912 variables in 201,917 participants, and we compare our results to the literature. Our framework provides a highly capable model-averaging approach that can be applied to the systematic evaluation of direct risk factors in biobank-scale resources.

Theory

We consider a general regression setting in which there are n observed outcomes $y_1 \ldots y_n$ and v variables (features) with regression coefficients $\beta_1 \ldots \beta_v$. The aim is to identify which variables directly influence the outcome, i.e. which of the regression coefficients are non-zero. In total there are 2^v hypotheses, ω_v, which we index with vector $\boldsymbol{v}$. The jth element of $\boldsymbol{v}$ indicates whether we are testing that variable j is zero ($v_j = 0$; $\beta_j = 0$) or not testing it ($v_j = 1$; $\beta_j = 0$ or $\beta_j \neq 0$).

In parallel, we define 2^v models with non-overlapping parameter spaces $\Theta_{\boldsymbol{s}}$, indexed by vector $\boldsymbol{s}$, the jth element of which indicates whether variable j is zero ($s_j = 0$; $\beta_j = 0$) or non-zero ($s_j = 1$; $\beta_j \neq 0$). Each null hypothesis ω_v is compatible with one or more models $\Theta_{\boldsymbol{s}}$, indexed by the set

$$\mathcal{O}_v = \{s: s_j = 0 \text{ for all } v_j = 0\},$$

and incompatible with all the other models, indexed by the complementary set $\mathcal{A}_v$. In the Bayesian setting, we reject null hypothesis ω_v if the posterior odds of models in $\mathcal{A}_v$ versus $\mathcal{O}_v$,

$$\text{PO}_{\mathcal{A}_v : \mathcal{O}_v} = \frac{\sum_{\boldsymbol{s} \in \mathcal{A}_v} \text{PO}_{\boldsymbol{s}}}{\sum_{\boldsymbol{s} \in \mathcal{O}_v} \text{PO}_{\boldsymbol{s}}}, \tag{1}$$

exceed some threshold τ. Here $\text{PO}_{\boldsymbol{s}}$ represents the posterior odds of model $\boldsymbol{s}$ versus the grand null model $\boldsymbol{0}$. The Bayesian false discovery rate (FDR), both local and global (19), is then controlled at or below $1/(1 + \tau)$, contingent on the prior.

Fryer, Arning and Wilson (18) showed that Bayesian hypothesis tests, like the above, inherently control the frequentist familywise error rate (FWER) in the strong sense because they are closed testing procedures (20). The FWER can be quantified, subject to further assumptions. Johnson (21, 22) developed an approach in which the posterior odds of model $\boldsymbol{s}$ versus model $\boldsymbol{0}$ can be approximated as

$$\text{PO}_{\boldsymbol{s}} \approx \left(\mu \sqrt{\frac{h}{n+h}}\right)^{|s|} R_{\boldsymbol{s}}^{n/(n+h)}, \tag{2}$$

where $R_{\boldsymbol{s}}$ represents the ratio of maximized likelihoods under model $\boldsymbol{s}$ versus model $\boldsymbol{0}$, $|s|$ gives the difference in the number of parameters to be estimated under model $\boldsymbol{s}$ versus model $\boldsymbol{0}$, μ represents the prior odds that a regression coefficient is non-zero, n is the sample size, and h represents the precision of the prior on the non-zero regression coefficients. The approximation is based on assumptions including a large sample size, independent observations and the following prior:

$$\boldsymbol{\theta}_{\boldsymbol{s}} \sim \text{Normal}\left(0, h^{-1} \mathcal{I}_{\boldsymbol{s}}^{-1}\right) \tag{3a}$$

$$s_j \sim \text{Bernoulli}\left(\frac{\mu}{1+\mu}\right), \quad j = 1 \ldots v \tag{3b}$$

Here $\boldsymbol{\theta}_{\boldsymbol{s}}$ represents the unconstrained parameters in model $\boldsymbol{s}$ (the β_j for which $s_j = 1$, and any nuisance parameters), and $\mathcal{I}_{\boldsymbol{s}}$ is the per-observation Fisher information matrix for model $\boldsymbol{s}$, evaluated at $\theta_{\boldsymbol{s}} = \boldsymbol{0}$. Fisher's information matrix has been used widely in the definition of reference priors (e.g. 23, 24), and to generate concordance between Bayesian and frequentist point and interval estimates (see Table 1 of

18). Johnson's approach converges on the Bayesian information criterion (BIC), which has been shown to reasonably approximate a wide range of posterior odds when n is large (25, 26). The strength of Johnson's approach is the ability, for a pair of nested models, to interconvert posterior odds and p-values based on the likelihood ratio test (27).

Doublethink (18) extends this approach to the multiple testing setting, in which there is model uncertainty. Using the theory of heavy-tailed random variables (28-31) and asymptotic likelihood theory (e.g. 32), we showed that rejecting all null hypotheses ω_v for which $\text{PO}_{\mathcal{A}_v:\mathcal{O}_v} > \tau$ controls the FWER in the strong sense at or below level

$$\alpha \sim \Pr\left(\chi_1^2 > 2\log\frac{\tau}{v\,\mu\left(\frac{h}{n+h}\right)^{1/2}}\right) \text{ as } n \to \infty. \tag{4}$$

The Bayesian procedure is equivalent to rejecting the null hypothesis ω_v when an asymptotic p-value, adjusted for multiple testing,

$$p^{\star}_{\mathcal{A}_v:\mathcal{O}_v} \sim \Pr\left(\chi_1^2 > 2\log\frac{\text{PO}_{\mathcal{A}_v:\mathcal{O}_v}}{v\,\mu\left(\frac{h}{n+h}\right)^{1/2}}\right) \text{ as } n \to \infty, \tag{5}$$

is smaller than threshold α. In general, the convergence in these asymptotic results is pointwise.

An equivalent interpretation of the results is that the test statistic $2\log\text{PO}_{\mathcal{A}_v:\mathcal{O}_v}/\left(v\,\mu\,\sqrt{h/(n+h)}\right)$, which follows a chi-squared distribution with one degree of freedom when large, represents a model-averaged deviance. Doublethink p-values cannot be trivially rescaled by the prior parameters μ and h because (i) the null distribution of the model-averaged deviance does not depend on them and (ii) the realized value depends on them only through weights. Therefore μ and h influence the power of the test, but not its large-sample theoretical distribution under the null hypothesis. This makes model-averaged hypothesis testing a workable frequentist method by facilitating a prior-agnostic approach to quantifying Bayesian significance thresholds in terms of frequentist FWER, for large samples. For further details, simulations, limitations of the approach such as inflation of the tests due to highly correlated variables, and mitigations such as testing groups of highly correlated variables, see (18).

Methods
In this study, we developed a Monte Carlo Markov Chain algorithm (33) in R (34) and Python (35) that implements the Doublethink approach for thousands of variables, in order to identify direct risk factors for COVID-19 hospitalization in UK Biobank. We followed the COVID-19 Host Genetics Initiative definition of COVID-19 hospitalization, as applied to UK Biobank.

Outcomes. Cases were identified from Public Health England's Second Generation Surveillance System (SGSS), the National Health Service's Hospital Episode Statistics (HES) and the National Health Service's death registry between January and December 2020 as PCR positive for SARS-CoV-2 in SGSS, and hospitalized with International Classification of Diseases, Tenth Revision (ICD-10) diagnosis code U07.1 or U07.2 in HES. Participants not identified as cases were considered controls. We excluded participants that died before 2020, non-England residents determined by assessment centre, and those that withdrew before the analysis. The total number of controls was down-sampled to 200,000 to speed computation. The total number of cases was 1,917.

Variables. We considered data fields approved for UK Biobank project 53100 'Microbiology, disease and genetics', across the categories Population characteristics, Assessment centre, Biological samples, Online follow-up, Additional exposures and Health-related outcomes. We excluded Compound, Date,

Text and Time variables, and variables concerning genetics and sampling processes. For repeated measures, we took the first instance. We excluded factors exceeding 50 levels, except self-reported illnesses, and variables missing in more than 15% of participants. Special values, including negative factor levels, were treated as missing. We imputed missing continuous and integer covariates taking the mean of non-missing values. Missing factor levels were treated as a separate level and excluded. We created binary variables for all levels of every factor observed with frequency above 0.2%. We created a binary variable for every ICD-10 code with frequency above 0.2% recorded before 2020 in HES. Overall, we analysed 184 covariates, binary variables encoding 865 levels across 193 factors, and 863 ICD-10 admission codes, a total of 1,912 variables (Supplementary Table S2).

Model. We fitted the data separately for each outcome via a logistic regression model implemented in R using the glm function, assuming an additive linear predictor with an intercept term. We assumed the prior odds of variable inclusion were $\mu = 0.0053$, independently for the $v = 1,912$ variables, implying a prior expectation of 10 variables in the model. We assumed a unit information prior ($h = 1$) for the regression coefficients (26). We disallowed the inclusion of collinear variables by defining a zero likelihood.

Implementation. In contrast to the Gibbs sampling approach of (36), we implemented a Metropolis-Hasting Markov Chain Monte Carlo (MCMC) sampler over the variable inclusion vector **s**. We ran 100 chains with 25,000 iterations of burn-in and 75,000 iterations of sampling. The average run-time per chain was 35 hours. Chains were initialized using a furthest neighbor algorithm to avoid including correlated variables. For initialization, we clustered variables into 200 groups with the scikit-learn-extra KMedoids algorithm, using rank correlation distance. Each chain was initialized with the medoid of one group, before adding nine more variables iteratively from the next-least correlated variables. Three Metropolis Hastings moves were implemented that respectively added, removed, or swapped pairs of variables with relative proposal probabilities 9:9:2. Variables were swapped preferentially for those with high squared correlation. We simulated regression coefficients directly from conditional Normal distributions by post-processing the MCMC iterations. We calculated posterior odds and parameter estimates by combining chains, computing Monte Carlo standard errors across independent chains.

Grouping variables. We were able to perform valid *post-hoc* variable grouping while controlling the FDR and FWER, which was useful since correlated variables reduce one-another's individual posterior inclusion probabilities. Moreover, tests of null hypotheses involving some but not all members of a cluster of highly correlated variables are liable to inflation (18). We clustered variables in two ways: (i) *A posteriori*, using the scikit-learn OPTICS algorithm (38) with distances defined by their posterior correlation in inclusion probabilities. This grouped the variables with the strongest negative correlations in inclusion probabilities. (ii) *A priori*, using the same algorithm applied to one minus the squared correlation between variables. We computed the posterior odds of including one or more of the variables in each group.

***p*-value calculation.** We used the chi-squared distribution to compute adjusted *p*-values using Equation 5. In the case of orthogonal variables with one degree-of-freedom, this is conservative for $p < 0.02$; see (18). Since the large sample size assumption implies interest in small significance thresholds, we report any *p*-value larger than 0.02 as n.s. (not significant) or '-'. Effectively, this makes Doublethink incompatible with any threshold exceeding $\alpha = 0.02$. In practice, we did not explicitly set a threshold, instead reporting adjusted *p*-values alongside posterior odds.

Literature review. We reviewed the variables included in published analyses of COVID-19 risk factors in UK Biobank using the query "UK Biobank" (Abstract) and "COVID" (Abstract) in

 on 19 September 2023. After excluding Review Articles and Editorial Material, this search returned 203 publications. We analysed a subset of 127 of these papers that quantified the effect of non-genetic risk factors on COVID-19 outcomes; this predominantly excluded papers reporting genetic risk factors, two-sample Mendelian randomization, and COVID-19 as an exposure for other outcomes (Supplementary Table S1). We manually categorized the variables analysed by these 127 papers into groups (Supplementary Table S3). We summarized the frequency with which each category of variable was included in the published analysis or abstract.

Results

We aimed to identify risk factors that directly influenced COVID-19 hospitalization in UK Biobank participants to understand the underlying processes. We used model-averaged hypothesis tests to account for uncertainty in variable selection and deplete for potential confounders. We assumed the relevant variables were measured in UK Biobank. We aimed to limit the impact of collider bias by focusing on exposure variables measured before 2020, and by comparing cases to the rest of the biobank. This compounded the case definition with any selection biases in the sampling process, for example access to testing, which may affect interpretation (9). We focused on risk factors for hospitalization with COVID-19, because there were more cases than critical illness, and less obvious selection bias than infection, since testing was more widely available in hospitals.

Figure 1 Individual variables (circles) and pre-defined groups of variables (vertical lines) with the strongest evidence of direct effects on the risk of COVID-19 hospitalization in UK Biobank. Evidence was quantified simultaneously by $\log_{10}$ posterior odds (left axis) and $-\log_{10}$ adjusted *p*-value (right axis) using Doublethink. Points and lines are coloured for legibility. Variables were ordered horizontally using OPTICS (38). Vertical lines show the boost in significance (if any) from the most significant individual variable per group to the significance of the whole group. Significance was truncated to $\log_{10}$ posterior odds between -6 and 6. Individual variables and groups significant at $\log_{10}$ posterior odds ≥ 1 are labelled (with the most significant variable per group in parentheses). Individual variables are named by UK Biobank field ID or, when prefixed by a letter, ICD-10 code. Refer to Table 1 for full names.

Doublethink facilitates joint Bayesian-frequentist model-averaged hypothesis tests

Figure 1 shows a Manhattan plot displaying the evidence that each of the 1,912 individual variables (circles) directly affected the risk of COVID-19 hospitalization in UK Biobank, averaged over

uncertainty in the effect of all other variables. Points are plotted against both the $\log_{10}$ posterior odds (left side) and the $-\log_{10}$ adjusted p-value from Equation 5 (right side). This interconversion allows a Bayesian or frequentist approach to evaluating the strength of evidence.

Comparison of the two vertical axis scales shows that in the Doublethink model, the model-averaged posterior odds and adjusted p-values are approximately linearly related, for small p-values. Significant variables are identified by applying a threshold to either the posterior odds or the adjusted p-value; this simultaneously controls the FDR (subject to the assumed prior) and the FWER (subject to the asymptotic approximation). For example, a Bayesian threshold of $\tau = 10$ would control the FDR at $1/(1 + \tau) = 0.091$ and the FWER at $\alpha = 10^{-3.3} = 0.00047$. The latter is much smaller than the conventional threshold of 0.05 because of the large sample size.

At a significance threshold of $\tau = 10$ and $\alpha = 10^{-3.3}$, nine variables were identified as individually exposome-wide significant (Table 1). The interpretation is that significant variables, such as the ICD-10 codes `F03 Unspecified dementia` and `J22 Unspecified acute lower respiratory infection`, directly affect risk of COVID-19 hospitalization, even after controlling for the effects of all other measured variables. This differs from the common practice of testing the significance of a variable in the context of a single model that controls for a limited set of other variables. Model averaging is important in biobank-scale data where correlation between variables is pervasive, and no single model has high posterior probability.

For several significant variables, the interpretation that they directly affect risk of COVID-19 would seem too literal, such as `137 Number of treatments/medications taken`, which is based on the recruitment interview, `41214 Carer support indicators : 1 : Yes`, which indicates a hospital record of past carer support, `R29.6 Tendency to fall, not elsewhere classified`, which indicates a history and future risk of falls, and `Z86.4 Personal history of psychoactive substance abuse`, which indicates a hospital record of past alcohol, tobacco or drug use. More plausibly, these variables represent or aggregate one or more (perhaps unmeasured) variables that directly affect risk of COVID-19, like aspects of general health or behaviour. The estimated direct effect of these proxies was to increase the risk of COVID-19 hospitalization in all cases (Table 1). In contrast, significant measures of educational attainment, `6138 Qualifications : 3 : O levels/GCSEs or equivalent`, and `6138 Qualifications : 1 : College or University degree`, had protective direct effects on risk of COVID-19 hospitalization.

The significance of some variables was, at first glance, unexpectedly low, such as the well-established risk factors `31 Sex : 1 : Male` (Posterior probability, $PP = 49.9$; $p^\star = 10^{-2.23}$; where posterior odds $= PP/(1-PP)$) and `34 Year of birth (years)` ($PP = 40.8$; $p^\star = 10^{-2.05}$; Table 1). This is explained by the inclusion in the data of the other very highly correlated variables `31 Sex : 0 : Female`, `21003 Age when attended assessment centre (years)` and `21022 Age at recruitment (years)`. Including variables that are correlated, whether strongly or weakly, often dilutes the significance of individual variables when testing for the existence of a direct effect, over and above all other variables. For age and sex, an obvious solution would be to exclude these correlated variables – it may seem absurd not to have done so. However, correlation is pervasive in biobank-scale data, it may not be obvious which variables to exclude, and the impact of excluding variables on the results is hard to anticipate. An alternative solution is to define groups of correlated variables and test whether one or more members of a group directly affect the outcome. A major strength of Doublethink is it allows arbitrary groups of variables to be tested in this way, while controlling the FDR and FWER.

Testing the significance of groups of variables reveals more signals

Nine groups of variables defined *a priori* were significant at $\tau = 10$ and $\alpha = 10^{-3.3}$; for four groups, none of the member variables were individually significant. In Figure 1, vertical lines illustrate the boost in the significance of groups of variables compared to their most significant member. The groups are numbered for cross-reference with Table 1. Reassuringly, the well-established risk factors age (Group 1; $PP = 100$; $p^\star < 10^{-5.95}$), indices of multiple deprivation (Group 2; $PP = 100$; $p^\star < 10^{-5.95}$), and sex (Group 3; $PP = 100$; $p^\star = 10^{-5.95}$) were significant despite containing no individually significant member variables. In these examples, testing groups of variables recovered signal that was diluted by the inclusion in the data of highly correlated variables.

Another group was significant, despite containing no individually significant variables, demonstrating the ability of combined tests to detect subtle signals. Group 8 ($PP = 99.5$, $p^\star = 10^{-4.71}$) comprised `z50.1 Other physical therapy` ($PP = 79.9$, $p^\star = 10^{-2.89}$) and `z50.7 Occupational therapy and vocational rehabilitation, not elsewhere classified` ($PP = 19.6$, $p^\star > 0.02$). These indicators of rehabilitation might represent or aggregate aspects of convalescence less well captured by the other 1,917 variables analysed. The analysis suggests this history of convalescence directly increased the risk of COVID-19 hospitalization, after controlling for all other measured variables.

Testing groups is useful but defining them *a priori* is not necessarily the most effective method of discovering signals, because the groupings might not be relevant to the outcome under investigation. For example, Group 8 also included variable `z50.5 Speech therapy`, which appeared to contribute nothing to the group's overall significance ($PP = 0.0$, $p^\star > 0.02$). Conversely, failure to group relevant variables together can cause signals to be overlooked, as we see next.

Doublethink allows arbitrary groups to be tested

Figure 2 Individual variables (circles) and post-hoc groups of variables (vertical lines) with the strongest evidence of direct effects on the risk of COVID-19 hospitalization in UK Biobank. The horizontal axis is truncated to show only the most significant variables and groups. Vertical lines show the boost in significance (if any) from individual member variables to the significance of the whole group. See Figure 1 legend for further details.

One of the advantages of the Doublethink approach is it motivates the testing of arbitrary groups of variables without inflating the FWER or FDR through a multiple testing 'fishing expedition'. This is because the thresholds of all possible tests are pre-defined in the closed testing procedure. Therefore we were free to search for the most significant groups of variables. To this end, we grouped variables post-hoc whose posterior inclusion probabilities (*PP*s) were negatively correlated, because this suggests they 'competed' for inclusion in the model.

Figure 2 and Table 2 show that post-hoc grouping revealed signals that were weaker in pre-defined groupings, such as post-hoc Group D (*PP* = 99.5, $p^\star = 10^{-4.75}$), which captured aspects of obesity by combining the individual variables `48 Waist circumference (cm)` (*PP* = 89.0, $p^\star = 10^{-3.22}$), `21001 Body mass index (BMI) (Kg/m2)` (*PP* = 5.5, $p^\star > 0.02$) and `23104 Body mass index (BMI) (Kg/m2)` (*PP* = 5.0, $p^\star > 0.02$). In contrast, prior Group 10 (*PP* = 89.0, $p^\star = 10^{-3.22}$) had only combined `48 Waist circumference (cm)` with the non-significant variables `21002 Weight (Kg)` and `23098 Weight (Kg)` (Table 1).

Some post-hoc groups overlapped the pre-defined groups but dropped non-significant variables that did not contribute to the significance of the group. For example, Group H (*PP* = 99.2, $p^\star = 10^{-4.48}$), contained only the three most significant deprivation scores of the eight members of Group 2. Other groups revealed new connections between variables, such as Group I (*PP* = 95.7, $p^\star = 10^{-3.69}$), which combined the individually non-significant `K59.0 Constipation` (*PP* = 66.1, $p^\star = 10^{-2.55}$) and `N39.0 Urinary tract infection, site not specified` (*PP* = 58.4, $p^\star = 10^{-2.40}$). The former may cause the latter, but it is unclear how a history of these conditions could predispose to COVID-19 hospitalization.

The post-hoc grouping of `41218 History of psychiatric care on admission : 8 : Not applicable` with `I10 Essential (primary) hypertension` was at first glance surprising from the field descriptions (Group G: *PP* = 99.3, $p^\star = 10^{-4.55}$). However, the former variable indicates a history of non-psychiatric hospital care. This suggests it may act, in a manner interchangeable with `I10`, as a proxy for a history of underlying poor physical health. The direct effect of both variables was to increase the risk of COVID-19 hospitalization (Table 2).

The ability to quantify the evidence for groups of variables offers an alternative to approaches such as pre-analysis selection of representative candidate variables among groups of correlated variables. Doublethink permits all and any groups of variables to be tested while controlling the FDR and FWER. This presents new possibilities for identifying significant groups, and the identification of these groups may help with the interpretation of the role in the individual variables in the outcome.

Comparison to the literature on COVID-19 outcomes in UK Biobank
Since early in the COVID-19 pandemic, before the discovery of effective treatments, there were intense research efforts to understand susceptibility to infection, disease and poor outcomes. Many studies focused on large established cohorts like UK Biobank that could rapidly link to data on SARS-CoV-2 testing (37), COVID-19 hospitalization (39) and mortality (40). Since then, many risk factors have been reported, including diabetes (41, 42, 43, 44), asthma (45, 46) and vitamin D (47, 48) as predisposing to worse outcomes. We compared our results to the literature on COVID-19 in UK Biobank to identify any differences to standard approaches and find new insights. At the time of analysis, we identified 127 comparable studies through Web of Science. We manually assigned the most common risk factors in published analyses of COVID-19 outcomes to larger categories for

comparison to the variables and groups listed in Tables 1 and 2, which we assigned to the same list of categories.

Table 3 shows the most common categories of risk factors included in published analyses of COVID-19 outcomes in the 127 UK Biobank studies. Two summaries are shown: the percentage of papers and the percentage of abstracts in which each category of risk factors appeared. Alongside we show the evidence from our analysis, with values of $PP < 50\%$ (corresponding to $p^\star > 10^{-2.20}$) omitted, since the Bayesian interpretation is that this represents evidence against a direct effect of those risk factors. An important caveat is that many studies focused on the total (direct and indirect) effects of candidate risk factors, which our method does not quantify. Therefore, the discovery of a risk factor absent from the literature is more interesting than the non-discovery of (a possibly indirect) effect reported in the literature.

Age, Sex, Obesity, Ethnicity, Socioeconomic status (including deprivation indices) and Smoking were included in 66-90% of published analyses, but mentioned in just 6-20% of abstracts. Our analysis supported direct effects of all these categories with $PP \geq 99.5\%$ and $p^\star \leq 10^{-4.75}$ except ethnicity (Supplementary Table S4, S5). Post-hoc Group J, which combined self-reported ethnicity and country of birth with geographic measures of pollution, had strong support ($PP = 94.1$, $p^\star = 10^{-3.54}$). However, pre-defined Group 11, which combined self-reported ethnicity and country of birth without pollution metrics, was not significant at $\tau = 10$ and $\alpha = 10^{-3.3}$ ($PP = 80.0$, $p^\star = 10^{-2.89}$).

Other reasonably common categories of risk factor for which our analysis found evidence of direct effects included Lung disease, Alcohol intake, General comorbidity, Kidney disease and Educational attainment. Risk factors in these categories featured in 28-46% of published analyses and 4-9% of abstracts. Our analysis supported these categories with $PP \geq 95.7\%$ and $p^\star \leq 10^{-3.69}$.

Many categories of risk factors that appeared commonly in published analyses received no significant support for direct effects in our analysis. Diabetes, Cardiovascular disease and Hypertension were notable for inclusion in 54-63% of published analyses, and 10-12% of abstracts. No variables or groups of variables corresponding to these categories received support for direct effects in our analyses ($PP < 50\%$, $p^\star > 10^{-2.20}$). However, no evidence of a direct effect does not imply no evidence of an effect. These common diseases contribute to a general decline of health, and it is possible that their effects were mediated through pathways better represented by variables or groups we categorised under General comorbidity, such as `137 Number of treatments/medications taken` and Group G (non-psychiatric hospital care/hypertension). Mediation is not the only explanation; the sparsity-favouring prior may have penalized the inclusion of direct effects of several variables in favour of an aggregate variable like `137 Number of treatments/medications taken` that captured the signal more parsimoniously.

Several notable categories of risk factor that we found to have significant direct effects were included infrequently in published analyses of COVID-19 outcomes in UK Biobank. Variables representing Psychiatric disorders, Dementia, Aging (over and above Age) and Infection were included in 9-15% of published analyses, and 2-9% of abstracts, whereas we found strong evidence of direct effects of variables we assigned to these categories ($PP \geq 99.4\%$ and $p^\star \leq 10^{-4.65}$), including `41214 Carer support indicators : 1 : Yes` (which we categorised under Psychiatric disorders), `F03 Unspecified dementia` (Dementia), `R29.6 Tendency to fall, not elsewhere classified` (Aging) and `J22 Unspecified acute lower respiratory infection` (Infection). These variables were significant even after accounting for all others, such as age and number of treatments/medications.

Therefore a model-averaging big data approach that accounts for widespread correlations among variables and uncertainty in variable selection can bring useful perspective on our understanding of well-studied health outcomes like COVID-19 hospitalization in UK Biobank.

Discussion

All exposome-wide significant direct effects we discovered had received some attention among the 127 UK Biobank studies in our literature review. However, we found several strongly significant signals in categories of variable overlooked by 85% of studies or more. Since our analysis investigated the same cohort, we drew on studies from other populations to assess the plausibility of these signals.

Psychiatric disorders were investigated by twelve out of 127 UK Biobank studies (9%), several of which reported an association with elevated risk of hospitalization and mortality (49-60). We found an exposome-wide significant signal that hospitalization risk was directly increased by a history of psychiatric hospital care, specifically among patients with a carer available at their usual place of residence. Supporting this, US studies found psychiatric diagnosis to be a risk factor for COVID-19 diagnosis (61, 62). In a matched case-control analysis of 3.5 million patients, diagnosis with a psychiatric disorder (psychotic disorder, mood disorder, anxiety disorder or insomnia) in the year before the outbreak increased the risk of COVID-19 diagnosis by 65% (61). Following infection, patients with recent diagnosis of mental disorders encompassing ADHD, bipolar disorder, depression and schizophrenia had 1.5-fold higher risk of hospitalization and 1.8-fold higher risk of mortality (62). A meta-analysis of 23 studies reported a two-fold increase in COVID-19 mortality risk in individuals with psychotic disorders (63). Proposed mechanisms include reduced adherence to social distancing, immune-inflammatory dysregulation in psychiatric conditions, and interactions with psychotropic medications (62, 63).

Dementia was investigated by 19/127 UK Biobank studies (15%), reporting increased risk of infection, hospitalization and mortality, with Alzheimer's disease showing the highest risk of COVID-19 diagnosis and mortality (42, 53, 58, 59, 64-78). We found exposome-wide significant evidence that prior hospitalization with unspecified dementia directly increased the subsequent risk of COVID-19 hospitalization. In the US, analyses of electronic health records found that dementia increased the risk of COVID-19 diagnosis two-fold, with the strongest effect (3.2-fold) for vascular dementia (79), and increased the risk of mortality with COVID-19 by 1.3-fold (80). Proposed mechanisms for a causal effect of dementia on COVID-19 outcomes include compromised social distancing, particularly in care home settings, challenges maintaining personal hygiene, physical frailty, dementia-associated inflammation and immune dysregulation, and direct aggravating effects of viral infection on cardiovascular and respiratory brain function. Interestingly, imaging of 785 UK Biobank participants revealed brain tissue damage following SARS-CoV-2 infection (81), and the ACE2 receptor, by which SARS-CoV-2 invades human cells, is reportedly upregulated in the brains of Alzheimer's patients (82).

Whereas age was investigated by 114/127 UK Biobank studies (90%), molecular and physical signs of aging were investigated by just 19/127 studies (15%). Phenotypic age acceleration (which estimates excess aging via blood biomarkers (83)), shorter leukocyte telomere length, physical frailty including falls, and slower walking pace were associated with worse COVID-19 outcomes (49, 50, 66, 72, 84-98). We found exposome-wide significant evidence that a prior hospital diagnosis of a tendency to fall directly increased the risk of COVID-19 hospitalization. Falls have been identified as an atypical presenting symptom of COVID-19 in some patients, and serve as a marker of underlying frailty (99). Frailty, frequently measured through a subjective clinical frailty score (100), has been reported as a risk factor for COVID-19 mortality in multiple countries, with three meta-analyses supporting the conclusion that frailty increases risk even after accounting for age (101-103).

Whereas lung disease, including prior pneumonia, was commonly investigated (58/127 UK Biobank studies; 46%), other markers of infection were investigated infrequently (13/127 studies; 10%). In those studies, elevated immune cell counts, infection-related biomarkers and virus antibody titres were associated with increased COVID-19 infection and severity (59, 66, 70, 76, 89, 104-110). We found exposome-wide significant evidence for an increased risk of COVID-19 hospitalization directly associated with prior hospital episodes of (i) unspecified acute lower respiratory infection and (ii) constipation or urinary tract infection. In other populations, prior pneumonia, lung disease and genetic susceptibility to lung disease have been identified as risk factors for COVID-19 (111-116). It is unclear why prior hospital episodes of constipation or urinary tract infection should increase the risk of subsequent COVID-19 hospitalization (117), although perhaps interestingly, ACE2 is reportedly expressed in kidney, bladder and intestine(118, 119). Pre-existing pathologies in these tissues might be exacerbated directly by the cell tropism of SARS-CoV-2 infection.

We did not find exposome-wide significant evidence for direct effects of some previously reported risk factors, like diabetes (41, 42, 43, 44), asthma (45, 46) and vitamin D (47, 48). This does not rule out indirect effects, but it highlights an important contrast with variables for which we did find significant evidence of direct effects, like the self-reported number of treatments/medications taken, prior hospital diagnosis of acute lower respiratory infection, and a history of medically-relevant psychoactive substance abuse (including alcohol and tobacco). There are several reasons we may not have detected direct risk factors. (i) The exposome-wide approach demands stringent multiple testing, reducing power. (ii) We employed a nominal FWER of $\alpha = 10^{-3.3}$, 100-fold more stringent than the conventional threshold 0.05. Our FWER threshold was interconverted from an FDR of 0.09 using Doublethink (18), but stringent FWER thresholds such as this improve replicability (120) and scaling the FWER with $1/\sqrt{n}$ ensures the consistency of hypothesis testing in large samples (121). (iii) The Doublethink framework incorporates a prior distribution with hyperparameters μ and h that affect inference. (iv) The Bayesian prior penalizes complex models, which may favor aggregate variables (like number of treatments/medications) over a set of individual risk factors (like diabetes and hypertension).

A definitive approach to detecting direct risk factors is unattainable, even for a systematic, agnostic scan, because direct effects are defined only relative to a fixed set of variables. A "direct" effect could be mediated through one or more downstream variables that were not measured. No method is free of subjectivity: we curated variables to uphold quality control, avoid reverse causation and avoid collider bias (e.g. by restricting analysis to pre-2020 exposures), and to fix outcome definitions (restricting attention to 2020 due to time-varying vaccine effects). We restricted analysis to 1,912 variables measured in 85% of more of the cohort. The methods chosen to impute missing values, handle repeat measures and encode factors further shaped our results.

Our work offers a new approach at a time when there are increasing calls for exposome-wide association studies (e.g. 10). Agnostic exposome-wide approaches to discovering new risk factors were rare among published studies of COVID-19 outcomes in UK Biobank (Table S1). A few studies applied machine learning and univariable scans (9, 12, 44, 71, 117-123), but lacked a principled framework to control false positives rates. In contrast, Bayesian model averaging simultaneously controls the Bayesian FDR and the frequentist FWER, at a level that can be quantified using Doublethink (18). It naturally incorporates uncertainty in which variables to include, which in high-correlation biobank settings can strongly influence the evidence of direct effects. In this study it allowed us to test null hypotheses concerning arbitrary groups of variables, which brought multiple advantages by (i) avoiding the need to manually remove correlated variables pre-analysis, (ii) improving power by combining signals and reducing the stringency of significance thresholds, (iii)

conferring flexibility to pursue significant signals *post-hoc* (124), thereby challenging existing notions of fishing for significance, data dredging and *p*-hacking (125).

Despite its strengths, our approach has limitations. We tested only for direct effects, not total effects. This is an important distinction: in many applications, it is necessary to estimate the total effect to understand the likely impact of an intervention on the outcome. The direct effect can differ in magnitude and direction to the total effect, and confusing the two is a pitfall known as the Table 2 fallacy (8). Our Monte Carlo Markov Chain approach was computationally expensive, requiring 3500 CPU hours. Its feasibility depended on asymptotic approximations and a computationally expedient prior (18). These demands limited our ability to model important phenomena like interactions between variables, non-linear effects such as time-since-exposure, or variables with sparse representation like medication use and occupation. Like other methods, Doublethink is subject to *p*-value inflation when testing subsets of highly correlated variables; this can be mitigated by testing groups of correlated variables together (18).

By advancing a more powerful and agnostic approach to identifying direct non-genetic risk factors, our approach has the potential to help advance scientific discovery and bring together the advantages of Bayesian and classical hypothesis testing in biobank-scale settings.

Acknowledgements

This research has been conducted using the UK Biobank Resource under Application Number 53100. We thank all UK Biobank participants, staff and contributors. We wish to thank Naomi Allen, Jeff Chen, David Eyre, Steven Lin, Gil McVean, Tim Peto and Sarah Walker for comments and advice, and the Mathematisches Forschungsinstitut Oberwolfach, organizers and participants of workshop 2308 *Design and Analysis of Infectious Disease Studies*.

Funding

This work was funded by the Robertson Foundation, the Wellcome Trust and the Royal Society (grant no. 101237/Z/13/B). This study was supported by the National Institute for Health Research (NIHR) Health Protection Research Unit in Healthcare Associated Infections and Antimicrobial Resistance (NIHR200915), a partnership between the UK Health Security Agency (UKHSA) and the University of Oxford. The views expressed are those of the author(s) and not necessarily those of the NIHR, UKHSA or the Department of Health and Social Care. The research was supported by the National Institute for Health Research (NIHR) Oxford Biomedical Research Centre (BRC). The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

Supplementary Material

Table S1 Literature review of published papers analysing COVID-19 outcomes in UK Biobank, containing fields from Web of Science and curated with information on the type of analysis, exclusion criteria, and outcomes and exposures included in the abstract and analysis.

Table S2 All UK Biobank fields included in the analysis, annotated by field or ICD-10 code, UK Biobank or ICD-10 description and (if a factor) level.

Table S3 Synonyms and categories of variables used in the interpretation of the literature review of published papers analysing COVID-19 outcomes in UK Biobank.

Table S4 Categories applied to results in Table 1, prior groupings, for comparison to the literature review

Table S5 Categories applied to results in Table 2, post hoc groupings, for comparison to the literature review

Literature cited

1. Sudlow C, et al. (2015) UK Biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS Med* **12**:e1001779.
2. Tan VY, Timpson NJ (2022) The UK Biobank: a shining example of genome-wide association study science with the power to detect the murky complications of real-world epidemiology. *Annu Rev Genomics Hum Genet* **23**:569–589.
3. Fachal L, Dunning AM (2015) From candidate gene studies to GWAS and post-GWAS analyses in breast cancer. *Curr Opin Genet Dev* **30**:32–41.
4. Marigorta UM, et al. (2018) Replicability and prediction: lessons and challenges from GWAS. *Trends Genet* **34**(7):504–517.
5. Duncan LE, Ostacher M, Ballon J (2019) How genome-wide association studies (GWAS) made traditional candidate gene studies obsolete. *Neuropsychopharmacology* **44**(9):1518–1523.
6. Pearl J, Glymour M, Jewell NP (2016) *Causal inference in statistics: a primer*. Wiley, Hoboken, NJ.
7. Vandenbrouck JP (2002) The history of confounding. *Soz Praventivmed* **47**:216–224.
8. Westreich D, Greenland S (2013) The table 2 fallacy: presenting and interpreting confounder and modifier coefficients. *Am J Epidemiol* **177**(4):292–298.
9. Griffith GJ, et al. (2020) Collider bias undermines our understanding of COVID-19 disease risk and severity. *Nat Commun* **11**:5749.
10. Ding E, Wang Y, Liu J, Tang S, Shi X (2022) A review on the application of the exposome paradigm to unveil the environmental determinants of age-related diseases. *Hum Genomics* **16**:1–16.
11. Madakkatel I, Zhou A, McDonnell MD, Hyppönen E (2021) Combining machine learning and conventional statistical approaches for risk factor discovery in a large cohort study. *Sci Rep* **11**:22997.
12. Wan TK, Huang RX, Tulu TW, Liu JD, Vodencarevic A, Wong CW, Chan KHK (2022) Identifying predictors of COVID-19 mortality using machine learning. *Life* **12**(4):547.
13. Zech JR, Badgeley MA, Liu M, Costa AB, Titano JJ, Oermann EK (2018) Variable generalization performance of a deep learning model to detect pneumonia in chest radiographs: a cross-sectional study. *PLoS Med* **15**(11):e1002683.
14. House of Commons Science, Innovation and Technology Committee (2023) The governance of artificial intelligence: interim report. Ninth Report of Session 2022–23. HC 1769. https://committees.parliament.uk/publications/41130/documents/205611/default/
15. Raftery AE (1995) Bayesian model selection in social research. *Sociol Methodol* 111–163.
16. Gelman A (2008) Objections to Bayesian statistics. *Bayesian Anal* **3**(3):445–450.
17. Gelman A, Shalizi CR (2012) Philosophy and the practice of Bayesian statistics. *Br J Math Stat Psychol* **66**:8–38.

18. Fryer HR, Arning N, Wilson DJ (2025) Doublethink: simultaneous Bayesian-frequentist model-averaged hypothesis testing. arXiv:2312.17566.
19. Efron B, Tibshirani R, Storey JD, Tusher V (2001) Empirical Bayes analysis of a microarray experiment. *J Am Stat Assoc* **96**(456):1151–1160.
20. Marcus R, Eric P, Gabriel KR (1976) On closed testing procedures with special reference to ordered analysis of variance. *Biometrika* **63**(3):655–660.
21. Johnson VE (2005) Bayes factors based on test statistics. *J R Stat Soc B* **67**:689–701.
22. Johnson VE (2008) Properties of Bayes factors based on test statistics. *Scand J Stat* **35**:354–368.
23. Zellner A (1986) On assessing prior distributions and Bayesian regression analysis with g-Prior distributions, in *Bayesian Inference and Decision Techniques: Essays in Honor of Bruno de Finetti*, Goel PK, Zellner A, eds. North-Holland, Amsterdam, pp. 233–243.
24. Liang F, Paulo R, Molina G, Clyde MA, Berger JO (2008) Mixtures of g priors for Bayesian variable selection. *J Am Stat Assoc* **103**:410–423.
25. Schwarz G, et al. (1978) Estimating the dimension of a model. *Ann Stat* **6**:461–464.
26. Kass RE, Wasserman L (1995) A reference Bayesian test for nested hypotheses and its relationship to the Schwarz criterion. *J Am Stat Assoc* **90**:928–934.
27. Wilks SS (1938) The large-sample distribution of the likelihood ratio for testing composite hypotheses. *Ann Math Stat* **9**:60–62.
28. Karamata J (1933) Sur un mode de croissance régulière. théorèmes fondamentaux. *Bull Soc Math France* **61**:55–62.
29. Nagaev SV (1965) Some limit theorems for large deviations. *Theory Probab Appl* **10**(2):219–233.
30. Davis RA, Resnick SI (1996) Limit theory for bilinear processes with heavy-tailed noise. *Ann Appl Probab* **6**(4):1191–1210.
31. Wilson DJ (2019) The harmonic mean p-value for combining dependent tests. *Proc Natl Acad Sci USA* **116**(4):1195–1200.
32. Cox DR, Hinkley DV (1974) *Theoretical Statistics*. Chapman and Hall, London.
33. Hastings WK (1970) Monte Carlo sampling methods using Markov chains and their applications. *Biometrika* **57**(1):97–109.
34. R Core Team (2023) *R: a language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/
35. Anaconda Inc (2023). Python. https://www.python.org
36. Hu J, Johnson VE (2009) Bayesian model selection using test statistics. *J R Stat Soc B* **71**:143–158.
37. Armstrong J, Rudkin JK, Allen N, Crook DW, Wilson DJ, Wyllie DH, O'Connell AM (2020) Dynamic linkage of COVID-19 test results between Public Health England's second generation surveillance system and UK Biobank. *Microb Genom* **6**(7):mgen000397.
38. Ankerst M, Breunig MM, Kriegel H-P, Sander J (1999) OPTICS: ordering points to identify the clustering structure. *ACM SIGMOD Rec* 28(2):49–60.
39. UK Biobank (2023) Hospital inpatient data. Version 4.0. https://biobank.ndph.ox.ac.uk/ukb/refer.cgi?id=138483
40. UK Biobank (2023) Mortality data: linkage to death registries. Version 3.0. https://biobank.ctsu.ox.ac.uk/crystal/refer.cgi?id=115559
41. Elliott J, et al. (2021) COVID-19 mortality in the UK Biobank cohort: revisiting and evaluating risk factors. *Eur J Epidemiol* **36**(3):299–309.
42. Atkins JL, et al. (2020) Preexisting comorbidities predicting COVID-19 and mortality in the UK Biobank community cohort. *J Gerontol A Biol Sci Med Sci* **75**(11):2224–2230.

43. Gao M, et al. (2022) Associations between body composition, fat distribution and metabolic consequences of excess adiposity with severe COVID-19 outcomes: observational study and Mendelian randomisation analysis. *Int J Obes* **46**(5):943–950.

44. Wong KCY, Xiang Y, Yin L, So HC (2021) Uncovering clinical risk factors and predicting severe COVID-19 cases using UK Biobank data: machine learning approach. *JMIR Public Health Surveill* **7**(9):e29544.

45. Zhu Z, Hasegawa K, Ma B, Fujiogi M, Camargo CA Jr, Liang L (2020) Association of asthma and its genetic predisposition with the risk of severe COVID-19. *J Allergy Clin Immunol* **146**:327–329.e4.

46. Lodge CJ, Doherty A, Bui DS, Cassim R, Lowe AJ, Agusti A, Dharmage SC (2021) Is asthma associated with COVID-19 infection? A UK Biobank analysis. *ERJ Open Res* **7**:00309–2021.

47. Ma H, Zhou T, Heianza Y, Qi L (2021) Habitual use of vitamin D supplements and risk of coronavirus disease 2019 (COVID-19) infection: a prospective study in UK Biobank. *Am J Clin Nutr* **113**:1275–1281.

48. Li S, Cao Z, Yang H, Zhang Y, Xu F, Wang Y (2021) Metabolic healthy obesity, vitamin D status, and risk of COVID-19. *Aging Dis* **12**:61–71.

49. Lassale C, Gaye B, Hamer M, Gale CR, Batty GD (2020) Ethnic disparities in hospitalisation for COVID-19 in England: the role of socioeconomic factors, mental health, and inflammatory and pro-inflammatory factors in a community-based cohort study. *Brain Behav Immun* **88**:44–49.

50. Batty GD, Deary IJ, Luciano M, Altschul DM, Kivimäki M, et al. (2020) Psychosocial factors and hospitalisations for COVID-19: prospective cohort study based on a community sample. *Brain Behav Immun* **89**:569–578.

51. van der Meer D, Pinzón-Espinosa J, Lin BD, Tijdink JK, Vinkers CH, et al. (2020) Associations between psychiatric disorders, COVID-19 testing probability and COVID-19 testing results: findings from a population-based study. *BJPsych Open* **6**:e87.

52. Yang H, Chen W, Hu Y, Chen Y, Zeng Y, et al. (2020) Pre-pandemic psychiatric disorders and risk of COVID-19: a UK Biobank cohort analysis. *Lancet Healthy Longev* **1**:e69–e79.

53. Wang Y, Yang Y, Ren L, Shao Y, Tao W, et al. (2021) Preexisting mental disorders increase the risk of COVID-19 infection and associated mortality. *Front Public Health* **9**:684112.

54. Batty GD, Gaye B, Gale CR, Hamer M, Lassale C (2022) Explaining ethnic differentials in COVID-19 mortality: a cohort study. *Am J Epidemiol* **191**:275–281.

55. Hassan L, Peek N, Lovell K, Carvalho AF, Solmi M, et al. (2022) Disparities in COVID-19 infection, hospitalisation and death in people with schizophrenia, bipolar disorder, and major depressive disorder: a cohort study of the UK Biobank. *Mol Psychiatry* **27**:1248–1255.

56. Chen W, Zeng Y, Suo C, Yang H, Chen Y, et al. (2022) Genetic predispositions to psychiatric disorders and the risk of COVID-19. *BMC Med* **20**:314.

57. Topless R, Green R, Morgan SL, Robinson P, Merriman T, et al. (2022) Folic acid and methotrexate use and their association with COVID-19 diagnosis and mortality: a case-control analysis from the UK Biobank. *BMJ Open* **12**:e062945.

58. Ma Y, Li S, Yang H, Zhang Y, Li H, et al. (2023) Effect of psychotropics on the risk of COVID-19 in middle-aged and older adults. *Eur Neuropsychopharmacol* **66**:67–77.

59. Fan X, Han J, Zhao E, Fang J, Wang D, et al. (2023) The effects of obesity and metabolic abnormalities on severe COVID-19-related outcomes after vaccination: a population-based study. *Cell Metab* **35**:585–600.e5.

60. Wauye VM, Ho FK, Lyall DM (2023) Psychosocial predictors of COVID-19 infection in UK Biobank (N = 104,201). *J Public Health (Oxf)* **45**:560–568.

61. Taquet M, Luciano S, Geddes JR, Harrison PJ (2021) Bidirectional associations between COVID-19 and psychiatric disorder: retrospective cohort studies of 62,354 COVID-19 cases in the USA. *Lancet Psychiatry* **8**:130–140.
62. Wang Q, Xu R, Volkow ND (2021) Increased risk of COVID-19 infection and mortality in people with mental disorders: analysis from electronic health records in the United States. *World Psychiatry* **20**:124–130.
63. Vai B, Mazza MG, Colli CD, Foiselle M, Allen B, Benedetti F, et al. (2021) Mental disorders and risk of COVID-19-related mortality, hospitalisation, and intensive care unit admission: a systematic review and meta-analysis. *Lancet Psychiatry* **8**:797–812.
64. Aung N, Khanji MY, Munroe PB, Petersen SE (2020) Causal inference for genetic obesity, cardiometabolic profile and COVID-19 susceptibility: a Mendelian randomization study. *Front Genet* **11**:586308.
65. Hilser JR, Han Y, Biswas S, Gukasyan J, Cai Z, et al. (2021) Association of serum HDL-cholesterol and apolipoprotein A1 levels with risk of severe SARS-CoV-2 infection. *J Lipid Res* **62**:100061.
66. Kuo CL, Pilling LC, Atkins JL, Masoli JAH, Delgado J, et al. (2021) Biological aging predicts vulnerability to COVID-19 severity in UK Biobank participants. *J Gerontol A Biol Sci Med Sci* **76**:e133–e141.
67. Topless RK, Phipps-Green A, Leask M, Dalbeth N, Stamp LK, et al. (2021) Gout, rheumatoid arthritis, and the risk of death related to coronavirus disease 2019: an analysis of the UK Biobank. *ACR Open Rheumatol* **3**:333–340.
68. Tahira AC, Verjovski-Almeida S, Ferreira ST (2021) Dementia is an age-independent risk factor for severity and death in COVID-19 inpatients. *Alzheimers Dement* **17**:1818–1831.
69. Julkunen H, Cichoska A, Slagboom PE, Würtz P (2021) Metabolic biomarker profiling for identification of susceptibility to severe pneumonia and COVID-19 in the general population. *eLife* **10**:e63033.
70. Fan X, Liu Z, Poulsen KL, Wu X, Miyata T, et al. (2021) Alcohol consumption is associated with poor prognosis in obese patients with COVID-19: a Mendelian randomization study using UK Biobank. *Nutrients* **13**:1383.
71. Xiang Y, Wong KC, So HC (2021) Exploring drugs and vaccines associated with altered risks and severity of COVID-19: a UK Biobank cohort study of all ATC level-4 drug categories reveals repositioning opportunities. *Pharmaceutics* **13**:1337.
72. Ren J, Pang W, Luo Y, Cheng D, Qiu K, et al. (2022) Impact of allergic rhinitis and asthma on COVID-19 infection, hospitalization, and mortality. *J Allergy Clin Immunol Pract* **10**:124–133.
73. Hu Y, Yang H, Hou C, Chen W, Zhang H, et al. (2022) COVID-19 related outcomes among individuals with neurodegenerative diseases: a cohort analysis in the UK Biobank. *BMC Neurol* **22**:15.
74. Liu Z, Luo Y, Su Y, Wei Z, Li R, et al. (2022) Associations of sleep and circadian phenotypes with COVID-19 susceptibility and hospitalization: an observational cohort study based on the UK Biobank and a two-sample Mendelian randomization study. *Sleep* **45**:zsac005.
75. Topless RK, Gaffo A, Stamp LK, Robinson PC, Dalbeth N, et al. (2022) Gout and the risk of COVID-19 diagnosis and death in the UK Biobank: a population-based study. *Lancet Rheumatol* **4**:e274–e281.
76. Xie J, Feng S, Li X, Gea-Mallorquí E, Prats-Uribe A, et al. (2022) Comparative effectiveness of the BNT162b2 and ChAdOx1 vaccines against COVID-19 in people over 50. *Nat Commun* **13**:1519.
77. Zhou L, Li H, Zhang S, Yang H, Ma Y, et al. (2023) Impact of ultra-processed food intake on the risk of COVID-19: a prospective cohort study. *Eur J Nutr* **62**:275–287.

78. Patel AP, Paranjpe MD, Kathiresan NP, Rivas MA, Khera AV (2020) Race, socioeconomic deprivation, and hospitalization for COVID-19 in English participants of a national biobank. *Int J Equity Health* 19:114.

79. Wang Q, Davis PB, Gurney ME, Xu R (2021) COVID-19 and dementia: analyses of risk, disparity, and outcomes from electronic health records in the US. *Alzheimers Dement* 17:1297–1306.

80. Harrison SL, Fazio-Eynullayeva E, Lane DA, Underhill P, Lip GYH (2020) Comorbidities associated with mortality in 31,461 adults with COVID-19 in the United States: A federated electronic medical record analysis. *PLoS Med* 17:e1003321.

81. Douaud G, Lee S, Alfaro-Almagro F, Arthofer C, Wang C, et al. (2022) SARS-CoV-2 is associated with changes in brain structure in UK Biobank. *Nature* 604:697–707.

82. Ding Q, Shults NV, Gychka SG, Harris BT, Suzuki YJ (2021) Protein expression of angiotensin-converting enzyme 2 (ACE2) is upregulated in brains with Alzheimer's disease. *Int J Mol Sci* 22:1687.

83. Liu Z, Kuo PL, Horvath S, Crimmins E, Ferrucci L, et al. (2018) A new aging measure captures morbidity and mortality risk across diverse subpopulations from NHANES IV: a cohort study. *PLoS Med* 15:e1002718.

84. Atkins JL, Masoli JAH, Delgado J, Pilling LC, Kuo CL, et al. (2020) Preexisting comorbidities predicting COVID-19 and mortality in the UK Biobank community cohort. *J Gerontol A Biol Sci Med Sci* 75:2224–2230.

85. Woolford SJ, D'Angelo S, Curtis EM, Parsons CM, Ward KA, et al. (2020) COVID-19 and associations with frailty and multimorbidity: a prospective analysis of UK Biobank participants. *Aging Clin Exp Res* 32:1897–1905.

86. Toh C, Brody JP (2020) Evaluation of a genetic risk score for severity of COVID-19 using human chromosomal-scale length variation. *Hum Genomics* 14:36.

87. Ho FK, Petermann-Rocha F, Gray SR, Jani BD, Katikireddi SV, et al. (2020) Is older age associated with COVID-19 mortality in the absence of other risk factors? General population cohort study of 470,034 participants. *PLoS One* 15:e0241824.

88. Petermann-Rocha F, Hanlon P, Gray SR, Welsh P, Gill JMR, et al. (2020) Comparison of two different frailty measurements and risk of hospitalisation or death from COVID-19: findings from UK Biobank. *BMC Med* 18:355.

89. Ho FK, Celis-Morales CA, Gray SR, Katikireddi SV, Niedzwiedz CL, et al. (2020) Modifiable and non-modifiable risk factors for COVID-19, and comparison to risk factors for influenza and pneumonia: results from a UK Biobank prospective cohort study. *BMJ Open* 10:e040402.

90. Yates T, Razieh C, Zaccardi F, Rowlands AV, Seidu S, et al. (2021) Obesity, walking pace and risk of severe COVID-19 and mortality: analysis of UK Biobank. *Int J Obes (Lond)* 45:1155–1159.

91. Batty GD, Deary IJ, Gale CR (2021) Pre-pandemic cognitive function and COVID-19 mortality: prospective cohort study. *Eur J Epidemiol* 36:559–564.

92. Veronese N, Smith L, Barbagallo M, Giannelli G, Caruso MG, et al. (2021) Neurological diseases and COVID-19: prospective analyses using the UK Biobank. *Acta Neurol Belg* 121:1295–1303.

93. Wang Q, Codd V, Raisi-Estabragh Z, Musicha C, Bountziouka V, et al. (2021) Shorter leukocyte telomere length is associated with adverse COVID-19 outcomes: A cohort study in UK Biobank. *EBioMedicine* 70:103485.

94. Chudasama YV, Zaccardi F, Gillies CL, Razieh C, Yates T, et al. (2021) Patterns of multimorbidity and risk of severe SARS-CoV-2 infection: an observational study in the U.K. *BMC Infect Dis* 21:908.

95. Schneider CV, Schneider KM, Teumer A, Rudolph KL, Hartmann D, et al. (2022) Association of telomere length with risk of disease and mortality. *JAMA Intern Med* 182:291–300.
96. Reeves J, Kooner JS, Zhang W (2022) Accelerated ageing is associated with increased COVID-19 severity and differences across ethnic groups may exist. *Front Public Health* 10:1034227.
97. Hamrouni M, Roberts MJ, Bishop NC (2023) High grip strength attenuates risk of severe COVID-19 in males but not females with obesity: A short communication of prospective findings from UK Biobank. *Obes Res Clin Pract* 17:82–85.
98. Harris WS, Tintle NL, Sathyanarayanan SP, Westra J (2023) Association between blood N-3 fatty acid levels and the risk of coronavirus disease 2019 in the UK Biobank. *Am J Clin Nutr* 117:357–363.
99. Chen T, Hanna J, Walsh EE, Falsey AR, Laguio-Vila M, Lesho E (2020) Syncope, near syncope, or nonmechanical falls as a presenting feature of COVID-19. *Ann Emerg Med* 76:115.
100. Rockwood K, Song X, MacKnight C, Bergman H, Hogan DB, et al. (2005) A global clinical measure of fitness and frailty in elderly people. *CMAJ* 173:489–495.
101. Zhang XM, Jiao J, Cao J, Huo XP, Zhu C, et al. (2021) Frailty as a predictor of mortality among patients with COVID-19: a systematic review and meta-analysis. *BMC Geriatr* 21:1–11.
102. Pranata R, Henrina J, Lim MA, Lawrensia S, Yonas E, et al. (2021) Clinical frailty scale and mortality in COVID-19: a systematic review and dose-response meta-analysis. *Arch Gerontol Geriatr* 93:104324.
103. Rottler M, Ocskay K, Sipos Z, Görbe A, Virág M, et al. (2022) Clinical Frailty Scale (CFS) indicated frailty is associated with increased in-hospital and 30-day mortality in COVID-19 patients: a systematic review and meta-analysis. *Ann Intensive Care* 12:17.
104. Dite GS, Murphy NM, Allman R (2021) An integrated clinical and genetic model for predicting risk of severe COVID-19: a population-based case-control study. *PLoS One* 16:e0247205.
105. Christensen RAG, Arneja J, St Cyr K, Sturrock SL, Brooks JD (2021) The association of estimated cardiorespiratory fitness with COVID-19 incidence and mortality: A cohort study. *PLoS One* 16(5):e0250508.
106. Dite GS, Murphy NM, Allman R (2021) Development and validation of a clinical and genetic model for predicting risk of severe COVID-19. *Epidemiol Infect* 149:e162.
107. Huang BH, Inan-Eroglu E, Shaban RZ, Hamer M, Britton A, et al. (2022) Alcohol intake and mortality risk of COVID-19, pneumonia, and other infectious diseases: An analysis of 437191 UK Biobank participants. *Prev Med Rep* 26:101751.
108. Willette AA, Willette SA, Wang Q, Pappas C, Klinedinst BS, et al. (2022) Using machine learning to predict COVID-19 infection and severity risk among 4510 aged adults: a UK Biobank cohort study. *Sci Rep* 12(1):7736.
109. Lin Y, Ma B, Yang Y, Chen Y, Huang J, et al. (2022) Impaired kidney function biomarkers and risk of severe COVID-19: Analysis of population-based cohort data. *Mol Genet Genomic Med* 10(11):e2047.
110. Yanik EL, Evanoff BA, Dale AM, Ma Y, Walker-Bone KE (2022) Occupational characteristics associated with SARS-CoV-2 infection in the UK Biobank during August–November 2020: a cohort study. *BMC Public Health* 22(1):1884.
111. Estiri H, Strasser ZH, Klann JG, Naseri P, Wagholikar KB, Murphy SN (2021) Predicting COVID-19 mortality with electronic medical records. *NPJ Digit Med* 4(1):15.

112. Aveyard P, Gao M, Lindson N, Hartmann-Boyce J, Watkinson P, et al. (2021) Association between pre-existing respiratory disease and its treatment, and severe COVID-19: a population cohort study. *Lancet Respir Med* 9(8):909–923.
113. Signes-Costa J, Nunez-Gil IJ, Soriano JB, Arroyo-Espliguero R, Eid CM, et al. (2021) Prevalence and 30-day mortality in hospitalized patients with Covid-19 and prior lung diseases. *Arch Bronconeumol* 57:13–20.
114. Gülsen A, König IR, Jappe U, Drömann D (2021) Effect of comorbid pulmonary disease on the severity of COVID-19: a systematic review and meta-analysis. *Respirology* 26(6):552–565.
115. Wang, Y, Hao Y, Hu M, Wang Y, Yang H (2022) Interstitial lung disease independently associated with higher risk for COVID-19 severity and mortality: A meta-analysis of adjusted effect estimates. *Int Immunopharmacol* 111:109088.
116. Kanai M, Andrews SJ, Cordioli M, Stevens C, Neale BM, et al. (2023) A second update on mapping the human genetic architecture of COVID-19. *Nature* 621(7977):E7–E26.
117. Dabbah MA, Reed AB, Booth ATC, Yassaee A, Despotovic A, et al. (2021) Machine learning approach to dynamic risk modeling of mortality in COVID-19: a UK Biobank study. *Sci Rep* 11(1):16936.
118. Hikmet F, Méar L, Edvinsson Å, Micke P, Uhlén M, Lindskog C (2020) The protein expression profile of ACE2 in human tissues. *Mol Syst Biol* 16(7):e9610.
119. Zou X, Chen K, Zou J, Han P, Hao J, Han Z (2020) Single-cell RNA-seq data analysis on the receptor ACE2 expression reveals the potential risk of different human organs vulnerable to 2019-nCoV infection. *Front Med* 14:185–192.
120. Benjamin DJ, et al. (2018) Redefine statistical significance. *Nat Hum Behav* 2(1):6–10.
121. Wagenmakers E-J, Ly A (2023) History and nature of the Jeffreys–Lindley paradox. *Arch Hist Exact Sci* 77(1):25–72.
122. Tangirala S, Tierney BT, Patel CJ (2023) Prioritization of COVID-19 risk factors in July 2020 and February 2021 in the UK. *Commun Med* 3(1):45.
123. Córdova-Palomera A, Siffel C, DeBoever C, Wong E, Diogo D, Szalma S (2023) Assessing the potential of polygenic scores to strengthen medical risk prediction models of COVID-19. *PLoS One* 18(5):e0285991.
124. Goeman JJ, Solari A (2011) Multiple testing for exploratory research. *Stat Sci* 26(4):584–597.
125. Andrade C (2021) HARKing, cherry-picking, p-hacking, fishing expeditions, and data dredging and mining as questionable research practices. *J Clin Psychiatry* 82(1):25941.

Tables

	Group PP (%)	Group -log₁₀ p*	Variable	PP (%)	-log₁₀ p*	Direct effect when included	Standard error when included
1	**100**	**>5.95**	34 Year of birth (years)	40.8	2.05	-0.03	0.00
			21003 Age when attended assessment centre (years)	30.0	1.82	0.03	0.10
			21022 Age at recruitment (years)	29.2	1.80	0.03	0.10
2	**100**	**>5.95**	26413 Health score (England)	77.7	2.83	0.26	0.03
			26412 Employment score (England)	10.9	-	2.79	0.37
			26410 Index of Multiple Deprivation (England)	9.9	-	0.01	0.00
			189 Townsend deprivation index at recruitment	2.7	-	0.05	0.02
			26414 Education score (England)	0.3	-	0.01	0.00
			26411 Income score (England)	0.3	-	1.70	0.34
			26417 Living environment score (England)	0.0	-	0.00	0.00
			26416 Crime score (England)	0.0	-	0.01	0.04
3	**100**	**5.95**	31 Sex : 0 : Female	50.1	2.24	-0.49	0.11
			31 Sex : 1 : Male	49.9	2.23	0.49	0.11
4	**100**	**5.80**	41214 Carer support indicators : 1 : Yes	**100**	**5.78**	0.56	0.08
			54 UK Biobank assessment centre : 11006 : Stoke	0.0	-	0.24	0.21
5	**99.9**	**5.70**	6138 Qualifications : 3 : O levels/GCSEs or equivalent	**96.7**	**3.82**	-0.29	0.05
			6138 Qualifications : 1 : College or University degree	**96.5**	**3.80**	-0.34	0.06
			6138 Qualifications : 2 : A levels/AS levels or equivalent	0.1	-	-0.29	0.09
-	**99.9**	**5.44**	Z86.4 Personal history of psychoactive substance abuse	**99.9**	**5.44**	0.37	0.06
6	**99.7**	**5.01**	F03 Unspecified dementia	**99.7**	**5.00**	0.94	0.15
			G30.9 Alzheimer's disease, unspecified	0.0	-	0.57	0.23
			F00.9 Dementia in Alzheimer's disease, unspecified	0.0	-	0.59	0.26
7	**99.7**	**4.89**	137 Number of treatments/medications taken	**99.7**	**4.89**	0.06	0.01
			135 Number of self-reported non-cancer illnesses	0.0	-	0.02	0.01
-	**99.6**	**4.82**	J22 Unspecified acute lower respiratory infection	**99.6**	**4.82**	0.51	0.09
8	**99.5**	**4.71**	Z50.1 Other physical therapy	79.9	2.89	0.53	0.09
			Z50.7 Occupational therapy and vocational rehabilitation, not elsewhere classified	19.6	-	0.61	0.11
			Z50.5 Speech therapy	0.0	-	0.34	0.22
-	**99.2**	**4.47**	R29.6 Tendency to fall, not elsewhere classified	**99.2**	**4.47**	0.63	0.11
9	**98.2**	**4.10**	41218 History of psychiatric care on admission : 8 : Not applicable	**98.2**	**4.10**	0.49	0.09
			41214 Carer support indicators : 99 : Not known	0.0	-	0.03	0.19
10	89.0	3.22	48 Waist circumference (cm)	89.0	3.22	0.02	0.00
			21002 Weight (Kg)	0.0	-	0.01	0.00
			23098 Weight (Kg)	0.0	-	0.00	0.00
-	84.1	3.02	J18.1 Lobar pneumonia, unspecified	84.1	3.02	0.49	0.09
11	80.0	2.89	21000 Ethnic background : 1001 : British	70.2	2.64	-0.40	0.08
			1647 Country of birth (UK/elsewhere) : 6 : Elsewhere	9.8	-	0.42	0.09
			1647 Country of birth (UK/elsewhere) : 1 : England	0.0	-	-0.18	0.07
			21000 Ethnic background : 1003 : Any other white background	0.0	-	-0.21	0.14
-	66.1	2.55	K59.0 Constipation	66.1	2.55	0.40	0.08
12	58.4	2.40	N39.0 Urinary tract infection, site not specified	58.4	2.40	0.40	0.08
			B96.2 Escherichia coli [E. coli] as the cause of diseases classified to other chapters	0.0	-	0.32	0.12
13	40.1	2.04	3063 Forced expiratory volume in 1-second (FEV1) (litres)	28.1	1.77	-0.19	0.04
			3062 Forced vital capacity (FVC) (litres)	12.1	-	-0.15	0.03
			3064 Peak expiratory flow (PEF) (litres/min)	0.0	-	0.00	0.00
14	40.1	2.04	2188 Long-standing illness, disability or infirmity : 0 : No	21.6	-	-0.25	0.06
			2188 Long-standing illness, disability or infirmity : 1 : Yes	18.5	-	0.25	0.06
15	38.3	2.00	24017 Nitrogen dioxide air pollution; 2006 (micro-g/m3)	15.5	-	0.01	0.00
			24018 Nitrogen dioxide air pollution; 2007 (micro-g/m3)	15.1	-	0.01	0.00
			24016 Nitrogen dioxide air pollution; 2005 (micro-g/m3)	7.7	-	0.01	0.00
-	31.5	1.86	L97 Ulcer of lower limb, not elsewhere classified	31.5	1.86	0.69	0.14
-	25.5	1.71	N18.9 Chronic renal failure, unspecified	25.5	1.71	0.48	0.10

Table 1 Doublethink allows the interconversion of model-averaged posterior odds and p-values for groups of variables, defined here a priori using variable correlation. The most significant groups are shown, alongside details of constituent variables. The most significant individual, ungrouped, variables are also shown. PP: posterior probability. p: adjusted p-value. Dashes (-) indicate p* > 0.02.*

	Group PP (%)	Group -log₁₀ p*	Variable	PP (%)	-log₁₀ p*	Direct effect when included	Standard error when included
A	**100**	**>5.95**	34 Year of birth (years)	40.8	2.05	-0.03	0.00
			21003 Age when attended assessment centre (years)	30.0	1.82	0.03	0.10
			21022 Age at recruitment (years)	29.2	1.80	0.03	0.10
B	**100**	**5.95**	31 Sex : 0 : Female	50.1	2.24	-0.49	0.11
			31 Sex : 1 : Male	49.9	2.23	0.49	0.11
-	**100**	**5.78**	41214 Carer support indicators : 1 : Yes	**100.0**	**5.78**	0.56	0.08
C	**99.9**	**5.50**	Z86.4 Personal history of psychoactive substance abuse	**99.9**	**5.44**	0.37	0.06
			20116 Smoking status : 0 : Never	0.0	-	-0.17	0.07
-	**99.7**	**5.00**	F03 Unspecified dementia	**99.7**	**5.00**	0.94	0.15
-	**99.7**	**4.89**	137 Number of treatments/medications taken	**99.7**	**4.89**	0.06	0.01
-	**99.6**	**4.82**	J22 Unspecified acute lower respiratory infection	**99.6**	**4.82**	0.51	0.09
D	**99.5**	**4.75**	48 Waist circumference (cm)	89.0	3.22	0.02	0.00
			21001 Body mass index (BMI) (Kg/m2)	5.5	-	0.04	0.01
			23104 Body mass index (BMI) (Kg/m2)	5.0	-	0.04	0.01
E	**99.5**	**4.71**	Z50.1 Other physical therapy	79.9	2.89	0.53	0.09
			Z50.7 Occupational therapy and vocational rehabilitation, not elsewhere classified	19.6	-	0.61	0.11
F	**99.4**	**4.65**	R29.6 Tendency to fall, not elsewhere classified	**99.2**	**4.47**	0.63	0.11
			W19.0 Home	2.4	-	0.59	0.15
G	**99.3**	**4.55**	41218 History of psychiatric care on admission : 8 : Not applicable	**98.2**	**4.10**	0.49	0.09
			I10 Essential (primary) hypertension	5.5	-	0.24	0.06
H	**99.2**	**4.48**	26413 Health score (England)	77.7	2.83	0.26	0.03
			26412 Employment score (England)	10.9	-	2.79	0.37
			26410 Index of Multiple Deprivation (England)	9.9	-	0.01	0.00
			54 UK Biobank assessment centre : 11011 : Bristol	3.9	-	-0.47	0.14
-	**96.7**	**3.82**	6138 Qualifications : 3 : O levels/GCSEs or equivalent	**96.7**	**3.82**	-0.29	0.05
-	**96.5**	**3.80**	6138 Qualifications : 1 : College or University degree	**96.5**	**3.80**	-0.34	0.06
I	**95.7**	**3.69**	K59.0 Constipation	66.1	2.55	0.40	0.08
			N39.0 Urinary tract infection, site not specified	58.4	2.40	0.40	0.08
J	**94.1**	**3.54**	21000 Ethnic background : 1001 : British	70.2	2.64	-0.40	0.08
			24017 Nitrogen dioxide air pollution; 2006 (micro-g/m3)	15.5	-	0.01	0.00
			24018 Nitrogen dioxide air pollution; 2007 (micro-g/m3)	15.1	-	0.01	0.00
			1647 Country of birth (UK/elsewhere) : 6 : Elsewhere	9.8	-	0.42	0.09
-	84.1	3.02	J18.1 Lobar pneumonia, unspecified	84.1	3.02	0.49	0.09
K	40.1	2.04	3063 Forced expiratory volume in 1-second (FEV1) (litres)	28.1	1.77	-0.19	0.04
			3062 Forced vital capacity (FVC) (litres)	12.1	-	-0.15	0.03
L	40.1	2.04	2188 Long-standing illness, disability or infirmity : 0 : No	21.6	-	-0.25	0.06
			2188 Long-standing illness, disability or infirmity : 1 : Yes	18.5	-	0.25	0.06
-	31.5	1.86	L97 Ulcer of lower limb, not elsewhere classified	31.5	1.86	0.69	0.14
-	25.5	1.71	N18.9 Chronic renal failure, unspecified	25.5	1.71	0.48	0.10

Table 2 Doublethink allows arbitrary groups of variables to be assessed for significance while simultaneously controlling the FWER and FDR. Here groups were defined a posteriori *by identifying variables whose PPs were negatively correlated. The most significant groups are shown, alongside details of constituent variables. The most significant individual, ungrouped, variables are also shown. PP: posterior probability. p*: adjusted p-value. Dashes (-) indicate p* > 0.02.*

Category	% Papers	% Abstracts	PP (%)	$-\log_{10} p*$
Age	90	11	100	>5.95
Sex	84	14	100	5.95
Obesity	78	20	99.5	4.75
Ethnicity	78	16	80	2.89
Socioeconomic status	68	13	100	>5.95
Smoking	66	6	99.9	5.5
Diabetes	63	10		
Cardiovascular disease	59	12		
Hypertension	54	11		
Lung disease	46	6	99.6	4.82
Alcohol intake	35	3	99.9	5.5
General comorbidity	31	9	99.7	4.89
Cancer	29	1		
Kidney disease	28	6	95.7	3.69
Educational attainment	28	4	99.9	5.7
Asthma	26	3		
Physical activity	24	6		
Neurological disease	21	3		
Liver disease	19	2		
Inflammatory disease	17	2		
Geographic region	17	0		
Aging	15	9	99.4	4.65
Dementia	15	2	99.7	5.01
Employment	13	3		
Immune disease	12	2		
Diet	11	6		
Depression	11	4		
Infection	10	3	99.6	4.82
Arthritis	10	3		
Other	9	2		
Sleep disturbance	9	6		
Psychiatric disorders	9	5	100	5.8
Mental health	8	4		
Vitamin D	8	4		
Lipid disorders	7	2		
Pollution	5	2		
Covid-19 related	4	3		
Vaccination	4	3		
Allergy	4	1		
Haematological disease	3	1		
Lifestyle	3	2		
Gastrointestinal disease	3	2		
Sex hormones	2	2		
Periodontal disease	2	2		

Table 3 Comparison of risk factors for COVID-19 outcomes in previous UK Biobank studies versus this study. The number of papers, out of 127, are shown. Categories were assigned manually from a literature review, and from Tables 1 and 2. When there were multiple matches in Tables 1 and 2, the maximum significance is given. PP: *posterior probability (only values above 50% are shown).* p*: *adjusted* p-value (only values below $10^{-2.2}$ are shown).