

1 Closing the gap towards a successful referral system. A
2 case study of a tertiary teaching and referral hospital,
3 Kenya: pre-posttest study design
4

5 Authors: Maxwell Philip Omondi^{1*}

6
7
8 ¹ Orthopedic Unit, Department of Surgery, University of Nairobi, Kenya
9

10 *Corresponding author

11 Email address: maxwellomondi@gmail.com
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

Abstract

Inappropriate utilization of higher-level health facilities and ineffective management of the referral processes in resource-limited settings is increasingly becoming a concern in health care management in developing countries. This is characterized by self-referrals and frequent bypassing of nearest health facilities. On 1st July 2021, Kenyatta National Hospital (KNH) did enforce the national referral guidelines that required patients have a formal referral letter to reduce the number of self-referrals, decongest KNH and allow KNH to function as a referral facility as envisioned by Kenya Health Sector Referral Implementation Guidelines of 2014, Kenya 2010 constitution and KNH legal statute of 1987. The purpose of this study was to analyse the effect of enforcement of referral guidelines on facility referrals to KNH. This was a pre-posttest study design. The study was conducted amongst the orthopedic facility referrals in 2021 with 222 and 246 before and after enforcement of referral guidelines respectively. Data collection was done through data abstraction. Data was analyzed using frequency distribution, pearson chi-square test and logistic regression. Nairobi County and its environs constituted over four-fifth of all facility referrals to KNH. Over two-thirds of the facility referrals to KNH were from government facilities. There was significant reduction in health facilities tiers 2 and 3 referring patients directly to KNH after enforcement of referral guidelines ($p=0.002$). About 43 health facilities ceased referring patients to KNH with over two-thirds of these health facilities being private facilities. The major facility and patient factors that were associated with facility referrals to KNH were human resource capacity and availability and patient's preference. In conclusion, enforcement of the referral guidelines significantly reduced the lower tiers health facilities referring to KNH. We recommend

52 having written standard operating procedures on referrals based on the national referral
53 guidelines with continued enforcement of the same to sustain the gains made.

54

55 Key words: referral guidelines, health facility referrals, Kenyatta National Hospital

56

57

58

59

Background

60 Tertiary hospitals in resource-limited countries treat patients referred but in most cases
61 are the first level of care for the vast majority of patients (1). One of the challenges in
62 health care delivery in resource-limited settings is inappropriate utilization of tertiary
63 health facilities that results in patients' congestion in referral hospitals with simple medical
64 conditions that can be effectively managed at the lower peripheral health facilities. The
65 majority of these patients are self-referred, bypassing lower-level health facilities in the
66 process (2-5).

67

68 A study done in Lusaka demonstrated how University Teaching and Referral Hospital is
69 congested due to bypassing of lower peripheral health facilities and as a result the tertiary
70 facility effectively functions as a primary health care facility. The urban phenomenon of
71 widespread self-referral is associated with low rates of formal referral from peripheral
72 health facilities (4, 6). The net result is simple medical conditions end up being managed
73 in high-cost referral health facilities leading to overcrowding, long waiting times, and
74 scarce staff time consumed by lesser medical conditions at the expense of complex
75 medical conditions. It is therefore imperative that attempts need to be made to ensure
76 patients make use of lower health facilities by creating disincentives for patients by-
77 passing these health facilities (1, 7). The bypassing of lower level health facilities also

78 cripples the primary health care system and this effectively ensures that primary health
79 care facilities remain underused and inefficient (8). Taken together, this jeopardizes the
80 appropriate delivery of primary, secondary, and tertiary health care

81
82 In 2008, an assessment done in South African Public health sector revealed that patients
83 were accessing the health system at inappropriate levels and bypassing primary health
84 care to attend to regional and tertiary hospitals as their initial visit leading to overcrowding
85 and unnecessary costs to the referral facilities (9). In cross-sectional study was done to
86 review the self-referrals to a District-Regional Hospital in KwaZulu Natal, South Africa
87 showed 36% were self-referrals. However, majority (64%) were appropriately referred
88 with written referral letters (9).

89 A study on referral system in Nigeria showed that 92.9% reported to tertiary hospital
90 directly without referral while 7.1% were referred. The result of this is overcrowding of the
91 tertiary facilities with conditions that can be managed at lower level health facilities (7).

92
93 In India health care system is characterised by overcrowding, lack of specialist doctors
94 and effective referral system. Despite having a referral guideline that details referral of
95 patients from lower levels to sub-district, district and then to tertiary hospitals, the actual
96 practices are that patients go to any level of health care system without any referral (10).
97 This creates burden on the secondary and tertiary level hospitals. In Taiwan where we
98 have a free-access systems, bypass of primary care and seeking secondary care through
99 self-referral is rampant despite Taiwan's government taking various initiatives to mitigate
100 bypass (11).

101

102 Kenya is a low-income middle-income country. Kenya's health sector is one of the 14
103 functions devolved to the 47 county governments as provided for in the Fourth Schedule
104 of the 2010 constitution (12). There are six different levels or tiers of health care in Kenya.
105 Tiers 1 -5 are managed by the county government and tier 6 by the national government.
106 In this system patients are required to move from one tier to the next using a formal
107 referral letter. Kenya has a pluralistic health sector consisting of private and non-
108 governmental service providers alongside the government-run health facilities that is
109 organized into six (6) tiers (Figure 1).

110

111

112 Figure 1: Six tiers of health care service delivery in Kenya (13)

113

114 Tiers 4 are Sub-County Hospitals, Nursing Homes and also include medium-sized private
115 hospitals. Tier 5 are County Teaching and Referral Hospitals, large private as well as
116 faith-based hospitals. Tier 6 are National Teaching and Referral Hospitals. In Kenya there
117 are four teaching and referral hospitals: a) Mathari Hospital, b) Kenyatta National
118 Hospital, c) Moi Teaching and Referral Hospital and d) National Spinal Injury Hospital.
119 These tier 6 offer range of services similar to Tier V but additionally they offer specialised
120 treatments in addition to having advanced infrastructure and are not only accessed by
121 Kenyans but so serve East Africa and Central Africa.

122

123 Kenyatta National Hospital (KNH) was established as a National Referral and Teaching
124 Hospital, to provide training and medical research. KNH was established in 1901 and
125 became a State Corporation in 1987 and sits at the peak of the health referral system in

126 Kenya (14). According to the KNH Board order of 1987 contained in the Legal Notice No.
127 109, the functions of KNH were spelled out as a) to receive patients on referral for
128 specialized health care; b) to provide facilities for medical education for the University of
129 Nairobi and other health allied courses; c) to contribute to national health planning (14).
130 This understanding has been reinforced by the Kenya Health Sector Referral
131 Implementation Guidelines, 2014, and the Constitution of Kenya 2010 which tasks KNH
132 with the responsibility for health policy formulation (12, 15).

133
134 Orthopaedic wards in KNH have consistently recorded the highest bed occupancy
135 percent for the last couple of years. In 2018, 2019 and 2020 it recorded bed occupancy
136 percent of 142.2%, 138.2% and 116.5% respectively against the KNH bed occupancy
137 percent of 106.2%, 113.4% and 91.5% (16). This coupled with low nurse-patient ratio of
138 1:10 compromises not only the quality of nursing care given to patients but also the ability
139 of KNH to effectively perform its statutory obligations. Various studies have demonstrated
140 that an appropriate nurse-patient ratio is associated with the reduction in medical errors,
141 decubitus ulcers, hospital-acquired infections, long duration of hospital stay, high
142 readmission rates, and compromised patient well-being and safety due to the nurse
143 burnout (17-19).

144
145 On 1st July 2021, KNH management made a decision to enforce the Kenya Health Sector
146 Referral Guidelines 2014 that places KNH at the tip of the health sector referral system.
147 This meant that patients will be seen based on referral letters from other health facilities
148 to reduce the number of walk-in patients who would have otherwise been appropriately

149 seen at the peripheral health facilities. This would then allow KNH to focus on the
150 management of complex medical conditions and allow uncomplicated medical conditions
151 to be managed at lower-level health facilities. The purpose of this study was to analyse
152 the effect of enforcement of referral guidelines on facility referrals to KNH. The findings
153 of this study will help formulate policy and guidelines on orthopaedic and trauma essential
154 care at the peripheral and tertiary health facilities in Nairobi Metropolitan Area and the
155 country at large. It will also provide evidence-based information that will contribute to the
156 review of Kenya Health Sector Referral Implementation Guidelines of 2014.

157

158 **Methods**

159 **Study design**

160 This was a pre-posttest study design. The national referral guideline was enforced on 1st
161 July, 2021. The pretest covered 5 months before enforcement of referral guidelines
162 (February 1, to June 30, 2021) while post-test covered 5 months after enforcement of the
163 referral guidelines (August 1, to December 31, 2021). The variables compared were
164 county, type of health facility (government or private) and health facility tier before and
165 after enforcement of the referral guidelines. Enforcement of referral guidelines required
166 that the referring health facility consults with KNH referral Office for concurrence before
167 patients are referred to KNH and that patients should come with written official referral
168 letters. This was to ensure only patients who require specialized orthopaedic and trauma
169 care not available at the peripheral health facilities get admitted to KNH.

170 **Study area**

171 KNH is the largest teaching and referral hospital in East and Central Africa. KNH
172 Orthopaedic Wards were the study area. KNH is based in Upperhill, Nairobi, the capital
173 city of Kenya. It is located along Hospital Road, about 5km from the city centre. KNH has
174 a bed capacity of 1,800, 6,000+ staff members, 50 wards, 22 out-patient clinics, 24
175 theaters (16 specialized) and Accident & Emergency Department (14). Of the 1800 bed
176 capacity, 96 beds are allocated to orthopaedic wards. KNH is a 10-floor storied building
177 complex and the Orthopaedic wards are located on the 6th floor but we also have
178 orthopaedic admissions in private wings on 9th and 10th floor. Orthopaedic patients with
179 other co-morbidities also get admitted to other wards in KNH.

180 **Study duration**

181 The study duration was from 1st February 2021 to 31st December 2021. The referral
182 guidelines were enforced from 1st July 2021. Data abstraction was conducted from 1st
183 January to 31st March, 2022.

184

185 **Study population**

186 Orthopaedic inpatient caseload before and after enforcement of referral guidelines.

187

188 **Eligibility criteria**

189 **Inclusion criteria**

190 All orthopaedic and trauma facility referrals to KNH during the study period.

191

192 **Sample size calculation**

193 Sample size was calculated using an adjusted Casagrande formula for calculating sample
194 sizes that compare two binomial distributions (20).

195 A sample size of 468 facility referrals were enrolled during the study period with 220 and
196 248 facility referrals before and after the enforcement of national referral guidelines.

197 **Recruitment and sampling procedures**

198 Three (3) research assistants (RAs) were recruited to collect and abstract patient data
199 from patient files. The RAs were health care workers with a diploma in Orthopedic
200 Trauma and with some experience in research data collection. The Principal Investigator
201 (PI) was the research coordinator for the data collection. The orthopedic and trauma
202 admissions with facility referrals were identified from the a) admission desk of Health
203 Information System at KNH Accident and Emergency Unit (A&E) b) KNH Orthopedic
204 Outpatient clinic records (OC) c) KNH Corporate Outpatient Care (COC). They were then
205 recorded in a logbook. This logbook served as a master register for all facility referred
206 patients admitted and therefore the sampling frame for the study. All facility admissions
207 were logged into the logbook from the admission books stationed in these three (3)
208 services points. Proportional Population to Size (PPS) was then used to decide on the
209 numbers to be sampled per month from each of these three services points so that the
210 sample size would be a representative of the admissions by month from each of these
211 three orthopedic admissions entry points (Table 1).

212

213

214 Table 1: Orthopaedic and trauma admissions to KNH stratified by point of admission,
215 2021

	Month of the year, 2021`	Point of admission			
		A&E	Clinic	COC	Total
Before	February	54	3	3	60
	March	40	1	3	44
	April	37	0	3	40
	May	33	2	2	37
	June	39	1	1	41
	Total	203	7	12	222
After	August	42	2	2	46
	September	52	3	4	59
	October	55	2	4	59
	November	20	0	12	32
	December	43	2	3	48
	Total	212	9	25	246

216
217 The three (3) RAs were reporting to and working under the direction of the PI. The RAs
218 were trained for two (2) days by the PI on the research protocol, data collection tools,
219 data collection procedures and that included pilot testing of the data collection tools as
220 well before the actual data abstraction.

221
222 A written Informed consent was obtained from KNH Medical Research Department to
223 have access to the patient's health records in the Health Information Office (Room 19).

224 **Variables**

- 225 a) Name of health facility County c) Facility Type (Government/Private) d) Health
226 Facility Level/Tier e) Admission date (dd/mm/yyyy)

228 **Data collection procedures**

229 Data collection was done through a data abstraction form from the patient files.

230 **Data Abstraction form** – The three (3) RAs were trained on the data abstraction using
231 a data abstraction form as per the research protocol. The PI reviewed all the filled-in
232 abstraction forms for completeness and accuracy daily during the entire data collection
233 period and providing regular feedback to the RAs on a timely manner to ensure data
234 quality and compliance to the study protocol. All the completed and verified data
235 abstraction forms were then collected and filed by the PI at the end of every week under
236 a lockable cabinet.

237

238 **Ethical considerations**

239 The study protocol was presented to UoN/KNH Ethics and Research Committee and was
240 granted ethical approval (ERC No: P852/10/2021). Administrative approval was also
241 granted by KNH Medical Research Department and KNH Orthopaedics Department.

242

243 **Data management, analysis, and presentation plan**

244 Data abstraction tool was designed to collect quantitative and qualitative data. For
245 anonymity and confidentiality purposes the data abstraction tool were marked only with
246 the participant study numbers and no names were used. However, the research
247 assistants and the PI had access to the patients charts and therefore could identify
248 individual patients during the data collection period. The data were entered into a
249 password-protected Redcap database kept by the KNH Medical Research Department.
250 The data were analyzed using SPSS version 21.0. Descriptive statistics such as

251 frequencies while inferential statistics using Pearson’s chi-squared tests, logistic
252 regression were used. The calculations were done at a 95% level of confidence.

253 **Findings**

254 **Basic Profile of the sample population**

255 A total of 468 charts were abstracted of which 220 (47%) were before and 248 (53%)
256 were after the enforcement of the referral guidelines.

257
258 Nairobi County constituted more than half of facility referrals to KNH. This rose to over
259 four – fifths of facility referrals with inclusion of the neighbouring counties namely Kajiado,
260 Kiambu and Machakos. Almost all the facility referrals were from within Kenya (Table 2).

261
262 About three-fifths of the facility referrals to KNH were government health facilities with
263 private health facilities comprising about two-fifths. About half of the facility referrals are
264 from Tiers 2 to 4 and with the remainder of facility referrals to KNH from Tier 5 and 6
265 (Table 2).

266
267

268
269

270
271

272
273

274

275

276 Table 2: Basic profile of the sample population, 2021.

Variable	Category	Frequency (n=222)	Percent	Cumulative percent	Frequency (n=246)	Percent	Cumulative percent
			Before			After	
County	Nairobi	133	59.9%	59.9%	136	55.3%	55.2%
	Kiambu	31	14.0%	73.9%	31	12.6%	67.8%
	Kajiado	17	7.7%	81.6%	26	10.5%	78.3%
	Machakos	4	1.8%	83.4%	4	1.6%	79.9%
	Nakuru	4	1.8%	85.2%	2	0.8%	80.7%
	Others	6	2.7%	87.9%	15	6.1%	86.8%
	Central						
	Others	22	9.9%	98.3%	22	8.9%	95.7%
	Eastern						
	Others*	5	2.2%	100.0%	10	4.0%	99.7%
Burundi	0	0.0%	100.0%	1	0.4%	100.1%	
Type of Health Facility	Government	140	63.1%	63.1%	146	59.3%	59.3%
	Private	82	36.9%	100.0%	100	40.7%	100.0%
Health Facility Tier	2	14	6.3%	6.3%	2	0.8%	0.8%
	3	23	10.4%	16.7%	15	6.1%	6.9%
	4	74	33.3%	50.0%	106	43.1%	50.0%
	5	102	45.9%	95.9%	109	44.3%	94.3%
	6	9	4.1%	100.0%	14	5.7%	100.0%

277 Others* include facility referrals from former Coast, North Eastern, Nyanza, Western and
278 Rift Valley regions of Kenya

279

280

281

282

283

284 Figure 2: The major health facilities referring to KNH before and after enforcement of
285 referral guidelines, 2021

286

287 Mama Lucy Kibaki Hospital, Mbagathi District Hospital and Thika Level 5 Hospital were
288 the major health facilities referring patients to KNH before and after enforcement of the
289 referral guidelines (Figure 2).

290 There were increases in the health facility referrals to KNH after the enforcement of the
 291 referral guidelines. However, these increases were not statistically significant differences
 292 (Table 3).

293 Table 3: Bivariable analysis of the type of health facilities referrals referring to KNH
 294 before and after enforcement of referral guidelines, 2021

		Before (n=222)	After (n=246)	p-value
Type of Health Facility	Government Facility referrals	140 (49.0%)	146 (51.0%)	p= 0.411
	Private Facility referrals	82 (45.1%)	100 (54.9%)	

295 Legend: Pearson χ^2 statistics test was used to test for statistical significance at 95% CI.

296
 297 There were statistically significant differences in health facilities referrals to KNH before
 298 and after the enforcement of the referral guidelines when stratified by the health facility
 299 tier (Table 4). There was significant reduction in Tiers II and III referring patients directly
 300 to KNH and an increase in the facility referrals to KNH from tiers IV, V and VI (Table 4).
 301 There were less tier II and III health facilities referring patients to KNH after enforcement
 302 of the referral guidelines. The facility referrals from Tier II were 90.8% less likely as
 303 compared to Tier III after enforcement of the referral guidelines (p=0.006) (Table 4).

304
 305 Table 4: Multivariable analysis of the health facilities referrals, stratified by health facility
 306 tier, referring to KNH before and after enforcement of referral guidelines, 2021

		Before (n=222)	After (n=246)	Chi-square; p-value	Logistic regression
Health Facility Tier	II	14 (87.5%)	2 (12.5%)	16.505; p=0.002	1.0
	III	23 (60.5%)	15 (39.5%)		0.092 (p=0.006)
	IV	74 (41.1%)	106 (58.9%)		0.419 (p=0.108)
	V	102 (48.3%)	109 (51.7%)		0.921 (p=0.856)
	VI	9 (39.1%)	14 (60.9%)		0.687 (p=0.301)

307 Legend: Pearson χ^2 statistics test was used to test for statistical significance at 95% CI
 308 and Odds Ratio was used to test the strength of association. 1.0 refers to reference group.

309
 310 While the study showed about 43 health facilities ceased referring patients to KNH with
 311 over two-thirds of these health facilities being private facilities (Table 5).

312 The study revealed that lower tiers health facilities namely tier II-IV ceased referring
 313 patients to KNH and this comprised over four-fifths of all health facilities that stopped
 314 referring to KNH after enforcement of the referral guidelines (Table 5).

315
 316 Table 5: Frequency distribution of type of health facilities that stopped referring, stratified
 317 by Type of Health Facility and Facility Tier, after enforcement of the referral guidelines,
 318 2021

	Categories	Frequency (n=43)	Percent	Cumulative percent
Type of Health Facility	Government	14	32.6%	32.6%
	Private	29	67.4%	100.0%
Facility Tier	II	7	16.3%	16.3%
	III	12	27.9%	44.2%
	IV	16	37.2%	81.4%
	V	6	14.0%	95.4%
	VI	2	4.7%	100.1%

319
 320 The major facility and patient factors that were associated with facility referrals to KNH
 321 were human resource capacity and availability, health facility infrastructure, Orthopaedic
 322 equipment's and implants availability, patient's preference, financial considerations
 323 (Table 6). It's worth noting that across both private and government health facilities the
 324 main factors for referral were inadequate human resource capacity and availability
 325 followed by availability of medical supplies and equipments and financial considerations
 326 (Table 6). Amongst government health facility referrals, whereas financial considerations
 327 was among the common factors for facility referrals to KNH, the patients actually were
 328 opting to be referred to KNH as opposed to a private health facility due to cost "...the

329 *patient needed a total knee replacement which could not be done in Nyahururu and they*
330 *did not have enough funds to take him to a private hospital so they preferred to KNH...”*

331 *‘... the private hospital was too expensive so they opted for KNH...’*

332 Patients’ preference was shown to be a significant factor associated with private health
333 facility referrals to KNH as opposed to government health facility referrals to KNH (Table
334 6).

335 Table 6: Table showing the frequency distribution of the major facility and patient related
336 factors to the orthopaedic and trauma admissions at the KNH

Factors	Private health facilities referrals (n=182)	Government health facilities referrals (n=286)
Human resource capacity and availability	105 (57.7%)	178 (62.2%)
Infrastructure	16 (8.8%)	25 (8.7%)
Medical supplies and equipment	47 (25.8%)	74 (25.9%)
Finance	42 (23.1%)	79 (27.6%)
Patient preference	72 (39.6%)	15 (5.2%)

337 Legend: This was a cumulative tally of the facility and patient related factors to the
338 orthopaedic and trauma admissions at the KNH as a proportion of the sample size (n).

339

340 **Discussions**

341 While orthopaedic facility referrals were spread across the country, over four – fifths of
342 these facility referrals were from Nairobi County and its environs. This is in tandem with
343 studies done at tertiary teaching and referral health facilities in Rwanda, Tanzania, Malawi
344 and Nigeria that revealed significant majority of the facility referrals are from within the
345 region where the health facilities are co-located (7, 21-24).

346 Most of the facility referrals to KNH are from government health facilities with private
347 health facilities comprising about one-third. This compares favourably with a study done

348 in Rwanda that showed approximately 85.7% of the facility referrals were from public
349 hospitals while 10.7% of the referrals were from private health facilities (25). Similarly a
350 prospective study done at Muhimbili National Hospital, Tanzania to examine the medical
351 referral pattern of patients in 2004, revealed that about 22.4% of admissions were from
352 public health facilities while only 4.2% were from private health facilities (26).

353 Mama Lucy Kibaki Hospital, Mbagathi District Hospital and Thika Level 5 Hospital were
354 the major health facilities referring patients to KNH before and after enforcement of the
355 referral guidelines and they all happen to be government health facilities – which comprise
356 majority of health facility referrals to KNH as compared to private health facility referrals.
357 This is probably because of inadequate human resource, inadequate infrastructure like
358 bed capacity, theatre space as well as lack of orthopaedic equipments and implants.
359 However, this contrasts with a study on referrals to a tertiary health facility in Nigeria that
360 showed over half of the admissions were from private health facilities (7). In addition, St
361 Peters Orthopaedic and Surgical Speciality Centre, St Francis Community Hospital, Arthi
362 River Shalom Community Hospital and St Mary’s Mission Hospital were the most
363 frequently referring private health facilities to KNH for orthopaedic admissions in 2021.
364 These were probably due to inadequate human capacity, patients’ preference and
365 financial considerations.

366

367 There were increases in the number of health facilities referring patients to KNH after the
368 enforcement of the referral guidelines. This is in tandem with a study done in Republic of
369 Honduras that showed the referral rate was observed to be higher when institutional
370 managers emphasized the importance of the referral system (4). The enforcement of the

371 national referral guidelines meant more awareness on the referral requirements and
372 importance of referrals of patients to the next level of care and this might have resulted in
373 increase in the number of health facilities referring patients.

374 The enforcement of the national referral guidelines was meant to streamline the referral
375 process from the peripheral health facilities to KNH and in the process allow KNH to
376 manage complex orthopaedic cases that cannot be handled at the lower-level health
377 facilities. All potential referring health facilities were required to refer patients to the next
378 tier of health facility instead of referring directly to KNH and the higher tier health facilities
379 were expected to seek concurrence before referring patients to KNH and that should be
380 accompanied with an official referring letter. Due to the enforcement of these referral
381 requirements, there was significant reduction in Tiers 2 and 3 referring patients directly to
382 KNH and an increase in the patients being referred to KNH from tiers 4, 5 and 6.

383
384 The study showed about 43 health facilities ceased referring patients to directly to KNH
385 with over two-thirds of these health facilities being private facilities. This is in tandem with
386 a study on Geographic accessibility to public and private health facilities done in Kenya
387 in 2021 that revealed the private health facilities' distribution was skewed toward the
388 urban counties (27). The implication of this is that a good number of private health facilities
389 were no longer referred patients directly to KNH.

390 Lower tiers health facilities namely tier 2 and 3 largely reduced patient referrals to KNH
391 and this together with tier 4 comprised over four-fifths of all health facilities that stopped
392 referring to KNH after enforcement of the referral guidelines. This is because these lower-
393 tier facilities were now required to refer patients to the nearest Tier 4 and 5 health facilities

394 as their first referral point. This means that the enforcement of the national referral
395 guidelines did reduce referrals from lower tier health facilities and probably were referring
396 them to the next tier facility. This may explain why the numbers of higher tiers referring
397 patients to KNH increased after the enforcement of the referral guidelines.

398 Human resource capacity remains a major factor in health facility referrals both for private
399 and government health facilities. Inadequate human resource does also comprise the
400 quality of care provided at the health facilities. This is in tandem with a study finding in
401 Kenya that revealed Human resources for health was inadequately financed with
402 insufficient number of health workers and there was maldistribution of staff in favor of
403 higher-level facilities and in urban cities and this resulted in unnecessary referrals to
404 higher level consequently compromised quality of primary healthcare (28-30). A number
405 of studies in Brazil, Japan, United States of America and in Low- and Middle-income
406 countries have also demonstrated inadequate human resource capacity and availability
407 as the main reasons for patient referrals (31-35). However, this contradicts a study done
408 in China on types of health care facilities and the quality of primary care revealed that
409 Community Health Centers offer quality health care services than the secondary and
410 tertiary health care hospitals (36). This was largely due to the fact that it was Chinese
411 deliberate government policy to support and improve the capacity of the of the Community
412 Health Centers.

413
414 Private health facility referrals to KNH were to significant extent dictated by patients'
415 preference. This could be due to perceived good quality of care but also due to perceived
416 low cost of medical care in KNH by virtue of it being a government national referral

417 hospital. Studies done in California, USA and Islamic Republic of Iran, Cape Town, South
418 Africa did show that the perception of good quality of care does influence choice of health
419 facility (3, 37-39).

420
421 The study had a few limitations. First, the effect of COVID 19 pandemic on facility referrals
422 of cases from peripheral health facilities and walk-in patients. This was mitigated by
423 ensuring the data collection period covered the covid period where intercounty movement
424 restrictions were lifted. secondly, this study is retrospective and a quasi-experimental
425 study design and hence weaker in determining causality. Despite these limitations, given
426 the paucity of published literature in this study topic, this study offers key information on
427 the effects of enforcement of referral guidelines in health facility referrals to KNH with
428 important lessons for Kenya and possible sub-Saharan Africa.

429

430 **Conclusions**

431

432 The enforcement of the national referral guidelines did reduce the number of facility
433 referrals from lower tier health facilities with a good number of private health facilities
434 stopped referring patients to KNH altogether. The major facility and patient factors that
435 were associated with facility referrals to KNH were human resource capacity and
436 availability, health facility infrastructure, Orthopaedic equipment's and implants
437 availability, patient's preference, financial considerations. Therefore, it is possible to
438 successfully enforce the implementation of referral guidelines.

439

440 **Recommendations**

441 **Recommendations to Policy Makers/ County Government**

- 442 1. Regularly sensitize the health facilities on the implementation of the national
443 referral guidelines 2014 for continued enforcement of the referral guidelines;
- 444 2. Strengthen the human resource capacity and availability to minimize on referrals
445 to KNH and this will consequently strengthen the primary and tertiary level
446 orthopaedic and trauma care and help decongest KNH and allow it to function as
447 a premier referral facility as per its statutory obligations;
- 448 3. Develop, disseminate and train health facilities on standard operating procedures
449 based on the national referral guidelines and ensure adherence to the same.

450 **Recommendations to KNH**

- 451 1. Educate and sensitize the health facilities on the role of KNH as a premier National
452 Teaching and Referral facility that is mandated to manage complex referrals and
453 should not be the first point of contact for patients seeking orthopaedic care;
- 454 2. Develop, disseminate and train KNH staff from A&E department, Orthopedic Clinic
455 and Corporate Outpatient Centre on the national referral guidelines and referral
456 standard operating procedures.

457 **Recommendations to Health Facilities**

- 458 1. Ensure adherence to the national referral guidelines of 2014 for an effective
459 referral process and to ensure optimum utilization of primary, secondary and
460 tertiary level of care.

461

462 **Data availability**

463

464 Data is available and has been uploaded in the submission portal.

465

466 **Conflict of interests**

467

468 The author has no conflict of interest to declare.

469

470 **Funding statements**

471

472 This study was partially funded by Kenyatta National Hospital RFA 2020/21.

473

474 **Acknowledgements**

475

476 We would like to sincerely acknowledge the work of Brian Okinyi and Micah J. Kipkemei

477

477 for commitment and assistance in data collection process.

478

479

479 **References**

480

480 1. Hensher M, Price M, Adomakoh S. Referral Hospitals. Disease Control Priorities
481 in Developing Countries 2nd edition Washington (DC): The International Bank for
482 Reconstruction and Development / The World Bank; 2006 2nd Edition ed. New York:
483 Oxford University Press; 2006. p. 1229 - 43.

484

484 2. Holdsworth G, Garner PA, Harphan T. Crowded outpatient departments in city
485 hospitals of developing countries: a case study from Lesotho. Int J Health Plann Manage.
486 1993 Oct-Dec;8(4):315-24.

487

487 3. London L, Bachmann OM. Pediatric utilization of a teaching hospital and a
488 community health center. Predictors of level of care used by children from Khayelitsha,
489 Cape Town. S Afr Med J. 1997 Jan;87(1):31-6.

490

490 4. Ohara K, Melendez V, Uehara N, Ohi G. Study of a patient referral system in the
491 Republic of Honduras. Health Policy Plan. 1998 Dec;13(4):433-45.

492

492 5. Sanders D, Kravitz J, Lewin S, McKee M. Zimbabwe's hospital referral system:
493 does it work? Health Policy Plan. 1998 Dec;13(4):359-70.

- 494 6. Nordberg E, Holmberg S, Kiugu S. Exploring the interface between first and
495 second level of care: referrals in rural Africa. *Trop Med Int Health*. 1996 Feb;1(1):107-11.
- 496 7. Akande T. Referral system in Nigeria: Study of a tertiary health facility. *Annals of*
497 *African Medicine*. 2004;3(3):130 - 3.
- 498 8. Stefanini A. District hospitals and strengthening referral systems in developing
499 countries. *World Hosp Health Serv*. 1994;30(2):14-9.
- 500 9. Pillay I, Mahomed OH. Prevalence and determinants of self-referrals to a District-
501 Regional Hospital in KwaZulu Natal, South Africa: a cross sectional study. *Pan Afr Med*
502 *J*. 2019; 33:4.
- 503 10. Godlee F. Operationalizing an effective referral system in India. *BMJ* Yale.
504 2015;351.
- 505 11. Liang L, Huang N, Shen Y, Chen A, Chou Y. Do patients bypass primary care for
506 common health problems under a free-access system? Experience of Taiwan. *BMC*
507 *Health Service Research* 2020;1050.
- 508 12. GoK. Article 6 of the Constitution of Kenya 2010. Nairobi, Kenya; 2010.
- 509 13. (NCAPD) NCAfPaD, [Kenya], Ministry of Health (MOH), Central Bureau of
510 Statistics (CBS), Macro. O. Kenya Service Provision Assessment Survey 2004.
- 511 14. MoH. Nairobi; 2021 [updated 2021; cited 2021 10th May]; Available from:
512 <https://knh.or.ke/index.php/history/>
- 513 15. MoH. Kenya Health Policy 2014 – 2030. Nairobi, Kenya; 2014.
- 514 16. KNH IN-PATIENT STATISTICS REPORT 2018, 2019 & 2020 [database on the
515 Internet]. 2020 [cited June 2020].

- 516 17. Duffield C, Diers D, O'Brien-Pallas L, Aisbett C, Roche M, King M, et al. Nursing
517 staffing, nursing workload, the work environment and patient outcomes. *Appl Nurs Res.*
518 2010 Nov;24(4):244-55.
- 519 18. Buerhaus JN. Policy Implications of Research on Nurse Staffing and Quality of
520 Patient Care Policy, Politics, & Nursing Practice. 2000.
- 521 19. Sharma SK, Rani R. Nurse-to-patient ratio and nurse staffing norms for hospitals
522 in India: A critical analysis of national benchmarks. *J Family Med Prim Care.* 2020 Jun
523 2020;9(6):2631-7.
- 524 20. Casagrande JT, Pike MC, Smith PG. An improved approximation formula for
525 calculating sample sizes for comparing two binomial distributions. *Biometrics* 1978;
526 34:483-6
- 527 21. Simba DO, Mbembati NA, Museru LM, Lema LE. Referral pattern of patients
528 received at the national referral hospital: challenges in low-income countries. *East Afr J*
529 *Public Health.* 2008 Apr;5(1):6-9.
- 530 22. Pittalis C, Brugha R, Bijlmakers L, Mwapasa G, Borgstein E, Gajewski J. Patterns,
531 quality and appropriateness of surgical referrals in Malawi. *Trop Med Int Health.* 2020
532 Jul;25(7):824-33.
- 533 23. Theoneste N, Gabriel T, Jackline O, Rebecca M, Robert R, Neil G, et al. Referral
534 patterns and predictors of referral delays for patients with traumatic injuries in rural
535 Rwanda. *Global Surgery.* 2016;160(6):1636-44.
- 536 24. Premkumar A, Massawe HH, Mshabaha DJ, R.Foran J, XiaohanYing, Sheth NP.
537 The burden of orthopaedic disease presenting to a referral hospital in northern Tanzania.
538 *Global Surgery.* 2015;2(1):70-5.

- 539 25. Theoneste Nkurunziza, Gabriel Toma, Jackline Odhiambo, Rebecca Maine,
540 Robert Riviello, Neil Gupta, et al. Referral patterns and predictors of referral delays for
541 patients with traumatic injuries in rural Rwanda. *Global Surgery*. 2016;160(6):1636-44.
- 542 26. Simba DO, Mbembati NAA, Museru LML, E.K. L. Referral Pattern of Patients
543 Received at the National Referral Hospital: Challenges in Low Income Countries. *East
544 African Journal of Public Health*. 2008;5(1):6-9.
- 545 27. Moturi AK, Suiyanka L, Mumo E, Snow RW, Okiro EA, Macharia PM. Geographic
546 accessibility to public and private health facilities in Kenya in 2021: An updated geocoded
547 inventory and spatial analysis. *Front Public Health*. 2022; 10:1002975.
- 548 28. Nyawira L, Tsofa B, Musiega A, Munywoki J, Njuguna RG, Hanson K, et al.
549 Management of human resources for health: implications for health systems efficiency in
550 Kenya. *BMC Health Serv Res*. 2022 Aug 16;22(1):1046.
- 551 29. Wisborg T, Montshiwa TR, Mock C. Trauma research in low- and middle-income
552 countries is urgently needed to strengthen the chain of survival. *Scand J Trauma Resusc
553 Emerg Med*. 2011 Oct 24; 19:62.
- 554 30. Gwatkin DR, Bhuiya A, Victora CG. Making health systems more equitable.
555 *Lancet*. 2004 Oct 2-8;364(9441):1273-80.
- 556 31. Viel IL, Moura BRS, Martuchi SD, de Souza Nogueira L. Factors Associated with
557 Interhospital Transfer of Trauma Victims. *J Trauma Nurs*. 2019 Sep/Oct;26(5):257-62.
- 558 32. Garwe T, Cowan LD, Neas B, Cathey T, Danford BC, Greenawalt P. Survival
559 benefit of transfer to tertiary trauma centers for major trauma patients initially presenting
560 to nontertiary trauma centers. *Acad Emerg Med*. 2010 Nov;17(11):1223-32.

- 561 33. Newgard CD, McConnell KJ, Hedges JR, Mullins RJ. The benefit of higher level of
562 care transfer of injured patients from nontertiary hospital emergency departments. J
563 Trauma. 2007 Nov;63(5):965-71.
- 564 34. Pittalis C, Brugha R, Gajewski J. Surgical referral systems in low- and middle-
565 income countries: A review of the evidence. PLoS One. 2019;14(9):e0223328.
- 566 35. Jumbam DT, Menon G, Lama TN, Lodge li W, Maongezi S, Kapologwe NA, et al.
567 Surgical referrals in Northern Tanzania: a prospective assessment of rates, preventability,
568 reasons and patterns. BMC Health Serv Res. 2020 Aug 8;20(1):725.
- 569 36. Hu R, Liao Y, Du Z, Hao Y, Liang H, Shi L. Types of health care facilities and the
570 quality of primary care: a study of characteristics and experiences of Chinese patients in
571 Guangdong Province, China. BMC Health Serv Res. 2016 Aug 2;16(a):335.
- 572 37. Luft HS, Garnick DW, Mark DH, Peltzman DJ, Phibbs CS, Lichtenberg E, et al.
573 Does quality influence choice of hospital? JAMA. 1990 Jun 6;263(21):2899-906.
- 574 38. Bahadori M, Teymourzadeh E, Ravangard R, Nasiri A, Raadabadi M,
575 Alimohammadzadeh K. Factors contributing towards patient's choice of a hospital clinic
576 from the patients' and managers' perspective. Electron Physician. 2016 May;8(5):2378-
577 87.
- 578 39. Nolan T, Angos P, Cunha AJ, Muhe L, Qazi S, Simoes EA, et al. Quality of hospital
579 care for seriously ill children in less-developed countries. Lancet. 2001 Jan
580 13;357(9250):106-10.


581
582

Captions of supporting Information files

583
584


S1 Supporting Information – Working dataset 04.12.2023

- 585 S2 Supporting Information – Facilities stopped referring
- 586 S3 Supporting Information – Reasons for Referral spreadsheet
- 587 S4 Supporting Information – Reasons for Referral Tally spreadsheet
- 588 S5 Supporting Information – Referrals spreadsheet
- 589 S6 Supporting Information – Referral facilities spreadsheet
- 590 Other - ERB approval letter - ERC APPROVAL LETTER.pdf
- 591 Other - PLOSONe_Human_Subjects_Research_Checklist.doc



medRxiv preprint doi: <https://doi.org/10.1101/2024.01.02.24300732>; this version posted January 3, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Figure 1



medRxiv preprint doi: <https://doi.org/10.1101/2024.01.02.24300732>; this version posted January 3, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Figure 2