1 **Understanding the Absolute Risk of Neonatal Respiratory Morbidity among Late Preterm**
2 **Births**
3

4 Mark A. CLAPP, MD MPH[1], Siguo LI, MS[1], Jessica L. COHEN, PhD[2], Cynthia GYAMFI-
5 BANNERMAN, MD, MS[3], Amy B. KNUDSEN, PhD[4], Scott A LORCH, MD, MSCE[5],
6 Tanayott THAWEETHAI, PhD[6], Jason D. WRIGHT, MD[7], Anjali J. KAIMAL, MD MAS[8] *,
7 Alexander MELAMED, MD MPH[1] *
8
[1] Department of Obstetrics, Gynecology, and Reproductive Biology, Massachusetts General
Hospital
[2] Department of Global Health and Population, Harvard T.H. Chan School of Public Health
[3] Department of Obstetrics, Gynecology, and Reproductive Sciences, University of California at
San Diego
[4] Institute of Technology Assessment, Department of Radiology, Massachusetts General Hospital
[5] Division of Neonatology, Children's Hospital of Philadelphia
[6] Division of Biostatistics, Massachusetts General Hospital
[7] Department of Obstetrics and Gynecology, Columbia University
[8] Department of Obstetrics and Gynecology, University of South Florida
19
20 * Drs. Kaimal and Melamed contributed equally.
21
Corresponding Author:
Mark A Clapp, MD MPH
Department of Obstetrics, Gynecology, and Reproductive Biology
Massachusetts General Hospital
55 Fruit St, Boston, MA 02114
Email: mark.clapp@mgh.harvard.edu
28

29 Word Count:
30   Main – 2,659
31   Abstract - 381
32

Disclosures: Dr. Clapp serves as a medical advisory board member with private equity in Delfina
Health, outside the submitted work. Dr. Melamed reports grants from the National Center for
Advancing Translational Sciences, the National Cancer Institute, the Conquer Cancer-The
ASCO Foundation, and the Department of Defense outside the submitted work. Dr. Melamed
has also served as an advisor for AstraZeneca. Dr. Wright has received research funding from
Merck and honoraria from UpToDate. Dr. Gyamfi-Bannerman reports grants from NHLBI,
NICHD, and NIHMD and has received research funding from HealthCore Inc/SERA
Prognostics, Inc. and MIRVIE, Inc.
Conflicts of interest: The authors declare no conflicts of interest or financial support.
Funding: None.
Data Access, Responsibility, and Analysis: Dr. Clapp had full access to all the data in the study
and takes responsibility for the integrity of the data and the accuracy of the data analysis.
Data Sharing: Data used in this study is publicly available at https://dash.nichd.nih.gov/.
Acknowledgments: The authors have no other acknowledgments.

Precis

The absolute risk of neonatal respiratory morbidity among patients at risk for late preterm

delivery varies considerably by gestational age and mode of delivery.

Abstract

Objective: To estimate the effect of late preterm antenatal steroids on the absolute risk of

respiratory morbidity among subgroups of patients based on the planned mode of delivery and

gestational age at presentation.

Methods: This was a secondary analysis of the Antenatal Late Preterm Steroid (ALPS) Trial, a

multicenter trial originally conducted within the NICHD's MFMU network of individuals with

singleton gestations and without pre-existing diabetes who were at high risk for late preterm

delivery (34-36 weeks of gestation). We fit binomial regression models to estimate the absolute

risk of respiratory morbidity, with and without steroid administration, by gestational age and

planned mode of delivery at the time of presentation. We assumed a homogenous effect of

steroids on the log-odds scale, as was reported in the ALPS trial. The primary outcome was

neonatal respiratory morbidity, as defined in the ALPS Trial

Results: The analysis included 2,825 patients at risk for late preterm birth. The risk of respiratory

morbidity varied significantly by planned mode of delivery (adjusted risk ratio (aRR) 1.90 (95%

confidence interval (CI) 1.55, 2.33) for cesarean compared to vaginal delivery) and day of

gestation at presentation (aRR 0.92 (95% CI 0.90, 0.94)). For those planning a cesarean delivery

and presenting in the 34th week of gestation, the risk of neonatal respiratory morbidity was

39.4% (95% CI 30.8, 47.9%) without steroids and 32.0% (95% CI 24.6, 39.4%) with steroids. In

contrast, for patients presenting in the 36th week and planning a vaginal delivery, the risk of

neonatal respiratory morbidity was 6.9% (95% CI 5.6, 8.6%) without steroids and 5.6% (95%

4.2, 7.0%) with steroids.

<u>Conclusion</u>: The absolute risk of neonatal respiratory morbidity among patients at risk for late

preterm delivery varies considerably by gestational age at presentation and planned mode of

delivery. As only communicating the relative risk reduction of antenatal steroids for respiratory

morbidity may lead to an inaccurate perception of benefit, individualized estimates of absolute

risk expected with and without treatment may inform shared decision-making.

Introduction

Neonatal respiratory morbidity, including respiratory distress syndrome and transient tachypnea of the newborn, is a common complication of prematurity.[1–3] Multiple patient factors, including gestational, mode of delivery, and fetal sex, influence the risk of respiratory morbidity.[2–7] There is a broad and longstanding consensus that antenatal steroids decrease the risk of neonatal respiratory morbidity for neonates born before 34 weeks of gestation.[8,9] More recently, the Antenatal Late Preterm Steroid (ALPS) Trial demonstrated that this benefit extends to fetuses at risk of delivery in the late preterm period (34-36 weeks of gestation): neonates exposed antenatally to betamethasone in the late preterm period had a 20% risk reduction of the composite respiratory outcome, without statistically conclusive evidence of treatment heterogeneity among pre-specified subgroups.[10]

The Society for Maternal-Fetal Medicine recommends that clinicians and patients engage in shared decision-making around steroid administration in the late preterm period.[11] However, the baseline risk of respiratory morbidity differs substantially across the late preterm period, making summary measures of absolute risk, as reported in the original trial, less informative for individualized counseling.[2] Furthermore, communicating only the relative risk may lead to an inaccurate perception of the benefit of antenatal steroids, especially for those with a small absolute risk.[12] Thus, our objective in this secondary analysis of data from the ALPS Trial was to estimate the absolute risk of respiratory morbidity with and without steroid administration, based on patients' planned mode of delivery and gestational age at presentation.[10]

Methods

This is a secondary analysis of the Antenatal Late Preterm Steroid (ALPS) Trial

conducted within 17 university-based hospitals within the Maternal-Fetal Medicine Units

(MFMU).[10] In the trial, patients with a singleton pregnancy between 34 weeks 0 days and 36

weeks 5 days of gestation and with a high chance of delivery in the late preterm period were

enrolled and randomized to receive an antenatal course of betamethasone or placebo. The

primary outcome was a composite of respiratory morbidity within 72 hours after delivery and

included the following: high-flow nasal cannula or continuous positive airway pressure (CPAP)

for at least 2 hours; supplemental oxygen use for at least 4 hours; extracorporeal membrane

oxygenation (ECMO); and/or mechanical ventilation. We also examined the secondary outcome

of severe respiratory complications, as defined in the ALPS Trial as "a composite outcome of

CPAP or high-flow nasal cannula for at least 12 continuous hours, supplemental oxygen with a

fraction of inspired oxygen of at least 0.30 for at least at least 24 continuous hours, ECMO or

mechanical ventilation, stillbirth, or neonatal death within 72 hours after delivery."[10] The ALPS

trial randomized 2,831 patients to betamethasone (n=1,429) or placebo (n=1,402).[10] This analysis

included patients randomized at or after 34 weeks of gestation who were not lost to follow-up.

We examined the risk of neonatal respiratory morbidity among two patient factors

strongly associated with the risk of respiratory morbidity: mode of delivery and gestational

age.[2,3,5] To generate results that could guide clinical decision-making and inform counseling, we

considered *planned* mode of delivery (not actual mode) and gestational age *at presentation* (not

delivery), information available to clinicians and patients when deciding about antenatal steroid

administration. We defined the planned mode of delivery as "cesarean" if 1) a cesarean delivery

was scheduled in the late preterm period at the time of presentation or 2) if no delivery was

scheduled at the time of presentation but a cesarean section was performed at any gestational age

for the following indications: placenta accreta, placenta previa, fetal anomaly, prior uterine

surgery, or prior cesarean delivery. This definition was chosen to capture patients who were

unlikely to attempt a trial of labor. The prevalence of these factors was compared between the

two groups to examine for potential group imbalances in the original trial.

We constructed a binomial regression model that included the treatment assignment

(placebo or betamethasone), planned mode of delivery, and gestational age in days and used a

log link to report risk ratios. Because the prespecified subgroup analysis in the original trial did

not detect signs of treatment effect heterogeneity, the model assumed a constant effect size (on

the log-odds scale) of the intervention across all groups. We used this regression model to

estimate the absolute risks of respiratory morbidity in both the treated and untreated groups. The

predicted (or estimated) absolute risks of this model were presented graphically by planned mode

of delivery and across days of gestation; for comparison, we also plotted the observed absolute

risks of the outcome with a corresponding 95% confidence interval by planned mode of delivery

and weeks of gestation. To examine the model's discrimination and calibration, we used k-fold

cross-validation (k=10) to calculate the area under the receiver operating characteristic (ROC)

curves (AUC) and generate a calibration plot for both outcomes.

Additional steps were performed to further facilitate counseling around the absolute risk

of respiratory morbidity and expected outcome by mode of delivery and gestational age. First,

the risks of the primary and secondary outcomes were estimated using the same logistic

regression model, though gestational age in weeks was used instead of gestational age in days;

this allowed for the output to be summarized in a compact, interpretable table. Second, we built

an online reference tool to display the estimated risk of the primary and secondary outcome

using a patient's precise gestational age in days at presentation and their planned mode of

delivery, which is available at https://mghinferencelab.shinyapps.io/alps_web_calc/.

Data was obtained with permission from the NICHD Data and Specimen Hub (DASH).[13]

The XXX Institution Review Board exempted this study. All analyses used Stata (Stata Corp,

College Station, TX). P-values <0.05 were considered statistically significant. The STROBE

reporting guidelines were followed.

Results

Of the original 2,831 participants, we excluded 2 patients randomized before 34 weeks of

gestation and 4 patients lost to follow-up (2.1%). eTable 1 in the Supplement shows the sample

sizes of the various strata used in the analysis. As expected, there were no differences in

distributions of the planned mode of delivery or gestational age at randomization between the

two groups (eTable 2 in the Supplement).

Figure 1 shows the predicted absolute risks of respiratory morbidity among patients

treated with steroids and those who did not by the day of gestation and the planned mode of

delivery at the time of presentation. The observed rates (not those predicted by the model) by

week of gestation are shown on the same graph for comparison and included in eTable 3 in the

Supplement. The risk of respiratory morbidity varied significantly across planned modes of

delivery (adjusted risk ratio (aRR) 1.90 (95% confidence interval (CI) 1.55, 2.33) for cesarean

compared to vaginal delivery) and day of gestation at presentation (aRR 0.92 (95% CI 0.90,

0.94)). The unadjusted and adjusted model outputs are shown in eTable 4 in the Supplement. The

10-fold cross-validated AUC was 0.669 (95% CI 0.639, 0.698). The ROC curve and calibration

plot is shown in eFigure 1 and eFigure 2 in the Supplement.

167  The predicted absolute risks with and without treatment are summarized by week of

168 gestation in Table 1. At baseline (i.e., among those treated with the placebo), neonates born to

169 individuals planning a cesarean delivery have a 36.7% (95% CI 29.8, 43.7%) risk of respiratory

170 morbidity when presenting in the 34th week of gestation; this risk decreases to 31.4% (95% CI

171 24.5, 37.9%) with antenatal steroids. By contrast, neonates who are born via a planned vaginal

172 delivery after presenting in the 36th week of gestation have a 6.8% (95% CI 5.1, 8.5%) risk of

173 respiratory morbidity without steroids and a 5.4% (95% CI 4.0, 6.9%) risk with steroid

174 treatment.

175  Figure 2 shows the predicted absolute risks of severe respiratory complications among

176 those who received and did not receive antenatal steroids by day of gestational age and planned

177 mode of delivery. The observed risks are shown on the same graphs by week of gestation and are

178 included in eTable 5 in the Supplement. Planned mode of delivery (aRR 2.16 (95% CI 1.72,

179 2.72) for cesarean vs. vaginal delivery) and gestational age (aRR 0.92 (95% CI 0.90, 0.94)) were

180 significantly associated with the risk of severe respiratory complications. The unadjusted and

181 adjusted model outputs are shown in eTable 6 in the Supplement. The 10-fold cross-validated

182 AUC was 0.679 (95% CI 0.651, 0.717); the ROC curve and calibration plots are shown in

183 eFigure 3 and eFigure 4 in the Supplement. These findings are summarized by week of gestation

184 in Table 2 which tabulates the absolute risk of severe respiratory complication, with and without

185 treatment, by planned mode of delivery and week of gestational age at presentation.

186  The results of these two models are available as an interactive calculator available at

187 https://mghinferencelab.shinyapps.io/alps_web_calc/. This tool displays the estimated absolute

188 risk of respiratory morbidity (and associated 95% CIs) with and without treatment as a function

189 of day of gestation at presentation and planned mode of delivery (example shown in Figure 3).

Discussion

In this secondary analysis of the ALPS Trial, we demonstrated that the absolute risks of

neonatal respiratory morbidity vary significantly by gestational age and planned mode of

delivery. Notably, infants born via a planned cesarean delivery are at a much higher risk of

respiratory morbidity than those having a vaginal delivery, especially at earlier gestational ages

(approximately 37% for planned cesarean delivery at 34 weeks vs. 7% for planned vaginal

delivery at 36 weeks). While these relationships were previously known, we quantified this risk

in the late preterm period using prospectively collected data as part of the ALPS Trial. Similarly,

we demonstrated the challenge of predicting these outcomes for patients at the time of

counseling and decision-making for antenatal steroids, even in the context of a clinical trial that

used a strict definition of "high risk for preterm delivery," as demonstrated by the ROC curves;

to convey the level of uncertainty around the estimates, all data are presented with 95%

confidence intervals. This information can inform shared decision-making about late preterm

steroids and allow clinicians to estimate the baseline risk of neonatal respiratory morbidity more

accurately. This absolute baseline risk of respiratory in the unexposed group can then be used to

interpret the expected relative risk reduction that was demonstrated uniformly across all enrollees

in the trial.

These findings must be interpreted in the context of the original ALPS Trial protocol and

the trial enrollees. The ALPS Trial only enrolled patients who were at high risk of a late preterm

delivery as determined by predefined clinical factors (rupture of membranes, preterm labor at ≥3

cm dilation or ≥75% effacement, or expected preterm delivery via induction or cesarean within

24 hours and 7 days) or provider judgment who had not previously received antenatal

corticosteroids or had pregestational diabetes.[10] As a result, nearly 40% of enrollees delivered

before receiving the second dose, and the median time from presentation to delivery was approximately 30 hours.[10] The absolute risks reported from this trial are most relevant to patients presenting with a similar risk of preterm delivery. For example, the benefit of antenatal steroids may be higher than reported in the trial for patients who are more likely to deliver after receiving a full course; in contrast, the benefit may be less if more patients received antenatal steroids but went on to delivery at later gestations or full term (i.e., a lower risk of preterm delivery than the trial). Studies of late preterm antenatal steroids outside of the context of a clinical trial are needed to better estimate the absolute risks of respiratory morbidity and the benefit of the intervention.

The original trial did include subgroup comparisons for several factors, including planned cesarean delivery and gestational age, though gestational age was analyzed as 34-35 vs. 36 weeks of gestation. The study reported no significant effects for the interaction terms for the primary outcome when each factor was analyzed individually; however, the trial was not powered to detect differences in the primary outcome within subgroups. The possibility of a differential impact of betamethasone in actual practice (i.e., treatment effect heterogeneity) remains plausible, especially considering the differences in the number of days between first dose and delivery and the number of patients who received a full course of betamethasone before delivery by planned mode of delivery and gestational age at presentation. Investigations of treatment effect heterogeneity should be a focus of future research to further individualize counseling on risks and benefits.

This study focused on counseling regarding the individualized benefits of antenatal steroids among two common and known clinical factors across which the baseline risk of respiratory morbidity varies widely. However, other factors may inform clinical decision-

making, especially the known and theoretical intervention risks. Hypoglycemia, which was more

likely in the treatment group in the ALPS Trial (RR 1.60 (95% CI 1.37-1.87)), also likely varies

among population subgroups (e.g., gestational age).[10,14,15] We could not investigate the absolute

risks of hypoglycemia across gestational age and mode of delivery because these data are not

publicly available due to ongoing data collection at the time of this analysis.[13] Furthermore, there

is no data available to inform individualized counseling on the associations between long-term

neurocognitive or developmental risks and antenatal steroids in the late preterm period. Available

evidence in other populations is mixed, though a single dose of antenatal steroids in the early

preterm period has not been shown to increase long-term neurodevelopment risks.[16–21]

The study is strengthened by its use of rigorously collected data as part of a large

randomized controlled trial, allowing us to estimate the rates of respiratory morbidity among

those exposed and unexposed to antenatal steroids in the late preterm period. However, it is

limited in that the trial enrollees may not reflect the general population at risk for late preterm

birth or considering antenatal steroids. As mentioned, the baseline risks of respiratory morbidity

are presented for individuals with a high likelihood of delivery in the late preterm period.[10] The

baseline risks may vary in a population without the same likelihood of preterm delivery.

Similarly, the original trial was not powered to detect differences within population subgroups.

As such, the number of patients within our strata varies, some of which are small, leading to

considerable uncertainty. To help quantify this uncertainty, 95% confidence intervals are

provided in Tables 1 and 2 and in the online calculator. Other factors may also contribute to a

neonate's risk for respiratory morbidity. However, this study focused on two patient-specific

factors that would be known at the time of presentation and have previously been shown to have

a significant effect on risk in the literature; future studies could try to improve the performance

of the predictive model for respiratory morbidity using other factors, though the baseline

challenge of predicting who and when a patient will deliver in the preterm period likely will

continue to limit model discrimination. Last, we acknowledge that many non-patient factors may

affect clinical decision-making surrounding ALPS (e.g., hospital resources, etc.); this information

and the visualization tool are intended to help conceptualize the absolute risk reduction of

respiratory morbidity with ALPS, which may inform one aspect of the shared decision-making.

265   In conclusion, the absolute risk of neonatal respiratory morbidity among patients at high

risk for late preterm delivery varies considerably by gestational age at presentation and planned

mode of delivery. Assuming no treatment effect heterogeneity (i.e., a constant relative risk

reduction among all subgroups), patients with a low baseline risk (e.g., those presenting in the

36[th] week and planning a vaginal delivery) can expect to obtain a considerably smaller absolute

benefit from steroid administration compared to patients at the highest risk of neonatal

respiratory morbidity (e.g., those presenting in the 34[th] week and planning a cesarean delivery). ;

Accurate understanding of these anticipated benefits and potential risks is critical to patients'

making decisions based on informed patient preferences. Studies adequately powered to compare

the risks and benefits of antenatal late preterm steroids among population subgroups, and their

effectiveness outside the context of a clinical trial, are needed to increase the precision of shared

decision-making and public policy surrounding late preterm steroids.

References

1. Mahoney AD, Jain L. Respiratory disorders in moderately preterm, late preterm, and early term infants. *Clin Perinatol*. 2013;40(4):665-678. doi:10.1016/j.clp.2013.07.004

2. Consortium on Safe Labor, Hibbard JU, Wilkins I, et al. Respiratory morbidity in late preterm births. *JAMA*. 2010;304(4):419-425. doi:10.1001/jama.2010.1015

3. De Luca R, Boulvain M, Irion O, Berner M, Pfister RE. Incidence of early neonatal mortality and morbidity after late-preterm and term cesarean delivery. *Pediatrics*. 2009;123(6):e1064-1071. doi:10.1542/peds.2008-2407

4. Mekic N, Selimovic A, Cosickic A, et al. Predictors of adverse short-term outcomes in late preterm infants. *BMC Pediatr*. 2023;23(1):298. doi:10.1186/s12887-023-04112-z

5. Berthelot-Ricou A, Lacroze V, Courbiere B, Guidicelli B, Gamerre M, Simeoni U. Respiratory distress syndrome after elective caesarean section in near term infants: a 5-year cohort study. *J Matern Fetal Neonatal Med*. 2013;26(2):176-182. doi:10.3109/14767058.2012.733743

6. Altman M, Vanpée M, Cnattingius S, Norman M. Risk factors for acute respiratory morbidity in moderately preterm infants. *Paediatr Perinat Epidemiol*. 2013;27(2):172-181. doi:10.1111/ppe.12035

7. Condò V, Cipriani S, Colnaghi M, et al. Neonatal respiratory distress syndrome: are risk factors the same in preterm and term infants? *J Matern Fetal Neonatal Med*. 2017;30(11):1267-1272. doi:10.1080/14767058.2016.1210597

8. Roberts D, Dalziel S. Antenatal corticosteroids for accelerating fetal lung maturation for women at risk of preterm birth. *Cochrane Database Syst Rev*. 2006;(3):CD004454. doi:10.1002/14651858.CD004454.pub2

9. Gilstrap LC, Christensen R, Clewell WH, et al. Effect of Corticosteroids for Fetal Maturation on Perinatal Outcomes: NIH Consensus Development Panel on the Effect of Corticosteroids for Fetal Maturation on Perinatal Outcomes. *JAMA*. 1995;273(5):413-418. doi:10.1001/jama.1995.03520290065031

10. Gyamfi-Bannerman C, Thom EA, Blackwell SC, et al. Antenatal Betamethasone for Women at Risk for Late Preterm Delivery. *New England Journal of Medicine*. 2016;374(14):1311-1320. doi:10.1056/NEJMoa1516783

11. Reddy UM, Deshmukh U, Dude A, Harper L, Osmundson SS. Society for Maternal-Fetal Medicine Consult Series #58: Use of antenatal corticosteroids for individuals at risk for late preterm delivery: Replaces SMFM Statement #4, Implementation of the use of antenatal corticosteroids in the late preterm birth period in women at risk for preterm delivery, August 2016. *American Journal of Obstetrics & Gynecology*. 2021;225(5):B36-B42. doi:10.1016/j.ajog.2021.07.023

12. Freeman ALJ. How to communicate evidence to patients. *Drug Ther Bull*. 2019;57(8):119-
124. doi:10.1136/dtb.2019.000008

13. NICHD DASH - Eunice Kennedy Shriver National Institute of Child Health and Human
Development Data and Specimen Hub. Accessed November 11, 2023.
https://dash.nichd.nih.gov/

14. Bromiker R, Perry A, Kasirer Y, Einav S, Klinger G, Levy-Khademi F. Early neonatal
hypoglycemia: incidence of and risk factors. A cohort study using universal point of care
screening. *The Journal of Maternal-Fetal & Neonatal Medicine*. 2019;32(5):786-792.
doi:10.1080/14767058.2017.1391781

15. Lubchenco LO, Bard H. INCIDENCE OF HYPOGLYCEMIA IN NEWBORN INFANTS
CLASSIFIED BY BIRTH WEIGHT AND GESTATIONAL AGE. *Pediatrics*.
1971;47(5):831-838. doi:10.1542/peds.47.5.831

16. Räikkönen K, Gissler M, Tapiainen T, Kajantie E. Associations Between Maternal Antenatal
Corticosteroid Treatment and Psychological Developmental and Neurosensory Disorders in
Children. *JAMA Network Open*. 2022;5(8):e2228518.
doi:10.1001/jamanetworkopen.2022.28518

17. Ninan K, Liyanage SK, Murphy KE, Asztalos EV, McDonald SD. Evaluation of Long-term
Outcomes Associated With Preterm Exposure to Antenatal Corticosteroids: A Systematic
Review and Meta-analysis. *JAMA Pediatrics*. 2022;176(6):e220483.
doi:10.1001/jamapediatrics.2022.0483

18. van der Merwe JL, Sacco A, Toelen J, Deprest J. Long-term neuropathological and/or
neurobehavioral effects of antenatal corticosteroid therapy in animal models: a systematic
review. *Pediatr Res*. 2020;87(7):1157-1170. doi:10.1038/s41390-019-0712-1

19. Thevathasan I, Said JM. Controversies in antenatal corticosteroid treatment. *Prenatal
Diagnosis*. 2020;40(9):1138-1149. doi:10.1002/pd.5664

20. Dalziel SR, Lim VK, Lambert A, et al. Antenatal exposure to betamethasone: psychological
functioning and health related quality of life 31 years after inclusion in randomised
controlled trial. *BMJ*. 2005;331(7518):665. doi:10.1136/bmj.38576.494363.E0

21. Sotiriadis A, Tsiami A, Papatheodorou S, Baschat AA, Sarafidis K, Makrydimas G.
Neurodevelopmental Outcome After a Single Course of Antenatal Steroids in Children Born
Preterm: A Systematic Review and Meta-analysis. *Obstet Gynecol*. 2015;125(6):1385-1396.
doi:10.1097/AOG.0000000000000748

Figure 1: Risk of respiratory morbidity among individuals exposed and not exposed to antenatal
steroids by gestational age and planned mode of delivery
349

350
351
The lines represent the estimated rates of respiratory morbidity among those exposed (+ ALPS)
and not exposed (- ALPS) that were derived from a logistic regression model that included
gestation age at presentation (in days), planned mode of delivery (vaginal vs. cesarean), and
treatment group. For comparison, the observed rates (i.e., "actual risk") with their corresponding
95% confidence interval (CI) by week (not day) of gestation are also plotted; of note, their
position on the x-axis does not correspond to a specific day within the week of gestation.
358

359   Figure 2: Risk of severe respiratory complications among individuals exposed and not exposed
360   to antenatal steroids by gestational age and planned mode of delivery
361

362
363
The lines represent the estimated rates of severe respiratory complications among those exposed
(+ ALPS) and not exposed (- ALPS) that were derived from a logistic regression model that
included gestation age at presentation (in days), planned mode of delivery (vaginal vs. cesarean),
and treatment group. For comparison, the observed rates (i.e., "actual risk") with their
corresponding 95% confidence interval (CI) by week (not day) of gestation are also plotted; of
note, their position on the x-axis does not correspond to a specific day within the week of
gestation.
371

372   Figure 3: Image of online calculator to estimate the absolute risks of respiratory morbidity by
373   planned mode of delivery and gestational age with and without antenatal steroids
374

Absolute Risk of Respiratory Morbidity for Neonates at High Risk for Late Preterm Birth

The following model estimates the absolute risk of respiratory outcomes for singleton gestations at high risk for late preterm birth using data from the ALPS Trial. In the trial, *high risk for preterm delivery* was definied by the following criteria: rupture of membranes, preterm labor at ≥3 cm dilation or ≥75% effacement, expected preterm delivery via induction or cesarean within 24 hours and 7 days, or provider judgment. Risk estimates are a function of the planned mode of delivery and gestational age (in days) at the time of presentation and are presented between the trial's control group ('Without Steroids') and intervention group ('With Steroids').

Please note: This tool is only for research purposes, and the findings have not been externally validated. Confidence intervals around the point estimates are shown to reflect the degree of uncertainty.

	Without Steroids	With Steroids
Respiratory Morbidity	16.3% (14.3% - 18.4%)	13.6% (11.7% - 15.6%)
Severe Respiratory Complication	13.7% (11.8% - 15.8%)	9.5% (7.9% - 11.4%)

Outcome Definitions

Respiratory Morbidity

Respiratory support <72 hours after birth is defined as any of the following: the use of CPAP or high-flow nasal cannula for ≥2 consecutive hours, supplemental oxygen with a FiO2 ≥0.30 for ≥4 continuous hours, ECMO, or mechanical ventilation.

Severe Respiratory Complication

Defined as any of the following: CPAP or high-flow nasal cannula for ≥ 2 continuous hours, supplemental oxygen with FiO2 ≥0.30 for ≥24 continuous hours, ECMO or mechanical ventilation, stillbirth, or neonatal death within 72 hours after delivery.

377 Table 1: Estimated risk of respiratory morbidity among neonates not exposed and exposed to

378 antenatal steroids, stratified by week of gestation and planned mode of delivery

Week of Gestation at Presentation	Planned Vaginal Delivery		Planned Cesarean Delivery	
	No ALPS % (95% CI)	ALPS % (95% CI)	No ALPS % (95% CI)	ALPS % (95% CI)
34 weeks	20.9 (17.5-24.4)	17.0 (13.9-20.2)	39.4 (30.8, 47.9)	32.0 (24.6-39.4)
35 weeks	11.8 (9.6-14.1)	9.6 (7.7-11.5)	22.3 (17.3-27.3)	18.1 (14.0-22.3)
36 weeks	6.9 (5.2-8.6)	5.6 (4.2-7.0)	12.9 (9.3-16.6)	10.6 (7.5-13.)

381 ALPS, antenatal late preterm steroids; CI, confidence interval.

384 Table 2: Estimated risk of severe respiratory complications among neonates not exposed and

385 exposed to antenatal steroids, stratified by week of gestation and planned mode of delivery

Week of Gestation at Presentation	Planned Vaginal Delivery		Planned Cesarean Delivery	
	No ALPS % (95% CI)	ALPS % (95% CI)	No ALPS % (95% CI)	ALPS % (95% CI)
34 weeks	17.3 (14.1-20.1)	11.6 (9.0-14.2)	37.0 (28.1-45.9)	24.8 (18.1-31.4)
35 weeks	9.4 (7.6-11.8)	6.5 (5.0-8.1)	20.8 (15.6-25.9)	13.9 (10.2-17.6)
36 weeks	5.0 (3.5-6.4)	3.3 (2.3-4.4)	10.6 (7.1-14.1)	7.1 (4.7-9.5)

388 ALPS, antenatal late preterm steroids; CI, confidence interval.