

1 **ABSTRACT (277 words)**

2

3 **Background.** Tuberculosis (TB) remains a significant global health challenge, particularly
4 with the rise of drug-resistant strains, such as Pyrazinamide (PZA)-resistant TB. This
5 resistance hampers treatment effectiveness. Currently, there's a lack of affordable and precise
6 quantitative tests for detecting PZA resistance, highlighting the need for accessible diagnostic
7 tools. Our study introduces a direct, accurate, and accessible susceptibility test for PZA
8 resistance by quantifying pirazinoic acid (POA).

9 **Methods.** We analyzed 264 TB-positive samples (MP and TBN) to assess PZA susceptibility
10 by measuring POA production. This was done using the MODS-Wayne qualitative assay and
11 our developed quantitative approach (MODS-WQ). For MODS-WQ, TBN samples were
12 processed in 7H9 media (MODS-WQ 7H9) and MP samples in citrate buffer (MODS-WQ
13 CB). We measured POA levels using spectrophotometry against a calibration curve, with
14 PZA susceptibility determined by a composite standard. We also investigated the correlation
15 between POA levels and pyrazinamidase mutations. Multidrug-resistant samples were
16 assessed using the MODS test, and statistical analyses were conducted.

17 **Results.** Approximately 23.48% of the samples showed PZA resistance, which included
18 62.8% of the multidrug-resistant (MDR) samples. The MODS-Wayne assay demonstrated
19 an 87.8% sensitivity and 95.9% specificity for MP samples, and 62.5% sensitivity with
20 93.7% specificity for TBN samples. The MODS-WQ identified specific cut-off points—
21 123.25 μM for MODS-WQ 7H9 and 664.7 μM for MODS-WQ CB—with corresponding
22 sensitivities of 81.25% and 92.31%, and specificities of 77.22% and 95.93%. Significantly
23 lower POA levels were observed in samples with mutations in the metal binding site
24 compared to mutations on the enzyme's periphery.

25 **Conclusions.** The citrate-buffered MODS-WQ demonstrates high sensitivity and specificity
26 in quantifying POA. It offers a significant improvement over qualitative methods, effectively
27 overcoming their subjectivity limitations in PZA resistance detection.

28

29

30

31

1 **BACKGROUND (576 words)**

2 Tuberculosis (TB) remains a major global health concern, particularly in low- and mid-
3 income countries where it has reached pandemic proportions. The emergence of drug-
4 resistant TB strains poses a significant challenge to eradication efforts [1]. Pyrazinamide
5 (PZA), a first-line anti-TB drug, is pivotal in shortening treatment duration to six months for
6 both drug-sensitive and multidrug-resistant (MDR) TB cases [2]. PZA acts as a prodrug,
7 which is converted into its active form, pyrazinoic acid (POA), by the enzyme
8 pyrazinamidase (PZase). The PZase enzyme is encoded by the *pncA* gene, playing a crucial
9 role in the drug's efficacy [3].

10 Resistance to Pyrazinamide (PZA_r) significantly worsens TB treatment outcomes and
11 increases mortality rates [4]. It's estimated that PZA_r occurs in about 16.2% of all TB cases
12 globally [5] and constitutes approximately 60% of multidrug-resistant (MDR) TB cases. The
13 lack of a standardized and reliable method for routinely detecting PZA_r presents a major
14 obstacle in TB management strategies [6].

15 Routine testing for PZA resistance (PZA_r) is often not performed due to various technical
16 challenges [7]–[9]. Existing phenotypic culture-based drug susceptibility testing (pDST)
17 methods, such as the BACTEC MGIT 960 system [10], the Wayne assay [11], the Nitrate
18 reductase assay [12], and the Microscopic Observation Drug Susceptibility (MODS) assay
19 [13], face several limitations. These include difficulties in maintaining the precise inoculum
20 amount and pH level (5.5-6.5) necessary for PZA activity, the risk of contamination with
21 other bacterial strains leading to an error rate of 10-15% [7], and prolonged detection times
22 [8]. Additionally, the Bactec MGIT 960 system, despite its high cost, is known to produce a
23 significant rate of false resistance results [9].

24 The Wayne assay, which assesses the enzymatic activity of pyrazinamidase (PZase), is an
25 economical colorimetric test that identifies pyrazinoic acid (POA) as a marker of
26 susceptibility using ferrous ammonium sulfate (SAF) [11]. Despite its cost-effectiveness, this
27 assay has limitations: it is qualitative and subjective, and it necessitates extended incubation
28 periods until bacterial growth is detectable [8].

29 Genotypic drug susceptibility testing (gDST) methods, such as Sanger sequencing and
30 whole-genome sequencing, are employed to identify specific genetic mutations in the *pncA*
31 gene linked to PZA resistance (PZA_r) [14], [15]. These methods can also detect mutations in

1 other genes, including *rpsA*, *panD*, and *clpC1* [16]–[19]. gDSTs are generally more sensitive
2 and specific than phenotypic methods, with the majority (70-97%) of PZA-resistant clinical
3 isolates showing genetic variations in the *pncA* gene's promoter region or coding sequence
4 [20], [21]. However, these techniques require significant technical expertise and are costly,
5 making them less accessible in low- and mid-income countries where TB is endemic [22].
6 Additionally, the variability in mutation patterns linked to PZA^r can differ regionally,
7 affecting the representativeness of these tests [23].

8 The Microscopic Observation Drug Susceptibility (MODS) assay is an effective, low-cost,
9 and simple method for diagnosing TB and MDR-TB directly from sputum samples, with a
10 short turnaround time of 5 to 21 days. However, its application for Pyrazinamide (PZA)
11 testing has been limited. To address this, Alcántara et al. [24] developed the MODS-Wayne
12 method, combining the MODS assay with the Wayne test. This adaptation is based on
13 detecting pyrazinoic acid (POA), the hydrolyzed product of PZA expelled into the
14 extracellular environment, via a colorimetric reaction with ferrous ammonium sulfate (SAF)
15 [11]. The MODS-Wayne assay demonstrated high sensitivity and specificity (92.7% and
16 99.3%, respectively). Nonetheless, its subjective nature may affect the reproducibility of
17 results.

18 In this study, we aim to refine the MODS-Wayne test by introducing a quantitative approach
19 (MODS-WQ) using sputum samples from TB patients. This new approach seeks to reduce
20 diagnosis time, eliminate the need for pre-isolation in solid cultures, and move beyond
21 qualitative judgments.

22

23

24 **MATERIALS AND METHODS (920 words)**

25 Samples and Study design

26 264 TB-positive samples from two studies conducting during 2015-2016 (coded as MP) and
27 2017-2018 (coded as TBN), obtained from patients enrolled in the National TB Strategy
28 treatment program at the Hospital Nacional Dos de Mayo, Lima, Peru, and from the Regional
29 Tuberculosis Reference Laboratory, Callao, Lima, Peru. Only basic data (age, sex, treatment)
30 was recorded to the sample anonymity. This work was approved by the Institutional
31 Committee on Research Ethics (CIEI) of the Universidad Peruana Cayetano Heredia (Code

1 331-34-21). For the quantitative MODS-WQ test, two variants were performed and
2 contrasted with the qualitative version MODS-Wayne. One variant (MP n=169), were carried
3 out in citrate buffer (pH 7.0) matrix, named as MODS-WQ BC, whereas the other (TBN
4 n=95) was coded as TBN, which were performed in the same media culture 7H9 (pH~6.8),
5 named as MODS-WQ 7H9 (**Figure 1**).

6 7 Multidrug resistant test: rifampicin (RIF) and isoniazid (INH) resistance

8 All sputum samples were previously decontaminated according to the protocol by Kent and
9 Kubica [25] and seeded in 7H9 Middlebrook (**See Supplementary methods**). MODS test
10 was carried out using the protocol implemented in our laboratory [26]. Samples were
11 incubated with INH at final concentration of 0.4 µg/ml or RIF at 1.0 µg/ml at 37°C for a
12 maximum of 21 days followed by evaluation using an inverted microscope.

13 14 MODS-Wayne qualitative assay for PZA susceptibility

15 This test was conducted simultaneously with the MODS-MDR assay (**Supp. Figure S1A**).
16 Samples were allocated into two wells: Control (PZA-C) and treatment (PZA-Wayne), then
17 incubated at 37°C. Upon observing growth in the PZA-C well, the date was recorded, and
18 incubation continued for an additional three days. Subsequently, PZA (800 µg/mL) was
19 introduced into the PZA-Wayne well and further incubated for three days. Next, 100 µL of
20 10% SAF was added and incubated for five minutes, with immediate color registration. A
21 red color indicated sensitivity, while the absence of color denoted resistance. Due to
22 variations in observed coloration among samples, an intensity scale from 0 to 3 was
23 established, where 0 represented resistance and 3 signified high sensitivity (**Supp. Figure**
24 **S1C-D**). For more details **see supplementary methods**.

25 26 MODS-WQ assay for PZA susceptibility

27 This assay included two variants: MODS-WQ BC (10 mM citrate pH 7.0) and MODS-WQ
28 7H9 (pH ~6.8), performed in separate 24-well plates (**Supp. Figure S1B**). Samples were
29 incubated as in MODS-Wayne assay. Following a similar incubation process to the MODS-
30 Wayne assay, upon observing positive growth in the PZA-C well on the third day, samples
31 underwent variant-specific treatments. For MODS-WQ 7H9, samples were completed to 1

1 mL by adding 100 μ L of 7H9 media to the PZA-C well and 100 μ L of PZA (800 μ g/mL final
2 concentration) to the PZA-WQ well, followed by an additional three-day incubation.
3 Conversely, in the MODS-WQ CB procedure, samples were centrifuged, the resulting pellet
4 resuspended in 900 μ L of citrate buffer and returned to a 24-well plate. Then, 100 μ L of
5 citrate buffer was added to the PZA-C well and 100 μ L of PZA (800 μ g/mL final
6 concentration) to the PZA-WQ well, also incubated for three days.
7 Post-incubation, 100 μ L of 10% SAF was added and incubated for 5 minutes. Subsequently,
8 500 μ L was transferred to cryotubes and stored at -80°C for subsequent analysis. During the
9 analysis, triplicates of samples (100 μ L) were transferred to a 96-well plate and measured at
10 a 450nm wavelength using a spectrophotometer. Notably, MODS-WQ 7H9 samples were
11 centrifuged (13000 rpm, 2 minutes) and supernatants were analyzed. Absorbance values were
12 then interpolated against standard POA curves, ranging from 31.25 μ M to 4000 μ M (Table
13 S1). Calibration curves with 7H9 matrix underwent also previous centrifugation (13000 rpm,
14 2 minutes), with subsequent measurement of supernatants. Mixture triplicates (100 μ L) were
15 analyzed on a spectrophotometer at a 450nm wavelength to determine absorbance values, for
16 calculating linear regression on standard curves). For more details **see supplementary**
17 **methods**.

18 19 Composite reference standard for PZA susceptibility determination

20 In the absence of a singular standard test, we worked with a composite standard, based on
21 three tests: (i) the BACTEC MGIT960 (ii) conventional Wayne test, and (iii) sequencing of
22 the *pncA* gene (**Supp. Figure S2A**). Each sample was coded as 0 (sensitive) or 1 (resistant)
23 based on its response to these three standard tests. Samples were then categorized as resistant
24 (summing up to 2 or 3) or sensitive (summing up to 0 or 1) within the composite standard
25 (**Supp. Figure S2B**). Mutations were classified according to WHO catalog[27] and the
26 SuspectPZA webtool: https://biosig.lab.uq.edu.au/suspect_pza/. Both MODS-Wayne and
27 MODS-WQ were evaluated against this composite standard. More details in **supplementary**
28 **methods**.

29 30 Evaluation of PZase mutations effect on POA production

1 To assess the impact of enzyme mutations on pyrazinoic acid (POA) production, POA values
2 (μM) were plotted per sample according to mutation sites, categorized as Active Catalytic
3 Site (ACS), Metal Binding Site (MBS), PZA Binding Site (PBS), Enzymatic Core (EC), and
4 the Enzyme's Periphery (PER). POA production was compared to that of the wild-type
5 enzyme to evaluate alterations caused by mutations.

6

7 Statistical analysis

8 Continuous variables were expressed as median \pm IC95% or median and interquartile range
9 (IQR). Sample comparisons were made using non-parametric Mann-Whitney test. The
10 sensitivity and specificity of qualitative MODS-Wayne was estimated using a 2-by-2
11 contingency table. The predictive value of MODS-WQ assay was evaluated with the logistic
12 regression model to calculate the Receiver Operating Characteristic (ROC) curve, sensitivity
13 and specificity of the test. Agreement between assays and the composite standard was
14 estimated using the Kappa index. Statistical tests were performed with STATA 14 and
15 GraphPad Prism 9.0.2. The analysis of mutations was performed with the Pymol 2.5.4
16 educational license software.

17

18 **RESULTS (778 words)**

19 **Study population and MDR characterization**

20 Patient characterization is shown in **Table 1**. The MP and TBN groups exhibited no
21 significant differences in age and gender distribution. In MP group only 11.00% of patients
22 had received prior treatment, while in the TBN group, 67.00% had received it. Among the
23 264 collected samples, 78 (29.54%) were identified as MDR samples. Within the non-MDR
24 samples ($n = 186$, 70.45%), 26 demonstrated monoresistance to INH (9.8%), 15 displayed
25 monoresistance to RIF (5.7%), and 145 samples exhibited pansusceptibility (55.30%).

26

27 **Composite reference standard tests for PZA susceptibility determination**

28 In the TBN group ($n = 95$), MGIT-PZA identified 80 isolates (84.2%) as sensitive and 15
29 (15.8%) as resistant. Conversely, in the MP group ($n = 169$), 122 isolates (72.2%) were
30 sensitive, while 47 (27.8%) were resistant. The conventional Wayne test showed sensitivity
31 in 83 (87.4%) isolates and resistance in 12 (12.6%) within the TBN group; for the MP group,

1 137 (81.1%) were sensitive, and 32 (18.9%) were resistant. The *pncA* sequencing revealed
2 187 wild-type isolates (TBN: n = 71, MP: n = 116) and 69 isolates with resistant mutations,
3 (undetermined in 8 isolates). Among the mutations (n = 19), 16 were point mutations, 2
4 deletions, and 2 insertions (Table 3). Notably, 1 (1.5%) affected the promoter site, 1 (1.5%)
5 affected the catalytic site, 4 (5.8%) targeted the metal binding site, 5 (7.3%) were located in
6 the periphery, and 6 (8.7%) affected the enzyme core (**Table 2**). Following the three tests'
7 susceptibility results, a composite susceptibility was calculated for each isolate (**Table 3**).

9 **MODS-Wayne qualitative assay performance**

10 In the MP group, based on the observed color scale, 24.26% of samples were classified as
11 PZA-resistant, while 75.74% were deemed sensitive. Conversely, in the TBN group, 15.79%
12 were identified as resistant, and 84.21% were labeled as sensitive to PZA (Table S4).

13 For assessing MODS-Wayne's sensitivity and specificity, the obtained proportions were
14 compared to those derived from the composite standard. The results indicated an 87.8%
15 sensitivity (95% CI: 0.7446 - 0.9468, Wilson-Brown test) and 95.9% specificity (95% CI:
16 0.9084 - 0.9825), accompanied by a kappa index of 0.83 (p=0.001) (**Table 4**).

18 **MODS-WQ assay performance**

19 The samples were categorized into sensitive and resistant groups based on the composite
20 standard. In MODS-WQ 7H9 samples, the median POA production according calibration
21 curve (**Figure 2A**) for the sensitive group was 274 μM (IQR = 373.3 μM), contrasting with
22 92 μM (IQR = 77.95 μM) for the resistant group (**Figure 2B**). The Mann-Whitney test
23 yielded a significant p-value (<0.0001) although no pronounced discrimination was
24 observed. In MODS-WQ BC samples, the median concentration for the sensitive group was
25 2105 μM (IQR = 1672 μM), whereas for the resistant group, it was 258 μM (IQR = 1383.3
26 μM) (**Figure 2B**), revealing a strong distinction with a p-value of <0.0001.

27 ROC curve analysis displayed good performance for both variants. However, the area under
28 the curve (AUC) for MODS-WQ BC was 97.57% (95% CI: 95.30% – 99.84%), significantly
29 surpassing (p-value= 0.0037) MODS-WQ 7H9 (AUC = 82.28%, 95% CI: 71.28% - 93.28%)
30 (**Figure 2C**), indicating superior discrimination for MODS-WQ CB. The optimal cut-off
31 points associated with the ROC curves were 126.70 μM for MODS-WQ 7H9, demonstrating

1 a sensitivity of 81.25% (95% CI: 56.99% - 93.41%) and a specificity of 77.22% (95% CI:
2 66.83% - 85.07%) (**Table 4-Figure 2D**). Conversely, for MODS-WQ CB, the optimal cut-
3 off was 664.7 μM , showcasing a sensitivity of 92.31% (95% CI: 79.68% – 97.35%) and a
4 specificity of 95.93% (95% CI: 90.84% - 98.25%) (**Table 4-Figure 2D**). The kappa index
5 resulted in 0.83 (p-value=0.001) for MODS-WQ 7H9 and 0.87 (p-value=0.001) for MODS-
6 WQ BC.

7

8 **Relationship between POA production and PZase mutations.**

9 The advantage of determining resistance quantifying POA production rather than relying
10 solely on qualitative determination (MODS-Wayne) lies in establishing resistance levels and
11 correlating them with enzymatic activity. Mutations in *pncA* gene can impact in the enzyme
12 activity, so that, we assessed POA production in isolates with mutations across different
13 enzyme regions (**Figure 3A**), excluding deletions and mutations at the promoter site. The
14 **Figure 3B**, shows that mutations within the EC (I6S, Q10P, Q10R, K48T, and H71R)
15 exhibited a median POA production of 252 μM . Mutations in the MBS (D49N, H51R, H57R)
16 caused a median of 266.4 μM . A single mutation in the ACS (D8E) detected in two isolates
17 demonstrated widely varying POA values of 59 μM and 847 μM (median: 453 μM). In
18 contrast, mutations in PER (P62S, F81S) revealed a median of 644.3 μM , close to the cut-
19 off. The median POA production in wild-type isolates was 2195 μM , indicating that
20 mutations in the EC and MBS regions decreased POA levels by approximately 88.5% and
21 88.0%, respectively. Meanwhile, mutations in PER decreased POA production by
22 approximately 70.6%.

23

24 **DISCUSSION (930 words)**

25 Prompt detection of PZAr is crucial for effective TB treatment. Traditional culturing
26 methods, however, face difficulties in maintaining appropriate pH levels and inoculum
27 concentrations. In our study of 264 samples, 78 (29.54%) were identified as multidrug-
28 resistant (MDR). Among these MDR samples, 49 (62.80%) exhibited resistance to PZA, as
29 determined by a composite standard. This finding aligns with other studies reporting PZAr
30 rates of 38% to 65% in MDR TB cases [28]–[30].

1 The citrate buffer variant of the MODS-Wayne quantitative (MODS-WQ CB) assay
2 demonstrated outstanding efficacy, with a sensitivity of 92.31%, specificity of 95.93%, and
3 a positive predictive value (PPV) of 88%. In contrast, the 7H9 matrix variant showed notably
4 lower performance, with a sensitivity of 81.3%, specificity of 77.2%, and a PPV of 42%.
5 The spectrophotometric measurement of POA in 7H9 media is complicated by the presence
6 of bovine serum albumin (BSA), which is added to foster mycobacterial growth [31]. BSA
7 exhibits a higher affinity for POA compared to PZA at concentrations greater than 0.4%,
8 significantly raising the POA minimum inhibitory concentration (MIC) [32]. Additionally,
9 BSA tends to interact with various ligands, particularly divalent metallic cations, including
10 copper and zinc. Notably, BSA shows a strong affinity for Fe²⁺ and Fe³⁺ ions at neutral pH,
11 particularly near tryptophan residues, similar to lactoferrin [33]–[35]. This affinity for POA
12 and iron could hinder the formation of the SAF-POA complex, leading to variable sensitivity
13 and specificity in our assay results.

14 The use of the citrate buffer (CB) matrix in our study not only simplified colorimetric
15 observation but also ensured more accurate absorbance detection by minimizing medium-
16 related biases. This contrasts with previous research by Meinzen et al. [36], which quantified
17 POA using similar matrices but relied on samples grown on agar and stored in a culture bank.
18 Our approach, in contrast, involves direct sampling from patient sputum. Meinzen et al.
19 reported sensitivity and specificity rates of 96.0% and 97.4%, respectively, using a PZA
20 concentration of 400 µg/mL, and 90.5% and 94.4% for 800 µg/mL PZA, achieving an area
21 under the curve (AUC) of over 93.0% for both concentrations.

22 While our study did not match the high sensitivity and specificity levels reported by Meinzen
23 et al., our approach, using direct patient samples, significantly reduced the diagnosis time.
24 On average, we achieved a 13.5-day reduction compared to traditional methods like the
25 Wayne Test (30-45 days) [24] and the MODS-Wayne (mean 19.5 days). Although the
26 performance of MODS-Wayne and our quantitative MODS-WQ were comparable (Kappa
27 indices of 0.83 and 0.87, respectively), it's crucial to note the variability in MODS-Wayne
28 results across different sample types (MP, TBN, and previous study), mainly due to observer
29 subjectivity. This variability could affect reproducibility [37], [38], an issue potentially
30 resolved by our quantitative MODS-WQ variant. Further studies focusing on reproducibility
31 are essential to validate our findings.

1 POA production is a key indirect marker of PZase enzymatic activity, which can be
2 influenced by protein mutations [39]–[41]. In our study, mutations within the PZase enzyme
3 were categorized into four areas: ACS, MBS, EC, and PER. We observed that mutations in
4 the EC and MBS regions led to significantly lower POA levels (252.7 μ M and 266.4 μ M,
5 respectively) compared to those in PER, which produced POA levels near the 665 μ M cut-
6 off ($p < 0.01$ and $p < 0.05$, respectively). Supporting our findings, Sheen et al. [39] reported that
7 mutations in MBS (H51R, D49N) resulted in a substantial decrease in enzymatic activity
8 (99%) in terms of POA production, compared to the wild-type H37Rv strain. Similarly,
9 mutations near active sites (core) led to a 72% reduction, while peripheral mutations caused
10 a 47% decrease in enzymatic activity.

11 Our study aligns with previous research, showing a marked decrease in POA production in
12 samples with mutations in the EC and MBS regions (88%) compared to wild-type samples.
13 In samples with PER mutations, this reduction was slightly lower at 70.6%. It's important to
14 distinguish our methodology from that of Sheen et al., who directly quantified POA
15 production in purified enzymes in vitro, free from external influences on the mycobacteria's
16 POA output.

17 The release of POA is influenced not only by enzymatic efficiency but also by external
18 factors, such as efflux pump activity, which plays a crucial role in POA transport. Zimic et
19 al. [42] highlighted a strong link between POA efflux rate and PZA resistance. Their study
20 showed that resistant isolates have lower efflux rates, while sensitive isolates display higher
21 rates [40], allowing for the prediction of resistance with high sensitivity and specificity.

22 Enzymatic activity alone accounts for only 27.3% of PZA resistance cases [39], while the
23 POA efflux rate contributes to around 51% [41]. Notably, 70-90% of PZA-resistant strains
24 show mutations in the *pncA* gene and its promoter region [43]–[45]. This suggests that a
25 significant portion of resistant strains may involve other, yet unidentified, mechanisms
26 contributing to PZA resistance.

27 Mutations in other genes, such as *panD* and *rpsA*, might also play a role in PZA resistance,
28 as could the presence of heteroresistant strains with varying susceptibility genotypes [28]–
29 [30], [46]. Interestingly, not all mutations in the *pncA* gene are linked to PZA resistance;
30 some result in PZA-sensitive strains [45]. This complexity is illustrated in Figure 4.
31 Consequently, assessing POA production in the culture medium offers a more comprehensive

1 understanding of PZA susceptibility compared to isolated enzymatic activity measurements
2 alone.

3 In conclusion, the MODS-Wayne quantitative (MODS-WQ) assay in citrate buffer, with its
4 exceptional predictive capacity (ROC: 97.6%, Kappa index: 0.87) and an average incubation
5 period of just 13 days from sputum sample culturing, represents a significant advancement
6 in TB diagnostics. This assay comprehensively accounts for factors like enzymatic efficiency
7 and efflux pump activity, crucial for determining POA production in the medium.
8 Additionally, its cost-effectiveness is particularly beneficial for resource-limited laboratories,
9 especially in developing regions.

10

11

12 ***Acknowledgments:*** The authors extend their gratitude to the Hospital Nacional Dos de Mayo,
13 Lima, Peru, and the Regional Tuberculosis Reference Laboratory, Callao, Lima, Peru for
14 providing the sputum samples.

15 ***Author contributions***

16 Contributors ETG, MZ and PS conceived and designed the study. ETG analyzed data,
17 generated figures and tables and wrote the manuscript. ETG, and RA collected clinical
18 samples and data. ETG, and RA performed experiments, and analyzed data. PS, RG, LG and
19 MZ contributed to data interpretation and critically reviewed the manuscript. All authors
20 approved the final manuscript for submission.

21 **The authors declare no conflicts of interest.**

22

23

24

25

26

27

28

29

30

31

1 REFERENCES

- 2 [1] World Health Organization, ‘Global tuberculosis report 2020’, Geneva, 2020.
- 3 [2] P. Sinha, K. R. Jacobson, C. R. Horsburgh Jr., and C. Acuña-Villaorduña, ‘At Long
4 Last: Short, All-Oral Regimens for Multidrug-Resistant Tuberculosis in the United
5 States’, *Open Forum Infectious Diseases*, vol. 10, no. 4, p. ofad177, Apr. 2023, doi:
6 10.1093/ofid/ofad177.
- 7 [3] H. Zhang *et al.*, ‘Characterization of Mycobacterium tuberculosis
8 nicotinamidase/pyrazinamidase’, *FEBS Journal*, vol. 275, no. 4, pp. 753–762, Feb.
9 2008, doi: 10.1111/j.1742-4658.2007.06241.x.
- 10 [4] Collaborative Group for the Meta-Analysis of Individual Patient Data in MDR-TB
11 treatment–2017 *et al.*, ‘Treatment correlates of successful outcomes in pulmonary
12 multidrug-resistant tuberculosis: an individual patient data meta-analysis’, *Lancet*, vol.
13 392, no. 10150, pp. 821–834, 08 2018, doi: 10.1016/S0140-6736(18)31644-1.
- 14 [5] M. G. Whitfield *et al.*, ‘A Global Perspective on Pyrazinamide Resistance: Systematic
15 Review and Meta-Analysis’, *PLOS ONE*, vol. 10, no. 7, p. e0133869, Jul. 2015, doi:
16 10.1371/journal.pone.0133869.
- 17 [6] S. Mok *et al.*, ‘Overcoming the Challenges of Pyrazinamide Susceptibility Testing in
18 Clinical Mycobacterium tuberculosis Isolates’, *Antimicrob Agents Chemother*, vol. 65,
19 no. 8, p. e0261720, Jul. 2021, doi: 10.1128/AAC.02617-20.
- 20 [7] G. L. Woods *et al.*, *Susceptibility Testing of Mycobacteria, Nocardiae, and Other*
21 *Aerobic Actinomycetes*, 2nd ed. in CLSI Standards: Guidelines for Health Care
22 Excellence. Wayne (PA): Clinical and Laboratory Standards Institute, 2011. Accessed:
23 Mar. 21, 2020. [Online]. Available: <http://www.ncbi.nlm.nih.gov/books/NBK544374/>
- 24 [8] Y. Zhang, S. Permar, and Z. Sun, ‘Conditions that may affect the results of susceptibility
25 testing of Mycobacterium tuberculosis to pyrazinamide’, *J. Med. Microbiol.*, vol. 51,
26 no. 1, pp. 42–49, Jan. 2002, doi: 10.1099/0022-1317-51-1-42.
- 27 [9] P. Chedore, L. Bertucci, J. Wolfe, M. Sharma, and F. Jamieson, ‘Potential for erroneous
28 results indicating resistance when using the Bactec MGIT 960 system for testing
29 susceptibility of Mycobacterium tuberculosis to pyrazinamide’, *J. Clin. Microbiol.*, vol.
30 48, no. 1, pp. 300–301, Jan. 2010, doi: 10.1128/JCM.01775-09.
- 31 [10] World Health Organization, ‘Technical manual for drug susceptibility testing of
32 medicines used in the treatment of tuberculosis.’, Switzerland, 2018.
- 33 [11] Wayne G. L., ‘Simple Pyrazinamidase and Urease Tests for Routine Identification of
34 Mycobacteria’, *Am Rev Respir Dis*, vol. 109, no. 1, pp. 147–151, Jan. 1974, doi:
35 10.1164/arrd.1974.109.1.147.
- 36 [12] A. Martin, A. Cubillos-Ruiz, A. Von Groll, P. Del Portillo, F. Portaels, and J. C.
37 Palomino, ‘Nitrate reductase assay for the rapid detection of pyrazinamide resistance in
38 Mycobacterium tuberculosis using nicotinamide’, *J Antimicrob Chemother*, vol. 61, no.
39 1, pp. 123–127, Jan. 2008, doi: 10.1093/jac/dkm418.

- 1 [13] L. Caviedes *et al.*, ‘Rapid, Efficient Detection and Drug Susceptibility Testing of
2 Mycobacterium tuberculosis in Sputum by Microscopic Observation of Broth Cultures’,
3 *J Clin Microbiol*, vol. 38, no. 3, pp. 1203–1208, Mar. 2000.
- 4 [14] D. C. Alexander, J. H. Ma, J. L. Guthrie, J. Blair, P. Chedore, and F. B. Jamieson, ‘Gene
5 Sequencing for Routine Verification of Pyrazinamide Resistance in Mycobacterium
6 tuberculosis: a Role for *pncA* but Not *rpsA*’, *J Clin Microbiol*, vol. 50, no. 11, pp. 3726–
7 3728, Nov. 2012, doi: 10.1128/JCM.00620-12.
- 8 [15] E. Sodja, S. Koren, N. Toplak, S. Truden, and M. Žolnir-Dovč, ‘Next-generation
9 sequencing to characterise pyrazinamide resistance in Mycobacterium tuberculosis
10 isolates from two Balkan countries’, *Journal of Global Antimicrobial Resistance*, vol.
11 29, pp. 507–512, Jun. 2022, doi: 10.1016/j.jgar.2021.09.019.
- 12 [16] S. J. Modlin, T. Marbach, J. Werngren, M. Mansjö, S. E. Hoffner, and F. Valafar,
13 ‘Atypical Genetic Basis of Pyrazinamide Resistance in Mono-resistant Mycobacterium
14 tuberculosis’, *Antimicrobial Agents and Chemotherapy*, vol. 65, no. 6, p.
15 10.1128/aac.01916-20, May 2021, doi: 10.1128/aac.01916-20.
- 16 [17] M. T. Khan, A. U. Rehman, M. Junaid, S. I. Malik, and D.-Q. Wei, ‘Insight into novel
17 clinical mutants of *RpsA*-S324F, E325K, and G341R of Mycobacterium tuberculosis
18 associated with pyrazinamide resistance’, *Computational and Structural Biotechnology
19 Journal*, vol. 16, pp. 379–387, Jan. 2018, doi: 10.1016/j.csbj.2018.09.004.
- 20 [18] S. Zhang, J. Chen, W. Shi, W. Liu, W. Zhang, and Y. Zhang, ‘Mutations in *panD*
21 encoding aspartate decarboxylase are associated with pyrazinamide resistance in
22 Mycobacterium tuberculosis’, *Emerging Microbes & Infections*, vol. 2, no. 1, pp. 1–5,
23 Jan. 2013, doi: 10.1038/emi.2013.38.
- 24 [19] P. Gopal *et al.*, ‘Pyrazinamide triggers degradation of its target aspartate
25 decarboxylase’, *Nat Commun*, vol. 11, no. 1, Art. no. 1, Apr. 2020, doi:
26 10.1038/s41467-020-15516-1.
- 27 [20] S. M. Ramirez-Busby *et al.*, ‘A Multinational Analysis of Mutations and Heterogeneity
28 in *PZase*, *RpsA*, and *PanD* Associated with Pyrazinamide Resistance in M/XDR
29 Mycobacterium tuberculosis’, *Sci Rep*, vol. 7, Jun. 2017, doi: 10.1038/s41598-017-
30 03452-y.
- 31 [21] P. Sheen *et al.*, ‘A multiple genome analysis of Mycobacterium tuberculosis reveals
32 specific novel genes and mutations associated with pyrazinamide resistance’, *BMC
33 Genomics*, vol. 18, p. 769, Oct. 2017, doi: 10.1186/s12864-017-4146-z.
- 34 [22] M. Shah, V. Chihota, G. Coetzee, G. Churchyard, and S. E. Dorman, ‘Comparison of
35 laboratory costs of rapid molecular tests and conventional diagnostics for detection of
36 tuberculosis and drug-resistant tuberculosis in South Africa’, *BMC Infect Dis*, vol. 13,
37 p. 352, Jul. 2013, doi: 10.1186/1471-2334-13-352.
- 38 [23] R. Baddam *et al.*, ‘Analysis of mutations in *pncA* reveals non-overlapping patterns
39 among various lineages of Mycobacterium tuberculosis’, *Scientific Reports*, vol. 8, no.
40 1, pp. 1–9, Mar. 2018, doi: 10.1038/s41598-018-22883-9.
- 41 [24] R. Alcántara *et al.*, ‘MODS-Wayne, a Colorimetric Adaptation of the Microscopic-
42 Observation Drug Susceptibility (MODS) Assay for Detection of Mycobacterium

- 1 tuberculosis Pyrazinamide Resistance from Sputum Samples’, *Journal of Clinical*
2 *Microbiology*, vol. 57, no. 2, Feb. 2019, doi: 10.1128/JCM.01162-18.
- 3 [25] P. T. Kent, *Public health mycobacteriology: a guide for the level III laboratory*.
4 Atlanta, Ga.: U.S. Dept. of Health and Human Services, Public Health Service, Centers
5 for Disease Control, 1985. Accessed: Jun. 15, 2023. [Online]. Available:
6 <https://search.library.wisc.edu/catalog/999574039202121>
- 7 [26] Universidad Peruana Cayetano Heredia, ‘MODS: A User Guide. Microscopic
8 Observation Drug Susceptibility Test’, Universidad Peruana Cayetano Heredia, Lima,
9 Perú, 2008.
- 10 [27] World Health Organization, ‘Catalogue of mutations in Mycobacterium tuberculosis
11 complex and their association with drug resistance.’, Geneva, 2021.
- 12 [28] K. Stoffels, V. Mathys, M. Fauville-Dufaux, R. Wintjens, and P. Bifani, ‘Systematic
13 Analysis of Pyrazinamide-Resistant Spontaneous Mutants and Clinical Isolates of
14 Mycobacterium tuberculosis’, *Antimicrob Agents Chemother*, vol. 56, no. 10, pp. 5186–
15 5193, Oct. 2012, doi: 10.1128/AAC.05385-11.
- 16 [29] R. I. Calderón *et al.*, ‘Prevalence of pyrazinamide resistance and Wayne assay
17 performance analysis in a tuberculosis cohort in Lima, Peru’, *The International Journal*
18 *of Tuberculosis and Lung Disease*, vol. 21, no. 8, pp. 894–901, Aug. 2017, doi:
19 10.5588/ijtld.16.0850.
- 20 [30] Y. Pang *et al.*, ‘Prevalence and molecular characterization of pyrazinamide resistance
21 among multidrug-resistant Mycobacterium tuberculosis isolates from Southern China’,
22 *BMC Infectious Diseases*, vol. 17, no. 1, p. 711, Nov. 2017, doi: 10.1186/s12879-017-
23 2761-6.
- 24 [31] M. Lynn, A. R. Wilson, and M. Solotorovsky, ‘Role of bovine serum albumin in the
25 nutrition of Mycobacterium tuberculosis.’, *Appl Environ Microbiol*, vol. 38, no. 5, pp.
26 806–810, Nov. 1979.
- 27 [32] Y. Zhang and D. Mitchison, ‘The curious characteristics of pyrazinamide: a review’,
28 *The International Journal of Tuberculosis and Lung Disease*, vol. 7, no. 1, pp. 6–21,
29 Jan. 2003.
- 30 [33] X. Xu, L. Zhang, D. Shen, H. Wu, and Q. Liu, ‘Oxygen-dependent oxidation of Fe(II)
31 to Fe(III) and interaction of Fe(III) with bovine serum albumin, leading to a hysteric
32 effect on the fluorescence of bovine serum albumin’, *J Fluoresc*, vol. 18, no. 1, pp. 193–
33 201, Jan. 2008, doi: 10.1007/s10895-007-0263-4.
- 34 [34] H. M. Ueno, H. Urazono, and T. Kobayashi, ‘Serum albumin forms a lactoferrin-like
35 soluble iron-binding complex in presence of hydrogen carbonate ions’, *Food Chem*, vol.
36 145, pp. 90–94, Feb. 2014, doi: 10.1016/j.foodchem.2013.07.143.
- 37 [35] S. Rudra, S. Dasmandal, C. Patra, A. Kundu, and A. Mahapatra, ‘Binding affinities of
38 Schiff base Fe(II) complex with BSA and calf-thymus DNA: Spectroscopic
39 investigations and molecular docking analysis’, *Spectrochimica Acta Part A: Molecular*
40 *and Biomolecular Spectroscopy*, vol. 166, pp. 84–94, Sep. 2016, doi:
41 10.1016/j.saa.2016.04.050.

- 1 [36] C. Meinzen *et al.*, ‘A quantitative adaptation of the Wayne test for pyrazinamide
2 resistance’, *Tuberculosis*, vol. 99, pp. 41–46, Jul. 2016, doi:
3 10.1016/j.tube.2016.03.011.
- 4 [37] P. Talkad and R. Metoyer, *Replication and Transparency of Qualitative Research from
5 a Constructivist Perspective*. OSF Preprints, 2019. doi: 10.31219/osf.io/6efvp.
- 6 [38] R. Tuval-Mashiach, ‘Is replication relevant for qualitative research?’, *Qualitative
7 Psychology*, vol. 8, no. 3, pp. 365–377, 2021, doi: 10.1037/qup0000217.
- 8 [39] P. Sheen *et al.*, ‘Effect of pyrazinamidase activity on pyrazinamide resistance in
9 Mycobacterium tuberculosis’, *Tuberculosis (Edinb)*, vol. 89, no. 2, pp. 109–113, Mar.
10 2009, doi: 10.1016/j.tube.2009.01.004.
- 11 [40] M. Zimic, P. Fuentes, R. H. Gilman, A. H. Gutiérrez, D. Kirwan, and P. Sheen,
12 ‘Pyrazinoic acid efflux rate in Mycobacterium tuberculosis is a better proxy of
13 pyrazinamide resistance’, *Tuberculosis (Edinb)*, vol. 92, no. 1, pp. 84–91, Jan. 2012,
14 doi: 10.1016/j.tube.2011.09.002.
- 15 [41] P. Sheen *et al.*, ‘pncA gene expression and prediction factors on pyrazinamide
16 resistance in Mycobacterium tuberculosis’, *Tuberculosis (Edinb)*, vol. 93, no. 5, pp.
17 515–522, Sep. 2013, doi: 10.1016/j.tube.2013.03.005.
- 18 [42] M. Zimic *et al.*, ‘A new approach for pyrazinamide susceptibility testing in
19 Mycobacterium tuberculosis’, *Microb. Drug Resist.*, vol. 18, no. 4, pp. 372–375, Aug.
20 2012, doi: 10.1089/mdr.2011.0207.
- 21 [43] M. T. Khan *et al.*, ‘Pyrazinamide resistance and mutations in pncA among isolates of
22 Mycobacterium tuberculosis from Khyber Pakhtunkhwa, Pakistan’, *BMC Infectious
23 Diseases*, vol. 19, no. 1, p. 116, Feb. 2019, doi: 10.1186/s12879-019-3764-2.
- 24 [44] P. W. Ei *et al.*, ‘Pyrazinamide resistance and pncA mutations in drug resistant
25 Mycobacterium tuberculosis clinical isolates from Myanmar’, *Tuberculosis*, vol. 125,
26 p. 102013, Dec. 2020, doi: 10.1016/j.tube.2020.102013.
- 27 [45] J. Shi *et al.*, ‘Pyrazinamide Resistance and Mutation Patterns Among Multidrug-
28 Resistant Mycobacterium tuberculosis from Henan Province’, *Infect Drug Resist*, vol.
29 13, pp. 2929–2941, Aug. 2020, doi: 10.2147/IDR.S260161.
- 30 [46] J. Werngren, E. Alm, and M. Mansjö, ‘Non-pncA Gene-Mutated but Pyrazinamide-
31 Resistant Mycobacterium tuberculosis: Why Is That?’, *Journal of Clinical
32 Microbiology*, vol. 55, no. 6, pp. 1920–1927, Jun. 2017, doi: 10.1128/JCM.02532-16.

33

34

35


36

37


38

1 FIGURES LEGENDS

2 **Figure 1. Study design.** Sputum samples were collected from two studies: MP and TBN.
3 Only samples positive for bacilloscopy were included. Samples were used in four
4 experimental assays: MODS (MDR characterization), MODS-Wayne (qualitative assay for
5 PZA resistance), MODS-WQ (quantitative variant) and three reference standards. The
6 diagnostic value for MODS-Wayne and MODS-WQ was calculated using the standard
7 composite.




1 **Figure 2. Performance of the MODS-WQ variants.** A) Standard curves of pyrazinoic acid
 2 dissolved in two types of matrices. Curves are result of 3 repetitions and their respective
 3 standard deviations. The equation for calculating POA concentration (POAc) is shown. B)
 4 Violin graph of the classification of TBN and CB samples according to the composite
 5 standard test and POA values in μM obtained with the MODS-WQ 7H9 and CB respectively.
 6 The medians of POA are shown for each group. C) ROC curves and p-values obtained with
 7 the two variants MODS-WQ BC and MODS-WQ 7H9, statistical differences are observed
 8 between both variants. D) Violin graph of classifieds according to MODS-WQ 7H9 and
 9 MODS-WQ BC showing the cut-off point for each variant.
 10 * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$
 11




12

1 **Figure 3. Effect of pyrazinamidase mutations on POA production.** A) 3-D structure of
 2 pyrazinamidase, the active sites are depicted in green, violet and red. Enzyme core area near
 3 to active sites and the periphery distant to active sites. Green: Binding site for PZA formed
 4 by the amino acids Phe13 and Trp68 (A.1) Red: Catalytic triad, composed of Asp8, Lys96,
 5 and Cys138 (A.2) Purple: Metal-binding site (Fe⁺), consisting of Asp49, His51, and His57
 6 (A.3). B) Distribution of POA values at each mutation site, as well as in wild type samples.
 7 The MODS-WQ cut-off (665 μM) is indicated by a dashed line. EC: Enzymatic Core, N=22;
 8 MBS: Metal-Binding Site, N=11; ACS: Active Catalytic Site, N=2; Periphery: N=4, WT:
 9 Wild type, N=116. Mann-Whitney test, * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001,
 10 ns: not significant.



11 **Figure 4. Complexity of PZA susceptibility phenotype.** Variability in susceptibility
 12 phenotype is result of different factors such as, lack of specific PZA target, the large number
 13 of potential mutations throughout the coding and promoter sequence of *pncA*, the diversity
 14 of phenotypic methods and errors that they entail, among others. In this figure we can observe
 15 the 4 types of isolates according to the susceptibility phenotype.
 16



17
 18

1 **TABLES**

	MP	TBN	Total
	N=169	N= 95	(N= 264)
Age (years), median (IQR)	36 (27.5)	29 (21)	33 (24.5)
Sex, n (%)			
Males	96 (57.0)	66 (69.0)	162 (61.4)
Females	54 (32)	27 (31)	81 (30.7)
N.D	19 (11)	2 (2)	21 (7.8)
Treatment, n (%)			
No treated	83 (49.1)	29 (30.5)	112 (42.4)
Treated	18 (10.7)	64 (67.4)	82 (31.1)
N.D	68 (40.2)	2 (2.1)	70 (26.5)
MDR	52 (30.8)	26 (27.4)	78 (29.5)
Mono-resistant	Rifampicin	15 (8.9)	0 (0.0)
	Isoniazid	20 (11.8)	6 (6.3)
Sensitive	82 (48.5)	63 (66.3)	145 (54.9)

2 **Table 1. Baseline characteristics of the samples**
3 Abbreviations: MP, samples collected in the study 2015-2016; TBN: Samples collected in the study 2017-
4 2018; IQR, Interquartile range; MDR, Multidrug resistance; N.D, no determined.

5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21

Table 2. Categorization and characterization of the mutations found in the study.

Mutation	N° Samples		Location in the enzyme	Susceptibility phenotype		Final susceptibility
	TBN	MP		WHO ^a	SuspectPZ A ^b	
I6S	-	2	EC	Uncertain	R	R
D8E	2	2	ACS	A w RI	R	R
Q10P	-	2	EC	A w R	R	R
Q10R	2	12	EC	A w R	R	R
A30T	1	-	PER	-	S	S
K48T	3	6	EC	Uncertain	R	R
D49N	2	1	MBS	A w RI	R	R
H51R	-	12	MBS	A w R	R	R
H57L	3	-	MBS	-	R	R
H57R	-	1	MBS	A w R	R	R
P62S	-	1	PER	A w RI	R	R
H71R	-	2	EC	A w R	R	R
F81S	5	2	PER	Uncertain	R	R
H82D	1	-	PER	Uncertain	R	R
V180F	3	-	PER	A w R	R	R
Prom A-11G	-	1	Promoter	A w R	-	R
407insA/ 461insA	1	-	EC	A w RI	-	R
Δ456-466	-	3	PER	A w RI	-	R
Δ375-389	-	1	PER	A w R	-	R
WT	71	116		S	S	S

R: Resistant; S: sensitive; Uncertain: Ambiguous association, A w R: associated with resistance, A w RI: associated with intermediate resistance

^a Mutation classification according to the WHO catalog

b Mutation classification according to the SuspectPZA web tool

c Mutation classification considering *a* and *b*

1
2
3
4
5
6
7
8

Table 3. PZA susceptibility results according to composite standard

Phenotype	Sum value	Standard composite				Total (264)
		MP (169)		TBN (95)		
		N (%)	N Subtotal	N (%)	N Subtotal	
Resistant	0 (%)	30 (17.8)	46 (27.2)	8 (8.4)	16 (16.8)	62 (23.5)
	1 (%)	16 (9.5)		8 (8.4)		
Sensitive	2 (%)	10 (5.9)	123 (72.8)	10 (10.5)	79 (83.2)	202 (76.5)
	3 (%)	113 (66.9)		69 (72.6)		


9
10
11
12
13
14
15
16
17
18
19





1



Table 4. Comparative performance of each MODS-Wayne and MODS-WQ methodology with their variants in CB and 7H9

Method/Sample		Ratio	Sensitivity [95% CI]	Ratio	Specificity [95% CI]	PPV	NPV	Ki
MODS-Wayne	MP	36/41	0.878 [0.745-0.947]	118/123	0.959 [0.901-0.983]	0.88	0.95	0.83
	TBN	10/16	0.625 [0.387-0.815]	74/79	0.937 [0.860-0.973]	0.67	0.93	0.57
MODS-WQ	MP-BC <665 µM	38/41	0.923 [0.797-0.974]	118/123	0.959 [0.908-0.983]	0.88	0.98	0.87
	TBN-7H9 <127 µM	13/16	0.813 [0.569-0.934]	61/79	0.772 [0.668-0.851]	0.42	0.95	0.43

Measures of accuracy (sensitivity, specificity, positive predictive value PPV, negative predictive value NPV) and confidence intervals were observed for each type of test evaluated.









a Resistant with mutations in *pncA*

b Resistant with wild-type *pncA*

c Sensitive with mutations in *pncA*

d Sensitive with wild-type *pncA*