Title: Exploring athlete pain assessment experiences and priorities; A two-part qualitative series of

athlete and physiotherapist interactions.

Part One. "Gauging and discerning" - Athlete & physiotherapist pain assessment experiences and

interactions.

Abstract

Objectives: To explore the content (subjective questions, objective tools and outcome measures) and

discuss the nature (qualitative elements and wider considerations) of the athlete pain assessment by

facilitating shared understandings of athlete and sports physiotherapists.

Design; Qualitative Research using a hermeneutic phenomenological approach.

Methods: We carried out focus groups comprising a deliberate criterion sample using a constructivist

perspective. We developed a topic guide and used reflexive thematic analysis. We developed codes,

candidate themes and finalised themes iteratively, and employed a critical friend to add depth to our

analysis. Our paper follows the consolidated criteria for reporting qualitative research (COREQ)

guidelines.

Results: We completed five focus groups comprising twelve athletes (five female, seven male) and

four sports physiotherapists (four male). Three final themes (and eight subthemes) were created; (i)

Measures, Scales and Dimensions (value and limitations of tools and scales, multidimensional

methods, making sense and interpreting), (ii) Connect, Listen and Learn (the pain interview and

athlete's story, forging the athlete-clinician connection), (iii) Lighthouse in the Storm: providing

direction for athletes in pain. (information overload and indecision, a beacon of direction; the role of

the physiotherapist, the burden of expectation; challenges for physiotherapists)

Conclusion: We described and explored the phenomena of pain assessment in sport including current

pain assessment strategies. Comprehensive multidimensional assessment methods that preserve the

athlete-clinician therapeutic relationship and facilitate optimal communication are priorities for future

research and practice.

Keywords

'Pain Measurement', 'Musculoskeletal Pain', 'Athletic Injuries' 'Athletic Performance' 'Focus Groups', 'Pain Management'.

Introduction

A comprehensive biopsychosocial pain assessment that includes affective (emotional), cognitive (understandings and appraisals) and socioenvironmental (sport, home, work/school, social, cultural and environmental) as well as the more frequently used neurophysiological (pain characteristics and qualities) and biomechanical local and wider biomechanical considerations) aspects has been proposed to capture the full extent of an athlete's pain experience. [1][2] Understanding the impact of these aspects on an athlete's sport and wider life facilitates the planning and delivery of more effective management strategies.[3] In a recent scoping review, we identified gaps in pain assessment practice specifically around the affective, cognitive and socioenvironmental aspects of pain assessment.[4] Contemporary models outline the impact of wider aspects such as stress, sports environment, support networks and coping strategies on athlete pain experience and injury prevention.[5][6] Clear and effective communication allows athletes to develop relationships with their support network and to access coping strategies they may not be able to otherwise. This is in line with contemporary research that highlights the importance of effective communication between all members of the athlete support staff in injury prevention.[7] In this series we explore one of the key relationships and communication opportunities in an athlete's pain journey, the athlete-physiotherapist pain assessment. To date, little has been published exploring athlete pain assessment, and to the best of our knowledge, this is the first study to combine both athlete and physiotherapist experiences and perspectives. Our objectives were to explore the content (subjective questions, objective tools and outcome measures) and discuss the nature (qualitative elements and wider considerations) of an assessment for athletes with upper and lower limb pain

Methods

We conducted a qualitative study with mixed focus groups that combined athletes and sports

physiotherapists and followed the consolidated criteria for reporting qualitative research (COREQ)

guidelines.[8] Ethical permission was granted for our study by the UCD Human Research Ethics

Committee (LS-22-40-Purcell-Caulfield).

Positionality statement and team background.

We viewed perceptions of athlete upper and lower limb pain and pain assessment from a

constructivist perspective taking a pragmatic approach and drawing on the relevant knowledge from

elements of empirico-analytical, interpretive and critical research paradigms.[9] As the primary

interviewer I (CP) recognise that being a sports physiotherapist and PhD athlete pain researcher

influences the dynamic of discussions, I am unavoidably enmeshed, and I embrace this to understand

athlete and clinician upper and lower limb pain experiences with a reflexive approach. [9][10]. GvO PhD

candidate and sports physiotherapist with seventeen years of experience acted as moderator for the

first two focus groups to ensure all voices were heard and took field notes of the setting and context

and IOC Athlete Pain domains discussed. [1] BC a Professor in physiotherapy with twenty years of

research experience acted as a neutral observer for the first session. CBW, Physiotherapist and PhD

researcher in the field of chronic conditions with experience in the conduct of focus groups, acted as a

critical friend for the qualitative analysis.

Participants

We recruited a deliberate criterion-based sample of: (i) athletes (of varying age, gender, sport, and

competition level) through university and local sports clubs and (ii) sports physiotherapists (with a

minimum of three years of postgraduate experience working with athletes as part of their weekly

caseload from a variety of sports and clinical settings) through the Irish Society of Chartered

Physiotherapists in Sports and Exercise Medicine group via email and other communication channels

and networks. All athletes had been assessed for pain in the upper or lower limb in the past year, and

all sports physiotherapists had assessed such patients also. To facilitate engagement we offered

participants their preference of face-to-face and online (zoom) focus group settings (which were run

separately). We used a reflexive sampling strategy with data collection and analysis following each

focus group informing the total sample size. We ceased sampling and recruitment when sufficient

depth of information on athlete pain experience and athlete-physiotherapist pain assessment

experience was gathered rather than seeking objective data saturation in line with our active

constructivist approach.[11]

Protocol

We sent participants a pre-participation questionnaire to capture demographic information including

their sports background and previous pain and injury history. We asked physiotherapists about their

experience working in sports, additional qualifications and any current or prior sports participation.

We developed a topic guide (Appendix A) using an established step-by-step approach; brainstorming,

ordering, timing and phrasing of questions, obtaining feedback from peers, revising and piloting.[12]

The topic guide moved from broad questions concerning athlete pain understandings and experiences

to reflections on current pain assessment practice before identifying priorities for an athlete upper and

lower limb pain assessment framework. We altered the sequencing of questions slightly in subsequent

focus groups in line with an iterative and reflexive approach.[10] We allocated two hours for the

sessions to ensure enough time for all components. We set ground rules establishing confidentiality,

equity and respect for diverse views and opinions before each group. We moderated the power

imbalance between physiotherapists and athletes by emphasising the importance of the athlete's voice

asking for athletes' opinions to be shared before clinicians. Where possible we ensured athletes and

physiotherapists who previously worked together were separated. Our initial successful pilot focus

group of two athletes was followed up with four focus groups combining athletes and physiotherapists

in line with methodological recommendations to facilitate interaction through dialogue and capture

various perspectives in a shared space.[13]

Analysis

We completed the transcription verbatim and transcripts were checked against the audio recordings

for accuracy. The full, uncoded and anonymised set of transcripts has been published in an online data

repository (10.17632/t47tw94mzd). We chose reflexive thematic analysis as it was congruent with

both our data collection and epistemological position.[10] Following initial observations and notes CP

carried out initial coding of the entire transcripts (both semantic and latent). During the data analysis

process, we assigned athletes alphanumeric participant IDs beginning with the letter "A" and

physiotherapists alphanumeric IDs beginning with ''P''. Athletes and physiotherapists were talking

about and experiencing the same concepts so we merged the data for coding and analysis. Participant

IDs were known only to CP, the lead researcher. Once participant IDs were allocated and analysis was

complete all records of participant information were deleted ensuring anonymity Athlete and

physiotherapist data were merged for coding and analysis. Subsequent analysis was completed to

identify codes that were present in (i) athletes only, (ii) physiotherapists only and (iii) codes that were

present in both athletes and physiotherapists. One-third of the transcripts were given to CBW to code

independently and we discussed, compared and refined initial codes using a critical friend approach to

incorporate breadth and variation in experiences and perspectives.[14] We updated coding where

necessary, with similar codes being highlighted and grouped to form rough clusters of candidate

themes. The team reviewed candidate themes in the context of the research question with

consideration given to the data available to support each theme. We merged some themes and

adjusted others at this refinement stage. We developed an initial thematic map of the refined themes

before we applied titles and created a final thematic map. (Appendix B) In this paper, Part One of this

series, we report on data primarily from questions focused on athlete pain assessment experience.

Results and Discussion

Table 1 – Participant Demographics inserted here

Participant demographics are summarised in Table 1. All final themes exploring athlete pain

assessment included a mixture of codes from all three categories (i, ii and iii in the data analysis

section above). Figure 1 displays the finalised themes and codes. Figure 2 is a thematic map of the

themes and subthemes relevant to athlete pain assessment.

Figures 1 – Themes (Measures, Scales & Dimensions; Connect Listen & Learn; Lighthouse in

the Storm: providing direction for athletes in pain) and codes inserted here.

Figure 2 – Themes & Subthemes (and thematic map) inserted here.

Theme 1. Measures Scales and Dimensions

Theme 1 includes three subthemes; 1.1 – value and limitations of tools and scales, 1.2 –

multidimensional methods, and 1.3 – making sense and interpreting. Despite positive advances in

identifying and managing sports-related pain the nebulous nature of pain experience proves

challenging when it comes to accurate and comprehensive assessment.[3] Participants in this study

contrasted the value of tangible measures and scales which add a level of objectivity to the subjective

experience of pain with the limitations of currently available tools and scales for assessing athlete

pain. Additionally, the need for wider biopsychosocial and multidimensional pain assessment

strategies is firmly established in the literature, something with which participants concurred but

rarely encountered.[2 4]

1.1 Value and limitations of tools and scales

Participants discussed aspects of a standard physiotherapy pain assessment when recalling previous

experiences, including observation of the athlete and measures of physical function such as range of

motion, strength and performance markers. Physiotherapists and athletes alike noted the importance

of pain provocation and localisation to identify the "athlete's pain" to validate the athlete's specific

pain experience which is promoted by contemporary pain standards.[15] Physiotherapists acknowledged

how criteria-based rehabilitation and pain management, guided by objective indicators can provide

clear goals for athletes and preferred this strategy to focusing on timelines which can be difficult to

predict. Additionally, athletes appreciated having tangible measures to both represent their pain and

gauge progress, a concept that is firmly established in sports-related injury and can also be applied to

the assessment of pain.[16]

"Mobility, and kind of, my range..strength on both legs, that would have been tested, and then

pinpointing the pain, physically." – A11

"I do a lot of like single leg hopping and stuff like that as I often say to a runner if you can't

single leg hop, you can't run, you kind of set your parameters early." – P01

" I appreciated that like there was an objective outcome .. (Physiotherapist) being able to see

if pain is actually affecting me. There was something that was tangible…I was quite happy

that meant he was going to take my pain serious(ly)." – A08

However, in cases where pain was more diffuse or difficult to pinpoint, the limitations of representing

pain in a clinical setting were a source of frustration for athletes. Additionally, some athletes

experienced negative consequences from pain provocation testing or repeatedly being asked about

symptoms, highlighting the need for the selective use of certain pain assessment tools. Choosing the

right time to assess pain response and ask about symptoms is an important aspect that has been

previously discussed by established pain clinician-researchers.[17]

"I couldn't tell where it was I couldn't point to it, I couldn't palpate it they couldn't locate

where it was .. which kind of frustrated me." – A03

"When I single leg hopped I then had seven out of ten pain for the next two days when I could

have told them that 'this is going to be really sore'." – A03

The development of technology in sport science and medicine has facilitated additional means for

capturing data that can be helpful in the assessment of pain in athletes including wearable devices for

physical performance and physiologic data, as well as smartphones and applications for assisting with

subjective health and wellness monitoring.[18] Athletes reported familiarity with physiological markers

such as heart rate metrics however, effective use requires planning, implementation and reporting and

at times athletes felt overwhelmed with the amount of information.

"I'm like a real techy person like I'm always on Garmin and Whoop and Auras and all the

different things…I went back to physios, went back to Doctors, because my resting heart rate

would have been normally you know low forties.. and then for a good two months, it was mid-

sixties." – A10

"Maybe it was a bad thing that I was keeping such a close eye on you know the data I was

collecting yes because I was probably forming some sort of psychological response."– A10

Questionnaires and pain diaries can be enlisted as a method to add a layer of objectivity to the

retrospective subjective reporting of pain.[19] Athletes, describing the individual nature of preferences

for communicating pain, noted how written diaries allow pain to be reported at multiple time points

throughout the day addressing the limitations of the once-off point-in-time assessments.

Physiotherapists also noted that written methods add an opportunity for athlete reflection. Conversely,

the time taken to complete them was seen as burdensome. Athletes felt they were repeating what had

already been expressed through conversation, and physiotherapists acknowledged that they are often

not appropriately analysed and utilised, citing time restrictions, and instead placed higher value on

objective measures.

"..where you kind of pull out a sheet and describe the pain and it kind of gets the flow of

thought going and you're better able to communicate it then to the physio.'' – A11

"Sometimes there's a period of time where like pain diaries and you know getting patients to

write down every day what the pain was and what they were doing." – P01

" A pain diary .., that takes persistence that I didn't have." –A08

Various pain scales are used to gauge the severity of pain an athlete feels during the clinical encounter

including during pain provocation tests, at specific time points throughout the day, and following

certain activities.[20] The numerical pain rating scale (NPRS), or 0-10 pain scale, was most frequently

encountered by participants who found it may be effective for repeated use with an athlete particularly

when tracking large changes in pain over time. However, both athletes and physiotherapists advised

comparisons between athletes should not be made. The benefits (simplicity, ability to track large

changes in pain in an athlete and widespread use) and drawbacks (once-off measure not representing

the 24-hour pain experience, floor/ceiling effects, subjective interpretation of scale) of the NPRS

described by participants are closely aligned with established pain measurement literature.[20]

"I often find with the 0-to-10 scale, someone can say a 5, doesn't really mean a whole lot. If

someone says a 9, it probably does, but if someone says a 3 or 4 that could be very different

to someone else's 3 or 4, ., I think it certainly gives me an insight into just that person, and if

your pain goes from a 6 to a 2, I know that that's an improvement for you." – P04

"So usually I get asked to describe it from nought to ten but I don't really get the opportunity

to describe the twenty-four-hour cycle of the pain, which I think is probably something maybe

more relevant ..." – A08

Athletes often struggled to represent their pain as a number and preferred descriptions that they could

attribute greater meaning to. Physiotherapists also used alternative pain scales offering different

means of measuring and interpreting pain severity including the traffic light system which indicates

when athletes must stop activities (red), proceed with caution (amber) or continue unabated (green),

an option that has been adopted widely in managing tendon pain.[21] Grading pain as mild, moderate or

severe was another option participants encountered.

"I find it very hard to say nought to ten so she would say mild, moderate or severe on the

pain scale like something that took it away from me giving it a number." – A08

1.2 Multidimensional methods

Athletes emphasised the importance of accurately representing the psychological/emotional aspects

(e.g. feelings and emotions related to pain and sport), wider biological and lifestyle factors (e.g. stress,

sleep and nutrition, age and time of a female athlete's menstrual cycle), and environmental factors

(e.g. pain culture within the sport, time of the competitive season, weather and playing conditions).

Athletes highlighted the profound impact these aspects have on pain perception, yet despite best

practice guidance, were rarely encountered.[24] Physiotherapists, acknowledging that their inclusion is

best practice, shared that some formal assessments of psychological and emotional aspects can be

lengthy and interrupt the flow of assessment in time-pressured sports settings where athletes may not

appreciate them. Instead, physiotherapists described how these aspects tend to be addressed more

informally as part of the pain interview. Techniques such as motivational interviewing and

psychologically informed practice are becoming standard practice for physiotherapists in the

general population and likely inform the assessment process for physiotherapists working in

sports also.[22]

It is worth highlighting how athletes in this study felt these aspects were often ignored and so

there may be a case for explicitly discussing these wider aspects to ensure athlete's pain

experience is validated. *"I know personally if my cortisol levels are higher, so if I'm stressed or*

lacking sleep that I nearly feel pain more and as a woman actually at different parts of your cycle." –

A08

"I think people probably didn't ask enough about the mental effects of injury until you are

visibly in a bad mood or something" – A05

"On the psychological stuff, I suppose the way I would normally do it is you kind of a little

more informal so you kind of be having a chat with them." – P02

1.3 Making sense and interpreting

Prior research has noted how physiotherapists can sometimes feel underequipped when assessing

pain, particularly with more complex and chronic presentations which has led to the recent

publication of pain education curricula at undergraduate and postgraduate levels by the European Pain

Federation.[22][23] Drawing on their experience of prior pain assessments, athletes highlighted the need to

go beyond the available pain scales and search for better ways to describe and represent pain

intensity. Physiotherapists also acknowledged how in their experience objective findings must be

correlated with athletes' subjective reports of pain to complete a comprehensive pain assessment that

is in line with best practice guidance and the recent pain education curricula.[2][23]

"Instead of like tell me what your pain is from nought to ten it's like tell me how it impacted

your ability to train, your ability to race… to live a normal life." – A08

"The outcome measures .. like handheld dynamometry and force plates .. you're trying to

correlate that with their own.. feedback on pain." –- P01

Theme 2. Connect, Listen & Learn

Theme 2 includes two subthemes; 2.1 – the pain interview and athlete's story and 2.2 – forging the

athlete-clinician connection. Every athlete has their own unique "pain story" and effectively

facilitating an athlete to describe the meaning their pain experiences holds for them is a cornerstone of

pain assessment.[17] This aspect of assessment was seen by athletes and physiotherapists as an

opportunity to develop a rapport and relationship with characteristics such as trust, authenticity and

empathy woven into the process. A well-conducted pain interview allowed athletes to feel listened to,

understood and validated, as prioritised in pain assessment guidelines.[15] In contrast, assessments in

sports settings are often time-pressured with establishing injury diagnosis being prioritised meaning

comprehensive pain assessment and optimal management cannot be achieved.[24]

2.1 The pain interview and athlete's story.

The pain interview allows the athlete and physiotherapist to explore the sensory and emotional aspects

of pain and their wider impact.[2] Athletes discussed being asked about aggravating and easing factors,

severity, irritability and nature of the pain and to describe it using various adjectives. The onset,

duration and preceding history of the pain experience were commonly explored. Athletes and

physiotherapists emphasized how the pain interview is an interactive process that is enhanced by the

physiotherapist exploring the context and history behind an athlete's pain experience. Athletes

gathered pain and injury experiences from their support network and teammates adding them to their

interpretations. Physiotherapists used the pain interview to guide the generation of hypotheses and the

selection of objective tests and measures.

"Because everyone has pain, no one goes through life without pain, and then it's individual to

that person so maybe help give them a bit of context to their own pain; this feels .. similar to

what I had before, .. to what this person told me they had, or... " – P04

"If we don't have an idea before they're on .. the treatment table, .. of two or three things that

might be going on here well then we either haven't asked the right question or we haven't

listened to them." –P01

Open-ended discussions allowed the athlete to go beyond the sensory or physical manifestations of

pain and begin to explore the emotional and multidimensional aspects. Open-ended questions aided

the clinician in gathering the necessary information needed to develop accurate hypotheses and

diagnoses in line with research findings of question styles used in other pain cohorts.[25] Conversely

participants found the judicious use of closed questions were helpful to focus on specific aspects of

the pain experience.

"Just letting someone talk to me about what their pain is, you learn a bit about the physical

side of it but then, you know, what does their pain means to them .. the emotion side of it and

the mental side of it as well, you get a bit more information out of it.., , .. a lot of the time

people will tell you exactly what's wrong with them" –P04

2.2 Forging the athlete-clinician connection

Athletes valued having an opportunity to give their unique and full perspective which lay the

foundation for a strong relationship with their physiotherapist. Listening to an athlete with empathy

provides validation and reassurance that they are being heard and understood enabling effective

therapeutic alliance and an effective working relationship.[15] Participants shared how truly

understanding an athlete's pain experience requires an investment of time and energy.

"I think it really matters how you talk to someone about their own pain you know if they think

that it's really important to them, it's really important." – A03

"This is the first time someone's actually listened to what's been going on and like. I think ..

you tend to get a little more insight into their pain." – P01

Athletes and physiotherapists alike stressed the importance of seeing the athlete as a whole person,

prioritising human interaction and connection and understanding their emotions, motivations and

behaviours. Trust plays an integral role in fostering the athlete-physiotherapist relationship and an

athlete's confidence in the diagnosis, pain explanation and prognosis they have received.[26] The

ambiguous nature of pain and the pressure to decide on whether or not to play and/or train through

pain in high-stakes sports settings can create barriers around forming strong relationships. This may

be due to sports physiotherapists' disclosure obligations to the management team or due to the

unwillingness of athletes to disclose the full extent of their pain. The athlete pain assessment was

described as a delicate balance between capturing sufficient objective data to structure the assessment

and management and dedicating time to nurture relationships and forge connections.

"I think it's all just about understanding people really and then obviously you have to have

the rationale to back up what you're going to do etc.." – P02

''There's a balance I suppose in terms of seeing people as a number on a sheet and also as a

human being. Sometimes ..with outcome measures. It ends up where people just don't be seen

at all as a person.'' – P03

Both athletes and physiotherapists being open to different methods of communication, assessment and

management is an important aspect previously highlighted in the literature that participants resonated

with and felt this helped achieve an effectively balanced and individualised assessment.[27]

"Understanding how people communicate is good because some people are, you know, would

be more comfortable writing, some people are comfortable with verbal, some people might be

more comfortable showing, you know.." – P04

'In an ideal world it's like every patient gets a very individual service I suppose... the person,

their previous experiences, you know their goals, their objectives this session like you know

so then you can (use) different parts of you know assessment tools and then you can go

dictate your treatments.'' – P03

Theme 3. Lighthouse In The Storm: providing direction for athletes in pain.

Theme 3 includes three subthemes; 3.1 – information overload and indecision, 3.2 – a beacon of

direction – the role of the physiotherapist and 3.3 – the burden of expectation – challenges for

physiotherapists. Understanding and making sense of pain can be challenging for athletes. They found

that the interaction with a physiotherapist helped them make informed decisions relating to their pain.

The role of providing clarity and direction for athletes is burdensome.[28] Time demands, high

expectations and work environment difficulties can be likened to the solitary and challenging role of

the lighthouse keeper. Research has highlighted how physiotherapists can sometimes feel challenged

in particular when managing chronic pain.[22]

Whilst it is important to seek support from other members of the interdisciplinary team when

appropriate, physiotherapists are core members of the pain team with expertise in exercise

prescription and psychologically informed care who are optimally positioned to guide the pain

management journey for athletes.

3.1 Information overload and indecision

Athletes described their frustration to accurately represent their pain during the assessment process.

Additionally, addressing the potential for pain hypervigilance, athletes noted that too much attention

amplified pain experiences and ruminating thoughts intruded. In line with previous findings, athletes

and physiotherapists noted the assessment of pain can be further complicated by cognitive overload

through an abundance of information gleaned from social media, support networks and various

clinicians that can leave an athlete feeling confused and frozen in indecision.[29]

"You kind of have that feeling of, you know, is that pain or is that, you know, your mind,

because you're overthinking when you're coming back ..that's something that I definitely

struggle with." – A12

"Like Instagram and people influencing .. I was changing exercises every single day trying to

solve my pains and everything .. the stress and the thought process and then doubting other

therapists was another huge thing so like ..there's a lot of therapists that clash and it just

doesn't help at all." – A07

3.2 A beacon of direction – the role of the physiotherapist

Athletes highlighted how a clear explanation of their pain presentation provides clarity and

reassurance helping them to accept their current situation and begin to plan their next steps. The pain

assessment is an opportunity to educate athletes about the causes and contributors to their pain

perception which can have a positive effect on pain experience and management and sports

performance.[3 5]

"I know exactly what it is, and I nearly come to terms with it an awful lot quicker whereas if it's an

injury that I'm not sure what it is sometimes I feel like my interpretation of that is worse. I feel like my

education around it has a huge impact so therefore that knowledge aspect is a drawback if I'm not

asked about it". – A08

"As a kind of full-circle to the person .. "this is what the pain is" or "this is what I think it is

and this is why I think it is" and "this is what we're going to do about it" and then if you understand

the first two then the third becomes a lot easier for the person to actually continue on because they

understand, you know" – P04

3.3 The burden of expectation – challenges for physiotherapists

Physiotherapists acknowledged the challenging nature of assessing and understanding pain and the

difficulty of choosing the appropriate pain assessment tools and measures. During the clinical

reasoning process, physiotherapists must be mindful of their communication strategy with athletes.

With due regard to the "information overload'' described by athletes, physiotherapists highlighted

how they were conscious not to cause further confusion in the pain assessment process, which can be

a challenge.

"Sometimes you come away from it and you wanted, why didn't I do that, why did I do this,

we should assess this, I didn't actually. You know, it can be hard." – P03

Striking the balance between assessment tools and diagnosis is a fine art, when not satisfied with the

pain explanation, diagnosis or prognosis provided by one clinician, athletes may seek additional

assessment, something clinicians were acutely aware of. Taking a health literacy-sensitive approach

is essential to ensure patients can be actively involved in the decision-making about their own

care. Health literacy impacts a variety of health outcomes in patients with chronic pain and

should also be prioritised with athlete cohorts to ensure optimal management.[30]

"All I can remember is I can't run for six weeks so that's not going to happen so I'll go to

somebody else and find the answer I want" – P01

Physiotherapists also described the challenges of completing pain assessments pitch side or in other

sports settings with conditions often less than optimal to conduct a thorough assessment. The timing

of the assessment is a key consideration with early and repeated assessments recommended in

musculoskeletal physiotherapy standards of practice.[30] This gives a clearer and more comprehensive

picture of pain. In contrast to these standards, the pain assessment practice described by athletes was

challenged by the lack of time available in both private practice and sports physiotherapy setting.

Additionally, the pressure to return to performance and minimise time out of sport was highlighted, an

aspect that has been previously discussed by sports physiotherapists.[28] Physiotherapists acknowledged

the difficulties of completing a thorough assessment and providing a diagnosis from the initial

assessment.

"Like if you're working with an athlete day-to-day then it's a lot easier to gauge pain versus

if I have someone that comes in off the street into the clinic that I've never met before, and you don't

know their tendencies or you don't know their history." – A12

"I think it's like, putting the pressure on yourself to know exactly what's wrong with someone

the first time you see them is unrealistic" –P04

Conclusion

We described and explored the phenomena of pain assessment in sport. Athletes and physiotherapists

described and critiqued the routine methods and measures used. We highlighted the value of the pain

interview and athlete's story and we discussed how combining objective findings with the athlete's

pain experience requires consideration and skill. Athletes shared their desire for direction in

understanding and managing pain and we highlighted the challenges this poses, particularly for

physiotherapists. Whilst every effort was made to collect experiences from a diverse range of athletes

and physiotherapists and variety was achieved in sport, competition level and practice setting the

experiences gathered from these focus groups may not apply to all athlete pain assessment settings.

Notably, participants were all recruited from Ireland and whilst some of the female athletes also had a

physiotherapy background, no female sports physiotherapists were available to participate.

Better, more comprehensive and multidimensional means to describe and assess pain are priorities for

research and practice. Additionally, improved communication strategies that facilitate more timely

and relevant pain information whilst preserving an effective athlete-physiotherapist relationship are

needed. Priorities and directions for pain assessment will be explored further in Part Two.

Practical Implications

- Objective assessment tools offer value in supporting the subjective nature of pain
experience and as indicators of progress.
- Some commonly used objective tools and scales have their limitations when it
comes to comprehensively assessing pain. The overuse of objective measures that
may not be relevant or helpful should be avoided.
- Multidimensional psychosocial assessment tools are underused and
Physiotherapists should consider integrating them into their athlete pain assessment
practice.
- Physiotherapists must place a high value on a well-conducted pain interview to (i)
facilitate the athlete to tell their pain story, (ii) educate the athlete regarding their
pain and (iii) as an opportunity to develop the therapeutic relationship.
- Physiotherapists must include multidimensional pain assessment measures and take
an individualised approach when interpreting pain assessment findings.

References

1. Hainline B, Derman W, Vernec A, et al. International Olympic Committee consensus statement on

pain management in elite athletes. *Br J Sports Med* 2017;51(17):1245-58. doi:

10.1136/bjsports-2017-097884 [published Online First: 2017/08/23]

2. Hainline B, Turner JA, Caneiro JP, et al. Pain in elite athletes-neurophysiological, biomechanical

and psychosocial considerations: a narrative review. *Br J Sports Med* 2017;51(17):1259-64.

doi: 10.1136/bjsports-2017-097890 [published Online First: 2017/08/23]

3. Hoegh M, Stanton T, George S, et al. Infographic. Pain or injury? Why differentiation

matters in exercise and sports medicine. *British Journal of Sports Medicine* 2022;56(5):299-

300. doi: 10.1136/bjsports-2021-104633

4. Purcell C, Duignan C, Fullen BM, et al. Comprehensive assessment and classification of upper and

lower limb pain in athletes: a scoping review. *British Journal of Sports Medicine*

2023:bjsports-2022-106380. doi: 10.1136/bjsports-2022-106380

5. Wouw Avd. Advocating a holistic approach for sport injury prevention and rehabilitation. *British*

Journal of Sports Medicine 2023;57(14):895-96. doi: 10.1136/bjsports-2022-105695

6. Titze C, Fett D, Trompeter K, et al. Psychosocial subgroups in high-performance athletes with low

back pain: eustress-endurance is most frequent, distress-endurance most problematic!

Scand J Pain 2021;21(1):59-69. doi: 10.1515/sjpain-2020-0053 [published Online First:

2020/09/07]

7. Tabben M, Verhagen E, Warsen M, et al. Obstacles and opportunities for injury prevention in

professional football in Qatar: exploring the implementation reality. *BMJ Open Sport &*

Exercise Medicine 2023;9(1):e001370. doi: 10.1136/bmjsem-2022-001370

8. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-

item checklist for interviews and focus groups. *International Journal for Quality in Health*

Care 2007;19(6):349-57. doi: 10.1093/intqhc/mzm042

9. Korstjens I, Moser A. Series: Practical guidance to qualitative research. Part 2: Context, research

questions and designs. *European Journal of General Practice* 2017;23(1):274-79. doi:

10.1080/13814788.2017.1375090

10. Braun V, Clarke V. Reflecting on reflexive thematic analysis. *Qualitative Research in Sport,*

Exercise and Health 2019;11(4):589-97. doi: 10.1080/2159676x.2019.1628806

11. Braun V, Clarke V. To saturate or not to saturate? Questioning data saturation as a useful

concept for thematic analysis and sample-size rationales. *Qualitative Research in Sport,*

Exercise and Health 2021;13(2):201-16. doi: 10.1080/2159676x.2019.1704846

12. Kallio H, Pietilä AM, Johnson M, et al. Systematic methodological review: developing a

framework for a qualitative semi-structured interview guide. *J Adv Nurs* 2016;72(12):2954-

65. doi: 10.1111/jan.13031 [published Online First: 2016/05/26]

13. Tausch AP, Menold N. Methodological Aspects of Focus Groups in Health Research. *Global*

Qualitative Nursing Research 2016;3(0):233339361663046. doi:

10.1177/2333393616630466

14. Smith B, McGannon KR. Developing rigor in qualitative research: problems and opportunities

within sport and exercise psychology. *International Review of Sport and Exercise Psychology*

2018;11(1):101-21. doi: 10.1080/1750984x.2017.1317357

15. Raja SN, Carr DB, Cohen M, et al. The revised International Association for the Study of Pain

definition of pain: concepts, challenges, and compromises. *Pain* 2020;161(9):1976-82. doi:

10.1097/j.pain.0000000000001939 [published Online First: 2020/07/23]

16 Mendiguchia J, Martinez-Ruiz E, Edouard P, et al. A Multifactorial, Criteria-based Progressive

Algorithm for Hamstring Injury Treatment. *Med Sci Sports Exerc* 2017;49(7):1482-92. doi:

10.1249/MSS.0000000000001241 [published Online First: 2017/03/10]

17 Wilson D, Williams M, Butler D. Language and the pain experience. *Physiother Res Int*

2009;14(1):56-65. doi: 10.1002/pri.424 [published Online First: 2008/11/15]

18 Seçkin AÇ, Ateş B, Seçkin M. Review on Wearable Technology in Sports: Concepts, Challenges and

Opportunities. *Applied Sciences* 2023;13(18):10399.

19. Haase I. Accuracy of retrospective pain measurement in patients with chronic pain. *Med Int*

(Lond) 2023;3(4):35. doi: 10.3892/mi.2023.95 [published Online First: 2023/07/14]

20. Correll DJ. Chapter 22 - The Measurement of Pain: Objectifying the Subjective. In: Waldman SD,

ed. Pain Management (Second Edition). Philadelphia: W.B. Saunders 2011:191-201.

21. Silbernagel KG, Thomeé R, Eriksson BI, et al. Continued sports activity, using a pain-monitoring

model, during rehabilitation in patients with Achilles tendinopathy: a randomized controlled

study. *Am J Sports Med* 2007;35(6):897-906. doi: 10.1177/0363546506298279 [published

Online First: 2007/02/20]

22. Cowell I, McGregor A, O'Sullivan P, et al. Physiotherapists' Approaches to Patients' Concerns in

Back Pain Consultations Following a Psychologically Informed Training Program. *Qualitative*

Health Research 2021;31(13):2486-501. doi: 10.1177/10497323211037651

23. European Pain Federation (EFIC). Core Pain Curriculum for Bachelor/ Pre Registration

Physiotherapy Programmes. 2023. Available online at:

https://europeanpainfederation.eu/education/pain-curricula/ Last Accessed 22.02.2024

24. Thorborg K, Rathleff MS, Petersen P, et al. Prevalence and severity of hip and groin pain in sub-

elite male football: a cross-sectional cohort study of 695 players. *Scand J Med Sci Sports*

2017;27(1):107-14. doi: 10.1111/sms.12623 [published Online First: 2015/12/09]

25. McDonald DD, M S, Rose L, et al. The effect of pain question phrasing on older adult pain

information. *J Pain Symptom Manage* 2009;37(6):1050-60. doi:

10.1016/j.jpainsymman.2008.06.008

26. David S, Hitchcock J. Understanding Patient Trust in the Athletic Setting through Interviews.

Internet Journal of Allied Health Sciences and Practice 2018 doi: 10.46743/1540-

580x/2018.1683

27. UK NGC. Evidence review for communication between healthcare professionals and people with

chronic pain (chronic primary pain and chronic secondary pain): Chronic pain (primary and

secondary) in over 16s: assessment of all chronic pain and management of chronic primary

pain: Evidence review B. . London National Institute for Health and Care Excellence (NICE)

2021.

28. Morgan N. Practical tips for a future in sports physiotherapy: Association of Chartered

Physiotherapists in Sport and Exercise Medicine blog series. In: Medicine BJoS, ed.: BMJ,

2023.

29. Gould D, Nalepa J, Mignano M. Coaching Generation Z Athletes. *Journal of Applied Sport*

Psychology 2020;32(1):104-20. doi: 10.1080/10413200.2019.1581856

30. Mackey LM, Blake C, Casey MB, et al. The impact of health literacy on health outcomes in

individuals with chronic pain: a cross-sectional study. *Physiotherapy* 2019(1873-1465

(Electronic)) doi: 10.1016/j.physio.2018.11.006. [published Online First: 2018]

31. Bishop A, Blackburn, J., Hallam, F., McComiskie, E., Rankin, G. Musculoskeletal

physiotherapy service standards; The delivery of musculoskeletal (MSK) physiotherapy

services in the UK for adults of 16 years and over. In: Physiotherapy TCSo, ed. London 2021.

PARTICIPANTS	N	FEMALE	MALE	SPORT (N)	COMPETITION LEVEL	PHYSIOTHERAPY BACKGROUND (N)	ATHLETE BACKGROUND
ATHLETES	12	5	7	Athletics (5) Tag Ruby (1) Field Hockey (1) Soccer (2), Cycling (1) Mixed Martial Arts (1) Gaelic Football	Professional/International (1) Semi-Professional (2) Elite Amateur (9)	Yes (5) No (7)	NA

567　**Table 1 – Participant Demographics**

			Athletics (1), Rugby (1), Gaelic Football & Hurling (2), Mixed Sports in Private Practice (3)	Professional/Inter national (2) Mixed Competition level in Private Practice (3)		Current Athletes (2) Retired Athletes (2)
			& Basketball (1)			
PHYSIOTHERAPIS TS	4	-	4		NA	

Figure Captions

Figure 1. Themes (Measures, Scales and Dimensions; Connect, Listen and Learn; Lighthouse in

the Storm: providing direction for athletes in pain) and codes. Dark shading – indicates codes

that were present in athletes and physiotherapists. Light shading – indicates codes that were present in

athletes only, No shading – indicates codes that were present in physiotherapists only

Figure 2. **Athlete pain assessment experience themes and subthemes.** The themes for each part of

this series represent a row in the priorities for the pain assessment pyramid, this paper presents the

bottom or foundation row of the pyramid. This comprises the three themes that describe the "athlete

pain assessment" experience alongside the constituent subthemes. In the top row, the "priorities and

directions for athlete pain assessment" themes and subthemes will be presented in Part Two and will

therefore build on the themes from this paper.

PARTICIPANTS	N	FEMALE	MALE	SPORT (N)	COMPETITION LEVEL	PHYSIOTHERAPY BACKGROUND (N)	ATHLETE BACKGROUND
ATHLETES	12	5	7	Athletics (5) Tag Ruby (1) Field Hockey (1) Soccer (2), Cycling (1) Mixed Martial Arts (1) Gaelic Football &	Professional/International (1) Semi-Professional (2) Elite Amateur (9)	Yes (5) No (7)	NA

			Basketball (1)			
PHYSIOTHERAPISTS	4	-	4	Athletics (1), Rugby (1), Gaelic Football & Hurling (2), Mixed Sports in Private Practice (3)	Professional/International (2) Mixed Competition level in Private Practice (3)	NA	Current Athletes (2) Retired Athletes (2)

Enhanced Communication & Pain Descriptions

- Pain and decision making – difficult
- Acute/clear diagnosis vs chronic/unclear pain
- Importance of physiotherapist providing clear direction
- Role of physiotherapist to make objective decisions
- Pre assessment opportunity for reflection
- Increased frequency of contextual pain assessment
- Descriptions, criteria & categories
- Integrative multidomain App
- Pain and rehab
- Conscious incompetence and understanding limitations
- A collaborative approach to pain and decision making

Integrating Sport Specific & Multidimensional Assessments

- Wider biological (sleep, stress and nutrition)
- More sport and context specific
- Pain medication
- Psychological aspects
- Psychological readiness to return to play
- Environmental factors
- Integrative Multidomain App

Measures, Scales & Dimensions

Objective assessment tools	Pain diaries/written descriptions	Numerical Pain Rating Scale	Limitations of numerical pain rating scale	Importance of objective indicators	Importance of identifying ''athlete's pain'' with objective tools	Clinicians and athletes value objective measures
Too much information	Psychological aspects of pain assessed informally	Importance of wider biological and lifestyle factors	Time of season is significant	Pain and culture	Increased frequencies of injury and pain with age	Limited value of subjective questionnaires and forms
Subjective questionnaires - adjunct to pain conversation and research	Correlating objective findings with subjective experience – considering multiple factors	Limitations to clinic based assessments	Lack of assessment and consideration of psychological aspects of pain	Pain and environment	Benefit of strategies to limit the subjective interpretation of pain	Wearable technologies and pain
Pain and physiological data	Regular pain and performance reporting	Mild, moderate and severe pain scale	Negative perceptions of pain assessment	Traffic light pain scale	Psychosocial and lifestyle factors	Timelines vs criteria – diagnosis and return to play decisions

Connect, Listen & Learn

Pain interview/ subjective assessment	Importance of being listened to and understood/ validating pain	Trust and honesty between athlete and physiotherapist	Value of developing relationship between athlete and physiotherapist	Human interaction – Understanding people and behaviour	Athlete values and preferences and an individualised pain assessment approach	Expectations and open-mindedness
Athlete's understanding of pain and pain context	Previous pain experience	Positive perceptions of pain assessment	Value of physiotherapist understanding sport and context	Importance of subjective assessment	Importance of letting the athlete tell their story over closed/leading questions or objective tests	Empathising, validating and seeking athlete experience
A collaborative approach to pain and decision making						

Lighthouse in the Storm

Persistent pain is challenging – frustration and doubt	Pain and decision making – difficult	Too much information	Influence of internet & social media	Importance of clear diagnosis & pain explanation	Importance of prognosis & timeline	Importance of physiotherapist providing clear direction
Role of physiotherapist to make objective decisions	Understanding and assessing pain is challenging	Hypervigilance to pain	Challenges of working in sport and private practice	Developing accurate diagnosis/prognosis can take time and/or experience	Acute/clear diagnosis vs chronic/unclear pain	Setting and environment of pain assessment
Access to physiotherapy – timeline and frequency	Role of physiotherapist to educate athlete	Pain and imaging	Time demands and pain assessment	Describing pain is difficult	Persisting pain beyond normal training/ competition triggering action	Negative perceptions of pain assessment
Overassessment	Clinician shopping	Conscious incompetence and understanding limitations	A collaborative approach to pain and decision making	Effects of physiotherapist's communication with athlete		