

Frequency of ocular manifestations in rheumatic autoimmune diseases (ARDs); a cross-sectional study.

AUTHOR:

Usama Sohail (Shaikh Zayed Hospital)

ABSTRACT:

Autoimmune disorders involve diverse symptomatic manifestations constituting an abnormal or exaggerated immune function and serve to attack self-antigens and result in tissue destruction. This study aims to comprehensively analyse the prevalence, type, associative factors, and the resultant impact of ocular symptoms on patients presented to rheumatology departments of multi-center tertiary care hospitals in Lahore. Methods. A total of 300 patients were studied during the time frame of Jan 2021-Dec 2022. Results. Rheumatoid Arthritis comprised 45.7 % (n=137) of patients, Systemic Lupus Erythematosus (SLE) was present in 34.3% (N=103) patients. Spondylarthritis 8.3%, Granulomatous Polyangiitis 3.3%, Juvenile idiopathic arthritis 0.7%, and Bechet Disease 2.3% were present. The total number of patients who developed eye manifestations was 18%. Keratoconjunctivitis Sicca was observed in 4 females and 2 males with overall frequency of 11.1%. Conclusions. Our study showed a surge in the cases of SLE diagnosed and witnessed during the period of 2022, but our study showed an overall decrease in the incidence of eye complications.

INTRODUCTION:

Autoimmune disorders involve diverse symptomatic manifestations constituting an abnormal or exaggerated immune function and serve to attack self-antigens and result in tissue destruction. (1) 80 different types of autoimmune disorders have been recognized. (2) A statistic from the United States of America showed an increased incidence of autoimmune disorders, around 5-7%, with females being the major targeted population. (3) Diseases with positive "Rheumatic Factor" are the main focus for the associated extra-articular effects and include a broad-spectrum range, including Bechet Disease, Systemic Lupus Erythematosus (SLE), Rheumatic Arthritis (RA), Dermatomyositis (DM), Vasculitis, Sjogren's Syndrome and Polymyositis (PM). (4) Autoimmune Rheumatic Diseases (ARDs) target people globally, with a count affecting millions of people. Worldwide prevalence of SLE, RA, DM/PM, SS, Sjogren's Syndrome and are 0.02–0.07% (5), (6) , 0.5–1.1% (7) ,0.02% (8), 0.02–0.04%, (9) 0.05–4.8%, (10) respectively. ARDs can involve many body organs outside the joints, including skin, heart, lungs, eyes, kidneys, etc. Our research purpose is to find a link between rheumatic autoimmune disease and their ocular manifestations (11). Ocular symptoms containing autoimmune diseases mainly consist of Rheumatoid Arthritis (12) , Sjogren Syndrome (13) , Spondylarthritis (14), (15), Juvenile Idiopathic Arthritis, Systemic Sclerosis Granulomatosis with Polyangiitis (16) , (17), Systemic Lupus Erythematosus associated with antiphospholipid syndrome (APA), Relapsing Polychondritis. (18) Inclusion of different eye tissues in patients with rheumatic diseases can lead to troublesome effects, of which the most distinctive ones are retinal vasculitis, silica, keratoconjunctivitis, glaucoma, and uveitis. The delicacy of the eye (19) as an organ can be indicated by the clinical signs that appear as the first indicator of any systemic distortion that leads to a change in its sensitive microenvironment. (20) A specialized sterile environment is present between

blood and eye compartments (containing aqueous and vitreous humor), and rheumatic autoimmune disease tends to disturb this equilibrium. (21) But ocular problems following rheumatic diseases can present during the active stage or even years after the initial diagnosis. The optimal and gold standard treatment regimens for RA diseases have been disease-modifying anti-rheumatic drugs or DMARDs, as they are involved in halting the long-term progression of the disease and the resultant joint manifestations. (22) But the downside to these drugs is the unpremeditated ocular toxicity, especially those caused by the prototypic agent Methotrexate, retinal cotton-wool spots, orbital non-Hodgkin's lymphoma, and ischemic neuropathy. (23) A study conducted in Mexico included 276 medical records in which the most common autoimmune rheumatic disease and ocular manifestation were spondylarthritis and keratoconjunctivitis sicca, respectively. (24) Another one done in a tertiary care setting in Lahore (Fatima Memorial Hospital) included 83 patients, in whom commonly occurring diseases were spondylarthritis and uveitis. (25) Hence, a detailed collaboration between the departments of rheumatology and ophthalmology is required for the ideal management of such patients. At best, routine-based checkups should be opted for as a planning methodology for diagnosing and treating such patients. As many diseases mentioned in this study result in acute visual complications, regular screening should be emphasized, even for asymptomatic patients. Till now, one such study has been conducted about this inter-relationship in our population, but more data is needed. This research aims to comprehensively analyze the prevalence, type, associative factors, and the resultant impact of ocular symptoms on patients presented to rheumatology departments of multi-center tertiary care hospitals in Lahore.

MATERIAL AND METHODS:

- Our study was retrospective cross sectional, carried out at Shaikh Zayed Hospital, Lahore. Data was taken between the time of Jan 2021 - Dec 2022. Data was included by probability method of simple random sampling. Out of total, a sample of 300 was taken. Sample included only those patients with autoimmune rheumatic diseases (RA factor +) and those who were diagnosed at the time by the rheumatologist. Our study was approved by the ethical review board and informed consent was taken to perform this study. Parameters assessed were demographics, presence of autoimmune disease and the resultant diagnosis of the eye disease. It also included the duration of the eye symptoms and the use of medication like steroids and the effect of medications.
- The data taken was analyzed by SPSS tool. Continuous variables like age were presented and the resultant frequency (percentage) was taken.

Results:

- A total of 300 patients were studied during the time frame of Jan 2021-Dec 2022. Average age ----- is taken. About 59.7% of the total data taken were females.
- Rheumatoid Arthritis comprised 45.7 % (n=137) of patients, SLE was present in 34.3% (n=103) patients. While Systemic Sclerosis 5.3% (n=16), Spondylarthritis 8.3% (n=25), Granulomatous Polyangiitis 3.3% (n=10), Juvenile idiopathic arthritis 0.7% (n=2) and Behcet Disease was present in 2.3% (n=7). The total number of patients who developed ocular manifestations was 18%. The individual ocular sub types of presents include, Keratoconjunctivitis Sicca 6% (n=18), Uveitis 5.3% (n=16), Scleritis 4% (n=12), and Peripheral ulcer keratitis 2% (n=6).

DISCUSSIONS:

Ophthalmological involvement is the main stay process we see in most autoimmune rheumatic processes. This study provides an overlook about the major eye complications in patients of Autoimmune Rheumatic Diseases. Most of the patients included in our study were cases of Uveitis and Keratoconjunctivitis Sicca.

Autoimmune Rheumatic Diseases:

During the time of our study (Jan 2021-Dec 2022). We took a total of 300 patients were taken. The data we had taken included the age between 36-45 years as the most prevalent age group. RA is the most common articular manifestation noticed in our study. It included 87 females and 50 males. Rheumatoid Arthritis was the most common prevalence seen during the period of 2021, but our retrospective study showed a surge in the cases of SLE diagnosed and witnessed during the time of 2022. Going further SLE was our second most common disease with a prevalence of 69 among females and 39 among males. Spondylarthritis was present in 10 females and 15 males. In reference to this study, one conducted in Fatima Memorial Hospital, Lahore demonstrated Spondylarthritis as the most common disease with the overall frequency of 38.6%. Out of 16 patients of SS, 10 females and 6 males were present. And lastly, GPA had a frequency of 4 males and 6 females.

Ocular Manifestations:


They constitute the most common extra-articular symptomology studied in autoimmune rheumatic spectrum. Our study included most of the cases including Uveitis and Keratoconjunctivitis Sicca as the ocular diseases. Keratoconjunctivitis sicca, also known as Dry syndrome, the most common etiological factor for which is Sjogren syndrome. In our clinical cases, diagnosis of KS is made by Schirmer's test with the help of Whitman filter paper. Our study included total 18 patients out of which there 12 females and 6 males with the overall ocular frequency 33.3%. In Uveitis cases, there were 7 females and 9 males with overall percentage 29.6%. Scleritis had 6 females and 6 males with cumulative frequency of 22.2%. While peripheral keratitis was present in 4 females and 2 males with overall frequency of 11.1%. The age group categorization of the study included most patients of autoimmune rheumatic cases in between 36 to 45 years. Late adult onset has some striking association with hormonal changes. Keratoconjunctivitis siccas were predominant in between the age of 27-36 years whereas uveitis and scleritis were found in predominance 36-45 years. Our study also postulated the linkage between the steroids and ocular manifestations, as 5 of the patients had steroid-onset ocular diseases and all those cases were of Keratoconjunctivitis Sicca.

Conclusion:

To summarize, in our retrospective cross-sectional study, the most prevalent rheumatic autoimmune disease was Rheumatoid Arthritis (RA). And the resultant ocular complication was Keratoconjunctivitis Sicca. In view of the studies done on the linkage between the rheumatic autoimmune diseases and the ocular diseases in Pakistan as well as in other countries, our study showed an overall decrease in the ocular involvement.

RHEUMATIC AUTOIMMUNE DISEASE:


		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Bechet Disease	7	2.3	2.3	2.3
	GPA	10	3.3	3.3	5.7
	JIA	2	.7	.7	6.3
	RA	137	45.7	45.7	52.0
	SLE	103	34.3	34.3	86.3
	SpA	25	8.3	8.3	94.7
	SS	16	5.3	5.3	100.0
	Total	300	100.0	100.0	




SEX * RAD Crosstabulation:

	Bechet Disease	GPA	JIA	RA	SLE	SpA	SS	Total
Female	7	4	0	87	69	2	10	179
Male	0	6	2	50	34	23	6	121

Total	7	10	2	137	103	25	16	300
-------	---	----	---	-----	-----	----	----	-----




DISTRIBUTION ACCORDING TO AGE:



OCCULAR DISEASES:

	Frequency	Percent	Valid Percent	Commutative Percent
--	-----------	---------	---------------	---------------------

Keratoconjunctivitis Sicca	18	6.0	6.0	88.0
Peripheral Ulcer Keratitis	6	2.0	2.0	90.0
Retinal Vasculitis	2	.7	.7	90.7
Scleritis	12	4.0	4.0	94.7
Uveitis	16	5.3	5.3	100.0
Total	300	100.0	100.0	



RHEUMATIC AUTOIMMUNE DISEASE:



1. RHEUMATOID ARTHRITIS (RA) 2. SYSTEMIC LUPUS ERYTHEMATOSUS (SLE) 3. MICTIGANITIS 4. GOUT 5. ANKYLOSING SPONDYLITIS 6. DERMATOMYOSITIS 7. POLYMYOSITIS 8. SCLERODERMA 9. MISCHELANEUS

Year	Rheumatoid Disease	Prevalence		Cumulative Prevalence	
		Prevalence	Prevalence	Prevalence	Prevalence
1974	RA	100	100	100	100
1975	RA	100	100	100	100
1976	RA	100	100	100	100
1977	RA	100	100	100	100
1978	RA	100	100	100	100
1979	RA	100	100	100	100
1980	RA	100	100	100	100
1981	RA	100	100	100	100
1982	RA	100	100	100	100
1983	RA	100	100	100	100
1984	RA	100	100	100	100
1985	RA	100	100	100	100
1986	RA	100	100	100	100
1987	RA	100	100	100	100
1988	RA	100	100	100	100
1989	RA	100	100	100	100
1990	RA	100	100	100	100
1991	RA	100	100	100	100
1992	RA	100	100	100	100
1993	RA	100	100	100	100
1994	RA	100	100	100	100
1995	RA	100	100	100	100
1996	RA	100	100	100	100
1997	RA	100	100	100	100
1998	RA	100	100	100	100
1999	RA	100	100	100	100
2000	RA	100	100	100	100
2001	RA	100	100	100	100
2002	RA	100	100	100	100
2003	RA	100	100	100	100
2004	RA	100	100	100	100
2005	RA	100	100	100	100
2006	RA	100	100	100	100
2007	RA	100	100	100	100
2008	RA	100	100	100	100
2009	RA	100	100	100	100
2010	RA	100	100	100	100
2011	RA	100	100	100	100
2012	RA	100	100	100	100
2013	RA	100	100	100	100
2014	RA	100	100	100	100
2015	RA	100	100	100	100
2016	RA	100	100	100	100
2017	RA	100	100	100	100
2018	RA	100	100	100	100
2019	RA	100	100	100	100
2020	RA	100	100	100	100
2021	RA	100	100	100	100
2022	RA	100	100	100	100
2023	RA	100	100	100	100
2024	RA	100	100	100	100
2025	RA	100	100	100	100
2026	RA	100	100	100	100
2027	RA	100	100	100	100
2028	RA	100	100	100	100
2029	RA	100	100	100	100
2030	RA	100	100	100	100
2031	RA	100	100	100	100
2032	RA	100	100	100	100
2033	RA	100	100	100	100
2034	RA	100	100	100	100
2035	RA	100	100	100	100
2036	RA	100	100	100	100
2037	RA	100	100	100	100
2038	RA	100	100	100	100
2039	RA	100	100	100	100
2040	RA	100	100	100	100
2041	RA	100	100	100	100
2042	RA	100	100	100	100
2043	RA	100	100	100	100
2044	RA	100	100	100	100
2045	RA	100	100	100	100
2046	RA	100	100	100	100
2047	RA	100	100	100	100
2048	RA	100	100	100	100
2049	RA	100	100	100	100
2050	RA	100	100	100	100


Frequency



Frequency (20%) Frequency (30%) Frequency (10%) Frequency (40%) Frequency (10%) Frequency (0%)

SEX * RAD Cross-tabulation:

	Secret Service	IRS	IRS	IRS	IRS	IRS	IRS	Total
Female	7	4	8	37	62	2	10	119
Male	0	6	2	59	34	10	1	111
Total		10	10	96	96	12	11	230



OCULAR DISEASES:

	Frequency	Percent	Valid Percent	Cumulative Percent
Nonbacterial conjunctivitis (NBA)	118	6.4%	6.8%	33.4%
Bacterial conjunctivitis (BAC)	4	2.2%	2.6%	36.0%
Other conjunctivitis	7	3.7%	4.1%	40.1%
Glaucoma	12	6.4%	6.8%	46.9%
Cataract	118	6.4%	6.8%	53.3%
Total	180	100.0%	100.0%	

Frequency



■ Nonbacterial conjunctivitis (NBA)
 ■ Bacterial conjunctivitis (BAC)
 ■ Other conjunctivitis
 ■ Glaucoma
 ■ Cataract