4   **Sex Differences in the Role of Sleep on Cognition in Older Adults**

7   Yumiko Wiranto[1]* and Amber Watts[1, 2]

12   [1]Department of Psychology, University of Kansas, Lawrence, Kansas, United States of America

13   [2]University of Kansas, Alzheimer's Disease Research Center, Fairway, Kansas, United States of

14   America

16   * Corresponding author

17   Email: yumiko.wiranto@ku.edu (YW)

Abstract

Advancing age is associated with decreased sleep quality and cognition. Previous work has shown differences in sleep patterns and perceptions of sleep quality between men and women. However, whether the effects of sleep quality on cognition differ between sexes in older populations remains unclear. We analyzed data from 207 cognitively intact and mildly cognitively impaired participants (89 men and 118 women) aged 60< to estimate the effects of subjective and objective measures of sleep on executive function, verbal memory, and attention using generalized additive models. Objective sleep was measured with the GT9X Link Actigraph, and subjective sleep was measured with the Pittsburgh Sleep Quality Index. We found that women showed optimal executive function with up to about 400 minutes of total sleep time, followed by a decline in performance ($p = 0.01$). Additionally, a longer total sleep time contributed to lower verbal memory in a linear manner ($p = 0.043$). In men, there was a marginally significant improvement in executive function with higher sleep efficiency ($p = 0.05$). Higher self-reported sleep complaints were associated with poorer executive function in cognitively intact women ($p = 0.024$). Our findings suggest that the effects of sleep quality on cognitive performance differ between older men and women. Furthermore, executive function seems to be the cognitive domain that was most closely related to objective and subjective sleep. This study indicates that interventions focusing on sleep to mitigate the risk of cognitive impairment in older adults may need to be sex-specific.

Keywords: sleep, cognition, aging, sex differences, actigraphy

Introduction

The impact of aging on sleep and cognition is well documented in the literature. Aging

has been associated with fragmented sleep and decreased sleep quality, which could exacerbate

cognitive decline that occurs with aging [1]. However, the literature on sex differences in sleep

and its interaction with cognition in older adults is still limited. Women experience dramatic

hormonal and physiologic developmental changes (puberty, pregnancy, and menopause) across

the lifespan that contribute to alterations in sleep patterns and a higher number of reported sleep

complaints [2]. Moreover, women have twice the risk of developing Alzheimer's disease (AD),

for which poor sleep is a risk factor [3,4]. Understanding how men and women differ in their

sleep health and the implications of these differences on cognition is vital for designing effective

interventions and prevention strategies for AD.

Sleep and Aging

Advancing age is associated with physiological changes, among which are the alterations

of sleep architecture (i.e., stages of sleep) and circadian rhythm. Older adults have a shorter

duration of deep sleep (i.e., slow wave and rapid eye movement sleep) and more frequent

awakenings during the night [5,6]. Additionally, they tend to go to bed earlier in the evening and

wake up earlier in the morning. These changes contribute to the high prevalence of sleep

disturbances experienced by this population. Previous studies reported that up to 75% of older

adults experienced sleep difficulty [7–9]. In addition to sleep issues, increasing age is associated

with the accumulation of beta-amyloid, a protein that is strongly implicated in cognitive decline

and AD [10,11]. Because one of the roles of sleep is to clear out neurotoxic waste products

produced during the day, poor sleep has been found to exacerbate beta-amyloid deposition and

cognitive aging [12].

The investigation of the interrelationships between aging, sleep, and cognition has been

relatively limited in the existing literature. Moreover, the results from these studies have been

inconclusive. In cognitively healthy older adults, both short (< 6 hours) and long (> 9 hours)

sleep durations have been linked to reduced global cognition and memory [12,13]. However,

Sabeti et al. [14] observed this association solely in cognitively impaired older adults. In a two-

year longitudinal study, it was found that older women who slept for only four hours exhibited a

more significant cognitive decline compared to those who slept for seven hours. Notably,

individuals who slept for nine hours did not experience any cognitive decline during the same

period [15]. Conversely, Faubel et al. [16] indicated that older adults who slept for 11 hours

demonstrated lower cognitive performance compared to those who slept for seven hours. In

addition to global cognition and memory, sleep disturbances have been reported to impact other

cognitive domains, including attention and executive function, in older populations. Specifically,

Blackwell et al. [17] and Wilckens et al. [18] revealed a negative correlation between frequent

nighttime awakenings and performance on executive function and attention. However, Wilckens

et al. [18] did not find a significant association between sleep duration and cognitive

performance in their study.

Sex Differences in Sleep

Biological sex is another major factor that influences sleep. Research investigating sex

differences in subjective sleep quality suggests that middle-aged and older women take longer to

fall asleep, have shorter sleep duration, and have lower sleep efficiency [19,20]. However, this

result is contradicted by studies looking at objective sleep quality between older men and women

using actigraphy and polysomnography (PSG). A meta-analysis including studies on sleep

characteristics across three countries reported that older women had longer sleep duration and

higher sleep efficiency based on actigraphy data [21]. Studies using PSG show that reduced

slow-wave sleep due to age is more pronounced in men than women [22,23]. The variation

between subjective and objective sleep quality may be attributed to sex-related differences in

sleep perception and symptom reporting. Recent research suggests a stronger link between

objective and subjective sleep quality in women compared to men [24,25], potentially indicating

that women have a more accurate perception of their sleep quality. Additionally, it raises

questions about whether the current criteria for defining good objective sleep quality apply

equally to both sexes, suggesting a need for distinct criteria for women [26,27].

98   Regarding the cognitive implications of poor sleep, limited research has been conducted

to investigate sex differences in this topic. Notably, it has been observed that sleep deprivation

affects working memory to a greater extent in young women compared to men [28,29].

However, whether the same findings apply to older populations is still not well understood. This

lack of research leaves an important gap in our understanding of how sleep impacts cognitive

function across the lifespan. Some possible mechanistic explanations for sex differences in this

relationship include depletion in sex steroids (estradiol) due to menopause that could adversely

affect cognition, specifically memory processing [30]. Moreover, since sleep is known to

influence memory performance, the connection between reduced estradiol levels due to

menopause and memory might be further intensified by poor sleep.

Objective

109   The present study investigated sex differences in the relationship between sleep quality

110   and cognitive performance among older individuals. We assessed sleep quality using subjective

111   (self-report) and objective (actigraphy) measures including total sleep time and sleep efficiency.

112   We hypothesized that sleep quality would be more significantly associated with cognitive

113   performance in older women than it would in older men.

Materials and Methods

Participants

Participants were recruited from the University of Kansas Alzheimer's Disease Research

Center (KU-ADRC) Clinical Cohort, an ongoing longitudinal observational study to support

regional and national research on aging, cognition, and AD. This study collects demographic,

medical, psychological, and cognitive data annually, and imaging and biospecimen data as

needed if participants opt in. In the second year of participation (the first follow-up visit),

participants could opt to partake in the Physical Activity and Sleep Study (PASS), a sub-study of

the Clinical Cohort initiated on July 29, 2015, that was repeated once every two years until May

27, 2021. To be eligible for the Clinical Cohort study, participants had to be at least 60 years old,

native English speakers, and not exhibit the following conditions: significant depressive

symptoms, untreated thyroid dysfunction, visual or auditory impairment, active (< 2 years)

ischemic heart disease, and uncontrolled insulin-dependent diabetes mellitus.

In the current analysis, we included 207 participants with normal cognition or mild

impairment as indicated by clinical dementia rating (CDR) scores of 0 and >0, respectively.

Among these participants, 161 exhibited normal cognitive function, while 46 had mild cognitive

impairment with a CDR of 1. The research was approved by the KU-ADRC's Institutional

Review Board, and all participants provided written informed consent separately for both the

KU-ADRC Clinical Cohort and the PASS study.

Sleep Quality Measures

Actigraphy

ActiGraph GT9X Link accelerometers (Pensacola, FL) are body-worn monitors that

measure objective sleep patterns. Our study used placement on the non-dominant wrist to

estimate sleep parameters. The GT9X devices were programmed to collect data at a sample rate

of 30 Hz. ActiLife software version 6.13.2 or 6.13.4 (ActiGraph, LLC) was used to process and

analyze the raw actigraphy data using the Cole-Kripke algorithm [31]. Wear time was measured

using the Choi algorithm, and a valid wear time included a minimum of ten hours per day, at

least one weekend day, and a minimum of four days [32].

We included sleep efficiency (SE) and total sleep time (TST) in this analysis due to the

prevalent occurrence of sleep disturbances and variations in sleep duration among older adults.

SE refers to the percentage of total time spent asleep during the total time in bed (Time asleep /

Time in Bed). TST refers to the time spent asleep in bed (in minutes). TST was calculated by

subtracting duration to fall asleep and awakenings from the total amount of time in bed (in

minutes; Out of Bed Time – In Bed Time). A higher TST means a longer sleep duration.

The Pittsburgh Sleep Quality Index

The Pittsburgh Sleep Quality Index (PSQI) is a self-report questionnaire consisting of 10

items measuring sleep quality over the past month. The overall score of the PSQI ranges from 0

("good sleep quality") to 21 ("poor sleep quality"). The PSQI is a reliable measure to distinguish

between good and poor sleepers (PSQI > 5) with a sensitivity of 89.6% and a specificity of

86.5% [33]. It is also valid to use for clinical and research purposes.

Cognitive Performance

As part of the Clinical Cohort study annual evaluation, participants completed a cognitive

test battery, specifically the National Alzheimer's Coordinating Center (NACC) Uniform Data

Set (UDS), and several additional tests. The UDS was designed to measure cognitive

performance in mild cognitive impairment or dementia due to AD by the Alzheimer's Disease

Centers Clinical Task Force established by the National Institute on Aging. The subtests included

in the present analyses were the Wechsler Adult Intelligence Scales-Revised (WAIS-R; Digit

Symbol Substitution Test) [34], two subtests from the Wechsler Memory Scales-Revised (WMS-

R; Letter Number Sequencing and Digit Span Forward and Backward) [35], the Stroop Test

(interference condition) [36], the Craft Story 21 Immediate and Delayed recall [37], the Free and

Cued Selective Reminding Test [38], two tests of semantic verbal fluency (animal and vegetable

naming) [39], and Trail Making Test B [40]. The administration procedures of the subtests are

described in their respective references.

The present study used cognitive factor scores derived from a confirmatory factor

analysis. This approach reduces type 1 error resulting from multiple testing and improves

measurement accuracy by combining common variance across multiple subtests while

accounting for measurement error. The details of this analysis can be found elsewhere [41]. The

result yielded three factors: verbal memory (immediate and delayed logical memory, selective

reminding test trials sum), attention (digits forward, digits backward, letter-number sequencing),

and executive function (category fluency sum of animals and vegetables, Stroop color word

interference, Trail Making Test B, and digit symbol substitution test).

Statistical Analysis

We implemented generalized additive models (GAMs) with the mgcv package in R to

analyze the relationship between sleep and each cognitive domain (executive function, verbal

memory, and attention). Because previous studies have suggested that men and women have

different sleep patterns and perceptions of sleep quality, we chose to perform the analyses

separately for men and women to allow for clearer interpretations. We selected GAMs because

previous research has shown that sleep and cognition were non-linearly correlated in older adults

[42]. GAMs are suitable for accounting for non-parametric and non-linear relationships. The

spline fitting, a method to find the best fit of the data points, was estimated using Restricted

Maximum Likelihood (REML). REML is particularly adept at reducing bias in the estimation of

variance components. It adjusts for the degrees of freedom used by the fixed effects, allowing for

more accurate and less biased estimates of the random effects variance. There was 16.4% ($n =$

34) of missing data in the PSQI variable and was imputed with multiple imputation chained

equation with the mice package in R. The imputation was performed by dividing the dataset into

two: cognitively healthy and cognitively impaired groups. This was done because previous

research has shown that cognitively impaired older adults have skewed awareness resulting in

inaccurate self-reporting [43]. Other variables, such as age, years of education, and CDR, that

affect cognitive performance were added as covariates.

193   Cognition ~ s(total sleep time) + s(sleep efficiency) + s(PSQI, by=CDR) +

194   Age + Education + CDR

Results

The summary of participants' demographics is presented in Table 1. Most participants

were White (93.70%), educated (16.4 years), and cognitively normal (i.e., CDR = 0) (77.78%).

Our results revealed that women outperformed men in verbal memory ($\beta = 0.330$, $p = 0.029$)

accounting for age, educational level, and CDR score. Consistent with previous findings, women

reported more sleep problems ($\beta = 0.995$, $p = 0.010$) on the PSQI after adjusting for age and

CDR score. We did not find significant differences in sleep duration and sleep efficiency

between older men and women as measured by the actigraphy.

Table 1. Descriptive Statistics on Demographics, Sleep, and Cognitive Performance

Participants characteristics	Men (*n* = 89)		Women (*n* = 118)		Total (*N*= 207)	
	M	*SD*	*M*	*SD*	*M*	*SD*
Age (years)	76.1	6.9	74	6.6	74.9	6.8
Education (years)	16.7	2.8	16.1	3	16.4	2.9
	N	*%*	*N*	*%*	*N*	*%*
Race						
White	86	96.6	108	91.5	194	93.7
African American	2	2.3	8	6.8	10	4.8
Asian	1	1.1	2	1.7	3	1.5
CDR = 0	63	70.8	98	83.1	161	77.78
	M	*SD*	*M*	*SD*	*M*	*SD*
TST (minutes)	409.7	68.3	419.8	60.4	415.5	64
PSQI	5.8 *	2.5	6.8 *	2.9	6.4	2.7
SE (%)	85.4	7	86.6	6	86.1	6.4
Verbal Memory	0.4 *	1.3	0.9 *	1.2	0.7	1.3
Executive Function	0.1	0.9	0.4	0.8	0.3	0.8
Attention	0.1	0.4	0.2	0.4	0.2	0.4

CDR = Clinical Dementia Rating. TST = total sleep time. PSQI = Pittsburgh Sleep Quality
Index. SE = sleep efficiency. M = mean. SD = standard deviation.
* Differences between men and women, *p* < 0.05

Sleep and Cognitive Function

The General Additive Models (GAMs) revealed significant correlations between sleep

parameters and cognitive function in older adults, after accounting for age, education, and CDR

score. In older women, total sleep time (TST) showed a non-linear association with executive

function (edf = 2.178, *p* = 0.01; Table 2). this relationship demonstrated stable executive

function performance up to a TST of 400 minutes, beyond which a decrease in executive

function was observed (Fig 1). Additionally, TST presented a negative linear connection with

verbal memory (edf = 1.0, *p* = 0.043) and attention (edf = 1.0, *p* = 0.078) as shown in Figs 2 and

3. This indicated that TST contributed to lower performance, although the relationship was not

significant in the case of attention. Contrarily, for older men, TST did not significantly influence

cognitive performance across the three domains: executive function (edf = 3.233; p = 0.167; Fig

4), verbal memory (edf = 1.885; p = 0.296; Fig 5), and attention (edf = 1.001; p = 0.789; Fig 6).

Table 2. Generalized Additive Models of the Relationship Between Sleep Quality (Total
Sleep Time, Sleep Efficiency, and PSQI) and Cognitive Performance (Executive Function,
Verbal Memory, and Attention)
220

	Executive Function		Verbal Memory		Attention	
	Women	Men	Women	Men	Women	Men
Spline fit	EDF	EDF	EDF	EDF	EDF	EDF
TST	2.178**	1.000	1.000*	1.000	1.000	1.000
SE	1.872	1.000*	1.000	1.000	2.257	1.000
PSQI (CDR =0)	1.000*	1.000	2.4	1.000	1.000	1.000
PSQI (CDR =1)	1.450	2.178***	1.000	2.102**	1.794	2.041
Covariate	**Estimate**	**Estimate**	**Estimate**	**Estimate**	**Estimate**	**Estimate**
Age	-0.021**	-0.031**	-0.062***	-0.034*	-0.017**	-0.016**
Education	0.045**	0.022	0.083**	0.004	0.019	0.015
CDR	-0.936***	-1.037***	-1.152***	-1.609***	-0.308**	-0.444***

221   Dependent variables: executive function, verbal memory, and attention. TST = total sleep time. SE =
222   sleep efficiency. CDR = Clinical Dementia Rating. EDF = effective degrees of freedom.
223   *p < 0.05, ** p<0.01, *** p<0.001
224

225   Sleep efficiency demonstrated non-linear associations with both executive function (edf =

3.233; p = 0.167) and attention (edf = 3.233; p = 0.167), while it was linearly related to verbal

memory (edf =1.000; p =0.0795) in older women (Figs 1-3). However, these associations did not

reach statistical significance. Intriguingly, in older men, sleep efficiency positively correlated

with executive function in a linear manner, a relationship that was marginally significant (edf =

1.000, p = 0.050; Fig 4). The same pattern was observed between sleep efficiency and both

verbal memory (edf = 1.000, p = 0.559) and attention (edf = 1.000, p = 0.123) in this group,

although these relationships were not significant (Figures 5-6).

233   For self-report measured by the PSQI, our findings suggest that total PSQI scores showed

non-linear associations with verbal memory (edf = 2.4; p = 0.159) in cognitively intact (CDR =

0) older women. This relationship was characterized by a relatively stable executive function up

to a PSQI score of 8, after which a decline in performance was observed. In addition, the PSQI

had a linear negative relationship with executive function (edf = 1.000; p = 0.024) and attention

(edf =1.000; p = 0.104; Figs 1-3). However, the relationship was only significant with executive

function indicating poorer subjective sleep quality, marked by a higher PSQI score, was

associated with lower performance. In men, the PSQI did not significantly correlate with

cognitive performance across the three domains: executive function (edf = 1.000; p = 0.625; Fig

4), verbal memory (edf = 1.000; p = 0.352; Fig 5), and attention (edf = 1.000; p = 0.929; Fig 6).

Remarkably, among cognitively impaired participants, non-linear positive relationships emerged

between PSQI scores and executive function (edf = 1.450; p = 0.143; Fig 1) and attention (edf =

1.794; p = 0.564; Fig 3) in women. The same pattern was observed in men's executive function

(edf = 2.178, p = 0.0002; Fig 4) and verbal memory (edf = 2.102, p = 0.002; Fig 5), with the

notable difference that these associations were statistically significant.

Discussion

Our study examined whether objective and subjective sleep quality, as measured by

actigraphy and self-report questionnaire, impact cognitive performance, specifically verbal

memory, executive function, and attention, within the older adult population. Concurrently, we

investigated potential sex-related disparities in this association. Our data revealed noteworthy

distinctions in sleep between sexes, with women reporting more sleep-related symptoms,

consistent with previous findings [20,44]. However, we did not find significant differences in

total sleep time and sleep efficiency between the two sexes. Further research is needed to

determine what constitutes optimal sleep for older women and men.

Based on our analysis, we found that total sleep time affected older women but not older

men in executive function and memory, while attention was not affected in both sexes. The

presence of both non-linear and linear relationships across various cognitive areas suggests that

sleep's influence on cognition is not uniform but rather specific to distinct cognitive domains.

This implies that each aspect of cognition may have its own threshold and optimal conditions for

sleep duration and quality. To elucidate the causes of these varied associations, more research is

required to unravel the complexities of how different cognitive functions are differentially

affected by sleep. Meanwhile, low sleep efficiency was associated with lower executive function

in older men. These findings suggest that sleep duration and sleep efficiency may have different

impacts on cognitive function between older women and men. Additionally, the results could

indicate that sleep could have more pronounced effects on higher-order cognitive processes, such

as memory and executive function [18] and less with attention that is considered a more basic

cognitive process. Prior studies reported that acute sleep loss impacts different types of attention

from the one focused on in our study, such as sustained attention, processing speed, and

vigilance [45,46]. It is noteworthy that our study concentrated on attention more closely aligned

with working memory and short-term attention span, which could contribute to different findings

from previous research.

We noted that men with Mild Cognitive Impairment (MCI) who reported more sleep

disturbances tended to exhibit better cognitive performance. This observation raises certain

considerations. Existing research has proposed that those with cognitive impairment might have

diminished self-awareness (i.e., anosognosia; [43], potentially leading to inaccuracies in their

self-reports. Additionally, previous research suggests that women generally have a more accurate

perception of their sleep quality, which may contribute to the sex differences we observed. This

cautions against relying solely on self-report data when assessing cognitively impaired

individuals and highlights the importance of obtaining collateral information from caregivers.

Another plausible explanation could be that sleep may no longer play as significant a role in

cognition because these individuals may have already surpassed a critical threshold in their

cognitive decline.

One main difference between our study and most studies that investigate sleep and

cognition is that prior research typically induces sleep loss to investigate the effects of acute or

chronic sleep loss on cognitive performance, resulting in a significant impact on memory and

executive function [28,29,47], as opposed to natural observations in our study. Furthermore, the

majority of our participants exhibited ideal sleep efficiency levels (around 85%). Studies that

include those with sleep disorders (e.g., insomnia, sleep apnea) may yield different results and

provide insights into the cognitive sequelae of clinically impaired sleep duration and efficiency.

Our study has several strengths. First, we used a comprehensive neuropsychological

battery to gauge participants' cognitive performance, opting for a more in-depth assessment

rather than relying on a simple test battery (e.g., the Mini Mental State Examination), which is

less sensitive to subtle cognitive impairment. Second, we included composite scores of cognitive

domains generated through confirmatory factor analysis to represent each cognitive domain

using multiple tests. This approach allows for better representation of each cognitive domain.

Third, we employed generalized additive models to capture non-linear relationships between

sleep quality and cognitive performance without imposing symmetrical curves. Moreover,

GAMs have the built-in capability to determine the appropriate level of smoothness for each

predictor, ensuring an optimal fit. Fourth, we objectively measured total sleep time and sleep

efficiency over multiple nights allowing for a more representative result of typical sleep and

compared it to self-reported measures to evaluate distinctions between the two common types of

measures.

Several limitations should also be acknowledged. First, the present study is limited by the

characteristics of the sample. The participants were primarily White, highly educated, relatively

healthy, and motivated to engage in research leading to limited generalizability of the findings.

Most of the participants in this study had relatively healthy sleep patterns. Second, our

participants did not undergo comprehensive check-ups to assess for sleep disorders, which are

prevalent in older adults. Third, we did not differentiate between how total sleep time and sleep

efficiency might distinctly impact cognitive performance in individuals with intact cognition

compared to those with mild cognitive impairment. This aspect remains unexplored in our study,

highlighting an area for further research to understand the nuanced effects of different sleep

parameters on cognitive functions across varying cognitive states.

Our study has yielded foundational evidence suggesting that sleep exerts differing effects

on cognitive health in older women and men. It also appears that sleep may impact distinct

cognitive domains. Nevertheless, we have yet to determine the optimal sleep characteristics

required to maintain cognitive health in individuals of different sexes. Future research should

explore the long-term effects of midlife hormonal changes on sleep and cognitive health in aging

populations since this period is notable for hormonal shifts. Midlife may serve as an optimal

window for intervention, as cognitive decline resulting from sleep deprivation tends to be less

severe at this stage. The implications of future findings have the potential to shape public health

policies and strategies for aging. Healthcare professionals should consider recommending regular

sleep assessments as part of a comprehensive check-up to promote cognitive well-being in aging

populations.

References

1.   Scullin MK, Bliwise DL. Sleep, Cognition, and Normal Aging: Integrating a Half-Century
of Multidisciplinary Research. Perspect Psychol Sci J Assoc Psychol Sci. 2015;10: 97–137.
doi:10.1177/1745691614556680

2.   Zhang B, Wing Y-K. Sex differences in insomnia: a meta-analysis. Sleep. 2006;29: 85–93.
doi:10.1093/sleep/29.1.85

3.   Rajan KB, Weuve J, Barnes LL, McAninch EA, Wilson RS, Evans DA. Population
estimate of people with clinical Alzheimer's disease and mild cognitive impairment in the
United States (2020–2060). Alzheimers Dement. 2021;17: 1966–1975.
doi:10.1002/alz.12362

4.   Spira AP, Chen-Edinboro LP, Wu MN, Yaffe K. Impact of Sleep on the Risk of Cognitive
Decline and Dementia. Curr Opin Psychiatry. 2014;27: 478–483.
doi:10.1097/YCO.0000000000000106

5.   Bliwise DL, Foley DJ, Vitiello MV, Ansari FP, Ancoli-Israel S, Walsh JK. Nocturia and
disturbed sleep in the elderly. Sleep Med. 2009;10: 540–548.
doi:10.1016/j.sleep.2008.04.002

6.   Ohayon MM, Carskadon MA, Guilleminault C, Vitiello MV. Meta-analysis of quantitative
sleep parameters from childhood to old age in healthy individuals: developing normative
sleep values across the human lifespan. Sleep. 2004;27: 1255–1273.
doi:10.1093/sleep/27.7.1255

7.   Ancoli-Israel S. Sleep and its disorders in aging populations. Sleep Med. 2009;10 Suppl 1:
S7-11. doi:10.1016/j.sleep.2009.07.004

8.   Li J, Gooneratne NS. Sleep and health in older adults. Sleep and health. San Diego, CA,
US: Elsevier Academic Press; 2019. pp. 31–43. doi:10.1016/B978-0-12-815373-4.00004-6

9.   Nguyen V, George T, Brewster GS. Insomnia in Older Adults. Curr Geriatr Rep. 2019;8:
271–290. doi:10.1007/s13670-019-00300-x

10.   Braak H, Braak E. Neuropathological stageing of Alzheimer-related changes. Acta
Neuropathol (Berl). 1991;82: 239–259. doi:10.1007/BF00308809

11.   Rodrigue KM, Kennedy KM, Devous MD, Rieck JR, Hebrank AC, Diaz-Arrastia R, et al.
β-Amyloid burden in healthy aging: regional distribution and cognitive consequences.
Neurology. 2012;78: 387–395. doi:10.1212/WNL.0b013e318245d295

12.   Winer JR, Deters KD, Kennedy G, Jin M, Goldstein-Piekarski A, Poston KL, et al.
Association of Short and Long Sleep Duration With Amyloid-β Burden and Cognition in
Aging. JAMA Neurol. 2021;78: 1187–1196. doi:10.1001/jamaneurol.2021.2876

13. Kondo R, Miyano I, Lee S, Shimada H, Kitaoka H. Association between self-reported night
sleep duration and cognitive function among older adults with intact global cognition. Int J
Geriatr Psychiatry. 2021;36: 766–774. doi:10.1002/gps.5476

14. Sabeti S, Al-Darsani Z, Mander BA, Corrada MM, Kawas CH. Sleep, hippocampal volume,
and cognition in adults over 90 years old. Aging Clin Exp Res. 2018;30: 1307–1318.
doi:10.1007/s40520-018-1030-x

15. Tworoger SS, Lee S, Schernhammer ES, Grodstein F. The association of self-reported sleep
duration, difficulty sleeping, and snoring with cognitive function in older women.
Alzheimer Dis Assoc Disord. 2006;20: 41–48. doi:10.1097/01.wad.0000201850.52707.80

16. Faubel R, López-García E, Guallar-Castillón P, Graciani A, Banegas JR, Rodríguez-
Artalejo F. Usual sleep duration and cognitive function in older adults in Spain. J Sleep
Res. 2009;18: 427–435. doi:10.1111/j.1365-2869.2009.00759.x

17. Blackwell T, Redline S, Ancoli-Israel S, Schneider JL, Surovec S, Johnson NL, et al.
Comparison of Sleep Parameters from Actigraphy and Polysomnography in Older Women:
The SOF Study. Sleep. 2008;31: 283–291. doi:10.1093/sleep/31.2.283

18. Wilckens KA, Woo SG, Kirk AR, Erickson KI, Wheeler ME. The role of sleep continuity
and total sleep time in executive function across the adult lifespan. Psychol Aging. 2014;29:
658–665. doi:10.1037/a0037234

19. Kim SJ, Lee DK, Jang YK, Jang H, Kim SE, Cho SH, et al. The Effects of Longitudinal
White Matter Hyperintensity Change on Cognitive Decline and Cortical Thinning over
Three Years. J Clin Med. 2020;9: 2663. doi:10.3390/jcm9082663

20. Morris JL, Rohay J, Chasens ER. Sex Differences in the Psychometric Properties of the
Pittsburgh Sleep Quality Index. J Womens Health. 2018;27: 278–282.
doi:10.1089/jwh.2017.6447

21. Kocevska D, Lysen TS, Dotinga A, Koopman-Verhoeff ME, Luijk MPCM, Antypa N, et al.
Sleep characteristics across the lifespan in 1.1 million people from the Netherlands, United
Kingdom and United States: a systematic review and meta-analysis. Nat Hum Behav.
2021;5: 113–122. doi:10.1038/s41562-020-00965-x

22. Hume KI, Van F, Watson A. A field study of age and gender differences in habitual adult
sleep. J Sleep Res. 1998;7: 85–94. doi:10.1046/j.1365-2869.1998.00103.x

23. Yoon JE, Oh D, Hwang I, Park JA, Im HJ, Lee SK, et al. Sleep structure and
electroencephalographic spectral power of middle-aged or older adults: Normative values
by age and sex in the Korean population. J Sleep Res. 2021;30. doi:10.1111/jsr.13358

24. Albinni B, Baker FC, Javitz H, Hasler BP, Franzen PL, Clark DB, et al. Morning perception
of sleep, stress, and mood, and its relationship with overnight physiological sleep: findings
from the National Consortium on Alcohol and Neurodevelopment in Adolescence
(NCANDA) study. J Sleep Res. 2023;32: e13886. doi:10.1111/jsr.13886

25.   Della Monica C, Johnsen S, Atzori G, Groeger JA, Dijk D-J. Rapid Eye Movement Sleep,
Sleep Continuity and Slow Wave Sleep as Predictors of Cognition, Mood, and Subjective
Sleep Quality in Healthy Men and Women, Aged 20-84 Years. Front Psychiatry. 2018;9:
255. doi:10.3389/fpsyt.2018.00255

26.   Groeger JA, Zijlstra FRH, Dijk D-J. Sleep quantity, sleep difficulties and their perceived
consequences in a representative sample of some 2000 British adults. J Sleep Res. 2004;13:
359–371. doi:10.1111/j.1365-2869.2004.00418.x

27.   Vitiello MV, Larsen LH, Moe KE. Age-related sleep change: Gender and estrogen effects
on the subjective-objective sleep quality relationships of healthy, noncomplaining older
men and women. J Psychosom Res. 2004;56: 503–510. doi:10.1016/S0022-3999(04)00023-
6

28.   Rångtell FH, Karamchedu S, Andersson P, Liethof L, Olaya Búcaro M, Lampola L, et al. A
single night of sleep loss impairs objective but not subjective working memory performance
in a sex-dependent manner. J Sleep Res. 2019;28: e12651. doi:10.1111/jsr.12651

29.   Santhi N, Lazar AS, McCabe PJ, Lo JC, Groeger JA, Dijk D-J. Sex differences in the
circadian regulation of sleep and waking cognition in humans. Proc Natl Acad Sci U S A.
2016;113: E2730-2739. doi:10.1073/pnas.1521637113

30.   Baker FC, Sattari N, de Zambotti M, Goldstone A, Alaynick WA, Mednick SC. Impact of
sex steroids and reproductive stage on sleep-dependent memory consolidation in women.
Neurobiol Learn Mem. 2019;160: 118–131. doi:10.1016/j.nlm.2018.03.017

31.   Cole RJ, Kripke DF, Gruen W, Mullaney DJ, Gillin JC. Automatic sleep/wake
identification from wrist activity. Sleep. 1992;15: 461–469. doi:10.1093/sleep/15.5.461

32.   Choi L, Liu Z, Matthews CE, Buchowski MS. Validation of accelerometer wear and
nonwear time classification algorithm. Med Sci Sports Exerc. 2011;43: 357–364.
doi:10.1249/MSS.0b013e3181ed61a3

33.   Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh sleep quality
index: A new instrument for psychiatric practice and research. Psychiatry Res. 1989;28:
193–213. doi:10.1016/0165-1781(89)90047-4

34.   Wechsler D. Wechsler Adult Intelligence Scale. 3rd (Ed.) San Antonio: The Psychological
Corporation.[Google Scholar]. 1997.

35.   Wechsler D. The Wechsler Memory Scale, San Antonio, Tex, Psychological Corp.
Harcourt; 1997.

36.   Stroop JR. Studies of interference in serial verbal reactions. J Exp Psychol. 1935;18: 643–
662. doi:10.1037/h0054651

37. Craft S, Newcomer J, Kanne S, Dagogo-Jack S, Cryer P, Sheline Y, et al. Memory
improvement following induced hyperinsulinemia in Alzheimer's disease. Neurobiol
Aging. 1996;17: 123–130. doi:10.1016/0197-4580(95)02002-0

38. Grober E, Buschke H, Crystal H, Bang S, Dresner R. Screening for dementia by memory
testing. Neurology. 1988;38: 900–903. doi:10.1212/wnl.38.6.900

39. Goodglass H, Kaplan E. The assessment of aphasia and related disorders. 2nd ed.
Philadelphia, PA: Lea & Febiger; 1983. Available:
https://cir.nii.ac.jp/crid/1130282270892335488

40. Reitan RM, Wolfson D. The Halstead-Reitan Neuropsychological Test Battery: Theory and
Clinical Interpretation. Neuropsychology Press; 1985.

41. Watts A. Creating cognitive factor scores from KU-ADC data. 2020.

42. Lucey BP, Wisch J, Boerwinkle AH, Landsness EC, Toedebusch CD, McLeland JS, et al.
Sleep and longitudinal cognitive performance in preclinical and early symptomatic
Alzheimer's disease. Brain. 2021;144: 2852–2862. doi:10.1093/brain/awab272

43. De Carolis A, Cipollini V, Corigliano V, Comparelli A, Sepe-Monti M, Orzi F, et al.
Anosognosia in People with Cognitive Impairment: Association with Cognitive Deficits
and Behavioral Disturbances. Dement Geriatr Cogn Disord Extra. 2015;5: 42–50.
doi:10.1159/000367987

44. Kim H, Kim R, Kim S, Kim S, Kim S, Lee S, et al. Sex differences in deterioration of sleep
properties associated with aging: A 12-year longitudinal cohort study. J Clin Sleep Med.
2021;17. doi:10.5664/jcsm.9072

45. Goel N, Dinges DF. Behavioral and Genetic Markers of Sleepiness. J Clin Sleep Med.
2011;7: S19–S21. doi:10.5664/JCSM.1348

46. Lo JC, Groeger JA, Santhi N, Arbon EL, Lazar AS, Hasan S, et al. Effects of partial and
acute total sleep deprivation on performance across cognitive domains, individuals and
circadian phase. PloS One. 2012;7: e45987. doi:10.1371/journal.pone.0045987

47. Hennecke E, Lange D, Steenbergen F, Fronczek-Poncelet J, Elmenhorst D, Bauer A, et al.
Adverse interaction effects of chronic and acute sleep deficits on spatial working memory
but not on verbal working memory or declarative memory. J Sleep Res. 2021;30: e13225.
doi:10.1111/jsr.13225

Fig 1. Partial effects of total sleep time, sleep efficiency, and Pittsburgh Sleep Quality Index (PSQI) on executive function in women.

Fig 1

Fig 2. Partial effects of total sleep time, sleep efficiency, and Pittsburgh Sleep Quality Index (PSQI) on verbal memory in women.

Fig 2

Fig 3. Partial effects of total sleep time, sleep efficiency, and Pittsburgh Sleep Quality Index (PSQI) on attention in women.

Fig 3

Fig 4. Partial effects of total sleep time, sleep efficiency, and Pittsburgh Sleep Quality Index (PSQI) on executive function in men.

Fig 4

Fig 5. Partial effects of total sleep time, sleep efficiency, and Pittsburgh Sleep Quality Index (PSQI) on verbal memory in men.

Fig 5

Fig 6. Partial effects of total sleep time, sleep efficiency, and Pittsburgh Sleep Quality Index (PSQI) on attention in men.

Fig 6