Sex Differences in the Role of Sleep on Cognition in Older Adults

Yumiko Wiranto[1], Catherine Siengsukon[2], Diego R. Mazzotti[3,4],

Jeffrey M. Burns[5], and Amber Watts[1,5]

[1]Department of Psychology, University of Kansas, Lawrence, Kansas, United States of America

[2]University of Kansas Medical Center, Department of Physical Therapy and Rehabilitation Science, Kansas City, KS USA

[3]Division of Medical Informatics, Department of Internal Medicine, University of Kansas Medical Center

[4]Division of Pulmonary Critical Care and Sleep Medicine, Department of Internal Medicine, University of Kansas Medical Center

[5]University of Kansas, Alzheimer's Disease Research Center, Fairway, Kansas, United States of America

* Corresponding author

Yumiko Wiranto

Department of Psychology, University of Kansas, 1415 Jayhawk Boulevard, Lawrence, KS 66044

Email: yumiko.wiranto@ku.edu

Phone: +1 785-864-4131

Abstract

Study Objectives: The study aimed to investigate sex differences in the relationship between sleep quality (self-report and objective) and cognitive function across three domains (executive function, verbal memory, and attention) in older adults.

Methods: We analyzed cross-sectional data from 207 participants with normal cognition or mild cognitive impairment (89 males and 118 females) aged over 60. The relationship between sleep quality and cognitive performance was estimated using generalized additive models. Objective sleep was measured with the GT9X Link Actigraph, and self-reported sleep was measured with the Pittsburgh Sleep Quality Index.

Results: We found that females exhibited stable performance of executive function with up to 400 minutes of total sleep time, with significant declines in performance ($p = 0.010$) when sleep duration was longer. Additionally, a longer total sleep time contributed to lower verbal memory in a linear manner ($p = 0.043$). Higher self-reported sleep complaints were associated with poorer executive function in females with healthy cognition ($p = 0.024$). In males, a positive linear relationship emerged between sleep efficiency and executive function ($p = 0.05$), while self-reported sleep was not associated with cognitive performance.

Conclusions: Our findings suggest that the relationships between sleep quality and cognition differ between older males and females, with executive function being the most influenced by objective and self-reported sleep. Interventions targeting sleep quality to mitigate cognitive decline in older adults may need to be tailored according to sex, with distinct approaches for males and females.

Keywords: Sleep, cognition, aging, sex differences, actigraphy

Introduction

The impact of aging on sleep and cognition is well documented in the literature. Aging has been associated with fragmented sleep and decreased sleep quality, which could exacerbate cognitive decline that occurs with aging [1]. However, the literature on sex differences in sleep and its interaction with cognition in older adults is still limited. Females experience dramatic hormonal and physiologic developmental changes (puberty, pregnancy, and menopause) across the lifespan that contribute to alterations in sleep patterns and a higher number of reported sleep complaints [2]. Moreover, females have twice the risk of developing Alzheimer's disease (AD), for which poor sleep is a risk factor [3,4]. Understanding how males and females differ in their sleep health and the implications of these differences on cognition is vital for designing effective interventions and prevention strategies for AD.

Advancing age is associated with physiological changes, among which are the alterations of sleep architecture (i.e., stages of sleep) and circadian rhythm. Older adults have a shorter slow-wave sleep duration and activity, as well as shorter rapid eye movement sleep and more frequent awakenings during the night [5,6]. Additionally, they tend to go to bed earlier in the evening and wake up earlier in the morning [7]. These changes contribute to the high prevalence of sleep disturbances experienced by this population. Previous studies reported that up to 75% of older adults experienced sleep difficulty [8–10]. Because one of the roles of sleep is to clear out neurotoxic waste products produced during the day, poor sleep has been found to exacerbate cognitive decline and neuropathological markers of AD, such as beta-amyloid and tau deposition [11–13].

The investigation of the interrelationships between aging, sleep, and cognition has been relatively inconclusive and marked by variability in findings, potentially due to methodological variations and differences in operationalizing sleep metrics across studies. In older adults with

normal cognition, both short (< 6 hours) and long (> 9 hours) sleep durations have been linked to reduced global cognition and memory [14,15]. However, Sabeti et al. [16] observed this association solely in older adults with cognitive impairment. In a two-year longitudinal study, it was found that older females who slept for only four hours exhibited a more significant cognitive decline compared to those who slept for seven hours. Notably, individuals who slept for nine hours did not experience any cognitive decline during the same period [17]. Conversely, Faubel et al. [18] indicated that older adults who slept for 11 hours demonstrated lower cognitive performance compared to those who slept for seven hours. In addition to global cognition and memory, sleep disturbances have been reported to impact other cognitive domains, including attention and executive function, in older populations. Specifically, Blackwell et al. [19] and Wilckens et al. [20] revealed a negative correlation between frequent nighttime awakenings and performance on executive function and attention. However, Wilckens et al. [20] did not find a significant association between sleep duration and cognitive performance in their study.

Biological sex is another major factor that influences sleep. Research investigating sex differences in self-reported sleep quality suggests that middle-aged and older females take longer to fall asleep, have shorter sleep duration, and have lower sleep efficiency [21,22]. However, this result is contradicted by studies looking at objective sleep quality between older males and females using actigraphy and polysomnography (PSG). A meta-analysis including studies on sleep characteristics across three countries reported that older females had longer sleep duration and higher sleep efficiency based on actigraphy data [23]. Studies using PSG show that reduced slow-wave sleep due to age is more pronounced in males than females [24,25]. The variation between self-reported and objective sleep quality may be attributed to sex-related differences in sleep perception and symptom reporting. Recent research suggests a stronger link between

objective and self-reported sleep quality in females compared to males [26,27], which may reflect a closer alignment between females' self-reported sleep experiences and objective sleep parameters. Additionally, it raises questions about whether the current criteria for defining good objective sleep quality apply equally to both sexes, suggesting a need for distinct criteria for females [28,29].

Regarding the cognitive implications of poor sleep, limited research has been conducted to investigate sex differences in this topic. Notably, it has been observed that sleep deprivation affects working memory to a greater extent in young females compared to males [30,31]. However, whether the same findings apply to older populations is still not well understood. This lack of research leaves an important gap in our understanding of how sleep impacts cognitive function across the lifespan. Some possible mechanistic explanations for sex differences in this relationship include depletion in sex steroids (estradiol) due to menopause that could adversely affect cognition, specifically memory processing [32]. Moreover, since sleep is known to influence memory performance, the connection between reduced estradiol levels due to menopause and memory might be further intensified by poor sleep.

The present study investigated sex differences in the relationship between sleep quality and cognitive performance in a clinical cohort of older individuals. We assessed sleep quality using a self-report questionnaire and actigraphy, measuring total sleep time and sleep efficiency. We hypothesized that sleep quality would be more highly associated with cognitive performance in older females than it would in older males.

Materials and Methods

Participants

Participants were recruited from the University of Kansas Alzheimer's Disease Research Center (KU-ADRC) Clinical Cohort, an ongoing longitudinal observational study to support regional and national research on aging, cognition, and AD. This study collects demographic, medical, psychological, and cognitive data annually, and imaging and biospecimen data as needed if participants opt in. Participants could opt to partake in the Physical Activity and Sleep Study (PASS), a sub-study of the Clinical Cohort initiated on July 29, 2015, in which participants completed self-report sleep questionnaires annually and wore an actigraphy every two years until May 27, 2021. During this time, the KU-ADRC Clinical Cohort enrolled over 1,000 participants. To be eligible for the Clinical Cohort study, participants had to be at least 60 years old, native English speakers, and not exhibit the following conditions: significant depressive symptoms, untreated thyroid dysfunction, visual or auditory impairment, active (< 2 years) ischemic heart disease, and uncontrolled insulin-dependent diabetes mellitus.

In the current analysis, we included 207 participants with normal cognition or mild impairment as indicated by clinical dementia rating (CDR) scores of 0 and >0, respectively. Among these participants, 161 exhibited normal cognitive function, while 46 had mild cognitive impairment with a CDR of 1. The research was approved by the KU-ADRC's Institutional Review Board, and all participants provided written informed consent separately for both the KU-ADRC Clinical Cohort and the PASS study.

Sleep Quality Measures

Actigraphy

ActiGraph GT9X Link accelerometers (Pensacola, FL) are body-worn monitors that measure objective sleep patterns. Our study used placement on the non-dominant wrist to estimate sleep parameters. The GT9X devices were programmed to collect data at a sample rate

of 30 Hz. ActiLife software version 6.13.2 or 6.13.4 (ActiGraph, LLC) was used to process and analyze the raw actigraphy data using the Cole-Kripke algorithm [33]. Wear time was measured using the Choi algorithm, and a valid wear time included a minimum of ten hours per day, at least one weekend day, and a minimum of four days [34]. Participants were also provided with a sleep diary to document the timing and length of their sleep during the day and at night. They were asked to note how long it took them to fall asleep, record any instances of waking up during the night, and track the time they woke up and got out of bed.

We included sleep efficiency (SE) and total sleep time (TST) in this analysis due to the prevalent occurrence of sleep disturbances and variations in sleep duration and sleep efficiency among older adults. SE refers to the percentage of total time spent asleep during the total time in bed (Time asleep / Time in Bed). TST refers to the time spent asleep in bed (in minutes) at night. TST was calculated by subtracting the duration of falling asleep and awakening from the total amount of time in bed (in minutes; Out of Bed Time – In Bed Time). A higher TST means a longer sleep duration. For each night's data, sleep intervals recorded by ActiGraph were compared with the participants' sleep diaries to check for discrepancies. When the sleep diary entries were closely aligned (within a 30-minute margin) with the ActiGraph-identified sleep period, the sleep period identified by ActiGraph was used. In cases where the discrepancy exceeded 30 minutes, the researchers manually evaluated the sleep data in ActiLife by looking for significant changes in activity levels. This manual assessment was also automatically initiated whenever participants did not submit or complete their sleep diaries.

The Pittsburgh Sleep Quality Index

The Pittsburgh Sleep Quality Index (PSQI) is a self-report questionnaire consisting of 10 items measuring sleep quality over the past month. The overall score of the PSQI ranges from 0

("good sleep quality") to 21 ("poor sleep quality"). The PSQI is a reliable measure to distinguish between good and poor sleepers (PSQI > 5) with a sensitivity of 89.6% and a specificity of 86.5% [35]. It is also valid to use for clinical and research purposes.

Cognitive Performance

As part of the Clinical Cohort study annual evaluation, participants completed a cognitive test battery, specifically the National Alzheimer's Coordinating Center (NACC) Uniform Data Set (UDS), and several additional tests. The UDS was designed to measure cognitive performance in mild cognitive impairment or dementia due to AD by the Alzheimer's Disease Centers Clinical Task Force established by the National Institute on Aging. The subtests included in the present analyses were the Wechsler Adult Intelligence Scales-Revised (WAIS-R; Digit Symbol Substitution Test) [36], two subtests from the Wechsler Memory Scales-Revised (WMS-R; Letter Number Sequencing and Digit Span Forward and Backward) [37], the Stroop Test (interference condition) [38], the Craft Story 21 Immediate and Delayed recall [39], the Free and Cued Selective Reminding Test [40], two tests of semantic verbal fluency (animal and vegetable naming) [41], and Trail Making Test B [42]. The administration procedures of the subtests are described in their respective references.

The present study used cognitive factor scores derived from a confirmatory factor analysis. This approach reduces type 1 error resulting from multiple testing and improves measurement accuracy by combining common variance across multiple subtests while accounting for measurement error. The details of this analysis can be found elsewhere [43]. The result yielded three factors: verbal memory (immediate and delayed logical memory, selective reminding test trials sum), attention (digits forward, digits backward, letter-number sequencing),

and executive function (category fluency sum of animals and vegetables, Stroop color word interference, Trail Making Test B, and digit symbol substitution test).

Statistical Analysis

We compared demographics (age and education) between males and females using an independent t-test. Analysis of covariance (ANCOVA) was used to compare the three cognitive domains and sleep variables (TST, SE, and PSQI) between sexes, adjusting for age, sex, and CDR, with education included as a covariate exclusively for the cognitive domains. We implemented generalized additive models (GAMs) with the mgcv package in R to analyze the relationship between sleep and each cognitive domain (executive function, verbal memory, and attention). Because previous studies have suggested that males and females have different sleep patterns and perceptions of sleep quality, we chose to perform the analyses separately for males and females to allow for clearer interpretations. We selected GAMs because previous research has shown that sleep and cognition were non-linearly correlated in older adults [44]. GAMs are suitable for accounting for non-parametric and non-linear relationships. The spline fitting, a method to find the best fit of the data points, was estimated using Restricted Maximum Likelihood (REML). REML is particularly adept at reducing bias in the estimation of variance components. It adjusts for the degrees of freedom used by the fixed effects, allowing for more accurate and less biased estimates of the random effects variance. There was 16.4% ($n = 34$) of missing data in the PSQI variable which was imputed with multiple imputation chained equation with the mice package in R. All variables were included in the imputation, except for the dependent variables (i.e., the three cognitive domains). The imputation was performed by dividing the dataset into two: cognitively healthy and cognitively impaired groups. This was done because previous research has shown that cognitively impaired older adults have skewed

awareness resulting in inaccurate self-reporting [45]. Other variables, such as age, years of

education, and CDR, that affect cognitive performance were added as covariates.

Cognition ~ s(total sleep time) + s(sleep efficiency) + s(PSQI, by=CDR) +

Age + Education + CDR

Results

The summary of participants' demographics is presented in Table 1. Most participants

were White (93.70%), college-educated (16.4 years), and cognitively normal (i.e., CDR = 0)

(77.78%). Our results revealed that females outperformed males in verbal memory (β = 0.329, p

= 0.029) accounting for age, educational level, and CDR score. Consistent with previous

findings, females reported more sleep problems (β = 0.994, p = 0.010) on the PSQI after

adjusting for age and CDR score. We did not find significant differences in sleep duration and

sleep efficiency between older males and females as measured by the actigraphy.

Sleep and Cognitive Function

The General Additive Models (GAMs) revealed significant correlations between sleep

parameters and cognitive function in older adults, after accounting for age, education, and CDR

score. In older females, total sleep time (TST) showed a non-linear association with executive

function (edf = 2.178, p = 0.010; Table 2). This relationship demonstrated stable executive

function performance up to a TST of 400 minutes, beyond which a decrease in executive

function was observed (Figure 1). Additionally, TST presented a negative linear relationship

with verbal memory (edf = 1.000, p = 0.042) and attention (edf = 1.000, p = 0.078) as shown in

Figs 2 and 3. This indicated that TST was associated with lower performance, although the

relationship did not reach statistical significance in the case of attention. Contrarily, among older

males, TST was not associated with cognitive performance across the three domains: executive

function (edf = 1.000; p = 0.248; Figure 4), verbal memory (edf = 1.000; p = 0.120; Figure 5),

and attention (edf = 1.000; p = 0.274; Figure 6).

Sleep efficiency demonstrated non-linear associations with both executive function (edf =

1.872; p = 0.201) and attention (edf = 2.257; p = 0.279), while it was linearly related to verbal

memory (edf =1.000; p =0.079) in older females (Figs 1-3). However, these associations did not

reach statistical significance. Intriguingly, in older males, sleep efficiency positively correlated

with executive function in a linear manner, a relationship that was marginally significant (edf =

1.000, p = 0.050; Figure 4). The same pattern was observed between sleep efficiency and both

verbal memory (edf = 1.000, p = 0.559) and attention (edf = 1.000, p = 0.123) in this group,

although these relationships were not significant (Figs 5-6).

Total PSQI scores showed non-linear associations with verbal memory (edf = 2.400; p =

0.158) in older females with normal cognition (CDR = 0). This relationship was characterized by

a relatively stable executive function up to a PSQI score of 8, after which a decline in

performance was observed. In addition, the PSQI had a linear negative relationship with

executive function (edf = 1.000; p = 0.023) and attention (edf =1.000; p = 0.103; Figs 1-3).

However, the relationship was only significant with executive function indicating poorer self-

reported sleep quality, marked by a higher PSQI score, was associated with lower performance.

In males with normal cognition, the PSQI did not significantly correlate with cognitive

performance across the three domains: executive function (edf = 1.000; p = 0.624; Figure 4),

verbal memory (edf = 1.000; p = 0.352; Figure 5), and attention (edf = 1.000; p = 0.929; Figure

6). Among female participants with cognitive impairment, a linear negative relationship emerged

between the PSQI and verbal memory (edf = 1.000; p = 0.398; Figure 5), and the PSQI exhibited

non-linear positive relationships with executive function (edf = 1.450; p = 0.143; Figure 1) and

attention (edf = 1.794; p = 0.564; Figure 3). Non-linear positive relationships were also observed in the PSQI with executive function (edf = 2.178, p = 0.0002; Figure 4), verbal memory (edf = 2.102, p = 0.002; Figure 5), and attention (edf = 2.041; p = 0.078; Figure 6) in males with cognitive impairment, with the notable difference that these associations were statistically significant, except for attention.

Discussion

Our study examined whether objective and self-reported sleep quality, as measured by actigraphy and a self-report questionnaire, were associated with cognitive performance, specifically verbal memory, executive function, and attention, within an older adult clinical population. Concurrently, we investigated potential sex-related disparities in this association. Our data revealed noteworthy distinctions in sleep between sexes, with females reporting more sleep-related symptoms, consistent with previous findings [22,46]. However, we did not find significant differences in total sleep time and sleep efficiency between the two sexes. Further research is needed to determine what constitutes optimal sleep for older females and males.

Based on our analysis, we found that total sleep time was a significant predictor, among older females but not older males, of executive function and memory, while attention was not associated in both sexes. The presence of both non-linear and linear relationships across various cognitive areas suggests that effects were not uniform but rather specific to distinct cognitive domains. This implies that each aspect of cognition may have its own threshold and optimal conditions for sleep duration and quality. To elucidate the sources of this heterogeneity, more research is required to unravel the complexities of how different cognitive functions are differentially affected by sleep and vice-versa. Meanwhile, low sleep efficiency was associated with lower executive function in older males. These findings suggest that sleep duration and

sleep efficiency may have different effects on cognitive function between older females and males. Additionally, the results could indicate that sleep could have more pronounced effects on higher-order cognitive processes, such as memory and executive function [20] and less with attention that is considered a more basic cognitive process. Prior studies reported that acute sleep loss impacts different types of attention from the one focused on in our study, such as sustained attention, processing speed, and vigilance [47,48]. It is noteworthy that our study concentrated on attention more closely aligned with working memory and short-term attention span, which could contribute to different findings from previous research.

We noted that males with mild cognitive impairment (MCI) who reported more sleep disturbances tended to exhibit better cognitive performance. This observation raises certain considerations. Existing research has proposed that those with cognitive impairment might have diminished self-awareness (i.e., anosognosia) [45], potentially leading to inaccuracies in their self-reports. Additionally, previous research suggests that females generally have a more accurate perception of their sleep quality [26,27], which may contribute to the sex differences we observed. This cautions against relying solely on self-report data when assessing individuals with cognitive impairment and highlights the importance of obtaining collateral information from caregivers. Another plausible explanation could be that sleep may no longer play as significant a role in cognition because these individuals may have already surpassed a critical threshold in their cognitive decline.

One main difference between our study and most studies that investigate sleep and cognition is that prior research typically induces sleep loss to investigate the effects of acute or chronic sleep loss on cognitive performance, resulting in a significant impact on memory and executive function [30,31,49], as opposed to real-world observations in our study. Furthermore, the

majority of our participants exhibited ideal sleep efficiency levels (around 85%). Studies that include those with lower sleep efficiency may yield different results and provide insights into the cognitive sequelae of clinically impaired sleep.

Our study has several strengths. First, we used a comprehensive neuropsychological battery to gauge participants' cognitive performance, opting for a more in-depth assessment rather than relying on a simple test battery (e.g., the Mini-Mental State Examination), which is less sensitive to subtle cognitive impairment. Second, we included composite scores of cognitive domains generated through confirmatory factor analysis to represent each cognitive domain using multiple tests. This approach allows for a better representation of each cognitive domain. Third, we employed generalized additive models to capture non-linear relationships between sleep quality and cognitive performance without imposing symmetrical curves. Moreover, GAMs have the built-in capability to determine the appropriate level of smoothness for each predictor, ensuring an optimal fit. Fourth, we objectively measured total sleep time and sleep efficiency over multiple nights allowing for a more representative result of typical sleep and compared it to self-reported measures to evaluate distinctions between the two common types of measures.

Several limitations should also be acknowledged. First, the present study is limited by the characteristics of the sample. The participants were primarily White, highly educated, relatively healthy, and motivated to engage in research, leading to limited generalizability of the findings. Most of the participants in this study had relatively healthy sleep patterns. Second, we did not include daytime napping in the total sleep time because previous research suggests that daytime and nighttime sleep have different impacts on cognitive processes [50]. Therefore, excluding daytime naps helped to isolate the specific effects of nighttime sleep, ensuring a clearer

understanding of how it relates to cognitive performance. Third, we acknowledged the potential

challenges of using the PSQI in individuals with MCI, particularly regarding their ability to

recall sleep details over the past 30 days. Despite these concerns, we found that the instrument

has been extensively used in research involving MCI populations as reported in a literature

review [51], which supports our decision to include participants with normal and mildly impaired

cognition in our analysis, allowing for a more comprehensive exploration of the relationship

between sleep quality and cognitive function across a spectrum of cognitive statuses. Moreover,

the potential influence of cognitive impairment on self-reported sleep quality was mitigated by

incorporating the clinical dementia rating (CDR) into the analysis as a covariate. We also

stratified the relationship between PSQI and cognitive outcomes by CDR levels.

Our study has yielded foundational evidence of the association between sleep and

cognitive health in older females and males, with effects across distinct cognitive domains. Our

findings suggest that studies focused on determining the optimal sleep characteristics required to

maintain cognitive health in individuals of different sexes are warranted. Future research should

also explore the long-term effects of midlife hormonal changes on sleep and cognitive health in

aging populations since this period is notable for hormonal shifts. Midlife may serve as an

optimal window for intervention, as cognitive decline resulting from sleep deprivation tends to

be less severe at this stage. Moreover, future research should extend these findings through

longitudinal analyses, examining how sleep quality and cognitive abilities evolve together over

time. Such studies could provide critical insights into the temporal dynamics of sleep and

cognitive decline, identifying potential periods of vulnerability and opportunity for preventive

interventions. The implications of future findings have the potential to shape public health

policies and strategies for aging. Healthcare professionals should consider recommending regular

sleep assessments as part of a comprehensive check-up to promote cognitive well-being in aging

populations.

sleep assessments as part of a comprehensive check-up to promote cognitive well-being in aging

Disclosure Statements

- Funding: This project was supported by the University of Kansas Alzheimer's Disease Research Center, the National Institute on Aging (grant nos. P30AG035982; R01AG033673), and the University of Kansas General Research Fund.

- Conflict of interest: CS is the owner and Chief Executive Officer of Sleep Health Education, LLC.

References

1. Scullin MK, Bliwise DL. Sleep, Cognition, and Normal Aging: Integrating a Half-Century of Multidisciplinary Research. *Perspect Psychol Sci J Assoc Psychol Sci*. 2015;10(1):97-137. doi:10.1177/1745691614556680

2. Zhang B, Wing YK. Sex differences in insomnia: a meta-analysis. *Sleep*. 2006;29(1):85-93. doi:10.1093/sleep/29.1.85

3. Rajan KB, Weuve J, Barnes LL, McAninch EA, Wilson RS, Evans DA. Population estimate of people with clinical Alzheimer's disease and mild cognitive impairment in the United States (2020–2060). *Alzheimers Dement*. 2021;17(12):1966-1975. doi:10.1002/alz.12362

4. Spira AP, Chen-Edinboro LP, Wu MN, Yaffe K. Impact of Sleep on the Risk of Cognitive Decline and Dementia. *Curr Opin Psychiatry*. 2014;27(6):478-483. doi:10.1097/YCO.0000000000000106

5. Bliwise DL, Foley DJ, Vitiello MV, Ansari FP, Ancoli-Israel S, Walsh JK. Nocturia and disturbed sleep in the elderly. *Sleep Med*. 2009;10(5):540-548. doi:10.1016/j.sleep.2008.04.002

6. Ohayon MM, Carskadon MA, Guilleminault C, Vitiello MV. Meta-analysis of quantitative sleep parameters from childhood to old age in healthy individuals: developing normative sleep values across the human lifespan. *Sleep*. 2004;27(7):1255-1273. doi:10.1093/sleep/27.7.1255

7. Ancoli-Israel S. Sleep and Aging: Prevalence of Disturbed Sleep and Treatment Considerations in Older Adults. *J Clin Psychiatry*. 2005;66:24.

8. Ancoli-Israel S. Sleep and its disorders in aging populations. *Sleep Med*. 2009;10 Suppl 1:S7-11. doi:10.1016/j.sleep.2009.07.004

9. Li J, Gooneratne NS. Sleep and health in older adults. In: *Sleep and Health*. Elsevier Academic Press; 2019:31-43. doi:10.1016/B978-0-12-815373-4.00004-6

10. Nguyen V, George T, Brewster GS. Insomnia in Older Adults. *Curr Geriatr Rep*. 2019;8(4):271-290. doi:10.1007/s13670-019-00300-x

11. Braak H, Braak E. Neuropathological stageing of Alzheimer-related changes. *Acta Neuropathol (Berl)*. 1991;82(4):239-259. doi:10.1007/BF00308809

12. Rodrigue KM, Kennedy KM, Devous MD, et al. β-Amyloid burden in healthy aging: regional distribution and cognitive consequences. *Neurology*. 2012;78(6):387-395. doi:10.1212/WNL.0b013e318245d295

13. Wang C, Holtzman DM. Bidirectional relationship between sleep and Alzheimer's disease: role of amyloid, tau, and other factors. *Neuropsychopharmacology*. 2020;45(1):104-120. doi:10.1038/s41386-019-0478-5

14. Kondo R, Miyano I, Lee S, Shimada H, Kitaoka H. Association between self-reported night sleep duration and cognitive function among older adults with intact global cognition. *Int J Geriatr Psychiatry*. 2021;36(5):766-774. doi:10.1002/gps.5476

15. Winer JR, Deters KD, Kennedy G, et al. Association of Short and Long Sleep Duration With Amyloid-β Burden and Cognition in Aging. *JAMA Neurol*. 2021;78(10):1187-1196. doi:10.1001/jamaneurol.2021.2876

16. Sabeti S, Al-Darsani Z, Mander BA, Corrada MM, Kawas CH. Sleep, hippocampal volume, and cognition in adults over 90 years old. *Aging Clin Exp Res*. 2018;30(11):1307-1318. doi:10.1007/s40520-018-1030-x

17. Tworoger SS, Lee S, Schernhammer ES, Grodstein F. The association of self-reported sleep duration, difficulty sleeping, and snoring with cognitive function in older women. *Alzheimer Dis Assoc Disord*. 2006;20(1):41-48. doi:10.1097/01.wad.0000201850.52707.80

18. Faubel R, López-García E, Guallar-Castillón P, Graciani A, Banegas JR, Rodríguez-Artalejo F. Usual sleep duration and cognitive function in older adults in Spain. *J Sleep Res*. 2009;18(4):427-435. doi:10.1111/j.1365-2869.2009.00759.x

19. Blackwell T, Redline S, Ancoli-Israel S, et al. Comparison of Sleep Parameters from Actigraphy and Polysomnography in Older Women: The SOF Study. *Sleep*. 2008;31(2):283-291. doi:10.1093/sleep/31.2.283

20. Wilckens KA, Woo SG, Kirk AR, Erickson KI, Wheeler ME. The role of sleep continuity and total sleep time in executive function across the adult lifespan. *Psychol Aging*. 2014;29(3):658-665. doi:10.1037/a0037234

21. Kim SJ, Lee DK, Jang YK, et al. The Effects of Longitudinal White Matter Hyperintensity Change on Cognitive Decline and Cortical Thinning over Three Years. *J Clin Med*. 2020;9(8):2663. doi:10.3390/jcm9082663

22. Morris JL, Rohay J, Chasens ER. Sex Differences in the Psychometric Properties of the Pittsburgh Sleep Quality Index. *J Womens Health*. 2018;27(3):278-282. doi:10.1089/jwh.2017.6447

23. Kocevska D, Lysen TS, Dotinga A, et al. Sleep characteristics across the lifespan in 1.1 million people from the Netherlands, United Kingdom and United States: a systematic review and meta-analysis. *Nat Hum Behav*. 2021;5(1):113-122. doi:10.1038/s41562-020-00965-x

24. Hume KI, Van F, Watson A. A field study of age and gender differences in habitual adult sleep. *J Sleep Res*. 1998;7(2):85-94. doi:10.1046/j.1365-2869.1998.00103.x

25. Yoon JE, Oh D, Hwang I, et al. Sleep structure and electroencephalographic spectral power of middle-aged or older adults: Normative values by age and sex in the Korean population. *J Sleep Res*. 2021;30(6). doi:10.1111/jsr.13358

26. Albinni B, Baker FC, Javitz H, et al. Morning perception of sleep, stress, and mood, and its relationship with overnight physiological sleep: findings from the National Consortium on Alcohol and Neurodevelopment in Adolescence (NCANDA) study. *J Sleep Res*. 2023;32(5):e13886. doi:10.1111/jsr.13886

27. Della Monica C, Johnsen S, Atzori G, Groeger JA, Dijk DJ. Rapid Eye Movement Sleep, Sleep Continuity and Slow Wave Sleep as Predictors of Cognition, Mood, and Subjective Sleep Quality in Healthy Men and Women, Aged 20-84 Years. *Front Psychiatry*. 2018;9:255. doi:10.3389/fpsyt.2018.00255

28. Groeger JA, Zijlstra FRH, Dijk DJ. Sleep quantity, sleep difficulties and their perceived consequences in a representative sample of some 2000 British adults. *J Sleep Res*. 2004;13(4):359-371. doi:10.1111/j.1365-2869.2004.00418.x

29. Vitiello MV, Larsen LH, Moe KE. Age-related sleep change: Gender and estrogen effects on the subjective-objective sleep quality relationships of healthy, noncomplaining older men and women. *J Psychosom Res*. 2004;56(5):503-510. doi:10.1016/S0022-3999(04)00023-6

30. Rångtell FH, Karamchedu S, Andersson P, et al. A single night of sleep loss impairs objective but not subjective working memory performance in a sex-dependent manner. *J Sleep Res*. 2019;28(1):e12651. doi:10.1111/jsr.12651

31. Santhi N, Lazar AS, McCabe PJ, Lo JC, Groeger JA, Dijk DJ. Sex differences in the circadian regulation of sleep and waking cognition in humans. *Proc Natl Acad Sci U S A*. 2016;113(19):E2730-2739. doi:10.1073/pnas.1521637113

32. Baker FC, Sattari N, de Zambotti M, Goldstone A, Alaynick WA, Mednick SC. Impact of sex steroids and reproductive stage on sleep-dependent memory consolidation in women. *Neurobiol Learn Mem*. 2019;160:118-131. doi:10.1016/j.nlm.2018.03.017

33. Cole RJ, Kripke DF, Gruen W, Mullaney DJ, Gillin JC. Automatic sleep/wake identification from wrist activity. *Sleep*. 1992;15(5):461-469. doi:10.1093/sleep/15.5.461

34. Choi L, Liu Z, Matthews CE, Buchowski MS. Validation of accelerometer wear and nonwear time classification algorithm. *Med Sci Sports Exerc*. 2011;43(2):357-364. doi:10.1249/MSS.0b013e3181ed61a3

35. Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh sleep quality index: A new instrument for psychiatric practice and research. *Psychiatry Res*. 1989;28(2):193-213. doi:10.1016/0165-1781(89)90047-4

36. Wechsler D. Wechsler Adult Intelligence Scale. 3rd (Ed.) San Antonio: The Psychological Corporation.[Google Scholar]. Published online 1997.

37. Wechsler D. The Wechsler Memory Scale, San Antonio, Tex, Psychological Corp. Published online 1997.

38. Stroop JR. Studies of interference in serial verbal reactions. *J Exp Psychol*. 1935;18(6):643-662. doi:10.1037/h0054651

39. Craft S, Newcomer J, Kanne S, et al. Memory improvement following induced hyperinsulinemia in Alzheimer's disease. *Neurobiol Aging*. 1996;17(1):123-130. doi:10.1016/0197-4580(95)02002-0

40. Grober E, Buschke H, Crystal H, Bang S, Dresner R. Screening for dementia by memory testing. *Neurology*. 1988;38(6):900-903. doi:10.1212/wnl.38.6.900

41. Goodglass H, Kaplan E. *The Assessment of Aphasia and Related Disorders*. 2nd ed. Lea & Febiger; 1983. Accessed December 17, 2023. https://cir.nii.ac.jp/crid/1130282270892335488

42. Reitan RM, Wolfson D. *The Halstead-Reitan Neuropsychological Test Battery: Theory and Clinical Interpretation*. Neuropsychology Press; 1985.

43. Watts A. Creating cognitive factor scores from KU-ADC data. Published online 2020.

44. Lucey BP, Wisch J, Boerwinkle AH, et al. Sleep and longitudinal cognitive performance in preclinical and early symptomatic Alzheimer's disease. *Brain*. 2021;144(9):2852-2862. doi:10.1093/brain/awab272

45. De Carolis A, Cipollini V, Corigliano V, et al. Anosognosia in People with Cognitive Impairment: Association with Cognitive Deficits and Behavioral Disturbances. *Dement Geriatr Cogn Disord Extra*. 2015;5(1):42-50. doi:10.1159/000367987

46. Kim H, Kim R, Kim S, et al. Sex differences in deterioration of sleep properties associated with aging: A 12-year longitudinal cohort study. *J Clin Sleep Med*. 2021;17. doi:10.5664/jcsm.9072

47. Goel N, Dinges DF. Behavioral and Genetic Markers of Sleepiness. *J Clin Sleep Med*. 2011;7(5 Suppl):S19-S21. doi:10.5664/JCSM.1348

48. Lo JC, Groeger JA, Santhi N, et al. Effects of partial and acute total sleep deprivation on performance across cognitive domains, individuals and circadian phase. *PloS One*. 2012;7(9):e45987. doi:10.1371/journal.pone.0045987

49. Hennecke E, Lange D, Steenbergen F, et al. Adverse interaction effects of chronic and acute sleep deficits on spatial working memory but not on verbal working memory or declarative memory. *J Sleep Res*. 2021;30(4):e13225. doi:10.1111/jsr.13225

50. van Schalkwijk FJ, Sauter C, Hoedlmoser K, et al. The effect of daytime napping and full-night sleep on the consolidation of declarative and procedural information. *J Sleep Res*. 2019;28(1):e12649. doi:10.1111/jsr.12649

51. Casagrande M, Forte G, Favieri F, Corbo I. Sleep Quality and Aging: A Systematic Review on Healthy Older People, Mild Cognitive Impairment and Alzheimer's Disease. *Int J Environ Res Public Health*. 2022;19(14):8457. doi:10.3390/ijerph19148457

Fig 1. Partial effects of total sleep time, sleep efficiency, and Pittsburgh Sleep Quality Index (PSQI) on executive function in females.

Fig 2. Partial effects of total sleep time, sleep efficiency, and Pittsburgh Sleep Quality Index (PSQI) on verbal memory in females.

Fig 3. Partial effects of total sleep time, sleep efficiency, and Pittsburgh Sleep Quality Index (PSQI) on attention in females.

Fig 4. Partial effects of total sleep time, sleep efficiency, and Pittsburgh Sleep Quality Index (PSQI) on executive function in males.

Fig 5. Partial effects of total sleep time, sleep efficiency, and Pittsburgh Sleep Quality Index (PSQI) on verbal memory in males.

Fig 6. Partial effects of total sleep time, sleep efficiency, and Pittsburgh Sleep Quality Index (PSQI) on attention in males.

Table 1. Descriptive Statistics on Demographics, Sleep, and Cognitive Performance

Participants characteristics	Males (*n* = 89)		Females (*n* = 118)		Total (*N*= 207)	
	M	*SD*	*M*	*SD*	*M*	*SD*
Age (years)	76.1*	6.9	74*	6.6	74.9	6.8
Education (years)	16.7	2.8	16.1	3	16.4	2.9
	N	%	*N*	%	*N*	%
Race						
White	86	96.6	108	91.5	194	93.7
African American	2	2.3	8	6.8	10	4.8
Asian	1	1.1	2	1.7	3	1.5
CDR = 0	63	70.8	98	83.1	161	77.78
	M	*SD*	*M*	*SD*	*M*	*SD*
TST (minutes)	409.7	68.3	419.8	60.4	415.5	64
PSQI	5.8 *	2.5	6.8 *	2.9	6.4	2.7
SE (%)	85.4	7	86.6	6	86.1	6.4
Verbal Memory	0.4 *	1.3	0.9 *	1.2	0.7	1.3
Executive Function	0.1	0.9	0.4	0.8	0.3	0.8
Attention	0.1	0.4	0.2	0.4	0.2	0.4

CDR = Clinical Dementia Rating. TST = total sleep time. PSQI = Pittsburgh Sleep Quality Index. SE = sleep efficiency. M = mean. SD = standard deviation.
* Differences between males and females, $p < 0.05$

Table 2. Generalized Additive Models of the Relationship Between Sleep Quality (Total Sleep Time, Sleep Efficiency, and PSQI) and Cognitive Performance (Executive Function, Verbal Memory, and Attention)

	Executive Function		Verbal Memory		Attention	
	Females	Males	Females	Males	Females	Males
Spline fit	**EDF**	**EDF**	**EDF**	**EDF**	**EDF**	**EDF**
TST	2.178**	1.000	1.000*	1.000	1.000	1.000
SE	1.872	1.000.	1.000	1.000	2.257	1.000
PSQI (CDR =0)	1.000*	1.000	2.400	1.000	1.000	1.000
PSQI (CDR =1)	1.450	2.178***	1.000	2.102**	1.794	2.041
Covariate	**Estimate**	**Estimate**	**Estimate**	**Estimate**	**Estimate**	**Estimate**
Age	-0.021**	-0.030**	-0.062***	-0.033*	-0.017**	-0.016**
Education	0.044**	0.021	0.083**	0.004	0.018	0.015
CDR = 1	-0.935***	-1.037***	-1.152***	-1.608***	-0.308**	-0.443***

Dependent variables: executive function, verbal memory, and attention. TST = total sleep time. SE = sleep efficiency. CDR = Clinical Dementia Rating. EDF = effective degrees of freedom (quantification of model's flexibility).

$p = 0.05$, $*p < 0.05$, $**p<0.01$, $***p<0.001$

Fig 6. Partial effects of total sleep time, sleep efficiency, and Pittsburgh Sleep Quality Index (PSQI) on attention in males.

Fig 1. Partial effects of total sleep time, sleep efficiency, and Pittsburgh Sleep Quality Index (PSQI) on executive function in females.

Fig 2. Partial effects of total sleep time, sleep efficiency, and Pittsburgh Sleep Quality Index (PSQI) on verbal memory in females.

Fig 3. Partial effects of total sleep time, sleep efficiency, and Pittsburgh Sleep Quality Index (PSQI) on attention in females.

Fig 4. Partial effects of total sleep time, sleep efficiency, and Pittsburgh Sleep Quality Index (PSQI) on executive function in males.

Fig 5. Partial effects of total sleep time, sleep efficiency, and Pittsburgh Sleep Quality Index (PSQI) on verbal memory in males.